



RESEARCH ARTICLE

REVISED **Dental health status of community-dwelling adults aged 50 years and over in Ireland. A cross-sectional analysis of the Wave 3 TILDA cohort. [version 3; peer review: 2 approved]**

Amara Naseer ¹, Jacinta McLoughlin¹, Orna A. Donoghue ², Rose Anne Kenny ^{2,3}, Brian O'Connell ¹

¹Dublin Dental University Hospital, Trinity College Dublin, Dublin, D02F859, Ireland

²The Irish Longitudinal Study on Ageing (TILDA), Trinity College Dublin, Dublin, D02PN40, Ireland

³Mercer's Institute for Successful Ageing, St James's Hospital, Dublin, Ireland

V3 **First published:** 14 Dec 2018, 1:26
<https://doi.org/10.12688/hrbopenres.12891.1>
Second version: 21 Sep 2021, 1:26
<https://doi.org/10.12688/hrbopenres.12891.2>
Latest published: 08 Feb 2022, 1:26
<https://doi.org/10.12688/hrbopenres.12891.3>

Abstract

Background: Little is known about the current oral health status of adults in Ireland. The aim of this study was to assess the dental health of community-dwelling adults aged 50 years and over in Ireland and to compare the current status to previous national surveys.
Methods: The Irish Longitudinal Study on Ageing (TILDA) Wave 3 assessed the dental health of a subset of participants. Respondents attending for health assessments were offered a dental examination. The World Health Organization examination criteria were used.
Results: Of the 3111 people who were offered the dental assessment, 2525 were examined. Adults below 50 years of age and respondents whose dental health data were unavailable at the time of analysis were omitted, giving a final sample of 2504. Among the dental assessment sample, 9.9% (249) were edentate. Of those aged 65 years and older, 15.6% were edentate while for the same age group 40.9% were edentate in the 2000-02 national survey. The mean number of teeth present in those aged 65 years or older was 14.9 for males and 14.2 for females, whereas in 2000-02 it was 9.9 and 7.4, respectively. 56.8% of the dentate sample had 10 or more tooth contacts. The mean DMFT of those aged 50 years or more was 18.5 and the Root Caries Index was 6.3. Between 2000-02 and 2014-5 (this study) in adults aged 65 years and over, the mean DMFT decreased from 25.9 to 20.1 and the Root Caries Index decreased from 11.6 to 9.1.
Conclusion: The results indicate the dental health of community-dwelling adults aged 50 years and over in Ireland improved since the

Open Peer Review

Approval Status

	1	2
version 3 (revision) 08 Feb 2022	 view	
version 2 (revision) 21 Sep 2021	 view	
version 1 14 Dec 2018	 view	 view

1. **W. Murray Thomson** , University of Otago, Dunedin, New Zealand

2. **Roseann Mulligan**, University of Southern California, Los Angeles, USA

Any reports and responses or comments on the article can be found at the end of the article.


previous survey of 2000-02. These improvements mean a change in the treatment needs of this age group and will require policy and service adjustments to meet these needs

Keywords

ageing demographics, cohort studies, oral health, health surveys, public health dentistry



This article is included in the [TILDA](#) gateway.



This article is included in the [Ageing Populations](#) collection.

Corresponding author: Brian O'Connell (brian.oconnell@dental.tcd.ie)

Author roles: **Naseer A:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **McLoughlin J:** Conceptualization, Investigation, Methodology, Project Administration, Supervision, Validation, Writing – Review & Editing; **Donoghue OA:** Conceptualization, Data Curation, Formal Analysis, Methodology, Project Administration, Supervision, Validation, Writing – Review & Editing; **Kenny RA:** Conceptualization, Funding Acquisition, Methodology, Supervision, Writing – Review & Editing; **O'Connell B:** Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Financial support for TILDA was provided by the Irish Government, the Atlantic Philanthropies and Irish Life plc. We also gratefully acknowledge financial support from the Department of Health for the oral health assessments and from the Dublin Dental Hospital Board for the analysis.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2022 Naseer A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Naseer A, McLoughlin J, Donoghue OA *et al.* **Dental health status of community-dwelling adults aged 50 years and over in Ireland. A cross-sectional analysis of the Wave 3 TILDA cohort. [version 3; peer review: 2 approved]** HRB Open Research 2022, 1:26 <https://doi.org/10.12688/hrbopenres.12891.3>

First published: 14 Dec 2018, 1:26 <https://doi.org/10.12688/hrbopenres.12891.1>

REVISED Amendments from Version 2

Previous surveys of Irish adults calculated the mean number of teeth in samples that included edentate individuals—this result is presented for comparison with the 2000–02 and 1989–90 surveys as in the last version of the text (Table 12). This second revision of the paper also recalculates the mean number of teeth in the age groups (50–64, 65–74 and 75 years+) in the TILDA dentate sample only, for the purposes of international comparison (Table 14).

The text was also edited for improvements in clarity and style.

Any further responses from the reviewers can be found at the end of the article

Introduction

The ageing population is one of the great challenges that will confront health services in developed countries in coming years. It is estimated that by 2050, the number of adults aged 60 years and older worldwide will increase from 901 million to 2.1 billion, and the adults called “oldest old” (80 years and over) will more than triple (125 to 434 million), from 2015¹. In 2011, in Ireland, adults aged 65 years and over comprised 11.4% of the total population and this proportion is predicted to reach 22.4% in 2041. The proportion of adults aged 80 years or older is predicted to be 7.5% of the total population in 2046^{2,3}.

The ageing population faces the challenge of various chronic diseases including physical and mental health related conditions. Maintaining good oral health may also be a challenge in older people and poor oral health has been called a silent epidemic⁴. Older adults may, as a result of reduced oral function, eat a poor quality diet and avoid social interaction and in this way poor oral health may adversely affect health and wellbeing. Similarly, loss of physical and cognitive function, along with increasing frailty, often result in less attention to oral self-care, while ill health or frailty may lead to reduced access to the professional care that is needed to maintain oral function.

Dental caries and periodontal disease are the most common chronic oral diseases affecting the adult population in Ireland and worldwide^{5,6}, and the loss of natural teeth is considered to be a key indicator of poor oral health in older people⁷. Although the complete loss of teeth with age has reduced dramatically in recent years, most oral health surveys have reported an age-related deterioration in oral health including an increase in tooth loss, poor periodontal health and increased tooth wear^{5,8,9}. The maintenance of a minimum number of functioning teeth, good periodontal health, absence of active dental caries and control of tooth wear are general indicators of satisfactory dental health in older adults.

In Ireland, the last national survey of the oral health status of adults was undertaken in 2000–02, so up-to-date information reflecting the current oral health status of adults is lacking. This makes it difficult to assess the needs of the population and to design services that maintain the oral health of older

adults. Wave 3 of The Irish Longitudinal Study on Ageing (TILDA) provided an opportunity to include a dental assessment of community dwelling adults aged 50 years and over¹⁰. The aim of this assessment was to provide an up-to-date picture of the dental health of older adults, and to compare the current data with previous data from Irish and international studies.

Methods**Ethical considerations and consent**

Ethical approval for this study was obtained from the Trinity College Dublin Faculty of Health Sciences Research Ethics Committee and participants provided written informed consent before the health assessment.

Study design

TILDA is a large-scale, nationally representative, comprehensive cohort study on ageing in Ireland. It was started in 2009 and at the time of writing had completed its fourth wave of data collection¹¹. The TILDA cohort consists of randomly selected community-dwelling adults aged 50 years and over, although partners or spouses of any age can also participate. There are three modes of data collection, a computer aided personal interview (CAPI), self-completion questionnaire (SCQ) and health assessments¹⁰.

In Wave 3 (March 2014 – December 2015), for the first time in the TILDA study a dental assessment was included as part of the health assessment conducted in the TILDA centre in Trinity College Dublin. The dental sample was a non-random convenience sample. As a dentist was not available during the full TILDA centre opening hours, only those participants who completed a health assessment during the dentist’s hours were invited to have the dental assessment; there were no other exclusion criteria. The respondents attending the TILDA centre in Dublin came from all over the country. Periodontal probing was omitted from the assessment of respondents at risk from bacteraemia. The examination criteria used in this study were the same as those used in previous Irish national oral health surveys and similar to those recommended by WHO^{5,12}. The examiners were trained by an experienced examiner (‘gold standard’--JMCL) from previous studies. A total of one trainer (‘gold standard’) and four assessors (including AN, BOC) completed the data collection. As the dental assessment was at the end of the health centre assessment (approximately three hours long), a maximum of 10 minutes was allocated for it. Because of the time constraint, it was not possible to perform duplicate examinations during the data collection. During the pilot phase, a calibration exercise was performed on non-participant volunteers, followed by dual assessment of study participants until any discrepancies between the trainer and assessors were resolved. For the dental examination, standardised equipment consisted of a dental chair with floor mounted Daray LED examination light (Model- XL200 LED examination light, 12–30v/5.8–8.2w), standard plane dental mirror and WHO recommended Community Periodontal Index of Treatment Need (CPITN) probe-E¹². Methods and equipment were the same for all examinations. Standard cross infection control measures were followed during all examinations.

All respondents attending for a health assessment were invited to participate in the OHA while an examining dentist was present. The data collected was; number of natural teeth, use of dentures, CPITN on index teeth, coronal caries at cavitation level and visual level (WHO and British Association for the Study of Community Dentistry-BASCD)¹³, Root Caries Index of Katz¹⁴, coronal tooth wear into dentine¹⁵ and tooth contacts between maxillary and mandibular teeth in the Maximum Intercuspal Position (MIP)^{16,17}. All criteria were based on visual examination and tactile sensing methods using a CPITN probe and no radiographs were taken. Data from the assessment was written on a paper form and then entered on a laptop computer and uploaded to the TILDA database.

For the purpose of the CPITN examination, the mouth was divided into sextants and the highest (worst) score in each sextant was recorded as the sextant score. The scores were; no disease (H), bleeding on examination (B), supra or sub gingival calculus present (C), pocket depth up to 4-5mm (P1), pocket depth >6mm (P2) and if no teeth were present in a sextant/unable to record (X)¹².

Tooth contacts in the Maximal Intercuspal Position (MIP) were recorded to evaluate the functional dentition and the need for replacement of teeth^{16,18-21}. To achieve MIP, participants were asked to swallow and keep their teeth closed together-- the number of mandibular occlusal units in contact with maxillary teeth was counted. An occlusal unit was considered to be a single anterior tooth or premolar, or half a molar tooth (mesial or distal)²¹. The percentages of dentate adults with fewer than 10 contacts, and 10 contacts or more, by age group and gender were calculated. Ten tooth contacts indicate approximately 20 teeth in occlusion, which is considered to be a minimal functioning dentition²¹.

The presence of root caries was recorded in all respondents and the Root Caries Index (RCI) was calculated among the dentate adults who had exposed roots due to gingival recession, as was done in previous Irish national surveys of adults¹⁴. This index gives the proportion of exposed roots with caries or restorations due to caries (RCI = mean decayed and filled roots/mean exposed roots %) in the population with exposed roots. Decayed and filled roots were recorded at tooth level rather than surface level, and so RCI was also calculated at tooth level.

Tooth wear was recorded by visual examination. The Bardsley tooth wear index was used in this study to record coronal tooth wear into dentine¹⁵. The mouth was divided into sextants and each sextant was individually scored. Tooth wear was recorded as; no wear, exposed dentine comprised <1/3 of worst surface of a tooth, exposed dentine comprised >1/3 of worst surface of a tooth, or the sextant was excluded, as no teeth were present in the sextant or unable to record a score. The worst tooth in a sextant was recorded as a sextant score. The highest score of dentine wear per person was recorded as a person's tooth wear level.

During the statistical analysis, some dental health indicators were calculated for the full dental assessment sample (denture

wear, number of teeth and mean number of decayed, missing or filled teeth (DMFT)) and other indicators (tooth contacts, RCI and periodontal health) were calculated for the dentate sample only, resulting in two bases for results. "base edentate/dentate" means the statistical analysis involved the full sample, including edentate and dentate respondents, whereas "base dentate" means statistical analysis was run only on the dentate sample.

TILDA is subject to the legislation under the Data Protection Act 1988 and the Data Protection (Amendment) Act 2003. All data protection protocols were followed during collection, processing, analysis and reporting on data²². Data analysis followed completion of data cleaning by accessing a TILDA hot desk at the TILDA research centre in Trinity College. STATA software (Stata 14.1 Stata Corp LLC Texas USA) was used for data analysis.

Results

Dental health assessment sample selection

The study sample was a sub-sample of the respondents who attended a TILDA health centre assessment at Wave 3. A total of 4309 respondents attended for the health assessment, of whom an opportunistic sample of 3111 (72.2%) were invited to have the dental assessment, and of these 2525 (81.1%) agreed to the assessment. Those aged less than 50 years (n=17) were omitted from this analysis. The full dental sample consisted of 2508 respondents, however, as the data for 4 of these respondents was not available at the time of the analysis, the results reported here are for 2504 respondents¹¹.

Dental health assessment sample vs TILDA sample (population sample)

The TILDA cohort is a nationally representative sample of adults aged 50 years and over (a population sample). However, the dental assessment was completed on a convenience sample of the TILDA cohort. Table 1 below reports the characteristics of the dental sample and a two-sample proportion test (Z test) for comparisons of the characteristics between the dental sample and the Wave 3 TILDA cohort. For the purpose of comparison between the different samples, 10 variables with 23 categories were selected. There was no difference between the samples (dental and TILDA) in 8 out of the 23 categories.

In summary, the dental sample was younger, more respondents were married, they were better educated, with good to excellent self-rated general health and more likely to be living in Dublin than the full TILDA sample.

Dental health assessment sample description

For the analysis, the dental sample was stratified into three age groups as recommended by the WHO¹²; 50–64 years old, 65–74 years old and 75 years and over (see Table 2 for gender breakdown). Almost half of the sample was aged 50–64 years, with 14.8% aged 75 years and over. Overall, the dental sample consisted of more females than males (55.3% vs 44.7%), but this trend was less marked in older respondents.

Dentate/edentate proportion

The adults with at least one natural tooth present were recorded as dentate. Table 3 shows that overall, 9.9% of the sample was edentate (no teeth).

Table 1. Comparison of the dental assessment sample with the TILDA sample by a two-sample proportion test (Z test). P values less than 0.05 are shown in bold.

Characteristic	Oral Health Assessment Sample (n=2508) n (%)	Population (TILDA sample) (n=6618) n (%)	Hypothesis test of proportions Population (TILDA) vs OHA
Age group			P value
50–64 years	1219 (48.6)	3036 (45.9)	0.0196
65–74 years	918 (36.6)	2110 (31.9)	< 0.001
≥75 years	371 (14.8)	1472 (22.2)	< 0.001
Female	1386 (55.3)	3679 (55.6)	0.7786
Education level			
Primary	478 (19.1)	1737 (26.3)	< 0.001
Secondary	1005 (40.1)	2610 (39.4)	0.5805
Tertiary/higher	1024 (40.8)	2269 (34.3)	< 0.001
Marital status			
Married	1889 (75.3)	4573 (69.1)	< 0.001
Never married	168 (6.7)	562 (8.5)	0.0048
Separated/divorced	192 (7.7)	469 (7.1)	0.3494
Widowed	259 (10.3)	1014 (15.3)	< 0.001
Locality			
Dublin	681 (27.1)	1592 (24.1)	0.0023
Other urban	661 (26.4)	1840 (27.8)	0.1664
Rural	1166 (46.5)	3186 (48.1)	0.1588
Grew up in rural area	1440 (57.4)	3891 (58.8)	0.2331
Never lived abroad	669 (26.7)	1534 (23.2)	0.0005
Current or former smoker	1302 (51.9)	3609 (54.5)	0.0251
No health insurance or medical card	209 (8.3)	591 (8.9)	0.3681
Self-reported health			
Excellent	394 (15.7)	921 (14.2)	0.0294
Very good	897 (35.8)	2169 (33.4)	0.0069
Good	851 (34.0)	2227 (34.2)	0.8001
Fair	312 (12.4)	970 (14.9)	0.0065
Poor	52 (2.1)	215 (3.3)	0.0029

Table 2. The dental assessment sample by gender and age group (n=2504).

Gender	Age groups			Total
	50-64 years	65-74 years	75 years and over	
Male	511	422	187	1120
	42.0%	46.1%	50.5%	44.7%
Female	707	494	183	1384
	58.0%	53.9%	49.5%	55.3%
Total Row %	1218	916	370	2504
	48.6%	36.6%	14.8%	100%

Table 3. Number and percentage of edentate/dentate sample by age group and gender (Base edentate/dentate, n=2504).

Age group	50-64 years		65-74 years		75 years & over		Total
	Male	Female	Male	Female	Male	Female	
Edentate	15	33	36	71	39	55	249
	2.9%	4.7%	8.5%	14.4%	20.9%	30.1%	9.9%
Dentate	496	674	386	423	148	128	2255
	97.1%	95.3%	91.5%	85.6%	79.1%	69.9%	90.1%

The proportion of edentate adults was higher in the older age group, and more females were edentate than males in all age groups. Overall, 9.9% of respondents were completely edentate, 14.0% were edentate in the upper arch only, and 19% were edentate in the lower arch only.

Denture wear

Table 4 shows that 46.9% of the sample had some type of removable denture. While 9.9% of the sample was edentate, only 9.1% was wearing complete dentures for both the upper and lower arches.

The percentage of adults wearing dentures was higher in the oldest age group irrespective of the type of denture. In those aged 75 years and over, 73% wore dentures and 24% wore complete dentures.

Number of teeth

Figure 1 shows the frequency distribution of the number of teeth present in the dental sample. It shows that 9.9% of adults had no teeth, and 54.3% of adults had 20 or more teeth. When the results in Table 5 are compared, it can be observed that although a higher proportion of females were edentate, females

in the youngest age group, who were dentate, had a similar mean number of teeth to males (21.5 vs 21.1). This suggests that the higher tooth loss in older women in Ireland may be reversing, although this would need to be substantiated by a longitudinal study.

Tooth contacts

The percentage of dentate adults with fewer than 10 occlusal contacts, and 10 contacts or more, by age group and gender was calculated. Figure 2 shows that 56.8% of the dentate sample had 10 or more tooth contacts. Notably, 13.6% of dentate adults had no contacts; these adults were edentate in one arch, wearing dentures, had cross bites, teeth not in contact with other teeth, or just roots remaining. Table 6 shows that the proportion of dentate adults with 10 or more contacts was less in the oldest age group and was higher in females, although the gender difference narrowed in the older age groups.

Decayed, missing and filled teeth (DMFT)

Dental caries was recorded at cavitation (DMFT-c) and at visual caries level (DMFT-v). As the results indicated a difference of 0.1 between DMFT-c and DMFT-v (DMFT-c= 18.5, DMFT-v =18.6), it was decided to only report DMFT-c.

Table 4. The proportion of denture wearers by age group and type of denture (Base edentate/dentate, n=2504).

Age group	No upper or lower denture		Complete upper and lower dentures		Complete upper and partial lower dentures		Complete upper denture only		All other combinations of complete and partial dentures		Total n
	n	%	n	%	n	%	n	%	n	%	
50-64 years	838	68.8	42	3.5	10	0.8	41	3.4	287	23.6	1,218
65-74 years	392	42.8	98	10.7	51	5.6	67	7.3	308	33.6	916
75 years and over	100	27.0	88	23.8	24	6.5	41	11.1	117	31.6	370
Total	1,330	53.1	228	9.1	85	3.4	149	5.9	712	28.4	2504




Figure 1. Percentage of adults by total number of teeth and the proportion with <20 teeth and ≥ 20 teeth. (Base edentate/dentate, n=2504). Mean number of teeth was 17.9, SD = 8.9, Median=21.

During the dental assessment it was possible to identify that some teeth were missing for reasons other than dental caries. These were missing premolars with no residual spaces (orthodontic extractions or congenital absence), third molars extracted due to impaction and teeth lost due to trauma. Where there was certainty about the reasons for loss these teeth they were recorded as missing for other reasons and not counted in the M component of DMFT. Where it was unclear, or there was

doubt about the reasons for tooth loss, these teeth were recorded as missing due to caries. Similarly, in the edentate group third molar teeth were not recorded as missing due to caries. For this reason, the maximum DMFT score for edentate adults is shown as 28 but for dentate adults is 32.

Figure 3 shows that overall, 10.2% of adults had a DMFT score of 28, which includes the 9.9% who were edentate.

The DMFT scores are negatively skewed with a greater percentage of adults having high values for DMFT. Only 1.8% of adults had a DMFT score of 1. Mean DMFT values by age group and gender are reported in Table 7. Overall,

females had a slightly higher mean DMFT (0.3) score than males, and this gender difference was present among all three age groups. Figure 4 shows the contribution of the decayed teeth, missing teeth, and filled teeth components of DMFT by age group and gender.

Table 5. Mean number of teeth per person by age group and gender (Base edentate/dentate, n=2504).

Age groups	Male	Female	Total
50-64 years	21.1	21.5	21.3
65-74 years	16.3	15.4	15.8
75 years & over	11.9	11.1	11.5
Total	17.7	17.9	17.9

Of the total mean DMFT of 18.5, the contribution of decayed teeth was 0.6 in males and 0.3 in females, and was almost the same in all age groups. When compared to men, women in all age groups had fewer decayed teeth, more filled teeth, and women aged 75 years and over had more missing teeth, which suggests that women access treatment more. It is notable that in the youngest age group, the proportion of missing teeth was much lower than in the two older age groups, with corresponding higher proportions of filled teeth and the same proportion of decayed teeth. It remains to be seen whether this younger cohort can maintain more of their natural teeth as they age, as this would represent a major shift in the dental health of older adults in Ireland.




Figure 2. Percentage of adults by total number of tooth contacts and with <10 tooth contacts and ≥10 contacts (Base dentate, n=2255). Mean 9.9, SD= 5.9, Median= 11.

Table 6. Number and percentage of adults with fewer than 10 tooth contacts, and equal to or more than 10 tooth contacts, by age group and gender (Base dentate, n=2255).

Age group	50-64 years		65-74 years		75 years & over		Total
	M	F	M	F	M	F	
<10 Contacts	172	179	214	225	101	84	975
	34.7%	26.6%	55.4%	53.2%	68.2%	65.6%	43.25%
≥ 10 Contacts	324	495	172	198	47	44	1280
	65.3%	73.4%	44.6%	46.8%	31.8%	34.4%	56.75%




Figure 3. Frequency distribution of the percentage of adults aged 50 years and over by DMFT score (Base edentate/dentate, n=2504). Mean= 18.5, SD= 6.3, Median=19.

Table 7. Mean DMFT level by age group and gender (Base edentate/dentate, n=2504).

Age groups	Male	Female	Total DMFT
50-64 years	16.4	17.0	16.7
65-74 years	19.4	19.6	19.5
75 years & over	21.3	22.0	21.6
Total	18.3	18.6	18.5

Root caries

Root caries is reported as the mean number of decayed and filled roots as a proportion of the mean number of roots with recession (Table 8). The RCI (4.3 vs 10.2) and the mean number of decayed/filled roots (0.5 vs 1.1) was higher, and the mean number of exposed roots (11.1 vs 10.3) was lower, in the oldest age group. In the lower mean number of exposed roots may be due to having fewer teeth with increasing age. Females aged less than 75 years had higher levels of root caries than males, but those aged 75 years and over had slightly lower levels than males.

Periodontal health

The CPITN was used for the periodontal health assessment. The severity of periodontal disease was reported by the

maximum CPITN score per person, and the extent as the mean number of sextants affected by the different scores for the dentate adults. Table 9 shows that the proportion of men and women with completely healthy periodontal tissues or bleeding gingivae or deep pockets was low, at less than 5%, in all age groups. The majority of respondents needed simple treatment for calculus and shallow pockets. The gender differences were small, with females tending to have better periodontal health status.

Table 10 shows the mean number of sextants per person affected by the different CPITN scores, stratified by age group and gender. The mean number of sextants with deep pockets is low (0.1-0.2). With respect to shallow periodontal pockets, the mean number of sextants affected was between 0.8 and 1.5, which suggests that pockets were not very extensive in this sample. The results from these two tables indicate that the periodontal treatment needs of this sample were neither complex nor extensive.

Tooth wear

Table 11 shows that in all age groups, fewer than 7% of respondents had no wear into dentine, while 50.6% had wear into dentine on less than one third of the worst surface. In all three age groups, there was more severe wear in males than females. In the age groups included in this study, some tooth wear would be considered to be physiological and for this reason, the low percentage with no wear is not unexpected.




Figure 4. Mean decayed, missing and filled teeth components of DMFT by age group and gender (Base edentate/dentate, n=2504).

Table 8. Mean number of exposed roots, mean decayed/filled roots and Root Caries Index (RCI) by age group and gender (Base dentate, n=2255).

Age groups	Mean decayed/filled roots (DFR)			Mean exposed roots (ER)			RCI		Total RCI
	Male	Female	Total	Male	Female	Total	Male	Female	
50-64 years	0.4	0.5	0.5	11.6	10.7	11.1	3.6	4.9	4.3
65-74 years	0.8	0.9	0.9	11.1	10.7	10.8	7.2	8.3	7.8
75 years & over	1.1	1.0	1.1	10.1	10.6	10.3	10.6	9.8	10.2

Table 9. Percentage and total number of adults with a maximum CPITN score of H (healthy), B (bleeding), C (calculus), P1 (shallow pocket), P2 (Deep pocket) and X (missing sextant) by age groups and gender (Base-Dentate n=2255).

Age groups	H%		B%		C%		P1%		P2%		X%		Total n
	M	F	M	F	M	F	M	F	M	F	M	F	
50-64 years	1.4	3.9	1.1	2.0	10.9	18.4	24.3	30.6	4.1	2.5	0.6	0.3	1170
65-74 years	2.4	3.5	0.9	2.7	14.3	19.3	24.9	23.9	2.7	2.2	2.6	0.7	809
75 years and over	3.6	4.7	3.3	1.8	15.9	19.9	22.1	17.4	2.5	1.5	6.2	1.1	276
Total n	45	86	29	50	288	426	546	599	77	51	45	13	2255

Table 10. Mean number of sextants per person affected by different CPITN score: H (healthy), B (bleeding), C (calculus), P1 (shallow pocket), P2 (Deep pocket) and X (missing sextant) among dentate sample by age group and gender (Base dentate, n=2255).

Age groups	H		B		C		P1		P2		X	
	M	F	M	F	M	F	M	F	M	F	M	F
50–64 years	1.6	2.2	0.6	0.7	1.4	1.3	1.5	1.2	0.2	0.1	0.7	0.5
65–74 years	1.4	1.9	0.5	0.5	1.4	1.3	1.1	0.8	0.1	0.1	1.5	1.4
75 years & over	1.4	1.7	0.4	0.4	1.1	1.2	0.9	0.8	0.1	0.1	2.2	1.8

Table 11. Percentage of adults with no tooth wear, dentine exposed less than 1/3 of worst surface, dentine exposed more than 1/3 of worst surface, or sextant excluded, among the dentate sample by age group and gender (Base dentate, n=2255).

Age groups	No wear		Wear <1/3 of dentine		Wear >1/3 of dentine		Excluded		Total
	M	F	M	F	M	F	M	F	
50–64 years	3.3	6.9	20.3	32.3	18.5	17.7	0.3	0.7	1170
65–74 years	3.3	5.4	20.2	28.1	23.9	18.4	0.5	0.4	809
75 years and over	2.5	4.7	23.9	25.4	26.1	16.3	1.1	0	276
Total n	73	138	465	675	482	401	10	11	2255

Discussion

In Ireland, there have been considerable improvements in the oral health status of adults from the previous Irish oral health surveys conducted in 1989–1990²³ and 2000–2002⁵. A summary of the principal oral health indicators in older adults in Ireland, from the last three surveys, is shown in Table 12. For national and international comparisons DMFT is reported by the WHO method of calculation (including teeth missing for all reasons).

Edentulism. Table 12 indicates that in Ireland, among adults aged 65 years and over, the prevalence of edentulism has reduced by more than two-thirds from 1989–90 to 2014–15 and most of that decrease has occurred since 2000–02. Though the prevalence of edentulism was reduced more dramatically in females than males, in 2014–5 the prevalence was still about 50% higher in females. A comparison of the Irish and international prevalence of edentulism is shown in Table 13.

There are similar age differences in the prevalence of edentulism in different countries, although the age groups and dates of the studies vary. For example, in the age group 75 years or older, the TILDA dental sample in 2014–15 had a similar prevalence of edentulism to the USA in 2012. For the group aged 65–74 years, edentulism in the TILDA dental sample

(11.7% in 2015) and in the UK (15.0% in 2009) are comparable taking account of the time difference between the studies.

Mean number of teeth. Table 12 shows that in Ireland there has been a doubling in the mean number of teeth in adults aged 65 years and older from 1989–90 to 2014–15 (7.3 to 14.6). This trend in females is particularly positive where the mean number of teeth almost doubled between 2000–02 and 2014–15. Although mean number of teeth is a crude measure of oral health status, as it gives no indication of the condition of these teeth, it is nonetheless a positive trend that natural teeth have been retained rather than extracted.

Table 14 shows that adults in Ireland, aged 65–74 years, and 75 years and over, had a lower mean number of teeth than the UK, New Zealand and Australia, despite the fact that the studies in these countries were completed 5–10 years before the TILDA study. Together, these findings suggest that in Ireland, fewer adults above 65 years may be completely edentate than in this group of countries, but the mean number of teeth per person is also less than the UK, Australia and New Zealand.

Dental caries. There was a decrease in mean DMFT from 1989–90 to 2014–15, for those aged 65 years and older in

Table 12. Changes in edentulism, mean number of teeth, DMFT (WHO calculation) and RCI among adults aged 65 years and over in Ireland from 1989–90²³ to 2000–02⁵ and in the current study, 2014–15.

Oral health indicators	Examination Year	Male	Female	Total
% Edentate	2014–15	12.3	18.6	15.6
	2000–02	34.6	45.6	40.9
	1989–90	33	61	48
Mean number of teeth	2014–15	14.9	14.2	14.6
	2000–02	9.9	7.4	8.5
	1989–90	10.1	4.9	7.3
Mean DMFT	2014–15	23.9	24.8	24.4
	2000–02	24.8	27	25.9
	1989–90	25.6	28.8	27.3
RCI	2014–15	9.1	9.1	9.1
	2000–02	12.7	10.6	11.6
	1989–90	20.9	14.9	18.5

Table 13. Percentage edentulism by age groups, country and examination year.

Country	Year of Examination	Age group	% Edentate
Ireland	2014–15	50–64 years	3.9
		65–74 years	11.7
		75 years and over	25.4
UK ²⁴	2009	55–64 years	5
		65–74 years	15
		75–84 years	30
		85 years and over	47
USA ²⁵	2012	65–74 years	13
		75 years and over	25.8
Australia ²⁶	2006	55–74 years	13.9
		75 years and over	35.7
New Zealand ²⁷	2009	55–64 years	14.5
		65–74 years	29.6
		75 years and over	39.6

Ireland (27.3 vs 24.4), as shown in Table 12. It is notable that most of this decrease in DMFT occurred since 2000–02. From 1989-90 to 2014–15, the gender difference in mean DMFT also reduced from 3.2 to 0.2.

These findings, along with the doubling of the mean number of teeth (Table 12), suggest that adults over 65 years are not only keeping more of their teeth, but these teeth are in a healthier state. It also appears that these adults have accessed dental care which is more oriented to the maintenance of teeth than in the past.

Table 12 also indicates that the Root Caries Index reduced by more than half in Ireland from 1989–90 to 2014–15 (18.5 vs 9.1).

The gender difference for RCI was reduced from 6 in 1989–90 to 0 in 2014–15. This is an encouraging finding, as it was thought that the prevalence of root caries in older people might increase with increased retention of natural teeth and a tendency to lose periodontal attachment with age, resulting in more exposed root surfaces that are vulnerable to the development of root caries.

Comparison of the mean DMFT among the TILDA dental sample and other countries is shown in Table 15. It should be noted that in both the Australian and New Zealand surveys these mean DMFT values were for the dentate population, as edentate people were excluded from the oral health assessments.

Table 14. Mean number of teeth per person by age groups, country and year of examination among Dentate Sample.

Country	Year of examination	Age group	Mean number of teeth
Ireland	2014–15	50–64 years	22.2
		65–74 years	17.9
		75 years and over	15.4
UK ²⁴	2009	55–64 years	23.2
		65–74 years	20.9
		75–84 years	17.1
		85 years and over	14.0
Canada (CHMS) ²⁸	2007–09	40–59 years	24.1
		60–79 years	19.4
New Zealand ²⁷	2009	55–64 years	24.0
		65–74 years	19.7
		75 years and over	18.1
Australia ²⁶	2004–06	65–74 years	21.8
		75 years and over	17.9

The TILDA dental sample had a lower mean DMFT than data from New Zealand and Australia, however, these studies took place 5–10 years before the Irish study which might account for some of this difference in mean DMFT. The inclusion of the edentate group in the TILDA calculation of mean DMFT would suggest that the difference in mean DMFT between Ireland and Australia and New Zealand may in fact be greater than it appears in this table.

Periodontal health. In Ireland, the changes over time in the proportion of adults aged 65 years and over with maximum CPITN scores from 2000–02 to 2014–15 are shown in Table 16. From 2000–02 to 2014–15, the proportion of adults with a CPITN score of ‘healthy’ had slightly reduced. There was a noticeable increase in the proportion of people with calculus and shallow periodontal pockets, but a substantial reduction in the proportion with deep periodontal pockets. It is also important to note the fall in the number of excluded sextants (X), which indicates that more sextants had the minimum number of teeth for the CPITN examination to be carried out. The increase in the proportion with calculus and shallow pockets probably also reflects increased tooth retention.

Data on the health of the population is a key part of identifying needs, planning public health strategies and assessing the effectiveness of public health policies. This is especially true of oral health, which is sensitive to socioeconomic conditions, dietary trends, lifestyle and access to care. At the same time, most dental disease is preventable so there is the potential for improvement in oral health at relatively low cost²⁹. The last national survey of adult oral health in Ireland was conducted in 2000–02 and the results were published in 2007. With the implementation of a new national oral health policy underway it is critical to understand the current oral health status of older adults in Ireland. The TILDA study was a valuable opportunity to examine the oral health of a nationally representative cohort of older Irish adults: this cohort has

been extensively characterised in terms of their physical and mental health, wellbeing, social interactions, and socioeconomic status.

The respondents who participated in the dental assessment were similar to the whole TILDA cohort in the key areas of gender, medical card status, urban-rural dwelling, and self-reported health status. The dental group had more respondents from Dublin, were younger, and were more likely to have tertiary/higher level education than the nationally representative TILDA cohort. Comparing this study to previous national surveys, the prevalence of edentulism has continued to decline rapidly in Irish adults, following the trend seen in many other English-speaking countries, though it was noted that older women are still more likely than men to have no natural teeth at all. The comparison also highlights economic, cultural, and historic differences between countries with respect to edentulism; for example New Zealand had a much higher prevalence of edentulism than the UK at the same time point, perhaps due to sociological reasons³⁰.

Having at least 20 natural teeth, or 10 pairs of contacting teeth, is considered a benchmark of a functional dentition, that is a dentition that will generally provide adequate functional capacity and may not require additional prosthetic teeth²¹. This study found that 56.8% of the dental sample had 10 or more pairs of contacting teeth, though as with other variables, there was a marked difference between the younger and older age groups. This means that there is a much larger need for intervention in the form of replacement teeth among older adults in Ireland, though a report on this cohort found that people with fewer teeth were actually less likely to access dental services³¹.

One of the concerns raised about maintaining oral health in older people is that, paradoxically, the retention of natural teeth incurs more ongoing care needs than a complete loss of teeth.

Table 15. Mean DMFT (WHO method) by age groups, country and examination year.

Country	Year of Examination	Age group	Mean DMFT
Ireland	2014-15	50-64 years	20.1
		65-74 years	23.7
		75 years and over	26.2
New Zealand* ²⁷	2009	55-64 years	21.7
		65-74 years	24.2
		75 years and over	24.8
Australia* ²⁶	2004-06	55-64 years	21.7
		65-74 years	23.2
		75 years and over	24.6

*Dentate sample only

Table 16. Changes over time in percentage of adults with maximum value of CPITN-severity score of H (healthy), B (bleeding), C (calculus), P1 (shallow pocket), P2 (deep pocket) and X (missing sextant) among dentate adults 65 years and over in Ireland (Base dentate).

Year of examination	H	B	C	P1	P2	X	Total (n)
2014-15	6.6	4.0	35.1	46.4	4.7	4.3	1085
2000-02	6.9	3.6	29.5	37.6	12.0	10.1	390

Without adequate regular plaque control, a healthy diet and some professional dental care, natural teeth can be susceptible to caries and periodontal disease which may lead to pain, infection and loss of function. The susceptibility of older adults to dental disease may be heightened due to age-related periodontal recession and an increase in medications that cause dry mouth. In Ireland, adults who become frail, dependent, or live in residential care will often lose access to regular dental services. This study found a low level of untreated crown and root caries, and mostly mild periodontal disease, which suggest that independently-living older adults can maintain their oral health fairly well. Nonetheless, the level of root caries in those aged 75 years or older was double that of the younger age groups. The challenge will be to provide adequate care for the growing number of frail older people who, in the future, will have a far greater number of natural teeth.

This study made a broad assessment of dental status, using a methodology that allows for comparison to previous national oral health surveys and similar international studies. Other strengths of the study design were the large sample size and the extensive information collected during the CAPI, SCQ and health assessment. This will be extremely useful in further analysis that will link the objective measures of dental health with other health outcomes. The dental sample also had similar characteristics to the TILDA cohort, suggesting that the findings should be representative of the overall sample. However, the study design had some limitations. The participants

in the dental assessment were recruited from those attending a health centre assessment; participants who had a home-based assessment or did not complete a health assessment were older and had poorer general health indicators than those who did attend a health centre³² so it is likely that the oral health status of these participants is also poorer than the dental sample. Therefore, while it was possible to generalise these results to the majority of the community-dwelling population aged 50 years and over, they may underestimate the prevalence and extent of oral health problems in adults with poorer health and those in residential care.

Conclusions

Data from this study shows that there has been considerable improvement in the retention of teeth among community dwelling adults aged 50 years and over in Ireland, since previous Irish national surveys. Since 1989-90, the proportion of edentate adults aged 65 years and over has decreased by two-thirds, and the mean number of natural teeth has more than doubled. However, tooth loss was still common among older adults, posing a challenge for our health services. Currently, there is little service provision for dentate older adults, especially those who become frail or dependent, suggesting the need for a qualitative change in the oral care for older people in Ireland.

If generalised to the whole population, these findings suggest that Ireland has more slowly followed international trends for improvement in the oral health status of older adults,

observed in other developed countries like the UK, USA, Australia and New Zealand. These trends are very positive but they indicate a requirement for more maintenance care, restorative and periodontal treatment, and less need for complete dentures than previously. The findings of this study should provide a valuable resource for oral health policy and planning of oral health services for older people in Ireland.

Data availability

The data presented in this report was collected during Wave 3 of TILDA. Wave 3 data is available from the Irish Social Science Data Archive (ISSDA):

ISSDA: Dataset 1. The Irish Longitudinal Study on Ageing (TILDA) Wave 3, 2014–2015. Study number: 0053-04. www.ucd.ie/issda/data/tilda/wave3³³.

Accessing the data

To access the data, please complete a [ISSDA Data Request Form for Research Purposes](#), sign it, and send it to ISSDA by email (issda@ucd.ie).

For teaching purposes, please complete the [ISSDA Data Request Form for Teaching Purposes](#), and follow the procedures, as above. Teaching requests are approved on a once-off module/workshop basis. Subsequent occurrences of the module/workshop require a new teaching request form.

Data will be disseminated on receipt of a fully completed, signed form. Requests to access the dental assessment data should be made directly to TILDA (tilda@tcd.ie)

Consent

Ethical approval for this study was obtained from the Faculty of Health Sciences Research Ethics Committee in Trinity College Dublin and participants provided written informed consent before the health assessment.

Acknowledgements

The authors would like to acknowledge the contribution of the TILDA participants and TILDA team, especially Margaret Foley, Ann Hever and Christine McGarrigle.

References

- United Nations: **World Population Ageing 2015**. United Nations. Department of Economic and Social Affairs Population Division: United Nations New York, 2015.
[Reference Source](#)
- Central Statistics Office: **Population and labour force projections 2016-2046**. Central Statistics Office, Editor. Published by the Stationery Office, Dublin, Ireland.: Dublin. 2013; 38.
[Reference Source](#)
- Central Statistics Office: **Census 2016 Summary Results - Part 1**. Central Statistics Office, Ireland: Ireland, 2017.
[Reference Source](#)
- Benjamin RM: **Oral health: the silent epidemic**. *Public Health Rep*. 2010; **125**(2): 158–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Whelton H, Crowley E, O'Mullane D, et al.: **Oral health of Irish adults 2000-2002: final report-April 2007**. Dublin: Department of Health and Children, 2007.
[Reference Source](#)
- Gerritsen S, Stefanogiannis N, Galloway Y, et al.: **A portrait of health: Key results of the 2006/07 New Zealand health survey**. Ministry of Health, 2008.
[Reference Source](#)
- Thorstensson H, Johansson B: **Why do some people lose teeth across their lifespan whereas others retain a functional dentition into very old age?** *Gerodontology*. 2010; **27**(1): 19–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Steele JG, Treasure ET, O'Sullivan I, et al.: **Adult dental health survey 2009: Transformations in British oral health 1968-2009**. *Br Dent J*. 2012; **213**(10): 523–527.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Polzer I, Schimmel M, Müller F, et al.: **Edentulism as part of the general health problems of elderly adults**. *Int J Epidemiol*. 2011; **40**(4): 877–884.
[PubMed Abstract](#)
- Kearney PM, Cronin H, O'Regan C, et al.: **Cohort profile: the Irish longitudinal study on ageing**. *Int J Epidemiol*. 2011; **40**(4): 877–884.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Donoghue OA, McGarrigle CA, Foley M, et al.: **Cohort Profile Update: The Irish Longitudinal Study on Ageing (TILDA)**. *Int J Epidemiol*. 2018; **47**(5): 1398–1398l.
[PubMed Abstract](#) | [Publisher Full Text](#)
- World Health Organization: **Oral Health Survey Basic Methods, 5th edition**. 2013.
[Reference Source](#)
- Pitts NB, Evans DJ, Pine CM: **British Association for the Study of Community Dentistry (BASCD) diagnostic criteria for caries prevalence surveys-1996/97**. *Community Dent Health*. 1997; **14 Suppl 1**: 6–9.
[PubMed Abstract](#)
- Katz RV, Hazen SP, Chilton NW, et al.: **Prevalence and intraoral distribution of root caries in an adult population**. *Caries Res*. 1982; **16**(3): 265–271.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bardsley PF: **The evolution of tooth wear indices**. *Clin Oral Investig*. 2008; **12 Suppl 1**: S15–19.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Witter DJ, van Palenstein Helderma WH, Creugers NH, et al.: **The shortened dental arch concept and its implications for oral health care**. *Community Dent Oral Epidemiol*. 1999; **27**(4): 249–258.
[PubMed Abstract](#)
- Kanno T, Carlsson GE: **A review of the shortened dental arch concept focusing on the work by the Käyser/Nijmegen group**. *J Oral Rehabil*. 2006; **33**(11): 850–862.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Armellini D, von Fraunhofer JA: **The shortened dental arch: a review of the literature**. *J Prosthet Dent*. 2004; **92**(6): 531–535.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Allen PF, Witter DF, Wilson NH, et al.: **Shortened dental arch therapy: views of consultants in restorative dentistry in the United Kingdom**. *J Oral Rehabil*. 1996; **23**(7): 481–485.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Khan S, Musekiwa A, Chikte UM, et al.: **Differences in functional outcomes for adult patients with prosthodontically-treated and -untreated shortened dental arches: a systematic review**. *PLoS One*. 2014; **9**(7): e101143.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Steele J, Sheiham A, Marcenes W, et al.: **National diet and nutrition survey: people aged 65 years and over**. Volume 2: Report of the oral health survey. 1998.
[Reference Source](#)
- Data Protection Commissioner: **Data Protection Acts 1988 and 2003. Informal Consolidation**. 1988 and 2003.
[Reference Source](#)
- O'Mullane D, Whelton H: **Oral health of Irish adults 1989-1990: a survey**

- conducted by the Oral Health Services Research Centre, University College Cork. Stationery Office, 1992.
[Reference Source](#)
24. O'Sullivan I, Lader D, Beavan-Seymour C, *et al.*: **Foundation Report: Adult Dental Health Survey 2009**. London: The Health and Social Care Information Centre. Dental and Eye Care Team, 2011.
[Reference Source](#)
 25. Dye BA, Thornton-Evans G, Li X, *et al.*: **Dental caries and tooth loss in adults in the United States, 2011-2012**. US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics, 2015.
[Reference Source](#)
 26. Slade GD, Spencer AJ, Roberts-Thomson KF: **Australia's dental generations: The National Survey of Adult Oral Health 2004-06**. A.J.S. Gary D Slade and a.K.F. Roberts-Thomson, Editors. Australian Institute of Health and Welfare, Australian Government, 2007.
[Reference Source](#)
 27. Haisman RJ, Mason K, Holmes E, *et al.*: **Our oral health: Key findings of the 2009 New Zealand Oral health Survey**. Ministry of Health, New Zealand.: PO Box 5013, Wellington 6145, New Zealand. 2010; Published in December 2010.
[Reference Source](#)
 28. Health Canada: **Summary Report on the Findings of the Oral Health Component of the Canadian Health Measures Survey, 2007-2009**. M.o.H. Canada., Editor. Ottawa, Ontario K1A 0K9, 2010.
[Reference Source](#)
 29. Righolt AJ, Jevdjevic M, Marcenes W, *et al.*: **Global-, regional-, and country-level economic impacts of dental diseases in 2015**. *J Dent Res*. 2018; **97**(5): 501–507.
[PubMed Abstract](#) | [Publisher Full Text](#)
 30. Ministry of Health New Zealand: **Our Oral Health: Key findings of the 2009 New Zealand Oral Health Survey**. Ministry of Health. PO Box 5013, Wellington 6145, New Zealand, 2010.
[Reference Source](#)
 31. Sheehan A, McGarrigle AC, O'Connell B: **Oral Health and Wellbeing of Older Irish Adults**. The Irish Longitudinal study on Ageing (TILDA), 2017.
[Publisher Full Text](#)
 32. Cronin H, O'Regan C, Finucane C, *et al.*: **Health and aging: development of the Irish Longitudinal Study on Ageing health assessment**. *J Am Geriatr Soc*. 2013; **61** Suppl 2: S269–S278.
[PubMed Abstract](#) | [Publisher Full Text](#)
 33. The Irish Longitudinal study on Ageing: **The Irish Longitudinal study on Ageing (TILDA) Wave 3, 2014-2015**. [dataset]. Version 3.1. Irish Social Science Data Archive. 2018.
<http://www.ucd.ie/issda/data/tilda/wave3/>

Open Peer Review

Current Peer Review Status:  

Version 3

Reviewer Report 09 February 2022

<https://doi.org/10.21956/hrbopenres.14705.r31432>

© 2022 Thomson W. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



W. Murray Thomson 

Department of Oral Sciences, Sir John Walsh Research Institute, Faculty of Dentistry, University of Otago, Dunedin, New Zealand

Congratulations to the authors for a fine report. I have no further comments or suggestions.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Dental epidemiology, gerodontology, health services research

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 2

Reviewer Report 24 September 2021

<https://doi.org/10.21956/hrbopenres.14597.r30347>

© 2021 Thomson W. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



W. Murray Thomson 

Department of Oral Sciences, Sir John Walsh Research Institute, Faculty of Dentistry, University of Otago, Dunedin, New Zealand

Overall, the paper is much improved – well done! I particularly like how the Introduction has improved.

I have only minor points for the authors:

- In the Discussion, “compared to” is still used in a number of places – surely “than” is a better word to use than that awful phrase. In the 3rd Discussion paragraph, the term “trend” is misused – “similar trends in the prevalence of edentulism with age” should read “similar age differences in edentulism prevalence”.
- Finally, the authors point out that it is difficult to compare their survey’s mean number of teeth with estimates from other national surveys because the latter have not included edentulous people in the denominator. Shouldn’t that be a very strong signal to them that one does not do that? That an estimate for the mean number of teeth should be just for the dentate sample?? I think they need to recalculate their mean number of teeth accordingly so that they are consistent with standard practice.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Dental epidemiology, gerodontology, health services research

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 05 Jan 2022

Brian O'Connell, Trinity College Dublin, Dublin, Ireland

- In the Discussion, “compared to” is still used in a number of places – surely “than” is a better word to use than that awful phrase. In the 3rd Discussion paragraph, the term “trend” is misused – “similar trends in the prevalence of edentulism with age” should read “similar age differences in edentulism prevalence”.

Thank you for spotting this. The recommended changes have been made in the attached revised version.

- Finally, the authors point out that it is difficult to compare their survey’s mean number of teeth with estimates from other national surveys because the latter have not included edentulous people in the denominator. Shouldn’t that be a very strong signal to them that one does not do that? That an estimate for the mean number of teeth should be just for the dentate sample?? I think they need to recalculate their mean number of teeth accordingly so that they are consistent with standard practice.

In both Irish adult surveys the mean number of teeth was calculated using the dentate and edentate samples. For that reason we calculated the mean number of teeth by the same method and reported this in the results and for national comparison (Table 12). However, as per your suggestion for international comparisons the mean number of teeth was calculated in the dentate sample only and the international comparison (Table 14) was updated.

Competing Interests: No competing interests were disclosed.

Version 1

Reviewer Report 31 January 2019

<https://doi.org/10.21956/hrbopenres.13960.r26445>

© 2019 Mulligan R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Roseann Mulligan**

Division of Dental Public Health and Pediatric Dentistry, Herman Ostrow School of Dentistry, University of Southern California, Los Angeles, CA, USA

Overview

There have been three waves of The Irish Longitudinal Study on Ageing (TILDA). In this work, the third wave is described during which time oral examinations were completed for a subset of participants aged 50 and over. The World Health Organization examination criteria was used to assess oral health as well as a number of previously used indices. Data on a variety of oral health indicators including number of retained teeth, number in occlusion, presence/absence of prostheses, caries, including root caries and periodontal disease indicators are presented and the results compared to previous Irish studies and findings from studies in other countries using similar criteria.

Overall comments:

There is a large amount of data of all types that was collected into this one paper. Typically the results of large studies are published in multiple papers that can orient the reader toward the findings in various discrete areas, such as edentulism, caries, periodontal findings, and allow discussion about each respective area. This paper has some of the characteristics of a monograph, displaying the results of all the data collected relating to all oral health findings and therefore the multitude of tables and figures is to be expected. I appreciate having all the OHA data in the TILDA displayed in one publication and suggest that the authors also provide an appendix of all the figures that one can readily negotiate when looking for a particular topic. However, unlike a monograph there is a lack of detailed discussion of findings relative to data about each specific finding. Usually only one reference is mentioned per topic and therefore the richness and meaning of the findings as compared to that found in other studies is often not expanded or expounded upon.

Comments on the Introduction:

While the sample size for the Oral Health Assessment (OHA) equaled 2504 individuals aged 50 and over, nearly ½ of the sample was made up of middle-aged adults from 50 to 64 year olds and those aged 75 and older only accounted for 14.8% (N = 371) of the entire OHA substudy. In most developed nations, 50 year olds are not considered elderly nor are their oral health findings that different from other middle-agers. Since TILDA is a Longitudinal Study on Aging, I was disappointed not to find more data about older adults, with details highlighting those 65+ and older by decades. Clearly the small sample size for the 75+ aged population precluded the ability of the investigators to analyze and present data. Therefore the authors' focus in the Introduction

that brings to the readers' attention the significant growth occurring in the 65 and older and 80 and older age groups is puzzling as it whets the appetite to learn new findings that are not in fact available or presented. I suggest rewriting most of the Introduction to remove this emphasis.

Comments about Methodology:

If the goal was to achieve robust cell sizes of older adults for analytic purposes and to make policy recommendations, I wonder why an oversampling methodology of the oldest categories was not used. Since this study appears to be a convenience sample of the overall TILDA pool of subjects attending a Health Center in Dublin, I have concern about its ability to serve as a representation of the general population of Ireland, which appears to be an objective of TILDA. Should there be future TILDA with OHA components, over sampling would be valuable of older age groups and I would strongly recommend such a strategy. If a longitudinal study is planned for the future, the current middle aged and newly aged elderly of the OHA substudy of TILDA 3 could serve as a baseline population for another wave. Still the subject selection would need to be improved upon by including cohorts outside of Dublin, so that the results have true national representativeness.

Comments on Results:

Displaying the large amounts of data is challenging, I can appreciate the authors attempts to include all of their findings in this manuscript. I find the tables much more helpful as they actually show the results with the sample sizes noted at the basic levels, and for the most part the data is not collapsed into large categories, thus making it easier to compare with other studies. For those who want to make future comparisons of this data to newer datasets, it will be the Tables that will provide the data cited. The Figures were more difficult to interpret even though visually more appealing with their display of comparative and trend data; they have less practicality or comparability. Figure 4 contains a highly pictorial and easy to grasp display of the separate components of the DMFT findings by three age adult categories. This is the figure I suggest the authors retain.

In the Results section there are many comparisons between previous findings in earlier Irish studies, however no statistical analysis were completed to determine if these differences were significant. Table 17 highlights the difficulty in this approach with the comparisons between the 2000-02 and 2014-15 studies described in the narrative as "slightly" and "substantial" to describe differences rather than indicating on the table whether or not there are statistical differences between the two studies using convention p value cut-offs.

While much is made of the valuable opportunity the TILDA study presents due to its extensive record of physical, mental and socioeconomic data on participants, few correlations were drawn between the oral findings outside of gender and middle to early older age groups due to sampling inadequacies (recruited small sample size of 75 and older individuals and the limited sample collection site.) The opportunity to better characterize those variables associated with individuals with the greatest oral health needs was missed.

Because there are so many tables, it is difficult to flip through different pages to find the table and narrative when they are not in proximity. While sometimes it is inevitable that tables and narratives are not on the same page, more effort needs to be made by the editor to place some of the smaller tables in close proximity to the relevant discussion.

Comments on Discussion:

I suggest that the authors be more careful about making generalizations about cause. In the discussion of the RCI findings on page 13, the authors state without citation that the explanation for an expected increase in root caries would be: “. . . because teeth tend to lose periodontal attachment with age and exposed root surfaces are vulnerable to the development of root caries.” While many would agree with the second statement, not all would agree with the first statement that periodontal disease as defined by attachment loss is an inevitable component of aging. Nor does this type of statement belong in the Results section for this discussion of potential causality should be covered in the Discussion.

I find statements such as, “The increase in the proportion with calculus and shallow pockets probably also reflects increased tooth retention.” If increasing tooth retention is aligned with calculus and shallow pockets, why not the other measured variables that trended in the same direction? If statistically significant comparisons were made and posted on the table, the readers would be able to determine if for example whether the increase in bleeding or the decrease in healthy tissue were significant? I again suggest that statistical analysis be run and the results indicated on the table so that readers can draw conclusions. While it might be difficult to compare results from various countries due to differing methodologies, where the CPITN was used in both Irish studies, one would expect some comparisons so that the importance or lack thereof could be appreciated.

It is in the Discussion section that one learns of the difficulties and differences in the OHA sub sample. The population sampled should have been described in the Methods section, not in the Discussion section. The characteristics defining the two different populations sampled may explain some of the significant jumps in improved health between the older Irish studies and the current studies in spite of the fact that the current OHA subgroup is representative of the TILDA study overall (Table 1). As a result the conclusions from this study may not be generalizable to the entire Irish population, especially those not living in the urban capital of the country.

Comments on Conclusions:

Conclusions should be based on the work and findings presented in this manuscript. Yet the authors say, “Currently, there is little service provision for dentate older adults, especially those who become frail or dependent, suggesting the need for a qualitative change in the oral care for older people in Ireland.” I am not sure why this statement is made in the Conclusion of the paper as this topic was not a part of the study and cannot be concluded from the data collected.

Closing Thoughts:

In the beginning of the article, the authors discuss the importance of this data to better inform policy makers in the development of a new national health care policy; however, little data was collected on the truly elderly population. This is a missed opportunity as limited health care dollars require that the at-risk population be studied to enable targeting of limited funds to cohorts at greatest need. I encourage future OHA studies, whether as part of TILDA or not, to do a better job of oversampling, especially the middle and oldest groups of elderly (75+ and 85+ years of age) and those who are living in a multitude of living arrangements and geographic settings, so that the most accurate picture of oral disease in Irish elderly is captured.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 09 January 2019

<https://doi.org/10.21956/hrbopenres.13960.r26444>

© 2019 Thomson W. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



W. Murray Thomson 

Department of Oral Sciences, Sir John Walsh Research Institute, Faculty of Dentistry, University of Otago, Dunedin, New Zealand

General comments

Paper reports the findings of a simple descriptive analysis of oral epi data from participants in a cohort study of adults aged 50+. The title and abstract refer to “oral health status”, but what is actually being described here is tooth retention, dental caries experience and a snapshot of periodontal status, rather than “oral health” per se. The presentation of findings is not as systematic and easy to follow as it should be, because the authors give the impression of not being sure whether this paper will be a simple overview of the oral characteristics of the sample or a comparison of changes over time – and the Results section reflects that, with data in the later stages being brought in from earlier Irish work and from overseas studies. I suggest that they confine the earlier Irish work and overseas data to the Discussion. A purely descriptive analysis like this one should be carefully and systematically presented, with a clear boundary between the study data and their comparison with earlier findings.

The authors insist on using the awful “compared with”; I suggest that they desist. Thus, “Of those

aged 65 years and older, 15.6% were dentate compared with 40.9% in 2000-02" should be written as "Of those aged 65 years and older, 15.6% were dentate, while it was 40.9% in 2000-02" (or somesuch).

There are too many acronyms – do we really need OHA and MIP?

There are far too many small Tables – the authors should concentrate on having larger, more efficient Tables where the sociodemographic (and other) associations are presented to the reader.

Section comments

Title

The title is unnecessarily long—a shorter and more appropriate one is indicated.

Abstract

This will need to be rewritten once the paper has been revised, but the authors should consider the following: do not leave the reader to calculate %; be consistent with the number of decimal places for the % that you do present (these should be to 1 decimal place); do not use a hackneyed phrase such as "the results suggest"; and the conclusion should be a valid conclusion (what's the take-home message from the study?).

Introduction

This is rather long and meandering. There are contentious assertions in paragraph 2 which should be omitted or tempered. The first clause of sentence 2 is vague – what do the AU mean here? Similarly, sentence 3 is unnecessary and can be contested – the great bulk of the evidence for the so-called "bidirectional relationship" is weak, fails to account for confounders, and is mostly the outcome of wishful thinking in interpreting cross-sectional data. Oral health is important enough in its own right without our having to try to justify investigating it using spurious associations with general health conditions.

In para 3, sentence 3 has structural problems and should be rewritten.

The last para of the Intro is actually Discussion and does not belong in the Intro.

Methods

There is insufficient information on the sampling procedures and what the sampling frame was. We are not told in the right place in Methods whether it is a simple random sample or a complex sample. Is it a national sample, or just people in and around Dublin? Also, these were people in Wave 3 – does this mean that they were also assessed earlier in Waves 1 and 2? If so, we need more information on that, and particularly on participation rates and loss to follow-up in earlier assessments.

Hyphenate "cross infection".

The assessment of tooth contacts in the dentition does not actually mention the maxillary teeth – I presume that they are referring to contacts between the maxillary and mandibular teeth. And the term "functionality of the dentition" is unusual.

The RCI is not a measure of root surface caries prevalence, despite Katz having used that term in his original paper. The prevalence of a condition is the proportion of the population who have the condition – that should be reported here, alongside the RCI and the mean number of root DFT. The RCI is a sort of “attack rate” indicator.

Use the term “denture wearing”, not “denture wear” – that’s something else entirely...

Results

As a general principle, have one paragraph of Results text per Table and do not repeat Table data in the accompanying text (instead, summarise). Do not discuss your findings in the Results section (e.g. Results para 3, sentence 2 is Discussion).

Re: the data in Table 1 – describe the Table as reporting the sample characteristics, etc – the actual statistical testing is secondary. While on the latter, the authors do themselves a disservice, I think – while a lot of those P values are <0.05, this is due to the large N rather than any meaningful differences in proportions. For example, 27.1% and 24.1% living in Dublin – each of those I would summarise as approximately 1 in 4. Doing the analyses by each category of age group (or education level, etc) means a lot of tests – why not one cross-tabulation for each domain rather than 3?

Results para 4, last sentence – you cannot say that the gender difference “decreased with increasing age” – it was less marked in those who were older; you did not see it decrease... This is actually a surprising finding, by the way – in most epi studies of older people, you will see higher proportions of females in the older age groups. Why is yours different?

Table 3 should also present the edentulous and dentate proportions for each age group (genders combined). Table 4 is not really a stand-alone Table – consider presenting that information in Table 3 as well – you are going to have to restructure it anyway. Again, you cannot say (as you do at the end of page 4) that the proportion of edentate adults increased with age – again, it was higher in those who were older.

Table 5 should refer to denture wearing (see above!). Again, why repeat the table data in the text? We can see the Table, so point out the highlights. Again, there is the problem phrase “increased with age” (see above)...

Figure 1 – is this necessary? (ditto Figures 2, 3 and 4 – especially Figure 4...). Whereas Tables are a very efficient way of presenting a lot of information, Figures are not, and that information can be summarised and presented efficiently in the Results text – or as a Table, with mean numbers and SDs presented by sociodemographic characteristics – that would certainly be more informative. The last sentence of the “number of teeth” section is actually Discussion – and two data points do not make a “trend”.

To the periodontal data – the CPI is a bad index to use anyway, and even worse when used in a longitudinal study. It’s too late now to point out that attachment loss data should have been collected. Again, the presentation of the data is not as informative as it might have been.

Discussion

See my earlier comments (page 1).

The last para of the Discussion misuses the term “extrapolated” – that should be “generalised”.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Dental epidemiology, gerodontology, health services research

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 18 Aug 2021

Brian O’Connell, Trinity College Dublin, Dublin, Ireland

General comments

Paper reports the findings of a simple descriptive analysis of oral epi data from participants in a cohort study of adults aged 50+. The title and abstract refer to “oral health status”, but what is actually being described here is tooth retention, dental caries experience and a snapshot of periodontal status, rather than “oral health” per se. The presentation of findings is not as systematic and easy to follow as it should be, because the authors give the impression of not being sure whether this paper will be a simple overview of the oral characteristics of the sample or a comparison of changes over time – and the Results section reflects that, with data in the later stages being brought in from earlier Irish work and from overseas studies. I suggest that they confine the earlier Irish work and overseas data to the Discussion. A purely descriptive analysis like this one should be carefully and systematically presented, with a clear boundary between the study data and their comparison with earlier findings.

Thank you for your helpful review and comments. We have done our best to reflect these

suggestions in the revised version of the paper. Throughout the paper and in the title 'oral health' has been changed to 'dental health' and we have limited comparisons with previous and international data to the Discussion section.

The authors insist on using the awful "compared with"; I suggest that they desist. Thus, "Of those aged 65 years and older, 15.6% were dentate compared with 40.9% in 2000-02" should be written as "Of those aged 65 years and older, 15.6% were dentate, while it was 40.9% in 2000-02" (or somesuch).

This has been amended throughout the document

There are too many acronyms – do we really need OHA and MIP?

We have removed OHA throughout the document and reduced the use of other acronyms as much as possible

There are far too many small Tables – the authors should concentrate on having larger, more efficient Tables where the sociodemographic (and other) associations are presented to the reader.

We have considered your suggestion about reporting the relationship between dental health and socio-economic status. This paper is already very long so we have decided to publish a separate paper on this topic by using socioeconomic variables from TILDA dataset. We have removed Table 4.

Section comments

Title

The title is unnecessarily long—a shorter and more appropriate one is indicated.

We understand the reviewer's comments but the publisher requires a full description of the study type in the title according to the STROBE guidelines

Abstract

This will need to be rewritten once the paper has been revised, but the authors should consider the following: do not leave the reader to calculate %; be consistent with the number of decimal places for the % that you do present (these should be to 1 decimal place);

The percentages reported have been checked and all are now at one decimal place.

do not use a hackneyed phrase such as "the results suggest"; and the conclusion should be a valid conclusion (what's the take-home message from the study?).

Changes has been made in the conclusions section to focus on the relevance of the findings of this cross-sectional study. We have removed the phrase 'the results suggest'

Introduction

This is rather long and meandering. There are contentious assertions in paragraph 2 which should be omitted or tempered. The first clause of sentence 2 is vague – what do the AU mean here? Similarly, sentence 3 is unnecessary and can be contested – the great bulk of the evidence for the so-called "bidirectional relationship" is weak, fails to account for confounders, and is mostly the outcome of wishful thinking in interpreting cross-sectional data. Oral health is important enough in its own right without our having to try to justify investigating it using spurious associations with general health conditions.

These suggestions have been included in the revisions. The focus in the introduction is on dental health indicators and any association between dental health and systemic health.

In para 3, sentence 3 has structural problems and should be rewritten.
This paragraph has been rewritten and is more focused.

The last para of the Intro is actually Discussion and does not belong in the Intro.
The last paragraph has been removed

Methods

There is insufficient information on the sampling procedures and what the sampling frame was. We are not told in the right place in Methods whether it is a simple random sample or a complex sample. Is it a national sample, or just people in and around Dublin? Also, these were people in Wave 3 –

We have given more information in the text to clarify these points.

does this mean that they were also assessed earlier in Waves 1 and 2? If so, we need more information on that, and particularly on participation rates and loss to follow-up in earlier assessments.

The dental health assessment was introduced in TILDA Wave 3 so we only have cross-sectional data. However, we are aiming to repeat dental assessment in the next TILDA wave to have longitudinal data.

Hyphenate “cross infection”.

This has been corrected.

The assessment of tooth contacts in the dentition does not actually mention the maxillary teeth – I presume that they are referring to contacts between the maxillary and mandibular teeth. And the term “functionality of the dentition” is unusual.

That is correct—tooth contacts (in Maximum Intercuspal Position) means contact between opposing pairs of teeth--this has been clarified in the text. The more usual term ‘functional dentition’ has been used to express the concept of a dentition that is adequate for normal function.

The RCI is not a measure of root surface caries prevalence, despite Katz having used that term in his original paper. The prevalence of a condition is the proportion of the population who have the condition – that should be reported here, alongside the RCI and the mean number of root DFT. The RCI is a sort of “attack rate” indicator.

Yes, we agree. It represents the proportion of exposed root surfaces with root caries and it was used to allow for comparisons to be made with previous Irish national surveys. The tables also report the mean number of teeth with decayed and filled roots. The text has been changed in the Methods section to clarify the meaning of the RCI.

Use the term “denture wearing”, not “denture wear” – that’s something else entirely...

The term has been changed to ‘denture wearing’.

Results

As a general principle, have one paragraph of Results text per Table and do not repeat Table data in the accompanying text (instead, summarise). Do not discuss your findings in the Results section (e.g. Results para 3, sentence 2 is Discussion).

The Results section has been revised as recommended.

Re: the data in Table 1 – describe the Table as reporting the sample characteristics, etc

The title of the table has been changed as suggested.

the actual statistical testing is secondary. While on the latter, the authors do themselves a disservice, I think – while a lot of those P values are <0.05 , this is due to the large N rather than any meaningful differences in proportions. For example, 27.1% and 24.1% living in Dublin – each of those I would summarise as approximately 1 in 4. Doing the analyses by each category of age group (or education level, etc) means a lot of tests – why not one cross-tabulation for each domain rather than 3?

In order to compare the characteristics of the two different-sized samples--dental health and total TILDA -- the two-sample proportion test was chosen (following the advice of two independent experienced biostatisticians).

For the cross tabulation we measured the relationship between different categories of two variables. Although the proportions were not much different between the full TILDA and the dental samples, the Z test showed similarities between the two samples for 8 out of 23 characteristics (Table 1).

Results para 4, last sentence – you cannot say that the gender difference “decreased with increasing age” – it was less marked in those who were older; you did not see it decrease...

We understand this wording was misleading so all wording in results has been changed from decreased with age to the reporting of comparison between two age groups.

This is actually a surprising finding, by the way – in most epi studies of older people, you will see higher proportions of females in the older age groups. Why is yours different?

This may be because the dental sample is a convenience sample, unlike the TILDA cohort, which is nationally representative.

Table 3 should also present the edentulous and dentate proportions for each age group (genders combined).

Given the finding in previous national surveys that the prevalence of edentulism was much higher in females (Table 12), it was considered notable to demonstrate the change in this trend over time and among the different age cohorts in the present study

Table 4 is not really a stand-alone Table – consider presenting that information in Table 3 as well –

Table 4 has been deleted and the main results given in the text.

you are going to have to restructure it anyway. Again, you cannot say (as you do at the end of page 4) that the proportion of edentate adults increased with age – again, it was higher in those who were older.

This wording has been changed throughout the document.

Table 5 should refer to denture wearing (see above!). Again, why repeat the table data in the text? We can see the Table, so point out the highlights. Again, there is the problem phrase “increased with age” (see above)...

As reported above, it has been changed

Figure 1 – is this necessary? (ditto Figures 2, 3 and 4 – especially Figure 4...). Whereas Tables are a very efficient way of presenting a lot of information, Figures are not, and that information can be summarised and presented efficiently in the Results text – or as a Table, with mean numbers and SDs presented by sociodemographic characteristics – that would certainly be more informative.

As the data was not categorical or normally distributed, we believe the figures give added information regarding the data distribution. This was limited to four figures, but can be reduced further if required.

We agree with the suggestion, also mentioned previously, to publish a separate paper on dental health according to sociodemographic characteristics.

The last sentence of the “number of teeth” section is actually Discussion

This has been removed – and two data points do not make a “trend”.

We understand the wording was misleading so it has been changed.

To the periodontal data – the CPI is a bad index to use anyway, and even worse when used in a longitudinal study. It’s too late now to point out that attachment loss data should have been collected. Again, the presentation of the data is not as informative as it might have been.

Yes, we understand CPI is not an ideal index for longitudinal studies but the dental assessment was allocated only 10 minutes in the study, which was not enough time to measure attachment loss. Also, there was only time to measure index teeth. We have requested a more comprehensive periodontal measurements in the next wave of TILDA, but as the whole health assessment takes three hours or more, this remains to be decided. We understand the onerous nature of a lengthy assessment for respondents, some of whom are frail and do not even understand why a dental assessment is part of the health assessment! But that is another matter.

Discussion

See my earlier comments (page 1).

The Discussion has been revised according to these suggestions

The last para of the Discussion misuses the term “extrapolated” – that should be “generalised”.

The wording has been changed

Thank you again for your in depth and valuable input. We hope the changes described above have improved the paper and addressed your significant reservations.

Competing Interests: No competing interests were disclosed.