

Recombinant *S. cerevisiae* expressing Old Yellow Enzymes from non-conventional yeasts: an easy system for selective reduction of activated alkenes

Romano *et al.*

RESEARCH

Open Access

Recombinant *S. cerevisiae* expressing Old Yellow Enzymes from non-conventional yeasts: an easy system for selective reduction of activated alkenes

Diego Romano², Martina Letizia Contente², Francesco Molinari², Ivano Eberini³, Erica Ruvutuso³, Cristina Sensi³, Alberto Amaretti¹, Maddalena Rossi¹ and Stefano Raimondi^{1*}

Abstract

Background: Old Yellow Enzymes (OYEs) are flavin-dependent enoate reductases (EC 1.6.99.1) that catalyze the stereoselective hydrogenation of electron-poor alkenes. Their ability to generate up to two stereocenters by the *trans*-hydrogenation of the C=C double bond is highly demanded in asymmetric synthesis. Isolated redox enzymes utilization require the addition of cofactors and systems for their regeneration. Microbial whole-cells may represent a valid alternative combining desired enzymatic activity and efficient cofactor regeneration. Considerable efforts were addressed at developing novel whole-cell OYE biocatalysts, based on recombinant *Saccharomyces cerevisiae* expressing OYE genes.

Results: Recombinant *S. cerevisiae* BY4741 Δ Oye2 strains, lacking endogenous OYE and expressing nine separate OYE genes from non-conventional yeasts, were used as whole-cell biocatalysts to reduce substrates with an electron-poor double bond activated by different electron-withdrawing groups. Ketoisophorone, α -methyl-*trans*-cinnamaldehyde, and *trans*- β -methyl- β -nitrostyrene were successfully reduced with high rates and selectivity. A series of four alkyl-substituted cyclohex-2-enones was tested to check the versatility and efficiency of the biocatalysts. Reduction of double bond occurred with high rates and enantioselectivity, except for 3,5,5-trimethyl-2-cyclohexenone. DFT (density functional theory) computational studies were performed to investigate whether the steric hindrance and/or the electronic properties of the substrates were crucial for reactivity. The three-dimensional structure of enoate reductases from *Kluyveromyces lodderae* and *Candida castellii*, predicted through comparative modeling, resulted similar to that of *S. cerevisiae* OYE2 and revealed the key role of Trp116 both in substrate specificity and stereocontrol. All the modeling studies indicate that steric hindrance was a major determinant in the enzyme reactivity.

Conclusions: The OYE biocatalysts, based on recombinant *S. cerevisiae* expressing OYE genes from non-conventional yeasts, were able to differently reduce the activated double bond of enones, enals and nitro-olefins, exhibiting a wide range of substrate specificity. Moreover whole-cells biocatalysts bypassed the necessity of the cofactor recycling and, tuning reaction parameters, allowed the synthetic exploitation of endogenous carbonyl reductases. Molecular modeling studies highlighted key structural features for further improvement of catalytic properties of OYE enzymes.

* Correspondence: stefano.raimondi@unimore.it

¹Dipartimento di Scienze della Vita, Università degli Studi di Modena e Reggio Emilia, Via G. Campi 183, 41125 Modena, Italy

Full list of author information is available at the end of the article

Background

Fine chemical industry is increasingly taking advantage of biocatalysts to produce high-value products because enzymes and whole-cell biocatalysts can be sometimes more convenient than synthetic chemistry [1]. Biotransformations occur under mild conditions of temperature, pressure and pH, and may display their activity with high chemo-, regio- and stereoselectivity [1,2]. Thus, biocatalysis has developed from a niche technology to a widely used manufacturing method, and the availability of new biocatalytic processes for the preparation of optically pure molecules is very attractive for organic chemists.

Due to the increasing demand of optically pure building blocks, considerable efforts are addressed at identifying, characterizing, improving, and developing Old Yellow Enzymes (OYE) biocatalysts, i.e. flavin-dependent enoate reductases (EC 1.6.99.1), that catalyze the chemo- and stereoselective hydrogenation of electron-poor alkenes [3,4]. Their biotechnological potential lies on the ability to generate up to two stereocenters by the stereoselective *anti*-hydrogenation of the activated C = C double bond, which is highly demanded in asymmetric synthesis, since conventional chemical methods mostly give *syn*-hydrogenation [5,6]. OYEs have been found in plants, bacteria, and fungi [7-10], OYE1 from *Saccharomyces pastorianus* being the first to be widely studied for structure, substrate selectivity and stereoselectivity [11-14]. In order to improve the tools for asymmetric reductions of activated olefins, the biodiversity of non-conventional yeasts was explored as well, and yielded several novel OYE genes [15,16]. OYE genes from unconventional yeasts were cloned and expressed in *Saccharomyces cerevisiae* which was used as a whole-cell biocatalyst, demonstrating the important role of unconventional yeasts as sources of novel biocatalysts and their potential for the expansion of known genetic biodiversity [16].

In the present study, recombinant strains of *S. cerevisiae* lacking endogenous OYE and expressing eight OYE genes from as many unconventional yeasts were used as whole-cell biocatalysts to reduce different substrates with an activated electron-poor double bond, in order to compare their catalytic and selectivity properties. It was investigated whether the electronic properties and the steric hindrance affected reactivity and/or enzyme stereoselectivity in the asymmetric reduction. The structure of three selected enzymes was predicted through comparative modeling and the active sites compared. The reactivity of the tested substrates, has been also studied by means of electronic computations, such as the lowest unoccupied molecular orbitals (LUMO) distribution and OYE-substrate interactions were analyzed to reveal the key features of the structure and the sequence patterns of these new enoate reductases.

Results and discussion


Asymmetric reductions with recombinant strains of *S. cerevisiae* expressing OYE genes.

The strain *S. cerevisiae* BY4741 Δ Oye2, that lacks endogenous *Oye2* gene and presents negligible overall OYE activity [16], was used as the host to express the OYE genes from *S. cerevisiae* BY4741 (OYE2), *S. cerevisiae* L12, and from the unconventional yeasts *Candida castellii* DBVPG3704, *Candida sake* DBVPG6162, *Kazachstania exigua* L10, *Kazachstania lodderae* DBVPG6308, *Kazachstania spencerorum* DBVPG6748, *Nakaseomyces bacillisporus* DBVPG6945, and *Naumovia castellii* DBVPG6298 [16]. Recombinant strains bearing heterologous OYE genes were used as whole-cells biocatalysts to reduce three molecules with electron-poor double bond activated by different electron-withdrawing groups (Figure 1), i.e. ketoisophorone (2,6,6-trimethylcyclohex-2-ene-1,4-dione, **1a**), α -methyl-*trans*-cinnamaldehyde ((*E*)-2-methyl-3-phenyl-2-propenal, **2a**), and *trans*- β -methyl- β -nitrostyrene ((*E*)-1-phenyl-2-nitropropene, **3a**).

The activity towards **1a** of the diverse recombinant *S. cerevisiae* was evaluated using resting cells. The conversion was at least 86% after 5 h and, for most of the strains, was nearly complete after 24 h (Table 1). However, even though high efficiency of **1a** transformation was observed, and the reaction proceeded with preference for the *R*-levodione (**1b**), the enantioselectivity generally decreased compared to that obtained with the corresponding wild type yeasts [16].

The biotransformation of **2a** by whole-cells of wild-type Baker's yeast has been reported to result in a mixture of saturated 2-methyl-3-phenylpropanol (**2c**) and unsaturated primary alcohol 2-methyl-3-phenyl-2-propenal (**2d**, Figure 1), with different ratios depending on the strain and the experimental conditions [17,18]. Consistently, the transformation of **2a** using the wild type *S. cerevisiae* BY4741 gave a mixture of **2c** (78%) and **2d** (22%). On the contrary, *S. cerevisiae* BY4741 Δ Oye2, that lacks endogenous *Oye2* gene, yielded **2d** as the sole detectable product, indicating the pivotal role of OYE2 protein in the enoate reductase activity of the parental strain [19]. Reduction of **2a** with most of the recombinant strains gave **2c** as the only observed product by means of the subsequent C = C and carbonyl reductions (Table 1). The enhanced OYE activity of the recombinant strains prevented the formation of **2d** observed using *S. cerevisiae* BY4741, with the exception of the strain expressing the *C. castellii* OYE gene that yielded a mixture of **2c** and **2d**. Interestingly, enantioselectivity ranged between 59 and > 98%, with four strains yielding (*S*)-**2c** as the sole detectable enantiomer.

The majority of recombinant strains exhibited a high rate in reducing **3a** to 1-phenyl-2-nitropropane (**3b**), but low e.e. (29-58%) was generally observed (Table 1).



Similar behavior was observed with different microbial whole-cells, where the scarce enantioselectivity was mostly due to the easy racemization of the saturated nitro-derivative [20].

The data herein reported proved that the OYE biocatalysts, based on recombinant *S. cerevisiae* BY4741 Δ Oye2 expressing OYE genes from non-conventional yeasts, were able to efficiently convert not only the enone **1a**, a well-established OYE substrate, but present a broader substrate specificities being able to reduce also the activated C = C double bond of enals (**2a**) and nitro-olefins (**3a**), showing that these systems seem to have a great potential for synthetic chemists.

Bioconversion of a series of alkyl-substituted cyclohex-2-enones

Recombinant strains were tested to reduce four alkyl-substituted cyclohex-2-enones (**4a**, **5a**, **6a**, and **7a**; Figure 1). The substrates were selected in order to investigate the versatility and efficiency of these biocatalysts, defining how the electronic properties and steric hindrance affect substrate reactivity and enzyme stereoselectivity in the asymmetric reduction of double bond.

All the recombinant yeasts totally reduced 2-cyclohexenone (**4a**) into cyclohexanone (**4b**) within 2 h (Table 2), yielding no traces of cyclohex-2-enol or cyclohexanol (**4c**), that could result from carbonyl reductases (CR) activity on the keto group of **4a** or **4b**, respectively.

The strains also reduced 4,4-dimethyl-2-cyclohexenone (**5a**) into 4,4-dimethyl-cyclohexanone (**5b**) at a good extent, with yields ranging between 76 and 97%. The strains carrying the genes from *Candida castellii* DBVPG3704 and *Kazachstania lodderae* DBVPG6308 showed the lowest conversions and produced also 4,4-dimethyl-2-cyclohexenol (**5d**; 20% and 15%, respectively) through carbonyl reduction of **5a**, indicating that CR became competitive when C = C double bond reduction was less effective.

The yeast bearing the OYE gene from *C. castellii* DBVPG3704 was the only ineffective in the reduction of 3-methyl-2-cyclohexenone (**6a**). All the other strains reduced this substrate with high stereoselectivity (>98%), with homogeneous stereobias and formation of (*S*)-**6b** as the sole detectable enantiomer. Small amounts of 3-methylcyclohexanol (**6c**, 3-6%) were observed after 48 h. Conversion yield of **6a** towards **6b** ranged from 23 to 75% after 72 h and was lower compared to the conversion of substrates **4a** and **5a**. Consistently, previous studies proved that **6a** is a 'difficult' substrate for enoate reductases, due to the presence of a substituent on the C3 position [21,22]. All the recombinant strains failed to transform α -isophorone (3,5,5-trimethyl-2-cyclohexenone, **7a**), neither the C = C or C = O double bonds being reduced after 72 h.

Table 1 Bioreductions of ketoisophorone (1a), α -methyl-*trans*-cinnamaldehyde (2a), and *trans*- β -methyl- β -nitrostyrene (3a), with recombinant *S. cerevisiae* BY4741 Δ Oye2 bearing heterologous OYE genes

OYE gene source	1a \rightarrow 1b*			2a \rightarrow 2c			3a \rightarrow 3b		
	Time	1b	(R)-1b	Time	2c	(S)-2c	Time	3b	(R)-3b
	h	%	e.e.%	h	%	e.e.%	h	%	e.e.%
<i>C. castellii</i> DBVPG3704	5	≥ 99	70	24	70 [†]	≥ 98	2	45	50
<i>C. sake</i> DBVPG6162	5	97	71	3	100	59	0.5	93	48
<i>K. exigua</i> L10	5	94	82	24	100	≥ 98	0.5	97	30
<i>K. lodderae</i> DBVPG6308	5	≥ 99	78	16	100	80	0.5	97	58
<i>K. spencerorum</i> DBVPG6748	5	96	65	3	100	≥ 98	0.5	94	50
<i>Nak. bacillisporus</i> DBVPG6945	5	98	68	2	100	95	0.5	92	40
<i>Nau. castellii</i> DBVPG6298	5	93	78	2	100	85	5	92	29
<i>S. cerevisiae</i> BY4741 (Oye2)	5	89	80	2	100	≥ 98	1	96	50
<i>S. cerevisiae</i> L12	5	86	79	2	100	72	2	95	44

Data are reported as molar conversion (%). *Results from Raimondi et al. 2010. [†]30% of compound 2d was also observed. The experiment were performed in triplicate, the yield standard deviations (SD) were always less than 4%, and the ee% SD were always less than 3%.

As a whole, most of the nine recombinant strains exhibited similar activity toward the tested cyclohexenone derivatives and a common stereobias was observed toward **6a**. Moreover, substrates **4a**, **5a**, **6a** and **7a** which are increasingly sterically hindered, were progressively less reactive in C = C reduction. The electronic features of the tested substrates and the structural properties of the investigated OYEs will be thoroughly described in a following section.

Computational analysis: evaluation of substrate electronic features, and OYE three-dimensional structure prediction through comparative modeling

The enoate reductases preferentially reduce substrates containing electron-poor double bonds [3]. In order to evaluate the electronic features of the tested substrates, a geometry optimization in the gas phase with an approach

based on density functional theory (DFT/B3LYP/6-31G*) was performed, resulting in prediction of the double bond lengths and of the electrostatic potential (ESP) derived charges for the C atom involved in the hydride attack of the substrates (Table 3). A further characterization has been carried out by the analysis of the lowest unoccupied molecular orbitals (LUMO) for the investigated substrates. The surfaces were rendered in order to encapsulate approximately 80% of the integral of the evaluated grid points (Figure 2). Based on the length of the double bonds, **4a**, **5a**, **6a**, and **7a** resulted very similar. The values 1.34 Å for substrates **4a** and **5a** and 1.35 Å for substrates **6a** and **7a**, typical of the sp² hybridization state, were not sufficiently different to explain a diverse reactivity based on electronic configurations. The comparison among the LUMO frontier orbitals revealed very similar configuration and

Table 2 Bioreductions of alkyl-substituted cyclohex-2-enones (substrates 4a, 5a, 6a, and 7a[#]) with recombinant *S. cerevisiae* BY4741 Δ Oye2 bearing heterologous OYE genes

OYE gene source	4a \rightarrow 4b		5a \rightarrow 5b		6a \rightarrow 6b [‡]		
	Time	Yield	Time	Yield	Time	Yield	(S)-6b
	h	%	h	%	h	%	e.e.%
<i>C. castellii</i> DBVPG3704	2	≥ 99	2	76 [†]	72	< 5	n.d.
<i>C. sake</i> DBVPG6162	1	≥ 99	1	95	48	75	≥ 98
<i>K. exigua</i> L10	2	≥ 99	2	92	72	23	≥ 98
<i>K. lodderae</i> DBVPG6308	2	≥ 99	2	84 [†]	72	25	≥ 98
<i>K. spencerorum</i> DBVPG6748	2	≥ 99	1	95	48	60	≥ 98
<i>Nak. bacillisporus</i> DBVPG6945	1	≥ 99	2	94	72	24	≥ 98
<i>Nau. castellii</i> DBVPG6298	2	≥ 99	2	95	48	53	≥ 98
<i>S. cerevisiae</i> BY4741 (Oye2)	1	≥ 99	2	96	48	45	≥ 98
<i>S. cerevisiae</i> L12	2	≥ 99	1	97	48	40	≥ 98

Data are reported as molar conversion (%). [#]No conversion of compound **7a** was observed in the first 72 h. [†]15-20% of compound **5d** was also observed; [‡]product **6c** was always less than 15%. The experiment were performed in triplicate, the yield standard deviations (SD) were always less than 4%, and the ee% SD were always less than 3%.

Table 3 Electronic features of the alkyl-substituted cyclohex-2-enones predicted by DFT calculations

Substrate	Double bond length	ESP charge
4a	1.34 Å	-0.003
5a	1.34 Å	-0.174
6a	1.35 Å	0.350
7a	1.35 Å	0.404

reactivity around the sp^2 carbons of all the substrates. Conversely, computed ESP charges suggested that substrates **5a**, **6a**, and **7a** were the most favored for the reduction of the double bond, outcome partially in contrast with bioconversion data indicating **6a** and **7a** as “difficult” substrates.

The data herein presented demonstrated that the different reactivity of these substrates cannot be ascribed to their electronic configurations. As a whole, the substrates which are increasingly sterically hindered, were progressively less reactive in double bond reduction and their electronic properties do not affect this behavior.

In order to evaluate the role of steric hindrance with respect to the substrate selectivity, the three-dimensional structures of the *S. cerevisiae* OYE2 and of the *K. lodderae* DBVPG6308 and *C. castellii* DBVPG3704 enoate reductases were predicted through comparative modeling, using the structure of the *S. pastorianus* OYE1 (RCSB PDB: 1OYA) as template [11]. These three enzymes were chosen because, when overexpressed in *S. cerevisiae*

BY4741ΔOye2, the corresponding recombinant strains exhibited different activity and/or enantioselectivity towards many of the tested substrates. The proteins and the template OYE1 share a very high level of identity, ranging from 73 to 91% (Table 4, Figure 3). The active sites are highly conserved and few differences in amino acid composition are present (Table 5). Furthermore, similar distances between the three core amino acids (His191, Asn194, and Tyr196) belonging to the catalytic site have been calculated (Table 6). The high identity between both the primary structures of the investigated OYEs and the low root mean square deviation (RMSD) values for the modeled structures suggested that the different transformation rates of the tested substrates can hardly be ascribed only to the structural diversity elucidated by our computational work.

The substrate α -isophorone (**7a**) was not reduced by OYEs, although electronic features suggested a good reactivity, likely because of a dramatic steric effect. A docking study of this molecule have been performed on the predicted structure of OYE2 protein. The repulsive van der Waals interactions between the alkylic groups of substrate **7a** and Trp116 of OYE2 confirmed that steric hindrance was present (Figure 4). Consistently, in a previous study, the interactions between substrates and Trp116 in the active site of *S. pastorianus* OYE1 influenced the catalytic activity of the enzyme and the stereochemistry of the products [23].

The analysis of the unfavorable contacts between the side chain of Trp116 and the substrates confirmed that the transformation efficiency on a series of alkyl-substituted cyclohex-2-enones was associated to the length of the alkyl-substituent. The larger was the substrate hindrance, the lower was the reactivity. Furthermore, the pivotal role of Trp116 on catalytic properties of OYE enzymes has been highlighted, suggesting this key structural features as an hot spot for modification and further improvement of the biocatalysts.

Whole-cell OYE biocatalysts: synthetic exploitation tuning reaction time

Biotransformation of $\alpha\beta$ -unsaturated carbonyls occurred with carbonyl reduction by endogenous CR as side-reaction. Thus, the biotransformations of substrates **4a**,




Table 4 OYE sequences alignment

	1OYA	OYE2	OYE from <i>K. lodderae</i>	OYE from <i>C. castellii</i>
1OYA	-	91.0	77.2	74.8
OYE2	91.0	-	77.7	73.2
OYE from <i>K. lodderae</i>	77.2	77.7	-	77.4
OYE from <i>C. castellii</i>	74.8	73.2	77.4	-

Sequence identity (%) among the template 1OYA and the three OYE proteins selected for the structure prediction through comparative modelling.




Figure 3 Sequence alignment of OYE proteins. Sequence alignment among the template 10YA from *S. pastorianus* and the three OYE proteins selected for the structure prediction through comparative modelling.

5a, 6a, and 8a were analyzed until prolonged reaction times, in order to obtain also the reduction of the carbonyl group. In all cases we observed a prevailing OYE activity in the first hours of reaction, while at longer times the carbonyl reductase activity led to the formation of the saturated alcohols (Figure 5). More in detail, using 4a as substrate, it was possible to quantitatively recover 4b after 2 hours of reaction. Later (144 h), the endogenous carbonyl reductase(s) led to the formation of 4c, with high yield (97%). The same behavior was observed with substrate 8a: at short reaction times (2 h) the OYE product (*R*)-8b was almost (99%) the only chemical species present, whereas a 1/10 ratio between 8b and 8c was observed after 168 h; the stereochemical analysis of 8c showed that the ratio (1*S*,2*R*)-8c/(1*R*,2*R*)-8c was 70/30. The bioreduction of 6a, that is a less reactive substrate, proceeded with the concomitant formation of 6b and 6c; 6b was obtained with high enantioselectivity (e.e. > 98%). The saturated alcohol (6c) was the minor product during the first hours of reaction and became the sole product after 120 h; 6c was produced with a *syn/anti* composition of 75/25. In the case of 5a, bioreductions proceeded faster: the saturated ketone 5b was produced at 95% molar conversion after 1 h while 5c reached 95% after 48 h of biotransformation.

Overall, our whole-cells system, compared to the use of purified enzymes, showed a peculiar synthetic advantage: with a unique biocatalyst, tuning the reaction time, it is possible to have access to different molecules, with high yields, starting from a single substrate.

Conclusions

S. cerevisiae is well suited for biotechnological applications owing to the wealth of the genetic techniques available, the ease of manipulation and the rapid growth rates. The expression in the host *S. cerevisiae* BY4741Δ*Oye2* of the OYE genes make possible the exploitation of these technological feature leading to the development of efficient biocatalysts base on yeast cells. The recombinant *S. cerevisiae*, showed a wide range of substrate specificity, being able to reduce the activated C = C double bond of enones, enals and nitro-olefins. Moreover, whole-cells biocatalysts bypassed the necessity of the cofactor recycling and, tuning reaction parameters, allowed the synthetic exploitation of the endogenous carbonyl reductases. Versatility and efficiency of these biocatalysts was analyzed revealing that the different reactivity was not ascribed to substrate electronic properties, but steric hindrance resulted the most relevant factor affecting the asymmetric reduction of double bond. Molecular modeling studies on OYE proteins highlighted the pivotal role of Trp116 both in activity and stereochemistry and it was established as a key structural features for further improvement of catalytic properties of OYE enzymes.

Methods

Strains and culture conditions

Recombinant *S. cerevisiae* BY4741Δ*Oye2* strains (bearing OYE genes from *Candida castellii* DBVPG3704, *Candida sake* DBVPG6162, *Naumovia castellii* DBVPG6298,

Table 5 Amino acid composition of OYE active sites

Position	OYE2	OYE from <i>K. lodderae</i>	OYE from <i>C. castellii</i>
82	TYR	TYR	TYR
116	TRP	TRP	TRP
117	VAL	VAL	VAL
118	LEU	LEU	LEU
119	GLY	GLY	GLY
120	TRP	ARG	ARG
121	ALA	GLN	GLN
122	ALA	GLY	GLY
191	HIS	HIS	HIS
192	SER	SER	SER
193	ALA	ALA	ALA
194	ASN	ASN	ASN
195	GLY	SER	SER
196	TYR	TYR	TYR
197	LEU	LEU	LEU
198	LEU	LEU	LEU
199	ASN	ASN	ASN
200	GLN	GLN	GLN
201	PHE	PHE	PHE
250	PHE	TYR	TYR
251	ASN	GLY	GLY
253	MET	MET	MET
295	PRO	SER	PRO
296	PHE	PHE	PHE
375	TYR	TYR	TYR

Differences are highlighted in bold.

Kazachstania exigua L10, *Kazachstania lodderae* DBVP G6308, *Kazachstania spencerorum* DBVPG6748, *Naka-seomyces bacillisporus* DBVPG6945, *Saccharomyces cerevisiae* BY4741 (Oye2), and *Saccharomyces cerevisiae* L12) were prepared as described earlier [16]. Strains were routinely maintained on YNB agar medium (Difco™ Yeast Nitrogen Base without aminoacids 6.7 g/L, glucose 20 g/L, histidine 0.15 g/L, leucine 0.73 g/L, methionine 0.15 g/L) and grown in YNB liquid medium, in baffled Erlenmeyer flasks on orbital shaker at 30°C and 150 rpm.

Table 6 OYEs catalytic triad

Protein	His-Asn	His-Tyr	Tyr-Asn
OYE2	6.5	6.8	5.1
<i>K. lodderae</i>	7.5	6.0	5.3
<i>C. castellii</i>	7.3	5.6	5.5

Distances between the core amino acids belonging to the catalytic site (His191, Asn194, and Tyr196) in the three modelled OYEs (Å).

Biotransformations

Recombinant cells grown for 24 h were centrifuged (5000 rpm for 10 min at 4°C), washed and resuspended at a concentration of 15 AU of OD₆₀₀ in 100 mM phosphate buffer at pH 7, containing 50 g/L glucose. Substrate stock solutions (200 g/L) prepared in ethanol were added to the reaction obtaining 1 g/L of substrate concentration. Biotransformations were carried out in baffled Erlenmeyer flasks on orbital shaker, incubated at 30°C and 150 rpm. Samples for analysis were extracted 1:1 with EtOAc containing 0.05% (v/v) of 1-phenylethanol as internal GC standard and dried over Na₂SO₄. At the end of the biotransformation, the whole reaction mixture was centrifuged and the aqueous supernatant extracted three times with EtOAc. The combined organic phases were dried over Na₂SO₄ and the residue purified by flash chromatography.

Analytical methods

Gas chromatographic analysis were performed on a Dani 86.10 HT instrument (Dani Instruments, Italy; carrier H₂, 0.6 bar, FID detector), using a MEGA-DEX DET-Beta (column A; Mega Snc, Legnano, Italy) chiral column or a SE30 polydimethylsiloxane non-chiral column (column B), under the following conditions. Bioreduction of **1a**: column A; 45°C for 2 min, 5° min⁻¹ rate to 130°C, 30° min⁻¹ to 200°C; retention times: **1a** 18.0 min, (*R*)-**1b** 19.2 min, (*S*)-**1b** 19.5 min. Bioreduction of **2a**: column A; 95°C, 1° min⁻¹ rate to 120°C, 30° min⁻¹ to 200°C; retention times: **2a** 31.0 min, (*R*)-**2b** 16.4 min, (*S*)-**2b** 16.7 min, (*R*)-**2c** 28.7 min, (*S*)-**2c** 29.2 min, **2d** 19.4 min. Bioreduction of **3a**: column B; 105°C for 5 min, 1° min⁻¹ rate to 115°C, 30° min⁻¹ to 200°C; retention times **3a** 20.4 min, (*R*)-**3b** 18.8 min, (*S*)-**3b** 19.0 min. Bioreduction of **4a**: column B; 55°C isotherm analysis; retention times **4a** 5.1 min, **4b** 4.1 min. Bioreduction of **5a**: column B; 60°C for 3 min, 1°C min⁻¹ rate to 140°C, 30° min⁻¹ to 200°C; retention times **5a** 14.6 min, **5b** 14.0 min, **5c** 16.9 min, **5d** 16.2 min. Bioreduction of **6a**: column A; 50°C, 1°C min⁻¹ rate to 150°, 30° min⁻¹ to 200°C; retention times **6a** 20.9 min, (*R*)-**6b** 13.0 min, (*S*)-**6b** 13.7 min. Column B; 40°C isotherm analysis; retention times **6a** 25.3 min, **6b** 11.4 min, **6c** 12.3 min. Bioreduction of **7a**: column A; 60°C for 5 min, 3°C min⁻¹ to 100°C, 30° min⁻¹ to 200°C; retention times **7a** 19.5 min. Bioreduction of **8a**: column A; 50°C for 5 min, 5°C min⁻¹ rate to 150°, 30° min⁻¹ to 200°C; retention times **8a** 19.8 min, (*R*)-**8b** 12.3, (*S*)-**8b** 13.0. Column B; 50°C for 5 min, 5°C min⁻¹ rate to 120°, 30° min⁻¹ to 200°C; retention times **8a** 8.70 min, **8b** 7.02 min, **8c** 7.35 min. Biotransformation products were identified by comparison with authentic reference materials. The absolute configuration and e.e. of **6b** and **8b** were determined by co-injection in chiral GC with reference materials of known absolute configuration [21 and references therein].




Figure 4 Repulsive van der waals interactions between the alkyl groups of α -isophorone (7a) and OYE2 Trp116.

Stereoisomeric composition of **6c** and **8c** were determined by ^1H NMR [24] after purification of the products.

Computational details

All the comparative modeling procedures were carried out with the Homology module of the Molecular Operating Environment 2011.10 (MOE; Chemical Computing Group

Inc., Montreal, QC, Canada). The alignment produced by the Align program of MOE with default parameters was manually adjusted. Comparative model building was carried out with the MOE Homology Model program. IOYA (*S. pastorianus* OYE1) was set as template for all the modeled sequences. For each sequence, ten independent models were built and refined, and then the highest




Figure 5 Bioreductions of alkyl-substituted cyclohex-2-enones (substrates **4a**, **5a**, **6a**, and **8a**) until prolonged reaction times. Data are reported as molar conversion (%). Blue line and circles: substrates. Red line and squares: OYE products; Green line and diamonds: OYE and CR products. The curves were representative of three independent experiments and the standard deviations were always less than 4%.

scoring intermediate model was submitted to a further round of energy minimization (EM). For both the intermediate and the final structures the refinement procedures consisted in EM runs based on the AMBER99 force field, with the reaction field solvation model.

Geometry optimization and electronic computations of the substrates were carried out in the gas phase through the MOE SCF Calculation interface to the GAMESS software [25,26]. The DFT calculations were based on the B3LYP hybrid functional with the 6-31G* basis set.

Abbreviations

OYE: Old Yellow Enzyme (enoate reductase); CR: Carbonyl reductase; **1a**: 2,6,6-trimethylcyclohex-2-ene-1,4-dione [ketoisophorone]; **1b**: 2,2,6-trimethylcyclohexane-1,4-dione [(6R)-levodione]; **1c**: 4-hydroxy-2,2,6-trimethylcyclohexanone [(4R, 6R)-actinol]; **2a**: (E)-2-methyl-3-phenyl-2-propenal [α -methyl-*trans*-cinnamaldehyde]; **2b**: 2-methyl-3-phenylpropanal; **2c**: 2-methyl-3-phenylpropanol; **2d**: 2-methyl-3-phenyl-2-propenol; **3a**: (E)-1-phenyl-2-nitropropene [*trans*- β -methyl- β -nitrostyrene]; **3b**: 1-phenyl-2-nitropropane; **4a**: 2-cyclohexenone; **4b**: Cyclohexanone; **4c**: Cyclohexanol; **5a**: 4,4-dimethyl-2-cyclohexenone; **5b**: 4,4-dimethylcyclohexanone; **5c**: 4,4-dimethylcyclohexanol; **5d**: 4,4-dimethyl-2-cyclohexenol; **6a**: 3-methyl-2-cyclohexenone; **6b**: 3-methylcyclohexanone; **6c**: 3-methylcyclohexanol; **7a**: 3,5,5-trimethyl-2-cyclohexenone [α -isophorone]; **8a**: 2-methyl-2-cyclohexenone; **8b**: 2-methylcyclohexanone; **8c**: 2-methylcyclohexanol; DFT: Density functional theory; ESP: Electrostatic potential; LUMO: Lowest Unoccupied Molecular Orbitals.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

DR and SR conceived the study, performed the experimental work and drafted the manuscript. MC performed bioreductions and chemical analysis. ER and CS contributed to the computational analysis. AA, FM, IE and MR accomplished data interpretation, participated in the design of the study, and writing of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors are most grateful to Dr. Luca Bertini from the Università degli Studi di Milano Bicocca for fruitful discussion on DFT application.

Author details

¹Dipartimento di Scienze della Vita, Università degli Studi di Modena e Reggio Emilia, Via G. Campi 183, 41125 Modena, Italy. ²Department of Food, Environmental and Nutritional Sciences (DeFENS), University of Milan, Via G. Celoria 2, 20133 Milan, Italy. ³Dipartimento di Scienze Farmacologiche e Biomolecolari, Università degli Studi di Milano, Via Trentacoste 2, 20134 Milan, Italy.

Received: 8 January 2014 Accepted: 15 April 2014

Published: 25 April 2014

References

1. Panke S, Held M, Wubbolts M: Trends and innovations in industrial biocatalysis for the production of fine chemicals. *Curr Opin Biotechnol* 2004, **15**:272–279.
2. Faber K: *Bioreductions in Organic Chemistry: A Textbook*. 6th edition. Berlin, Heidelberg: Springer; 2011.
3. Stuermer R, Hauer B, Hall M, Faber K: Asymmetric bioreduction of activated C = C bonds using enoate reductases from the old yellow enzyme family. *Curr Opin Chem Biol* 2007, **11**:203–213.
4. Williams RE, Bruce NC: "New uses for an Old Enzyme" – the Old Yellow Enzyme family of flavoenzymes. *Microbiology* 2002, **148**:1607–1614.
5. Blaser HU, Malan C, Pugin B, Spindler F, Steiner H, Studer M: Selective hydrogenation for fine chemicals: recent trends and new developments. *Adv Synth Catal* 2003, **345**:103–151.
6. De Wildeman SMA, Sonke T, Schoemaker HE, May O: Biocatalytic reductions: from lab curiosity to "first choice". *Acc Chem Res* 2007, **40**:1260–1266.

7. Strassner J, Fürholz A, Macheroux P, Amrhein N, Schaller A: A homolog of old yellow enzyme in tomato. Spectral properties and substrate specificity of the recombinant protein. *J Biol Chem* 1999, **274**:35067–35073.
8. Brigé A, Van Den Hemel D, Carpentier W, De Smet L, Van Beeumen JJ: Comparative characterization and expression analysis of the four Old Yellow Enzyme homologues from *Shewanella oneidensis* indicate differences in physiological function. *Biochem J* 2006, **394**:335–344.
9. Kataoka M, Kotaka A, Thiwthong R, Wada M, Nakamori S, Shimizu S: Cloning and overexpression of the old yellow enzyme gene of *Candida macedoniensis*, and its application to the production of a chiral compound. *J Biotechnol* 2004, **114**:1–9.
10. Arnone A, Cardillo R, Nasini G, Depava OV: Enantioselective reduction of racemic absicic acid by *Aspergillus niger* cultures. *J Chem Soc Perkin Trans 1* 1990, **11**:3061–3063.
11. Fox KM, Karplus PA: Old yellow enzyme at 2 Å resolution: overall structure, ligand binding, and comparison with related flavoproteins. *Structure* 1994, **2**:1089–1105.
12. Meah Y, Massey V: Old yellow enzyme: stepwise reduction of nitro-olefins and catalysis of aci-nitro tautomerization. *Proc Natl Acad Sci* 2000, **97**:10733–10738.
13. Swiderska MA, Stewart JD: Stereoselective enone reductions by *Saccharomyces carlsbergensis* old yellow enzyme. *J Mol Catal B: Enzym* 2006, **42**:52–54.
14. Vaz AD, Chakraborty S, Massey V: Old Yellow enzyme: aromatization of cyclic enones and the mechanism of a novel dismutation reaction. *Biochemistry* 1995, **34**:4246–4256.
15. Raimondi S, Roncaglia L, Amaretti A, Leonardi A, Buzzini P, Forti L, Rossi M: Rapid method for screening enoate reductase activity in yeasts. *J Microbiol Methods* 2010, **83**:106–110.
16. Raimondi S, Romano D, Amaretti A, Molinari F, Rossi M: Enoate reductases from non conventional yeasts: bioconversion, cloning, and functional expression in *Saccharomyces cerevisiae*. *J Biotechnol* 2010, **156**:279–285.
17. Fardelone LC, Augusto J, Rodrigues R, Moran PJ: Baker's yeast mediated asymmetric reduction of cinnamaldehyde derivatives. *J Mol Catal B: Enzym* 2004, **29**:41–45.
18. Fuganti C, Chiringhelli D, Grasselli P: Stereochemical course of the reduction of cinnamaldehyde and cinnamyl alcohol to 3-phenylpropanol by fermenting Baker's yeast. *J Chem Soc Chem Commun* 1975, **20**:846–847.
19. Contente M, Granato T, Remelli W, Zambelli P, Raimondi S, Rossi M, Romano D: Complementary microbial approaches for the preparation of optically pure aromatic molecules. *Ann Microbiol* 2013, **63**(3):1021–1027.
20. Mori A, Ishiyama I, Akita H, Suzuki K, Mitsuoka T, Oishi T: Reduction of nitroolefin using microorganisms. *Chem Pharm Bull* 1990, **38**:3449–3451.
21. Hall M, Stueckler C, Ehammer H, Pointner E, Oberdorfer G: Asymmetric bioreduction of C = C bonds using enoate reductases OPR1, OPR3 and YqjM: enzyme-based stereocontrol. *Adv Synth Catal* 2008, **350**:411–418.
22. Hall M, Stueckler C, Hauer B, Stuermer R, Friedrich T, Breuer M, Kroutil W, Faber K: Asymmetric bioreduction of activated c = c bonds Using *Zymomonas mobilis* NCR Enoate reductase and Old Yellow Enzymes OYE 1–3 from yeasts. *Eur J Org Chem* 2008, **9**:1511–1516.
23. Padhi SK, Bougioukou DJ, Stewart JD: Site-saturation mutagenesis of tryptophan 116 of *Saccharomyces pastorianus* old yellow enzyme uncovers stereocomplementary variants. *J Am Chem Soc* 2009, **131**:3271–3280.
24. Solladié-Cavallo A, Barama A, Choucaira E, Norouzi-Arabis H, Schmit M, Garin F: Heterogeneous hydrogenation of substituted phenols over Al2O3 supported ruthenium. *J Mol Catal A Chem* 2007, **273**:92–98.
25. Schmidt MW, Baldrige KK, Boatz JA, Elbert ST, Gordon MS, Jensen JH, Koseki S, Matsunaga N, Nguyen KA, Su S, Windus TL, Dupuis M, Montgomery JA: General atomic and molecular electronic structure system. *J Comput Chem* 1993, **14**:1347–1363.
26. Gordon MS, Schmidt MW: Chapter 41 - Advances in electronic structure theory: GAMESS a decade later. In *Theory and Applications of Computational Chemistry*. Edited by Dykstra CE, Frenking G, Kim KS, Scuseria GE. Amsterdam: Elsevier; 2005:1167–1189.

doi:10.1186/1475-2859-13-60

Cite this article as: Romano et al.: Recombinant *S. cerevisiae* expressing Old Yellow Enzymes from non-conventional yeasts: an easy system for selective reduction of activated alkenes. *Microbial Cell Factories* 2014 **13**:60.