

# Complete Genome Sequence of “*Candidatus Liberibacter africanus*,” a Bacterium Associated with Citrus Huanglongbing

Hong Lin,<sup>a</sup> Gerhard Pietersen,<sup>b</sup> Cliff Han,<sup>c</sup> David Alan Read,<sup>b</sup> Binghai Lou,<sup>d</sup> Goutam Gupta,<sup>c</sup> Edwin L. Civerolo<sup>a</sup>

USDA-ARS San Joaquin Valley Agricultural Sciences Center, Parlier, California, USA<sup>a</sup>; University of Pretoria, Pretoria, South Africa<sup>b</sup>; Los Alamos National Laboratory, Los Alamos, New Mexico, USA<sup>c</sup>; Guangxi Citrus Research Institute, Guilin, China<sup>d</sup>

**We report here the complete genome sequence of “*Candidatus Liberibacter africanus*” strain PTSAPSY. The 1,192,232-bp genome with 34.5% G+C content comprises 1,017 open reading frames, 44 tRNAs, and three complete rRNAs in a circular chromosome.**

Received 27 May 2015 Accepted 15 June 2015 Published 16 July 2015

**Citation** Lin H, Pietersen G, Han C, Read DA, Lou B, Gupta G, Civerolo EL. 2015. Complete genome sequence of “*Candidatus Liberibacter africanus*,” a bacterium associated with citrus huanglongbing. *Genome Announc* 3(4):e00733-15. doi:10.1128/genomeA.00733-15.

**Copyright** © 2015 Lin et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Hong Lin, hong.lin@ars.usda.gov.

Huanglongbing (HLB), also known as greening disease, is one of the most devastating diseases of citrus in the world (1). The disease is associated with three phloem-restricted insect-transmitted, Gram-negative, and fastidious species of alphaproteobacteria in the genus “*Candidatus Liberibacter*.” The name of each species was based on its presumptive origin; “*Candidatus Liberibacter asiaticus*” is believed to have originated in Asia, “*Ca. Liberibacter americanus*” in the Americas, and “*Ca. Liberibacter africanus*” in Africa (1–4). Among these three species, “*Ca. Liberibacter asiaticus*” is the most widely distributed and transmitted by the Asian citrus psyllid *Diaphorina citris* worldwide, while “*Ca. Liberibacter africanus*” is found only in Africa and the Mascarene Islands and is naturally transmitted by the psyllid *Trioza erythrae*. In contrast to “*Ca. Liberibacter asiaticus*” and “*Ca. Liberibacter americanus*,” “*Ca. Liberibacter africanus*” is heat sensitive and often suppressed at elevated temperatures of >30°C (1). In South Africa, a disease was observed as a yellow shoot or greening resembling mineral toxicity in 1928 and was confirmed to be a biotic disease in 1965 (1). While still a serious disease prevalent in many parts of South Africa (5), citriculture has flourished in spite of greening due to efficient vector control, inoculum removal strategies, and the milder nature of the disease caused by “*Ca. Liberibacter africanus*.”

We completed the genome sequence of a “*Ca. Liberibacter africanus*” strain, PTSAPSY. Genomic DNA was isolated from individual psyllids (*T. erythrae*) collected from Pretoria, South Africa. DNA samples containing higher titers of “*Ca. Liberibacter africanus*,” estimated by real-time PCR, were selected for whole-genome amplification. Amplified “*Ca. Liberibacter africanus*” genomic DNA was used to construct a sequencing library. The complete genome sequence of the strain was obtained by an Illumina HiSeq 2000 sequencing run with a 300-bp paired-end library, which achieved an average coverage of 60 to 80×. *De novo* assembly was conducted using the Velvet assembler version 1.1.04. “*Ca. Liberibacter africanus*” contigs were identified via BLASTn and BLASTx analyses against genomes of “*Ca. Liberibacter asiaticus*” (6), “*Ca. Liberibacter americanus*” (7, 8), “*Ca.*

*Liberibacter solanacearum*” (9), and the phylogenetically related bacteria *Agrobacterium tumefaciens* and *Sinorhizobium meliloti*. All identified contigs were reconfirmed by PCR. The contig gaps were then closed by PCR-based primer walking (9), and gap-closing PCR products were resequenced using the Sanger sequencing method. Final annotation was performed by using the NCBI Prokaryotic Genomes Annotation Pipeline (PGAP) (<http://www.ncbi.nlm.nih.gov/genomes/static/Pipeline.html>).

The genome of strain PTSAPSY comprises 1,192,232 nucleotides, 34.5% G+C content, 1,017 predicted coding sequences, 44 tRNAs, three complete copies of rRNA genes (16S, 23S, and 5S), and 279 hypothetical genes. Like “*Ca. Liberibacter americanus*” and “*Ca. Liberibacter solanacearum*,” two tandemly aligned prophage segments were identified in the “*Ca. Liberibacter africanus*” genome. The average G+C content in this prophage region is 42.1%, deviating significantly from that of the core genome. Genome information will contribute to a deeper understanding of the genomic evolution of liberibacters, permit comparative genomic analysis with other “*Ca. Liberibacter*” species, and may shed light on the molecular mechanisms associated with pathogenicity.

**Nucleotide sequence accession number.** The completed genome sequence of “*Ca. Liberibacter africanus*” strain PTSAPSY has been deposited in the GenBank database under the assigned accession no. CP004021. The version described in this paper is the first version.

## ACKNOWLEDGMENTS

This study was supported by U.S. Department of Agriculture, Agricultural Research Service and Citrus Research International, South Africa.

We thank Parminder Sahota for technical support.

Trade names or commercial products in this publication are mentioned solely for the purpose of providing specific information and do not imply recommendation or endorsement by the U.S. Department of Agriculture.

## REFERENCES

1. Bové JM. 2006. Huanglongbing: a destructive, newly-emerging, century-old disease of citrus. *J Plant Pathol* 88:7–37.
2. Jagoueix S, Bové JM, Garnier M. 1994. The phloem-limited bacterium of greening disease of citrus is a member of the  $\alpha$  subdivision of the *Proteobacteria*. *Int J Syst Bacteriol* 44:379–386. <http://dx.doi.org/10.1099/00207713-44-3-379>.
3. Teixeira Ddo C, do Carmo SC, Eveillard S, Danet JL, Ayres AJ, Bové JM, Bové J. 2005. “*Candidatus Liberibacter americanus*,” associated with citrus huanglongbing (greening disease) in São Paulo State, Brazil. *Int J Syst Evol Microbiol* 55:1857–1862. <http://dx.doi.org/10.1099/ijs.0.63677-0>.
4. Texeira DC, Ayres J, Kitajima EW, Danet L, Jagoueix-Eveillard S, Sailard C, Bové JM. 2005. First report of a huanglongbing-like disease of citrus in São Paulo State, Brazil and association of a new *Liberibacter* species, “*Candidatus Liberibacter americanus*”, with the disease. *Plant Dis* 89:107. <http://dx.doi.org/10.1094/PD-89-0107A>.
5. Pietersen G, Arrebola E, Breytenbach JHJ, Korsten L, le Roux HF, la Grange H, Lopes SA, Meyer JB, Pretorius MC, Schwerdtfeger M, van Vuuren SP, Yamamoto P. 2010. Survey for “*Candidatus Liberibacter*” species in South Africa confirms the presence of only “*Ca. L. africanus*” in commercial citrus. *Plant Dis* 94:244–249. <http://dx.doi.org/10.1094/PDIS-94-2-0244>.
6. Duan Y, Zhou L, Hall DG, Li W, Doddapaneni H, Lin H, Liu L, Vahling CM, Gabriel DW, Williams KP, Dickerman A, Sun Y, Gottwald T. 2009. Complete genome sequence of citrus huanglongbing bacterium, “*Candidatus Liberibacter asiaticus*” obtained through metagenomics. *Mol Plant Microbe Interact* 22:1011–1020. <http://dx.doi.org/10.1094/MPMI-22-8-1011>.
7. Lin H, Coletta-Filho HD, Han CS, Lou B, Civerolo EL, Machado MA, Gupta G. 2013. Draft genome sequence of “*Candidatus Liberibacter americanus*” bacterium associated with citrus huanglongbing in Brazil. *Genome Announc* 1(3):e00275-13. <http://dx.doi.org/10.1128/genomeA.00275-13>.
8. Wulff NA, Zhang S, Setubal JC, Almeida NF, Martins EC, Harakava R, Kumar D, Rangel LT, Foissac X, Bové JM, Gabriel DW. 2014. The complete genome sequence of “*Candidatus Liberibacter americanus*”, associated with citrus huanglongbing. *Mol Plant Microbe Interact* 27:163–176. <http://dx.doi.org/10.1094/MPMI-09-13-0292-R>.
9. Lin H, Lou B, Glynn JM, Doddapaneni H, Civerolo EL, Chen C, Duan Y, Zhou L, Vahling CM. 2011. The complete genome sequence of “*Candidatus Liberibacter solanacearum*”, the bacterium associated with potato zebra chip disease. *PLoS One* 6:e19135. <http://dx.doi.org/10.1371/journal.pone.0019135>.