




Usefulness of red dichromatic imaging for peroral cholangioscopy



► **Fig. 1** Observation of bile duct cancer using white-light imaging (WLI), narrow-band imaging (NBI), and red dichromatic imaging (RDI). **a** A clear image of the tumor was obtained using WLI. **b** But in the presence of bile juice, the view became unclear. **c** NBI showed a red image from the effects of bile juice. **d** RDI showed a clear image even in the presence of bile juice.



► **Fig. 2** A biopsy of the lesion was performed with RDI-3. It was possible to accurately recognize the lesion, even in the presence of bile juice.



► **Video 1** Performance of peroral cholangioscopy using red dichromatic imaging (RDI) with a new-generation endoscopy system.

Peroral cholangioscopy (POCS) was developed as a diagnostic modality to directly observe bile duct lesions that are difficult to visualize by cholangiography. However, one of its main issues is that the images are strongly affected by bile juice. Red dichromatic imaging (RDI) is a new image-enhancing technology using light tones of red, amber, and green. RDI is part of a new-generation endoscopy system (EVIS X1; Olympus Medical, Tokyo, Japan) and consists of three modes [1–4]. Among the three available modes, RDI-3 is most suitable for observing the

bile duct using POCS, as yellows can be excluded with minimal alteration of reds and greens [2]. We herein report a case in which RDI-3 was useful for both the observation and biopsy of biliary cancer under POCS.

A 65-year-old woman was admitted to our hospital for an examination of wall thickening of the bile duct after gallbladder cancer surgery. Contrast-enhanced computed tomography showed enhancement of the bile duct in the perihilar area. Endoscopic retrograde cholangiography

and mother-baby POCS were performed using a TJF-Q290V and CHF-B290 scope (Olympus Medical) with the EVIS X1 system. After saline injection and washing out the bile juice, observation was started. A continuous, flat, elevated lesion was noted in the perihilar area, and a good image was initially obtained with white-light imaging (WLI) and narrow-band imaging (NBI). However, bile juice overflowing from the proximal side made it difficult to obtain clear images. Therefore, WLI was switched to RDI-3,


and a good-quality image was obtained by excluding the yellow color of the bile juice (► Fig. 1). A biopsy from the lesion was then performed, and we were able to accurately recognize the lesion even in the presence of bile juice (► Fig. 2, ► Video 1). The pathological finding was malignancy.

Endoscopy_UCTN_Code_CCL_1AZ_2AN

Competing interests

The authors declare that they have no conflict of interest.

The authors

Kazuyuki Matsumoto , **Yuki Fujii**, **Daisuke Uchida**, **Shigeru Horiguchi**, **Koichiro Tsutsumi**, **Hironari Kato**

Department of Gastroenterology and Hepatology, Okayama University Hospital, Okayama, Japan

Corresponding author

Kazuyuki Matsumoto, MD

Department of Gastroenterology and Hepatology, Okayama University Hospital, 2-5-1 Shikata-cho, Okayama 700-8558, Japan
Fax: +81-86-225-5991
matsumoto.k@okayama-u.ac.jp

References

- [1] Yahagi N, Fujimoto A, Horii J et al. Dual red imaging: a novel endoscopic imaging technology visualizing thick blood vessels in the gastrointestinal wall. *Endosc Int Open* 2019; 7: E1632–E1635
- [2] Koiwai A, Hirota M, Satoh K. Observations on the presence of bile using red dichromatic imaging 3 in peroral cholangioscopy. *Dig Endosc* 2022; 34: 649
- [3] Inoue T, Ibusuki M, Kitano R et al. Usefulness of red dichromatic imaging for post-endoscopic sphincterotomy bleeding. *Endoscopy* 2022. doi:10.1055/a-1806-1605
- [4] Toyonaga H, Kin T, Hayashi T et al. The application of texture and color enhancement imaging in transpapillary biliary cannulation. *Endoscopy* 2022. doi:10.1055/a-1806-1913

Bibliography

Endoscopy 2023; 55: E264–E265
DOI 10.1055/a-1966-0534
ISSN 0013-726X
published online 25.11.2022
© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS <https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>