

POSTER PRESENTATION

Open Access

Effect of different pressure-targeted modes of ventilation on transpulmonary pressure and inspiratory effort

N Rittayamai^{1,2,3*}, F Beloncle^{1,2,4}, R Waheed^{1,2}, L Chen^{1,2}, M Rauseo^{1,2}, G Chen^{1,2}, EC Goligher^{2,5,6}, JCM Richard⁷, L Brochard^{1,2,3}

From ESICM LIVES 2015

Berlin, Germany. 3-7 October 2015

Introduction

Spontaneous breathing during mechanical ventilation improves gas exchange and might prevent ventilator-induced diaphragm dysfunction. In pressure-targeted modes, transpulmonary pressure (P_L) is the sum of pressure generated by the ventilator and muscular pressure. When inspiratory effort increases, P_L and tidal volume (V_T) increase, potentially resulting in lung injury. This effect depends on the degree of inspiratory synchronization (i-sync); pressure-targeted modes can be classified into fully, partially, and non i-sync modes. A bench study [1] demonstrated that non-i-sync mode resulted in lower P_L and V_T than other modes, protecting the lungs from injury. We undertook to assess the effect of varying synchronization during pressure-targeted ventilation in critically ill patients.

Objectives

To compare V_T , P_L , inspiratory effort (esophageal pressure-time product, PTP_{eso}) and respiratory drive (airway occlusion pressure, $P_{0.1}$) during three pressure-targeted modes with different degrees of i-sync.

Methods

We conducted a randomized cross-over physiological study in spontaneously breathing ventilated patients. Twelve patients were enrolled (1 subsequently withdrew). Three pressure-targeted modes (Evita XL, Draeger, Germany) including fully (PC-CMV), partially (PC-SIMV), and non i-sync (APRV) modes were sequentially applied for 20 minutes in random order using the same driving

pressure, PEEP and inspiratory time. Airway, esophageal, and gastric pressures, $P_{0.1}$, and flow were recorded along with gas exchange and hemodynamics. P_L and PTP_{eso} were calculated.

Results

11/12 patients successfully completed the study. V_T was significantly lower during non i-sync mode than fully i-sync mode (Table 1, $p = 0.003$) and V_T variability increased from 13 % to 35 % with decreasing inspiratory synchronization. Maximal P_L was significantly lower in non-i-sync mode than in partially or fully i-sync modes ($p = 0.008$). There were no significant differences in gas exchange and hemodynamic parameters between modes. PTP_{eso} was significantly higher with non i-sync modes (Table 1, $p = 0.047$). This increase in PTP_{eso} was observed in the 6 patients who were not receiving intravenous sedation; no increase was observed in the 5 patients receiving continuous intravenous sedation (Figure 1).

Conclusions

Non synchronized pressure-targeted ventilation lowers V_T and P_L in comparison to fully and partially synchronized modes in spontaneously breathing ventilated patients, even with the same driving pressure. Appropriate sedation may be important to alleviate increased patient effort during such modes.

Grant Acknowledgment

This study was supported by Dr. Brochard Laboratory Research Funding and by a grant from Siriraj Hospital in Bangkok, Thailand.


¹Li Ka Shing Knowledge Institute and Critical Care Department, St. Michael's Hospital, Toronto, Canada

Full list of author information is available at the end of the article

Table 1

	PC-CMV	PC-SIMV	APRV
Tidal volume per predicted body weight (mL/kg)	7.1 ± 1.0	6.5 ± 0.8	5.6 ± 1.2*
Maximal P _L (cmH ₂ O)	14.3 ± 4.5	14.0 ± 5.2	12.4 ± 4.8* [#]
Minute ventilation (L/min)	10.4 ± 2.3	9.8 ± 2.0	9.9 ± 2.2
Breathing frequency (breaths/min)	21.6 ± 2.9	22.5 ± 3.9	26.9 ± 7.1*
PaO ₂ /FiO ₂ ratio	221 ± 65	231 ± 57	227 ± 64
PaCO ₂ (mmHg)	48 ± 11	49 ± 12	50 ± 11
P _{0.1} (cmH ₂ O)	2.6 ± 1.9	2.7 ± 1.7	3.9 ± 2.9
PTP _{eso} (cmH ₂ O*sec/min)	129.6 ± 107.1	130.2 ± 91.4	209.0 ± 174.9*

* $p < 0.05$, PC-CMV vs APRV; [#] $p < 0.05$, PC-SIMV vs APRV



Authors' details

¹Li Ka Shing Knowledge Institute and Critical Care Department, St. Michael's Hospital, Toronto, Canada. ²Interdepartmental Division of Critical Care Medicine, University of Toronto, Toronto, Canada. ³Division of Respiratory Diseases and Tuberculosis, Department of Medicine, Faculty of Medicine Siriraj Hospital, Bangkok, Thailand. ⁴Department of Intensive Care and Hyperbaric Medicine, Angers University Hospital, Angers, France. ⁵Department of Medicine, Division of Respiriology, University Health Network, Toronto, Canada. ⁶Department of Physiology, University of Toronto, Toronto, Canada. ⁷Emergency Department, General Hospital of Annecy, Annecy, France.

Published: 1 October 2015

Reference

- Richard JC, Lyazidi A, Akoumianaki E, Mortaza S, Cordioli RL, Lefebvre JC, *et al*: Potentially harmful effects of inspiratory synchronization during pressure preset ventilation. *Intensive Care Med* 2013, **39**(11):2003-10.

doi:10.1186/2197-425X-3-S1-A825

Cite this article as: Rittayamai *et al*: Effect of different pressure-targeted modes of ventilation on transpulmonary pressure and inspiratory effort. *Intensive Care Medicine Experimental* 2015 **3**(Suppl 1):A825.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com