

Complete genome sequence of *Brachyspira murdochii* type strain (56-150^T)

Amrita Pati¹, Johannes Sikorski², Sabine Gronow², Christine Munk³, Alla Lapidus¹, Alex Copeland¹, Tijana Glavina Del Tio¹, Matt Nolan¹, Susan Lucas¹, Feng Chen¹, Hope Tice¹, Jan-Fang Cheng¹, Cliff Han^{1,3}, John C. Detter^{1,3}, David Bruce^{1,3}, Roxanne Tapia³, Lynne Goodwin^{1,3}, Sam Pitluck¹, Konstantinos Liolios¹, Natalia Ivanova¹, Konstantinos Mavromatis¹, Natalia Mikhailova¹, Amy Chen⁴, Krishna Palaniappan⁴, Miriam Land^{1,5}, Loren Hauser^{1,5}, Yun-Juan Chang^{1,5}, Cynthia D. Jeffries^{1,5}, Stefan Spring², Manfred Rohde⁶, Markus Göker², James Bristow¹, Jonathan A. Eisen^{1,7}, Victor Markowitz⁴, Philip Hugenholtz¹, Nikos C. Kyripides¹, and Hans-Peter Klenk^{2*}

¹ DOE Joint Genome Institute, Walnut Creek, California, USA

² DSMZ – German Collection of Microorganisms and Cell Cultures GmbH, Braunschweig, Germany

³ Los Alamos National Laboratory, Bioscience Division, Los Alamos, New Mexico, USA

⁴ Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, California, USA

⁵ Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

⁶ HZI – Helmholtz Centre for Infection Research, Braunschweig, Germany

⁷ University of California Davis Genome Center, Davis, California, USA

*Corresponding author: Hans-Peter Klenk

Keywords: host-associated, non-pathogenic, motile, anaerobic, Gram-negative, *Brachyspiraceae*, *Spirochaetes*, GEBA

Brachyspira murdochii Stanton et al. 1992 is a non-pathogenic host-associated spirochete of the family *Brachyspiraceae*. Initially isolated from the intestinal content of a healthy swine, the 'group B spirochaetes' were first described under as *Serpulina murdochii*. Members of the family *Brachyspiraceae* are of great phylogenetic interest because of the extremely isolated location of this family within the phylum '*Spirochaetes*'. Here we describe the features of this organism, together with the complete genome sequence and annotation. This is the first completed genome sequence of a type strain of a member of the family *Brachyspiraceae* and only the second genome sequence from a member of the genus *Brachyspira*. The 3,241,804 bp long genome with its 2,893 protein-coding and 40 RNA genes is a part of the *Genomic Encyclopedia of Bacteria and Archaea* project.

Introduction

Strain 56-150^T (= DSM 12563 = ATCC 51284 = CIP 105832) is the type strain of the species *Brachyspira murdochii*. This strain was first described as *Serpulina murdochii* [1,2], and later transferred to the genus *Brachyspira* [3]. The genus *Brachyspira* consists currently of seven species, with *Brachyspira aalborgi* as the type species [4,5]. The genus *Brachyspira* is the only genus in the not yet formally described family '*Brachyspiraceae*' [6,7]. The generic name derives from 'brachys', Greek for short, and 'spira', Latin for a coil, a helix, to mean 'a short helix' [5]. The species name for *B. murdochii* de-

rives from the city of Murdoch, in recognition of work conducted at Murdoch University in Western Australia, where the type strain was identified [1]. Some species of the genus *Brachyspira* cause swine dysentery and porcine intestinal spirochetosis. Swine dysentery is a severe, mucohemorrhagic disease that sometimes leads to death of the animals [1]. *B. murdochii* is generally not considered to be a pathogen, although occasionally it has been seen in association with colitis in pigs [3,8], and was also associated with clinical problems on certain farms [9-11].

In 1992, a user-friendly and robust novel PCR-based restriction fragment length polymorphism analysis of the *Brachyspira nox*-gene was developed which allows one to identify with high specificity members of *B. murdochii* using only two restriction endonucleases only [12]. More recently, a multi-locus sequence typing scheme was developed that facilitates the identification of *Brachyspira* species and reveals the intraspecies diversity of *B. murdochii* [13] (see also <http://pubmlst.org/brachyspira/>).

Only one genome of a member of the family '*Brachyspiraceae*' has been sequenced to date: *B. hyodysenteriae* strain WA1 [14]. It is an intestinal pathogen of pigs. Based on 16S rRNA sequence this strain is 0.8% different from strain 56-150T. Here we present a summary classification and a set of features for *B. murdochii* 56-150T, together with the description of the complete genomic sequencing and annotation.

Classification and features

Brachyspira species colonize the lower intestinal tract (cecum and colons) of animals and humans [6]. The type of *B. murdochii*, 56-150T, was isolated from a healthy swine in Canada [1,15]. Other isolates have been obtained from wild rats in Ohio, USA, from laboratory rats in Murdoch, Western Australia [16], and from the joint fluid of a lame

pig [17]. Further isolates have been obtained from the feces or gastrointestinal tract of pigs in Canada, Tasmania, Queensland, and Western Australia [2,15]. The type strains of the other species of the genus *Brachyspira* share 95.9-99.4% 16S rRNA sequence identity with strain 56-150T. GenBank contains 16S rRNA sequences for about 250 *Brachyspira* isolates, all of which share at least 96% sequence identity with strain 56-150T [18]. The closest related type strain of a species outside of the *Brachyspira*, but within the order *Spirochaetales*, is *Turneriella parva* [19], which exhibits only 75% 16S rRNA sequence similarity [18]. 16S rRNA sequences from environmental samples and metagenomic surveys do not exceed 78-79% sequence similarity to strain 56-150T, with the sole exception of one clone from a metagenomic analysis of human diarrhea [20], indicating that members of the species, genus and even family are poorly represented in the habitats outside of various animal intestines screened thus far (status March 2010).

Figure 1 shows the phylogenetic neighborhood of *B. murdochii* 56-150T in a 16S rRNA based tree. The sequence of the single 16S rRNA gene in the genome sequence is identical with the previously published 16S rRNA gene sequence generated from DSM 12563 (AY312492).




Figure 1. Phylogenetic tree highlighting the position of *B. murdochii* 56-150T relative to the other type strains within the genus and to the type strains of the other genera within the class *Spirochaetes* (excluding members of the *Spirochaetaceae*). The tree was inferred from 1,396 aligned characters [21,22] of the 16S rRNA gene sequence under the maximum likelihood criterion [23] and rooted in accordance with the current taxonomy. The branches are scaled in terms of the expected number of substitutions per site. Numbers above branches are support values from 1,000 bootstrap replicates if [24] larger than 60%. Lineages with type strain genome sequencing projects registered in GOLD [25] are shown in blue, published genomes in bold.

The cells of *B. murdochii* 56-150^T were 5 - 8 by 0.35 - 0.4 μm in size (Table 1 and Figure 2), and each cell possessed 22 to 26 flagella (11 to 13 inserted at each end) [1]. In brain/heart infusion broth containing 10% calf serum (BHIS) under an N₂-O₂ (99::1) atmosphere, strain 56-150^T had optimum growth temperatures of 39 to 42°C (shortest population doubling times and highest final population densities) [1]. In BHIS broth at 39°C, the doubling times of strain 56-150^T were 2 to 4 h, and the final population densities were 0.5 x 10⁹ to 2.0 x 10⁹ cells/ml. Strain 56-150^T did not grow at 32 or 47°C [1].

Substrates that support growth of strain 56-150^T in HS broth (basal heart infusion broth containing 10% fetal calf serum) include glucose, fructose, sucrose, N-acetylglucosamine, pyruvate, L-fucose, cellobiose, trehalose, maltose, mannose, and lactose, but not galactose, D-fucose, glucosamine, ribose, raffinose, rhamnose, or xylose [1]. In HS broth supplemented with 0.4% glucose under an N₂-O₂ (99::1) atmosphere, the metabolic end products of strain

56-150^T are acetate, butyrate, ethanol, CO₂, and H₂. Strain 56-150^T produces more H₂ than CO₂ [1], which is indicative of NADH-ferredoxin oxidoreductase reaction [6]. The ethanol is likely to be formed from acetyl-CoA by the enzymes acetaldehyde dehydrogenase and alcohol dehydrogenase [6]. Strain 56-150^T is weakly hemolytic, negative for indole production, does not hydrolyze hippurate, is negative for α -galactosidase and α -glucosidase activity, but positive for β -glucosidase activity [1]. Strain 56-150^T is anaerobic but aerotolerant [1].

Minimal inhibitory concentrations have been determined for strain 56-150^T for tiamulin hydrogen fumarate, tylosin tartrate, erythromycin, clindamycin hydrochloride, virginiamycin, and carbadox [36]. Several strains of *B. murdochii* have been described to be naturally resistant against rifampicin [7,32]. Also, a ring test for quality assessment for diagnostics and antimicrobial susceptibility testing of the genus *Brachyspira* has been reported [37].




Figure 2. Scanning electron micrograph of *B. murdochii* 56-150^T

Chemotaxonomy

At present, there are no reports on the chemotaxonomy of *B. murdochii*. However, some data are available for *B. innocens* (formerly classified as *Treponema innocens* [6]), the species that is currently most closely related to *B. murdochii* [13]. *B. innocens* cellular phospholipids and glycolipids were found to

contain acyl (fatty acids with ester linkage) with alkenyl (unsaturated alcohol with ether linkage) side chains [6,38]. The glycolipid of *B. innocens* contains monoglycosyldiglyceride (MGDG) and, in most strains, acylMGDG is also found, with galactose as the predominant sugar moiety [38].

Table 1. Classification and general features of *B. murdochii* 56-150 according to the MIGS recommendations [26]

MIGS ID	Property	Term	Evidence code
Current classification	Domain	<i>Bacteria</i>	TAS [27]
	Phylum	<i>Spirochaetes</i>	TAS [28]
	Class	<i>Spirochaetes</i>	TAS [28]
	Order	<i>Spirochaetales</i>	TAS [29,30]
	Family	<i>Brachyspiraceae</i>	TAS [31]
	Genus	<i>Brachyspira</i>	TAS [5]
	Species	<i>Brachyspira murdochii</i>	TAS [1]
MIGS-22	Type strain	56-150	TAS [1]
	Gram stain	negative	TAS [1]
	Cell shape	helical cells with regular coiling pattern	TAS [1]
	Motility	motile (periplasmic flagella)	TAS [1]
	Sporulation	non-sporulating	TAS [1]
	Temperature range	does not grow at 32°C or 47°C	TAS [1]
	Optimum temperature	39°C	TAS [1]
	Salinity	unknown	TAS
	Oxygen requirement	anaerobic, aerotolerant	TAS [1]
MIGS-6	Carbon source	soluble sugars	TAS [1]
	Energy source	chemoorganotrophic	TAS [1]
	Habitat	animal intestinal tract	TAS [6]
MIGS-15	Biotic relationship	host-associated	TAS [32]
MIGS-14	Pathogenicity	no	TAS [33]
MIGS-4	Biosafety level	1	TAS [34]
	Isolation	swine	TAS [15]
	Geographic location	Quebec, Canada	TAS [15]
MIGS-5	Sample collection time	1992	TAS [15]
MIGS-4.1	Latitude	52.939	TAS [1]
MIGS-4.2	Longitude	-73.549	TAS [1]
MIGS-4.3	Depth	not reported	TAS
MIGS-4.4	Altitude	not reported	TAS

Evidence codes - IDA: Inferred from Direct Assay (first time in publication); TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from of the Gene Ontology project [35]. If the evidence code is IDA, then the property was directly observed by one of the authors or an expert mentioned in the acknowledgements.

Genome sequencing and annotation

Genome project history

This organism was selected for sequencing on the basis of its phylogenetic position [39], and is part of the *Genomic Encyclopedia of Bacteria and Archaea* project [40]. The genome project is deposited in the Genome OnLine Database [25] and the com-

plete genome sequence is deposited in GenBank Sequencing, finishing and annotation were performed by the DOE Joint Genome Institute (JGI). A summary of the project information is shown in Table 2.

Table 2. Genome sequencing project information

MIGS ID	Property	Term
MIGS-31	Finishing quality	Finished
	Libraries used	Four genomic libraries: two Sanger 6kb and 8 kb pMCL200 library, one fosmid library, one 454 standard library
MIGS-28		
MIGS-29	Sequencing platforms	ABI3730, 454 GS FLX
MIGS-31.2	Sequencing coverage	19.7× Sanger; 48.9× pyrosequence
MIGS-30	Assemblers	Newbler version 1.1.02.15, phrap
MIGS-32	Gene calling method	Prodigal 1.4, GenePRIMP
	INSDC ID	CP001959
	Genbank Date of Release	May 13, 2010
	GOLD ID	Gc01276
	NCBI project ID	29543
	Database: IMG-GEBA	2502422316
MIGS-13	Source material identifier	DSM 12563
	Project relevance	Tree of Life, GEBA

Growth conditions and DNA isolation

B. murdochii, strain 56-150^T, DSM 12563, was grown anaerobically in DSMZ medium 840 (*Serpulina murdochii* medium) [41] at 37°C. DNA was isolated from 0.5–1 g of cell paste using Qiagen Genomic 500 DNA Kit (Qiagen, Hilden, Germany) with lysis modification st/L according to Wu *et al.* [40].

Genome sequencing and assembly

The genome was sequenced using a combination of Sanger and 454 sequencing platforms. All general aspects of library construction and sequencing performed can be found at the JGI website (<http://www.jgi.doe.gov/>). In total, 861,386 Pyrosequencing reads were assembled using the Newbler assembler version 1.1.02.15 (Roche). Large Newbler contigs were broken into 3,554 overlapping fragments of 1,000 bp and entered into assembly as pseudo-reads. The sequences were assigned quality scores based on Newbler consensus q-scores with modifications to account for overlap redundancy and adjust inflated q-scores. A hybrid 454/Sanger assembly was made using the parallel phrap assembler (High Performance Software, LLC). Possible misassemblies were corrected with Dupfinisher or transposon bombing of bridging clones [42]. A total of 300 Sanger finishing reads were produced to close gaps, to resolve repetitive regions, and to raise the quality of the finished sequence. The error rate of the completed genome sequence is less than 1 in

100,000. Together, the combination of the Sanger and 454 sequencing platforms provided 68.6× coverage of the genome. The final assembly contains 79,829 Sanger reads and 861,386 pyrosequencing reads.

Genome annotation

Genes were identified using [Prodigal](#) [43] as part of the Oak Ridge National Laboratory genome annotation pipeline, followed by a round of manual curation using the JGI [GenePRIMP](#) pipeline [44]. The predicted CDSs were translated and used to search the National Center for Biotechnology Information (NCBI) nonredundant database, UniProt, TIGRFam, Pfam, PRIAM, KEGG, COG, and InterPro databases. Additional gene prediction analysis and functional annotation was performed within the Integrated Microbial Genomes - Expert Review (IMG-ER) platform [45].

Genome properties

The genome is 3,241,804 bp long and comprises one main circular chromosome with an overall GC content of 27.8% (Table 3 and Figure 3). Of the 2,893 genes predicted, 2,853 were protein-coding genes, and 40 RNAs. A total of 44 pseudogenes were identified. The majority of the protein-coding genes (66.2%) were assigned a putative function while those remaining were annotated as hypothetical proteins. The distribution of genes into COGs functional categories is presented in Table 4.

Table 3. Genome Statistics

Attribute	Value	% of Total
Genome size (bp)	3,241,804	100.00%
DNA coding region (bp)	2,841,470	87.65%
DNA G+C content (bp)	899,647	27.75%
Number of replicons	1	
Extrachromosomal elements	0	
Total genes	2,893	100.00%
RNA genes	40	1.38%
rRNA operons	1	
Protein-coding genes	2,893	98.62%
Pseudo genes	44	1.52%
Genes with function prediction	1,914	66.16%
Genes in paralog clusters	610	21.09%
Genes assigned to COGs	1,815	62.74%
Genes assigned Pfam domains	1,973	68.20%
Genes with signal peptides	577	19.94%
Genes with transmembrane helices	737	25.48%
CRISPR repeats	2	


**Figure 3.** Graphical circular map of the genome. From outside to the center: Genes on forward strand (color by COG categories), Genes on reverse strand (color by COG categories), RNA genes (tRNAs green, rRNAs red, other RNAs black), GC content, GC skew.

Table 4. Number of genes associated with the general COG functional categories

Code	value	%age	Description
J	134	6.6	Translation, ribosomal structure and biogenesis
A	1	0.0	RNA processing and modification
K	81	4.0	Transcription
L	104	5.2	Replication, recombination and repair
B	0	0.0	Chromatin structure and dynamics
D	20	1.0	Cell cycle control, cell division, chromosome partitioning
Y	0	0.0	Nuclear structure
V	44	2.2	Defense mechanisms
T	116	5.8	Signal transduction mechanisms
M	143	7.1	Cell wall/membrane/envelope biogenesis
N	100	5.0	Cell motility
Z	0	0.0	Cytoskeleton
W	0	0.0	Extracellular structures
U	51	2.5	Intracellular trafficking secretion, and vesicular transport
O	62	3.1	Posttranslational modification, protein turnover, chaperones
C	111	5.5	Energy production and conversion
G	143	7.1	Carbohydrate transport and metabolism
E	185	9.2	Amino acid transport and metabolism
F	56	2.8	Nucleotide transport and metabolism
H	67	3.3	Coenzyme transport and metabolism
I	53	2.6	Lipid transport and metabolism
P	99	4.9	Inorganic ion transport and metabolism
Q	20	1.0	Secondary metabolites biosynthesis, transport and catabolism
R	286	14.2	General function prediction only
S	143	7.1	Function unknown
-	1,078	37.3	Not in COGs

Acknowledgements

We would like to gratefully acknowledge the help of Sabine Welnitz for growing *B. murdochii* cells and Susanne Schneider for DNA extraction and quality analysis (both at DSMZ). This work was performed under the auspices of the US Department of Energy Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory under con-

tract No. DE-AC02-05CH11231, Lawrence Livermore National Laboratory under Contract No. DE-AC52-07NA27344, Los Alamos National Laboratory under contract No. DE-AC02-06NA25396, UT-Battelle, and Oak Ridge National Laboratory under contract DE-AC05-00OR22725, as well as German Research Foundation (DFG) INST 599/1-1 and SI 1352/1-2.

References

1. Stanton TB, Fournie-Amazouz E, Postic D, Trott DJ, Grimont PAD, Baranton G, Hampson DJ, Saint Girons I. Recognition of two new species of intestinal spirochetes: *Serpulina intermedia* sp. nov. and *Serpulina murdochii* sp. nov. *Int J Syst Bacteriol* 1997; **47**:1007-1012. [PubMed doi:10.1099/00207713-47-4-1007](#)
2. Lee JL, Hampson DJ. Genetic characterisation of intestinal spirochaetes and their association with disease. *J Med Microbiol* 1994; **40**:365-371. [PubMed doi:10.1099/00222615-40-5-365](#)
3. Hampson DJ, La T. Reclassification of *Serpulina intermedia* and *Serpulina murdochii* in the genus *Brachyspira* as *Brachyspira intermedia* comb. nov.

- and *Brachyspira murdochii* comb. nov. *Int J Syst Evol Microbiol* 2006; **56**:1009-1012. [PubMed doi:10.1099/ijss.0.64004-0](#)
4. Euzéby JP. List of bacterial names with standing in nomenclature: A folder available on the Internet. *Int J Syst Bacteriol* 1997; **47**:590-592. [PubMed doi:10.1099/00207713-47-2-590](#)
5. Hovind-Hougen K, Birch-Andersen A, Henrik-Nielsen R, Orholm M, Pedersen JO, Teglbaerg PS, Thaysen EH. Intestinal spirochetosis: morphological characterization and cultivation of the spirochete *Brachyspira aalborgi* gen. nov., sp. nov. *J Clin Microbiol* 1982; **16**:1127-1136. [PubMed](#)
6. Stanton TB. 2006. The genus *Brachyspira*. In M Dworkin, S Falkow, E Rosenberg, KH Schleifer E Stackebrandt (eds), *The Prokaryotes*, 3. ed, vol. 7. Springer, New York, p. 330-356.
7. Paster BJ, Dewhirst FE. Phylogenetic foundation of spirochetes. *J Mol Microbiol Biotechnol* 2000; **2**:341-344. [PubMed](#)
8. Weissenböck H, Maderner A, Herzog AM, Lussy H, Nowotny N. Amplification and sequencing of *Brachyspira* spp. specific portions of *nox* using paraffin-embedded tissue samples from clinical colitis in Austrian pigs shows frequent solitary presence of *Brachyspira murdochii*. *Vet Microbiol* 2005; **111**:67-75. [PubMed doi:10.1016/j.vetmic.2005.09.002](#)
9. Stephens CP, Hampson DJ. Prevalence and disease association of intestinal spirochaetes in chickens in eastern Australia. 1999; **28**:447-454.
10. Stephens CP, Oxberry SL, Phillips ND, La T, Hampson DJ. The use of multilocus enzyme electrophoresis to characterise intestinal spirochaetes (*Brachyspira* spp.) colonising hens in commercial flocks. *Vet Microbiol* 2005; **107**:149-157. [PubMed doi:10.1016/j.vetmic.2005.01.011](#)
11. Feberwee A, Hampson DJ, Phillips ND, La T, van der Heijden HMJF, Wellenberg GJ, Dwars RM, Landman WJM. Identification of *Brachyspira hyodysenteriae* and other pathogenic *Brachyspira* species in chickens from laying flocks with diarrhea or reduced production or both. *J Clin Microbiol* 2008; **46**:593-600. [PubMed doi:10.1128/JCM.01829-07](#)
12. Rohde J, Rothkamp A, Gerlach GF. Differentiation of porcine *Brachyspira* species by a novel *nox* PCR-based restriction fragment length polymorphism analysis. *J Clin Microbiol* 2002; **40**:2598-2600. [PubMed doi:10.1128/JCM.40.7.2598-2600.2002](#)
13. Råsbäck T, Johansson KE, Jansson DS, Fellstrom C, Alikhani MY, La T, Dunn DS, Hampson DJ. Development of a multilocus sequence typing scheme for intestinal spirochaetes within the genus *Brachyspira*. *Microbiology* 2007; **153**:4074-4087. [PubMed doi:10.1099/mic.0.2007/008540-0](#)
14. Bellgard MI, Eanchanthuek P, La T, Ryan K, Moolhuijzen P, Albertyn Z, Shaban B, Motro Y, Dunn DS, Schibeci D, et al. Genome sequence of the pathogenic intestinal spirochaete *Brachyspira hyodysenteriae* reveals adaptations to its lifestyle in the porcine large intestines. *PLoS ONE* 2009; **4**:e4641. [PubMed doi:10.1371/journal.pone.0004641](#)
15. Lee JI, Hampson DJ, Lymbery AJ, Harders SJ. The porcine intestinal spirochaetes: identification of new genetic groups. *Vet Microbiol* 1993; **34**:273-285. [PubMed doi:10.1016/0378-1135\(93\)90017-2](#)
16. Trott DJ, Atyeo RF, Lee JI, Swayne DA, Stouteburg JW, Hampson DJ. Genetic relatedness amongst intestinal spirochaetes isolated from rats and birds. *Lett Appl Microbiol* 1996; **23**:431-436. [PubMed doi:10.1111/j.1472-765X.1996.tb01352.x](#)
17. Hampson DJ, Robertson ID, Oxberry SL. Isolation of *Serpulina murdochii* from the joint fluid of a lame pig. *Aust Vet J* 1999; **77**:48. [PubMed doi:10.1111/j.1751-0813.1999.tb12430.x](#)
18. Chun J, Lee JH, Jung Y, Kim M, Kim S, Kim BK, Lim YW. EzTaxon: a web-based tool for the identification of prokaryotes based on 16S ribosomal RNA gene sequences. *Int J Syst Evol Microbiol* 2007; **57**:2259-2261. [PubMed doi:10.1099/ijss.0.64915-0](#)
19. Levett PN, Morey RE, Galloway R, Steigerwalt AG, Ellis WA. Reclassification of *Leptospira parva* Hovind-Hougen et al. 1982 as *Turneriella parva* gen. nov., comb. nov. *Int J Syst Evol Microbiol* 2005; **55**:1497-1499. [PubMed doi:10.1099/ijss.0.63088-0](#)
20. Finkbeiner SR, Allred AF, Tarr PI, Klenin EJ, Kirkwood CD, Wang D. Metagenomic analysis of human diarrhea: viral detection and discovery. *PLoS Pathog* 2008; **4**:e1000011. [PubMed doi:10.1371/journal.ppat.1000011](#)
21. Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 2000; **17**:540-552. [PubMed](#)

22. Lee C, Grasso C, Sharlow MF. Multiple sequence alignment using partial order graphs. *Bioinformatics* 2002; **18**:452-464. [PubMed](#) [doi:10.1093/bioinformatics/18.3.452](https://doi.org/10.1093/bioinformatics/18.3.452)
23. Stamatakis A, Hoover P, Rougemont J. A Rapid Bootstrap algorithm for the RAxML web servers. *Syst Biol* 2008; **57**:758-771. [PubMed](#) [doi:10.1080/10635150802429642](https://doi.org/10.1080/10635150802429642)
24. Patterson ND, Alipour M, Bininda-Emonds ORP, Moret BME, Stamatakis A. How many bootstrap replicates are necessary? *Lect Notes Comput Sci* 2009; **5541**:184-200. [doi:10.1007/978-3-642-02008-7_13](https://doi.org/10.1007/978-3-642-02008-7_13)
25. Liolios K, Chen IM, Mavromatis K, Tavernarakis N, Hugenholtz P, Markowitz VM, Kyriakis NC. The Genomes On Line Database (GOLD) in 2009: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res* 2010; **38**:D346-D354. [PubMed](#) [doi:10.1093/nar/gkp848](https://doi.org/10.1093/nar/gkp848)
26. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Anjiouli SV, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547. [PubMed](#) [doi:10.1038/nbt1360](https://doi.org/10.1038/nbt1360)
27. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci USA* 1990; **87**:4576-4579. [PubMed](#) [doi:10.1073/pnas.87.12.4576](https://doi.org/10.1073/pnas.87.12.4576)
28. Garrity GM, Lilburn TG, Cole JR, Harrison SH, Euzéby J, Tindall BJ. Taxonomic outline of the Bacteria and Archaea, Release 7.7 March 6, 2007. Part 11 - The Bacteria: Phyla "Planctomycetes", "Chlamydiae", "Spirochaetes", "Fibrobacteres", "Acidobacteria", "Bacteroidetes", "Fusobacteria", "Verrucomicrobia", "Dictyoglomi", "Gemmatimonadetes", and "Lentisphaerae". <http://www.taxonomicoutline.org/index.php/toba/article/viewFile/188/220> 2007.
29. Skerman VBD, McGowan V, Sneath PHA. Approved Lists of Bacterial Names. *Int J Syst Bacteriol* 1980; **30**:225-420. [PubMed](#) [doi:10.1099/00207713-30-1-225](https://doi.org/10.1099/00207713-30-1-225)
30. Buchanan RE. Studies in the nomenclature and classification of Bacteria. II. The primary subdivisions of the Schizomycetes. *J Bacteriol* 1917; **2**:155-164. [PubMed](#)
31. Paster BJ, Dewhurst FE. Phylogenetic foundation of spirochetes. *J Mol Microbiol Biotechnol* 2000; **2**:341-344. [PubMed](#)
32. Imachi H, Sakai S, Hirayama H, Nakagawa S, Nunoura T, Takai K, Horikoshi K. *Exilispira thermophila* gen. nov., sp. nov., an anaerobic, thermophilic spirochaete isolated from a deep-sea hydrothermal vent chimney. *Int J Syst Evol Microbiol* 2008; **58**:2258-2265. [PubMed](#) [doi:10.1099/ijss.0.65727-0](https://doi.org/10.1099/ijss.0.65727-0)
33. Stanton TB, Postic D, Jensen NS. *Serpulina alvinipulli* sp. nov., a new *Serpulina* species that is enteropathogenic for chickens. *Int J Syst Bacteriol* 1998; **48**:669-676. [PubMed](#) [doi:10.1099/00207713-48-3-669](https://doi.org/10.1099/00207713-48-3-669)
34. Classification of Bacteria and Archaea in risk groups. www.baua.de TRBA 466.
35. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene Ontology: tool for the unification of biology. *Nat Genet* 2000; **25**:25-29. [PubMed](#) [doi:10.1038/75556](https://doi.org/10.1038/75556)
36. Karlsson M, Fellstrom C, Gunnarsson A, Landen A, Franklin A. Antimicrobial susceptibility testing of porcine brachyspira (*Serpulina*) species isolates. *J Clin Microbiol* 2003; **41**:2596-2604. [PubMed](#) [doi:10.1128/JCM.41.6.2596-2604.2003](https://doi.org/10.1128/JCM.41.6.2596-2604.2003)
37. Råsbäck T, Fellström C, Bergsjø B, Cizek A, Collin K, Gunnarsson A, Jensen SM, Mars A, Thomson J, Vyt P, et al. Assessment of diagnostics and antimicrobial susceptibility testing of *Brachyspira* species using a ring test. *Vet Microbiol* 2005; **109**:229-243. [PubMed](#) [doi:10.1016/j.vetmic.2005.05.009](https://doi.org/10.1016/j.vetmic.2005.05.009)
38. Matthews HM, Kinyon JM. Cellular lipid comparisons between strains of *Treponema hydysenteriae* and *Treponema innocens*. *Int J Syst Bacteriol* 1984; **34**:160-165. [PubMed](#) [doi:10.1099/00207713-34-1-160](https://doi.org/10.1099/00207713-34-1-160)
39. Klenk HP, Göker M. En route to a genome-based classification of Archaea and Bacteria? *Syst Appl Microbiol* (In press).
40. Wu D, Hugenholtz P, Mavromatis K, Pukall R, Dalin E, Ivanova NN, Kunin V, Goodwin L, Wu M, Tindall BJ, et al. A phylogeny-driven genomic encyclopaedia of Bacteria and Archaea. *Nature* 2009; **462**:1056-1060. [PubMed](#) [doi:10.1038/nature08656](https://doi.org/10.1038/nature08656)
41. List of growth media used at DSMZ: http://www.dsmz.de/microorganisms/media_list.php
42. Sims D, Brettin T, Detter J, Han C, Lapidus A, Copeland A, Glavina Del Rio T, Nolan M, Chen F, Lucas S, et al. Complete genome sequence of *Ky-*

- tococcus sedentarius* type strain (541^T). *Stand Genomic Sci* 2009; **1**:12-20. [doi:10.4056/sigs.761](https://doi.org/10.4056/sigs.761)
43. Hyatt D, Chen GL, Locascio PF, Land ML, Larmier FW, Hauser LJ. Prodigal Prokaryotic Dynamic Programming Genefinding Algorithm. *BMC Bioinformatics* 2010; **11**:119. [PubMed](https://pubmed.ncbi.nlm.nih.gov/20471210/) [doi:10.1186/1471-2105-11-119](https://doi.org/10.1186/1471-2105-11-119)
44. Pati A, Ivanova N, Mikhailova N, Ovchinkova G, Hooper SD, Lykidis A, Kyrpides NC. GenePRIMP: A Gene Prediction Improvement Pipeline for microbial genomes. *Nat Methods* (Epub). [doi:10.1038/nmeth.1457](https://doi.org/10.1038/nmeth.1457)
45. Markowitz VM, Ivanova NN, Chen IMA, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 2009; **25**:2271-2278. [PubMed](https://pubmed.ncbi.nlm.nih.gov/19624073/) [doi:10.1093/bioinformatics/btp393](https://doi.org/10.1093/bioinformatics/btp393)