



Review

Silver Nanoparticle's Toxicological Effects and Phytoremediation

Muhammad Ihtisham ^{1,†}, Azam Noori ^{2,†}, Saurabh Yadav ^{3,†}, Mohammad Sarraf ^{4,†}, Pragati Kumari ⁵, Marian Brestic ^{6,7}, Muhammad Imran ⁸, Fuxing Jiang ¹, Xiaojun Yan ^{1,*} and Anshu Rastogi ^{9,10,*}

- ¹ College of Landscape Architecture, Sichuan Agricultural University, Chengdu 611130, China; ihtisham@sicau.edu.cn (M.I.); bnjfx@sicau.edu.cn (F.J.)
 - ² Department of Biology, Merrimack College, North Andover, MA 01845, USA; nooria@merrimack.edu
 - ³ Department of Biotechnology, Hemvati Nandan Bahuguna Garhwal (Central) University, Garhwal, Srinagar 246174, Uttarakhand, India; saurabh Yadav40@rediffmail.com
 - ⁴ Department of Horticulture Science, Shiraz Branch, Islamic Azad University, Shiraz 71987-74731, Iran; sarraf.science@gmail.com
 - ⁵ Scientist Hostel-S-02, Chauras Campus, Garhwal, Srinagar 246174, Uttarakhand, India; pragati27@gmail.com
 - ⁶ Department of Plant Physiology, Slovak University of Agriculture, A. Hlinku 2, 94976 Nitra, Slovakia; marian.brestic@uniag.sk
 - ⁷ Department of Botany and Plant Physiology, Faculty of Agrobiology, Food and Natural Resources, Czech University of Life Sciences Prague, 16500 Prague, Czech Republic
 - ⁸ State Key Laboratory for Conservation and Utilization of Subtropical Agro-Bioresources, College of Agriculture, South China Agricultural University, Guangzhou 510642, China; muhammadimran@scau.edu.cn
 - ⁹ Laboratory of Bioclimatology, Department of Ecology and Environmental Protection, Poznan University of Life Sciences, Piątkowska 94, 60-649 Poznan, Poland
 - ¹⁰ Faculty of Geo-Information Science and Earth Observation (ITC), University of Twente, 7500 AE Enschede, The Netherlands
- * Correspondence: daniel@sicau.edu.cn (X.Y.); anshu.rastogi@up.poznan.pl (A.R.)
† These authors have contributed equally to the work.



Citation: Ihtisham, M.; Noori, A.; Yadav, S.; Sarraf, M.; Kumari, P.; Brestic, M.; Imran, M.; Jiang, F.; Yan, X.; Rastogi, A. Silver Nanoparticle's Toxicological Effects and Phytoremediation. *Nanomaterials* **2021**, *11*, 2164. <https://doi.org/10.3390/nano11092164>

Academic Editor: Alexey Pestryakov

Received: 27 May 2021

Accepted: 19 August 2021

Published: 24 August 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The advancement in nanotechnology has brought numerous benefits for humans in diverse areas including industry, medicine, and agriculture. The demand in the application of nanomaterials can result in the release of these anthropogenic materials into soil and water that can potentially harm the environment by affecting water and soil properties (e.g., soil texture, pH, organic matter, and water content), plants, animals, and subsequently human health. The properties of nanoparticles including their size, surface area, and reactivity affect their fate in the environment and can potentially result in their toxicological effects in the ecosystem and on living organisms. There is extensive research on the application of nano-based materials and the consequences of their release into the environment. However, there is little information about environmentally friendly approaches for removing nanomaterials from the environment. This article provides insight into the application of silver nanoparticles (AgNPs), as one of the most commonly used nanomaterials, their toxicological effects, their impacts on plants and microorganisms, and briefly reviews the possibility of remediation of these metabolites using phytotechnology approaches. This article provides invaluable information to better understand the fate of nanomaterials in the environment and strategies in removing them from the environment.

Keywords: phytoremediation; silver nanoparticles; toxicological effects; environmental sustainability

1. Introduction

Nanotechnology is a growing and advanced area of science and engineering that focuses on the synthesis, and application of matters on the nano scale with at least one dimension less than 100 nm [1]. The characteristics of nanomaterials are mainly derived from their small size, high surface area/volume ratio, and high stability [2]. Compared with conventional materials, the properties of engineered nanomaterials have improved their

application [3,4]. They have physical properties such as high uniformity, high conductivity, or special optical properties, making them popular in chemistry, physics, and biology [5].

Nanotechnology is used in various areas [6,7], including agriculture to improve crop yield by offering innovative agrochemical formulations and delivery methods, which in turn reduces the need for pesticide application [8]. Nanoparticles are also used in precision farming [9] to achieve sustainable agriculture, as these particles are attributed to altering critical plant life events and are used in agriculture for a variety of purposes such as minimizing nutritional loss, reducing various environmental stresses, and increasing crop yield [10,11]. In addition, nanoparticles are used in agriculture as nanofertilizers, nanopesticides, or nanosensors to protect plants against pathogens and improve productivity [12].

Extensive research on the impacts of nanoparticles on living organisms suggests that nanoparticles have “grey shade” by resulting in both pros and cons effects [13]. Metal-based nanoparticles are among the most recent types of anthropogenic materials that can potentially harm the ecosystem if used in a high concentration [14]. Nanoparticles can be released into the soil during production, agricultural or industrial applications, or accidental spillage [15]. Nanoparticles can also increase the bioavailability of metals in the soil and potentially result in higher environmental risk [16].

Different types of engineered NPs are synthesized for various applications. Silver nanoparticles (AgNPs) are among the most commonly used engineered nanomaterials with medicinal, industrial, and agricultural applications [17]. Considering the vast usage of AgNPs, there is a possibility of their release into the environment, and their potential toxicological effects on plants and animals. Apart from using the particulate form of silver, AgNPs may be transformed to silver oxide or silver sulfide via oxidation or sulfidation, respectively, and these ones impact the soil and living organisms in a variety of ways [18]. Therefore, it is critical to address the behavior of nanoparticles in the environment and possible methods for their removal. This review focuses on three objectives to discuss this issue including: the possible pathways for the release of AgNPs into the environment; the toxicological effects of AgNPs on plants and microorganisms; and the recommended phytoremediation approaches. This review provides invaluable information for a more sustainable application of AgNPs.

2. Applications of Silver Nanoparticles

The application of silver particles has a centuries-old history due to their therapeutic nature in medicine and storage vessel for beverages [19]. AgNPs are among the most commonly used nanoparticles, broadly due to their numerous uses. In 2015, the AgNPs market was reported to be \$1 billion, which is expected to rise to \$3 billion by 2024 [20,21]. Due to the antibacterial, antifungal, antiseptic, and antiviral properties, AgNPs are of prime importance in medical applications and were broadly used as an antimicrobial agent before the discovery of antibiotics [19,22,23]. Up until now, AgNPs have been utilized in a diverse range of applications in many fields, with some examples of health (immunity-based food supplements, pharmaceuticals, disinfectants, burn treatment products, and wound healing/dressings, radiotherapy), other biomedical products, optics, biosensing, environmental remediation, food industries, cleaning, water and air disinfection, electronics, textile, packaging, skincare products, detergents, plastics, paints, and even children toys [21,24–27]. AgNPs have also been used in agriculture as plant growth promoter/fertilizer, fruits ripening and preservation agent, plant disease control fungicide, and insecticide [25,28]. AgNPs also have several distinctive physicochemical properties, such as high thermal and electrical conductivity, improved surface Raman scattering, catalytic activity, and non-linear optical behavior [29]. This widespread use of AgNPs ensures particle release into the environment and makes it one of the most exploited nanomaterials [30,31] resulting in unknown impacts on plants, microorganisms, animals, and humans in the exposed environment.

AgNPs have the average production of 500 tons per year [32] and are known for their widespread application which has made their entry into many commercial products [33–35]. They have diverse applications and are becoming more common in industrial

processes [36]. The high usage of AgNPs can result in the direct or indirect release of these nanomaterials into the environment and can potentially cause toxicity for various organisms including aquatic organisms, plants, and humans. The direct transformation includes aerial deposition, run-off, or sewage discharges (release from industrial and household products into water bodies). The indirect release includes discharges from organic or inorganic fertilizers and engineered plant growth substances used as fertilizer substitutes that can be accumulated in soil and underground water, accidental spills during manufacturing and transport, and biosphere pollutions through atmospheric emissions from smelting, coal combustion, and cloud seeding [37–41]. AgNPs can be oxidized in the environment and be transformed into the ionic form of the silver (Ag^+) which can be more toxic than the particulate form of silver [42]. AgNPs and Ag^+ can make their way into the water bodies and soils during or after the lifetime of the product and eventually affect the ecosystem [30,43,44].

Prior to evaluating plant's potential to remove nanomaterials from the environment, it is critical to understand the environmental toxicology effects of these materials and their effects on plants and microorganisms.

3. Environmental and Toxicological Effects of AgNPs

The heavy inclusion of AgNPs in today's world has not been assessed and correlated with environmental risks [45,46] and balance between nanoparticles and biodiversity. Recently, the application of nanomaterials has gained more popularity in different areas [47,48] which raises the risk of their release into the environment and potential toxicological effects. AgNPs are considered as enormously toxic on the basis of L(E) C_{50} values (the amount which causes 50% death of tested animals when the material was given at once) for environmentally related organisms at the levels of L(E) C_{50} b 0.1 mg L^{-1} [32], while on the other hand, the predicted environmental concentration (PECs) of AgNPs in the environment ranges between 0.03 to 0.08 mg L^{-1} [49]. As per the World Health Organization (WHO) and Environmental Protection Agency (EPA), the maximum contamination limit (MCL) of toxic range of silver is 0.1 mg L^{-1} [50–54], although naturally occurring silver concentrations are generally low in the environment (surface waters) but are continuously increasing at higher levels due to runoff and wastewater from urban and industrial areas [50]. AgNPs released into the environment can be oxidized and generate the ionic form of silver that is more reactive than the particulate form. The high concentration of AgNPs and their potential to be oxidized in the environment can cause toxicity for living organisms. Consequently, this dilemma, if not addressed on time, will more negatively affect the ecosystem also by inhibiting growth of plants, polluting drinking water, and causing harm to human health. This will ultimately curtail the fauna and flora. Therefore, sustainable methods for the remediation of contaminated soil with AgNPs must be investigated.

Since the 1980s, the U.S. Environmental Protection Agency (EPA) regulated AgNPs usage and release into the environment to limit their impacts on the ecosystem [55]. To curb their impacts on living organisms, the concentration of total silver in aquatic systems is limited to 1.2–13 ppb (depending on CaCO_3 concentration) [55]. Over the last decade, the vast application of AgNPs called for examination of these particles to determine their long-lasting effects on ecosystems. The occurrence of silver species in the environment, particularly the water supply, can result in bioaccumulation across several trophic levels with potentially severe toxic effects [56]. Due to the increasing consumption of these materials, the aquatic environments are prone to silver contamination. The released AgNPs into the environment can be transformed into more toxic forms such as chloride, nitrate, etc. [57]. Among these silver species, silver nitrate is the most toxic due to dissociation into Ag^+ and nitrate interaction with other elements. Therefore, it is crucial to increase our understanding about the toxicity effects of AgNPs [56].

Silver is considered as the second most toxic metal to aquatic organisms after mercury [58]. The toxicity mechanisms of AgNPs and Ag^+ are different but both are fatal to a variety of organisms including bacteria, animals, and plants [37,59]. The cellular

structure and organelles of these organisms are affected by silver species through unfavorable binding interactions [60]. Silver cations can potentially enter cells through metal transporters such as copper ion transporters or transmembrane potassium channels [24]. Moreover, silver species translocation is attributed to variations in toxicity among Ag^+ , AgNPs, and insoluble silver salts [37,61]. The potential pathway for AgNPs released into the environment and its remediation is illustrated in Figure 1.




Figure 1. Potential pathway for AgNPs leakage into the environment and its remediation.

3.1. Effects of AgNPs on Plants

Unsustainable application of nanoparticles and their release into the environment can also negatively affect the economy by reducing the quality of crops and affecting human health. AgNPs released in waste or ground water can end up in plant growth environment. The small size of nanoparticles enables them to enter plant roots and travel short distances cell by cell or long distances, roots to shoots, via xylem cells [62]. AgNPs accumulated in plant tissues can subsequently enter the human body via the food chain. Numerous studies have reported the toxicological effects of AgNPs on plants (Table 1). The level of toxicity and the impact of AgNPs on plants depend on various factors namely the size, type, and concentration of nanoparticles as well as plant species, soil type, hydraulic conductivity, and the availability of essential nutrients in soil [63,64]. As small as the NPs are, their toxicity is still significant since they can be dissolved easier in water and enter living organisms [65]. As mentioned above, AgNPs have a tendency to be oxidized in the environment and form the ionic form of silver (Ag^+). Therefore, plants exposed to AgNPs are also affected by Ag^+ that is more interactive and toxic than the particulate form. Noori, et al. [21] detected both the ionic and particulate forms of silver in leaves of tomatoes (*Lycopersicon esculentum* L.) exposed to 10, 20, 30 mg L^{-1} of 20 nm AgNPs for 7 days via roots. It confirms that AgNPs can be directly taken up by plants and be stored in tissues in the form of particulates. The presence of AgNPs or Ag^+ in cells affects cellular components, metabolism, and plant growth at the morphological, molecular, and physiological level. At the morphological level, AgNPs affect seed germination, root devel-

opment, and cellular compartments [66,67]. The study by Qian et al. [68] showed that the thylakoid membrane and the structure of chloroplasts were affected in *A. thaliana* exposed to AgNPs. This impacts photosynthesis, metabolism, and plant growth rate. The study on *Oryza sativa* seedlings exposed to 0.5–1 mg L⁻¹ of AgNPs showed significant lowering of mitochondrial membrane potential [69]. A significant decrease in the photosynthetic rate, CO₂ assimilation, and plant growth is also reported in tomatoes exposed to 10 mg/kg of 7–14 nm AgNPs for 72 weeks [70]. Plant responses at the physiological level are modified by the expression of genes and proteins. Kaveh, et al. [71] reported the effects of AgNPs on the expression of over 300 genes. They reported the upregulation of 286 genes and downregulation of 81 genes in *Arabidopsis thaliana* seedlings exposed to up to 20 mg/L of 20 nm AgNPs for 10 days. Proteomics studies also show that AgNPs affect the expression of proteins involved in signal transduction, defense, and oxidative stress responses [72,73]. It implies that the effect of AgNPs on plants is related to the production of ROS and increased production of antioxidant enzymes which further affect the decrease in shoot and root growth and other pigments. This decreases photosynthesis, plant biomass and crop productivity.

Table 1. The effects of AgNPs on plants.

Species	Size (Diameter in nm)	Concentration mg L ⁻¹	Impacts	References
<i>Solanum tuberosum</i> L.	20	2, 10, 20	Increase in superoxide anion (O ₂ ⁻) and reactive oxygen species (ROS); Significant induction in the activities of superoxide dismutase (SOD), ascorbate peroxidase (APX), glutathione reductase (GR), and catalase (CAT)	[74]
<i>Oryza sativa</i> L.	20	0.2, 0.5, 1	Significant decrease in fresh weights, root elongation, shoot and root, carotenoids contents, total chlorophyll; Increased level of hydrogen peroxide (H ₂ O ₂) and lipid peroxidation (MDA) in shoots and roots, increased foliar proline accumulation, and decreased sugar contents	[69]
<i>Wolffia globosa</i>	10	1, 2, 5, 8, 10	Elevated level of malondialdehyde (MDA), ROS content, and SOD activity. Decrease in chlorophyll a, carotenoids, and soluble protein	[75]
<i>Lycopersicon esculentum</i>	10–15	100, 1000	Reduced in fruits productivity, significant decreases in root growth, chlorophyll contents, and increased activity of higher SOD content	[76]
<i>Lycopersicon esculentum</i>	20	10, 20, 30	Decrease in biomass, increased oxidative stress indicators content (H ₂ O ₂ and MDA), induced antioxidative stress responses such as flavonoids, anthocyanins, CAT, peroxidase (POD), reduced chlorophyll content, upregulation of the expression of membrane transporters and xylem cells enlargement	[24]
<i>Lycopersicon esculentum</i>	7–14	10	Lower fruit production, induction of oxidative stress, decrease in photosynthetic rate, elevated activity of antioxidant enzymes	[70]
<i>Arabidopsis thaliana</i>	10	0.2, 0.5, 3	Root growth inhibition, decreased in chlorophyll content and disruption of the thylakoid membrane structure; alteration of transcription of antioxidant and aquaporin channels	[68]
<i>Arabidopsis thaliana</i>	20	10–150	Inhibition of <i>Arabidopsis</i> root gravitropism, lower auxin accumulation in root tips, also, downregulation of the expression of auxin receptor-related genes	[77]

Table 1. Cont.

Species	Size (Diameter in nm)	Concentration mg L ⁻¹	Impacts	References
<i>Arabidopsis thaliana</i> <i>Populus deltoids</i> × <i>nigra</i> , DN-34	5, 10, 25	0.01–100	Toxicity of AgNPs increased with decreasing nanoparticles size; however, the stimulatory effect on fresh weight, evapotranspiration, and root elongation at sublethal concentrations	[78]
<i>Arabidopsis thaliana</i>	10	12.5	delay in flowering, decrease in petal development and vegetative growth, and pollen viability, downregulation of genes involved in floral development	[79]
<i>Lemna minor</i>	10–80	0.005–0.04	number of fronds decreased, growth reduction, chlorosis in leaves	[80]
<i>Cymodocea nodosa</i>	35	0.0002–0.2	length of leaves decreased, induction of oxidative stress indicator (H ₂ O ₂), and antioxidative enzymes activity, less actin and tubulin filaments	[81]
<i>Zea mays</i> (seedlings)	49	7.5	Inhibition of root and leaf growth, induction of O ₂ ⁻ , H ₂ O ₂ , and MDA, increased activity of antioxidative enzymes (SOD, GR, APX),	[82]
<i>Vicia faba</i>	25, 50, 75	100	Size dependent growth decrease, leaf necrosis and damage, and stomatal conductivity decrease	[83]

In addition, several studies have reported a significant increase in the concentration of hydrogen peroxide (H₂O₂), hydroxyl radical (^oOH), and malondialdehyde (MDA) as oxidative stress indicators upon exposure to AgNPs [81,84]. It means exposure to AgNPs can result in oxidative stress in plants. Oxidative stress increases ROS in plants under various environmental stresses, which is then countered by enzymatic and non-enzymatic molecules [85–87]. The level of oxidative stress and plant's ability to cope with NPs varies based on plant species, the size of nanoparticles, and environmental conditions [64,65,80]. Plants that can tolerate NPs in their environment have higher potential to synthesize antioxidants. Antioxidative metabolites (e.g., flavonoids, anthocyanins, or phenols) and enzymes (e.g., catalase, peroxidase, and superoxide dismutase) improve plants ability to reduce the oxidative stress induced by AgNPs.

Despite the extensive research on the toxicity effects of AgNPs on plants, there are reports on induced seed germination and plant growth upon exposure to AgNPs [88,89]. This could be due to AgNPs' role in improving water and nutrient uptake [90]. It is suggested that high concentration of NPs in the environment results in the aggregation of particles and creates larger particles (>100 nm). The aggregated particles do not easily enter plant cells and are less toxic than smaller particles [91]. On the other hand, AgNPs, as a type of oxidative stressor, can induce plant antioxidative and defense responses and improve plant tolerance in stressed condition. Kruszka et al. [92] reported upregulation of secondary metabolites involved in defense responses in *A. thaliana* exposed to 0.5–5 mg L⁻¹ AgNPs. They suggested that AgNPs have a role in improving the bioavailability of nitrogen. Elevated level of antioxidant enzymes and metabolites induce plant's ability to tolerate other environmental stresses as well. Khan et al. [93] reported that exposure to 30 mM AgNPs induced antioxidative responses of *Pennisetum glaucum* L. exposed to 150 mM NaCl. It suggests that plants with potential to tolerate AgNPs in their environment can be considered for phytotechnology approaches to simultaneously remove AgNPs and other contaminants from the environment. In summary, the effects of AgNPs on plants can be either induction or inhibition of growth, development, or defense responses based on various factors such as the type, size, concentration of AgNPs, as well as plant species and the period of exposure. The plant microbe interaction and the effects of AgNPs on microorganisms should be taken into consideration in phytotechnology approaches.

3.2. The Effects of AgNPs on Soil Microorganisms

The antimicrobial activities of AgNPs affect soil-borne microorganisms including pathogenic and beneficial bacteria and fungi [94,95]. The release of AgNPs in soil decreases soil microorganisms' population [96,97] and results in higher toxicity in soil. In addition to the effects on the soil microbial community, AgNPs also affect the activity of soil microorganisms [27,98]. AgNPs can both induce or inhibit enzymatic activities of soil microbes [99,100] depending on the type of the microorganisms, soil texture, osmotic potential, size, shape, and the concentration of AgNPs [101]. Jain et al. [102] reported that the enzymatic activity of decomposing bacteria and fungi depends on the concentration of NPs. They reported that exposure to 2.5 μM AgNPs induced the enzymatic activity of several fungi and bacteria, while 25 and 50 μM exposure had the opposite effect. Tripathi, et al. [103] summarized the effects of AgNPs on soil microorganisms. They highlighted that AgNPs and Ag^+ affect cell membrane integrity and enter the bacterial cell by disrupting the structure of the cell membrane. The impact on the membrane is the key in the antibacterial properties of AgNPs [104]. Exposure to AgNPs also interrupts the activity of membrane transporters and the transportation of essential elements in cells such as potassium and chloride. AgNPs and Ag^+ entered bacterial cells interfere with DNA replication [105] and bacterial growth. Based on recent studies, it is suggested that the antibacterial properties of AgNPs are mostly related to their impacts on the structure of membrane and cellular division. Similar to plants and bacteria, other studies on the interaction of AgNPs and fungi suggest that the size of AgNPs play a vital role in this interaction [25,103,106]. The presence of chitin gives a semipermeable structure to the fungal cell wall and reduces the penetration of larger size AgNPs into fungal cells [107]. In addition to soil bacteria and fungi, AgNPs can also affect decomposers. The study on plant decomposing invertebrate *Limnephilus* sp. showed that the leaf shredding behavior of this organism is affected by the size and concentration of AgNPs [108]. Invertebrates play a vital role in soil structure and are important in inducing plant defense responses. It is important to note that changes in soil microbial population and activity affect soil properties and subsequently plant growth, physiological, and molecular responses. These impact the plant's potential to remove nanoparticles from the environment. Table 2 lists the impacts of AgNPs on soil microorganisms.

Table 2. Overview on phytotoxicity functions of AgNPs on soil species.

Species	Size	Concentration	Major Functions	References
<i>Escherichia coli</i> and Nitrifying bacteria	16 nm	0.1–1 mg L ⁻¹	AgNPs inhibit respiration and nitrification (lack of change in dissolved oxygen); the effect varies depending on the size and bioavailability of the NPs.	[109]
Ammonia-oxidizing bacteria (AOB)	118 ± 11 nm	0.5–50 mg L ⁻¹	Silver treatments affect ammonia-oxidizing bacteria (AOB), reducing Nitrification potential rates.	[110]
<i>Bacillus subtilis</i>	27 nm	0.5–50 mg L ⁻¹	The growth of <i>B. subtilis</i> is affected by AgNPs depending on the size of the AgNPs which probably plays a role in toxicity differences.	[110]
<i>Nitrosomonas europaea</i>	AgNPs (35 nm)/AgNO ₃	0.075–0.75 mg L ⁻¹	The oxidation and production of NO ₂ by regulating the gene expression (nitric oxide reductase, ammonia monooxygenase, and nitrite reductase).	[111]
<i>Nitrosomonas europaea</i>	AgNPs (25.5 nm)/Ag ⁺	0.1 mL/h of solution, 0.05–2 ppm Ag ⁺ , 1.5–20 ppm AgNPs	Low concentrations of AgNPs cause enzymatic inhibition (of ammonia monooxygenase enzyme) and high concentrations cause cell death.	[112]
Soil microbial community	BAM-N001 (20 nm)/AgNO ₃	0.01 mg/kg	The toxicity of AgNPs increases over time (possibly due to Ag ⁺ release). AgNPs cause a decrease in biomass and the activity of soil microorganisms.	[97]

Table 2. Cont.

Species	Size	Concentration	Major Functions	References
Soil bacterial phyla	BAM-N001 (20 nm)/AgNO ₃	0.01 mg/kg	Long-term exposure to AgNPs cause a decrease in several phyla of bacteria, affecting important functions of soil like nitrification or organic carbon transformation.	[113]
Bacterial and fungal assemblages	AgNPs (20 nm)/AgMPs (3000 nm)	0.066% and 6.6%	Small particles are more toxic (cause a decrease in respiration, signature bacterial fatty acids, changes in richness and evenness in bacterial and fungal DNA sequence assemblages).	[114]
Heterotrophic bacterial and nitrifying communities	NM-300K (15 nm)	1.67 and 5 mg/kg supplied in one or three applications	Single application has a stronger effect on potential nitrification than split doses (i.e., same dose applied in 3 doses), whereas of respiration an opposite pattern is observed.	[115]
<i>Azotobacter vinelandii</i>	10 and 50 nm	2 and 10 mg L ⁻¹ for nano-Ag 10; 10 and 100 mg L ⁻¹ for nano-Ag 50	10 and 50 nm AgNPs induced apoptosis by 20.23% and 3.14%, reduced cell number, structural damage, inhibition of biological nitrogen fixation (BNF), ROS generation	[116]
<i>Nitrosomonas europaea</i> ATCC-19718	7 ± 3 (PVA doped) 40 ± 14 (Na ₂ ATP doped)	1, 5 and 10 mg L ⁻¹	Capping and size dependent decrease in NH ₃ oxidation, cell wall damage, and disintegrated nuclei	[117]
Ammonia oxidizing microorganisms	15 nm	1, 10, and 100 µg g ⁻¹ dry soil	10 and 100 µg g ⁻¹ AgNPs significantly inhibited soil urease activity and nitrification	[118]
Soil microbial activity	20.4 and 10 nm	0.1, 1, and 10 mg kg ⁻¹ soil	Decreased soil microbial metabolic activity, nitrification ability, and the abundances of ammonia-oxidizing bacteria at 0.1–10 mg kg ⁻¹ AgNPs	[119]
Mycorrhizal clover (<i>Trifolium repens</i>)	20.6 ± 3.1 (AgNPs)	0.01–1 mg kg ⁻¹	Drastic decrease in biomass of mycorrhizal clover, root nutrient acquisition of AMF, and glomalin content	[120]
<i>Glomus aggregatum</i> -Faba bean	5–50	800 µg kg ⁻¹ sandy soil-loam mixture	Lowered mycorrhizal colonization, glomalin content, and mycorrhizal responsiveness	[121]
AMF (unspecified)-Tomato	2 and 15 nm	12–36 mg kg ⁻¹ soil	Dose dependent AgNPs decrease in AMF colonization	[25]
Soil microbial activity	20 nm ± 10	50 mg kg ⁻¹	Decrease in urease and dehydrogenase activity, bacterial and archaeal amoA gene abundance in soil	[122]
Soil microbial activity	2–50 nm (average 35 nm)	550 mg/pot	Pyrosequence analysis showed no significant effect on soil microbial richness; however, individual analysis affected bacterial groups	[123]
Soil microbial activity	15–20 nm	220 mg kg ⁻¹	Decrease in C and N biomass and modification of microbial community structure	[124]
Soil microbial activity	21 ± 17 nm	0.14 mg kg ⁻¹	AgNPs caused a modification in the bacterial community	[125]
Soil microbial activity	10 and 50 nm	1600, or 3200 µg Ag kg ⁻¹ dry soil	Decreases in enzymatic activities due to AgNPs	[126]
Soil microbial activity	20.08 nm ± 2.24	1–1000 mg kg ⁻¹	AgNPs induced effects on enzymes	[99]

The consequences of the release of AgNPs in the environment and their toxicological effects on organisms in their vicinity provokes the need to consider sustainable approaches to reduce their harmful effects on living organisms. Phytotechnology, the application of plants in removing or sequestering pollutants, is a promising and environmentally friendly method that can be applied in environments contaminated with NPs.

4. Nanoparticle's Phytotechnology

Many soil remediation technologies (physical and chemical methods) have been established to reduce soil contamination [127,128]. However, most of these methods are laborious, time-consuming, and costly. Therefore, such technologies are not best for combating environmental pollution and remediation strategies for a long period [127]. Phytoremediation is an alternate technology applied for the remediation of pollutants using plants [129]. Plants being indispensable elements of ecosystems play a crucial role in the uptake, accumulation, and transport of elements including metal-based nanoparticles [130].

Phytotechnology refers to a technology that uses plants to remove, uptake, absorb, transform, transfer, attenuate, accumulate, degrade, or metabolite organic, inorganic, metallic, or metalloid contaminants from soil, water, or air [131–135]. Phytotechnology consists of several sub-methods such as rhizosphere biodegradation, phytoextraction, phytosequestration, phytovolatilization, phytodegradation, or phytoremediation. Although the phrase phytotechnology is used interchangeably with phytoremediation, the latter generally refers to a method of phytotechnology that removes pollutants from the environment. Phytoremediation is a natural, simple, cost effective, and widespread bioremediation technology that works on the principle of plant's metabolic system to clean, recover, and remediate contaminated sites by storing pollutants in plant biomass to ensure environmental safety [129,136]. Phytoremediation strategies [136] are namely (a) phytostabilization—when plants reduce bioavailability of pollutants in soil (b) phytovolatilization—when pollutants are converted as volatile compounds by plants (c) phytoextraction—when plants take out pollutants from soil (d) phytofiltration—when cultured plants absorb pollutants from water or waste materials. There are many advantages of such phytoremediation strategies, viz. phytostabilization temporarily counter the hazardous materials as compared to phytoextraction which is a permanent one, phytovolatilization converts to gaseous compounds and is more suitable [136].

In comparison to other methods, phytoremediation provides aesthetic appearance, less destruction, and high public acceptance [137–141]. Since the 1980s, phytoremediation has been significantly studied, practiced, and used in field studies at contaminated sites with heavy metals, metalloids, radionuclides, oil spills, fertilizers, pesticides, chlorinated solvents, and explosives [142,143]. In Table 3, we have showed some examples of AgNPs phytoremediation description and its effects.

Table 3. Phytoremediation of AgNPs.

Plant	Type of NPs	Description	Effect	Reference
<i>Phragmites australis</i>	AgNPs	It accumulated silver only in roots and in leaves and stems there was no metal accumulation	Phytoremediation in estuarine areas, Phytostabilization	[49]
<i>Pistia stratiotes</i>	AgNPs	Extracted silver from source in short period of time, Easy handling and can be used in polyculture	Phytoremediation of water source	[51]
<i>Egeria densa</i>	AgNPs	Plants absorbed AgNPs at concentrations as low as 5 ppm, bioaccumulation proportional to concentration of NPs	Phytoremediation of water source	[56]
<i>Phanerochaete chrysosporium</i>	AgNPs	Uptake of NPs in presence of cysteine amino acids	Phytoremediation of aquatic source	[144]
<i>Zea mays</i>	AgNPs	NPs increase the bioremediation potential of three PGPRs isolated from municipal wastewater	Phytoremediation of municipal wastewater	[134]
<i>Ipomoea carnea</i> , <i>Plantago major</i> , <i>Camellia sinensis</i>	AgNPs	Green synthesized NPs from few medicinal plants displayed bioremediation potential	Fipronil (an insecticide) contaminated water	[145]
<i>Lagerstroemia speciosa</i>	AgNPs	methyl orange and methylene blue showing 310- and 290-min degradation time, respectively	methyl orange and methylene blue dyes	[146]
<i>Aloe barbedensis</i> , <i>Azadirachta indica</i> and <i>Coriandrum sativum</i>	AgNPs and CuNPs	Green synthesized NPs from three plants decontaminated naphthalene in water	Wastewater remediation	[147]

Plant remediation technologies are recommended as a viable option for maintaining environmental sustainability among the various options for repairing these pollutants [148]. Recent advances in phytoremediation, molecular and metabolic engineering and nanotechnology have opened up new avenues for the effective handling of emerging organic and inorganic contaminants [149–151]. Plants with high potential to take up contaminants and store them in their tissues are considered as “hyperaccumulators”. These hyperaccumulators accumulate metals or metalloids at a greater level than other plants. These hyperaccumulators are found in metalliferous soils rich in any particular metals [152]. Some examples are, *Thlaspi caerulescens* [153], *Arabidopsis halleri* [154], *Pteris vittata* [155] *Pteris vittata* accumulates arsenic and three others namely *Thlaspi rotundifolium*, *Thlaspi ochroleucum*, *Thlaspi goesingense* accumulates zinc, lead and nickel, respectively [156]. There is a freely available global database containing information about hyperaccumulators (www.hyperaccumulators.org, accessed on 20 July 2021) [152].

These types of plants are favorable in phytoremediation studies especially in sites contaminated with heavy metals. In addition to heavy metals, most hyperaccumulators are also able to tolerate pollutants such as petroleum-based contaminants, explosives, or pesticides that contain a variety of heavy metals and nanoparticles [157]. Among different types of plants used in phytoremediation techniques, hyperaccumulators are best used in phytoextraction projects. AgNPs are metal based nanoparticles that can be extracted from the environment and be taken up by hyperaccumulators. To successfully implant a phytoremediation project, understanding the mechanism of uptake and translocation of contaminants, in this case metal-based nanomaterials, is required. Many studies by different groups have tried to understand the mechanism of uptake and translocation of metal NPs in plants [4,158], where the authors have indicated the size and chemical properties including zeta potential to be an important factor for initial penetration of NPs to the plant. Once the NPs enter the plants they may move through endocytosis or through symplastic transport to different plant tissues [158]. Understanding the molecular mechanism of hyperaccumulators enables scientists to use genetic engineering approaches to strengthen the remediation of environmental pollutants [159] including metal-based nanoparticles. Biotechnology is used to manipulate the expression of genes to improve hyperaccumulators potential to remove contaminants from the environment. In addition to hyperaccumulators, some plants that do not usually have the potential to remove contaminants can alter their physiology and biochemistry to release metabolites and hormones into their environment to sequester nanomaterials and reduce their reactivity in the environment [160]. This method is known as phytosequestration. To apply phytotechnology approaches in removing or sequestering NPs from the environment, several factors such as the phytotoxicity effects of NPs, plant species, weathering, soil structure and organic matter, soil microorganisms, and interaction with other chemicals should be considered [161]. The interaction of engineered NPs with soil organic matter can result in a more stable form of NPs and subsequently result in the release of the ionic form of NPs that are more interactive than the particulate form and have a higher chance to enter living organisms [161]. In addition, the hydrophobic properties of engineered NPs affect their interaction with organic matter and living organisms facilitating their uptake by plants [162]. In addition to above-mentioned factors, the chemical properties of NPs, their concentration, the ratio between the ionic and the particulate form of metal-based nanomaterials impact the phytotoxicity effects of engineered NPs and plants, potentially removing them from the environment. The study on the potential of salt marsh plant, *Phragmites australis*, in removing AgNPs and Ag⁺ showed that both AgNPs and Ag⁺ were accumulated in plant roots [49]. They emphasized that AgNPs interaction with the soil microbial community can interfere with the phytoremediation of Ag in either particulate or ionic form. Yang, et al. [163] also reported the accumulation of Ag in roots of rice exposed to up to 20 mg L⁻¹ AgNPs and AgNO₃ for five days. This study showed that uptake of AgNPs was more efficient than Ag⁺. They suggested that the high reactivity of Ag⁺ in the growth environment and generation of AgCl resulted in its lower uptake rate compared with the particulate form of Ag. The researcher has

also found the accumulation of AgNPs in *Lemna gibba* after its exposure to AgNPs at a concentration of 0.01 to 10 mg L⁻¹ for 7 days [164]. The authors have observed an accumulation from 7.7 to 17.5 µg/mg of AgNPs/plant dry mass. Considering the characteristics of engineered nanoparticles, it is suggested that phytoextraction and phytosequestration methods are promising approaches of phytoremediation in removing AgNPs from the environment [51,165]. The toxicity of AgNPs to plants and associated microorganisms is an important factor which is needed to be considered while the selection of plants for phytoremediation purposes. Along with the phytotoxicity effects of AgNPs and plant's potential to remove them from the environment, various environmental and biological factors should be also considered in selecting phytotechnology as an environmentally friendly method to remove these metal-based NPs from the environment. Due to the colloidal and dynamic properties of AgNPs, rhizosphere biodegradation can prove to be an important method where the plants release certain compounds which can lead to an enhancement in microbial activity resulting in the degradation of NPs. For example, *Bacillus subtilis* can colonize the rhizosphere and get nutrition from plants. In return, it provides several benefits to plants discussed in Hashem et al. [166]. It has been reported that during wastewater treatment, the AgNPs can be partially or fully sulfidized which results in much less toxic compounds than AgNPs in its original form [167]. The authors achieved a full sulfidization with the application of *B. subtilis* indicating an important role of *B. subtilis* in the processing of AgNPs to its less toxic form. Still, the AgNPs contamination has not been identified as an immediate threat to agriculture and only a few studies have been performed which are discussed above. But the increasing use of AgNPs in day-to-day life is making it a potential threat in the near future. Thus, more investigation is needed to better use the plants as a tool to extract/destroy the AgNPs from the agricultural field.

5. Conclusions

This review provides invaluable information about the consequences of the release of nanomaterials in the environment. Nanomaterials, especially AgNPs affect soil properties, microorganisms, and plants and can therefore cause toxicity for living organisms including humans. Although phytoextraction is suggested as a promising approach in removing metal-based nanomaterials, several factors such as the size, concentration, and type of AgNPs as well as soil structure, soil microbial community, and plant species should be considered to select the appropriate method of phytoremediation. In addition, molecular and physiological analysis that improves understanding about the involved transporters and metabolites in removing AgNPs from the environment will help scientists to improve the success of phytoremediation of these materials by altering the expression of related genes using plant biotechnology methods.

Despite the growth in biotechnology, phytotechnology, and nanotechnology, there is still very scarce information regarding sustainable removal of toxic nanoparticles from the environment. Further research is recommended to better understand the mechanism of AgNPs uptake by plants. The scientists must evaluate a small modeling kind of approach to see the effects on ecosystem and also 'lab to land' approach where there will be evaluation of small-scale toxicological data and its removal must be correlated at field level. Another area must be related to seeing the molecular changes upon nanoparticle exposure. Few of these efforts will be helpful in impact assessment of nanoparticles on biological organisms.

Author Contributions: M.I. (Muhammad Ihtisham) prepared original draft, writing—review and editing, and revision; A.N. prepared original draft, and revision; S.Y. revision and review and editing overall manuscript; M.S. prepared original draft, and revision; P.K. writing—review and editing, figure preparation; M.B. writing—review and editing, supervision; M.I. (Muhammad Imran) data curation; F.J. data curation; X.Y. data curation, funding acquisition; A.R. conceptualized the work, supervision, validation, formal analysis, arranged funding, preparation of figure, writing—review and editing. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: We thank the editors and anonymous reviewers for their precious time and comments.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Hochella, M.F.; Lower, S.K.; Maurice, P.A.; Penn, R.L.; Sahai, N.; Sparks, D.L.; Twining, B.S. Nanominerals, mineral nanoparticles, and earth systems. *Science* **2008**, *319*, 1631–1635. [CrossRef]
2. Keerthana, P.; Vijayakumar, S.; Vidhya, E.; Punitha, V.; Nilavukkarasi, M.; Praseetha, P. Biogenesis of ZnO nanoparticles for revolutionizing agriculture: A step towards anti-infection and growth promotion in plants. *Ind. Crop. Prod.* **2021**, *170*, 113762.
3. Dhawan, A.; Shanker, R.; Das, M.; Gupta, K.C. Guidance for safe handling of nanomaterials. *J. Biomed. Nanotechnol.* **2011**, *7*, 218–224. [CrossRef] [PubMed]
4. Rastogi, A.; Zivcak, M.; Sytar, O.; Kalaji, H.M.; He, X.; Mbarki, S.; Brestic, M. Impact of metal and metal oxide nanoparticles on plant: A critical review. *Front. Chem.* **2017**, *5*, 78. [CrossRef]
5. Dillon, A.D.; Ghidui, M.J.; Krick, A.L.; Griggs, J.; May, S.J.; Gogotsi, Y.; Barsoum, M.W.; Fafarman, A.T. Highly conductive optical quality solution-processed films of 2D titanium carbide. *Adv. Funct. Mater.* **2016**, *26*, 4162–4168. [CrossRef]
6. Chandra, H.; Patel, D.; Kumari, P.; Jangwan, J.; Yadav, S. Phyto-mediated synthesis of zinc oxide nanoparticles of *Berberis aristata*: Characterization, antioxidant activity and antibacterial activity with special reference to urinary tract pathogens. *Mater. Sci. Eng. C* **2019**, *102*, 212–220. [CrossRef]
7. Chandra, H.; Kumari, P.; Bontempi, E.; Yadav, S. Medicinal plants: Treasure trove for green synthesis of metallic nanoparticles and their biomedical applications. *Biocatal. Agric. Biotechnol.* **2020**, *24*, 101518. [CrossRef]
8. Chinnamuthu, C.; Boopathi, P.M. Nanotechnology and agroecosystem. *Madras Agric. J.* **2009**, *96*, 17–31.
9. Servin, A.; Elmer, W.; Mukherjee, A.; De la Torre-Roche, R.; Hamdi, H.; White, J.C.; Bindraban, P.; Dimkpa, C. A review of the use of engineered nanomaterials to suppress plant disease and enhance crop yield. *J. Nanopart. Res.* **2015**, *17*, 92. [CrossRef]
10. Kah, M.; Kookana, R.S.; Gogos, A.; Bucheli, T.D. A critical evaluation of nanopesticides and nanofertilizers against their conventional analogues. *Nat. Nanotechnol.* **2018**, *13*, 677–684. [CrossRef]
11. Gogos, A.; Knauer, K.; Bucheli, T.D. Nanomaterials in plant protection and fertilization: Current state, foreseen applications, and research priorities. *J. Agric. Food Chem.* **2012**, *60*, 9781–9792. [CrossRef] [PubMed]
12. Servin, A.D.; De la Torre-Roche, R.; Castillo-Michel, H.; Pagano, L.; Hawthorne, J.; Musante, C.; Pignatello, J.; Uchimiya, M.; White, J.C. Exposure of agricultural crops to nanoparticle CeO₂ in biochar-amended soil. *Plant Physiol. Biochem.* **2017**, *110*, 147–157. [CrossRef] [PubMed]
13. Singh, R.P.; Handa, R.; Manchanda, G. Nanoparticles in sustainable agriculture: An emerging opportunity. *J. Control. Release* **2020**, *329*, 1234–1248. [CrossRef] [PubMed]
14. Samal, A.K.; Polavarapu, L.; Rodal-Cedeira, S.; Liz-Marzaán, L.M.; Peérez-Juste, J.; Pastoriza-Santos, I. Size Tunable Au@ Ag core-shell nanoparticles: Synthesis and surface-enhanced raman scattering properties. *Langmuir* **2013**, *29*, 15076–15082. [CrossRef] [PubMed]
15. Simonin, M.; Martins, J.M.; Uzu, G.; Spadini, L.; Navel, A.; Richaume, A. Low mobility of CuO and TiO₂ nanoparticles in agricultural soils of contrasting texture and organic matter content. *Sci. Total Environ.* **2021**, *783*, 146952. [CrossRef] [PubMed]
16. Fang, J.; Zhang, K.; Sun, P.; Lin, D.; Shen, B.; Luo, Y. Co-transport of Pb²⁺ and TiO₂ nanoparticles in repacked homogeneous soil columns under saturation condition: Effect of ionic strength and fulvic acid. *Sci. Total Environ.* **2016**, *571*, 471–478. [CrossRef]
17. Courtois, P.; de Vaufléury, A.; Grosser, A.; Lors, C.; Vandembulcke, F. Transfer of sulfidized silver from silver nanoparticles, in sewage sludge, to plants and primary consumers in agricultural soil environment. *Sci. Total Environ.* **2021**, *777*, 145900. [CrossRef] [PubMed]
18. Torrent, L.; Marguí, E.; Queralt, I.; Hidalgo, M.; Iglesias, M. Interaction of silver nanoparticles with mediterranean agricultural soils: Lab-controlled adsorption and desorption studies. *J. Environ. Sci.* **2019**, *83*, 205–216. [CrossRef] [PubMed]
19. De Leersnyder, I.; Rijckaert, H.; De Gelder, L.; Van Driessche, I.; Vermeir, P. High variability in silver particle characteristics, silver concentrations, and production batches of commercially available products indicates the need for a more rigorous approach. *Nanomaterials* **2020**, *10*, 1394. [CrossRef] [PubMed]
20. Market Watch. Available online: <https://www.marketwatch.com/press-release/silvernanoparticles-market-future-scope-demands-and-projectedindustry-growth-to-2024-2019-05-09> (accessed on 20 October 2019).
21. Noori, A.; Ngo, A.; Gutierrez, P.; Theberge, S.; White, J.C. Silver nanoparticle detection and accumulation in tomato. *J. Nanopart. Res.* **2020**, *22*, 131. [CrossRef]
22. Wahab, M.A.; Li, L.; Li, H.; Abdala, A. Silver Nanoparticle-Based Nanocomposites for Combating Infectious Pathogens: Recent Advances and Future Prospects. *Nanomaterials* **2021**, *11*, 581. [CrossRef]
23. Peharec Štefanić, P.; Košpić, K.; Lyons, D.M.; Jurković, L.; Balen, B.; Tkalec, M. Phytotoxicity of silver nanoparticles on tobacco plants: Evaluation of coating effects on photosynthetic performance and chloroplast ultrastructure. *Nanomaterials* **2021**, *11*, 744. [CrossRef] [PubMed]
24. Noori, A.; Donnelly, T.; Colbert, J.; Cai, W.; Newman, L.A.; White, J.C. Exposure of tomato (*Lycopersicon esculentum*) to silver nanoparticles and silver nitrate: Physiological and molecular response. *Int. J. Phytoremediat.* **2020**, *22*, 40–51. [CrossRef]

25. Noori, A.; White, J.C.; Newman, L.A. Mycorrhizal fungi influence on silver uptake and membrane protein gene expression following silver nanoparticle exposure. *J. Nanopart. Res.* **2017**, *19*, 66. [[CrossRef](#)]
26. Rajput, V.D.; Minkina, T.; Sushkova, S.; Tsitsuashvili, V.; Mandzhieva, S.; Gorovtsov, A.; Nevidomskyaya, D.; Gromakova, N. Effect of nanoparticles on crops and soil microbial communities. *J. Soils Sediments* **2018**, *18*, 2179–2187. [[CrossRef](#)]
27. Courtois, P.; Rorat, A.; Lemiere, S.; Guyoneaud, R.; Attard, E.; Levard, C.; Vandebulcke, F. Ecotoxicology of silver nanoparticles and their derivatives introduced in soil with or without sewage sludge: A review of effects on microorganisms, plants and animals. *Environ. Pollut.* **2019**, *253*, 578–598. [[CrossRef](#)] [[PubMed](#)]
28. Ali, S.M.; Yousef, N.M.; Nafady, N.A. Application of biosynthesized silver nanoparticles for the control of land snail *Eobania vermiculata* and some plant pathogenic fungi. *J. Nanomater.* **2015**, *2015*, 218904. [[CrossRef](#)]
29. Capek, I. Preparation of metal nanoparticles in water-in-oil (w/o) microemulsions. *Adv. Colloid Interface Sci.* **2004**, *110*, 49–74. [[CrossRef](#)]
30. Reidy, B.; Haase, A.; Luch, A.; Dawson, K.A.; Lynch, I. Mechanisms of silver nanoparticle release, transformation and toxicity: A critical review of current knowledge and recommendations for future studies and applications. *Materials* **2013**, *6*, 2295–2350. [[CrossRef](#)]
31. Dobias, J.; Bernier-Latmani, R. Silver release from silver nanoparticles in natural waters. *Environ. Sci. Technol.* **2013**, *47*, 4140–4146. [[CrossRef](#)]
32. Burić, P.; Jakšić, Ž.; Štajner, L.; Sikirić, M.D.; Jurašin, D.; Cascio, C.; Calzolari, L.; Lyons, D.M. Effect of silver nanoparticles on Mediterranean sea urchin embryonal development is species specific and depends on moment of first exposure. *Mar. Environ. Res.* **2015**, *111*, 50–59. [[CrossRef](#)] [[PubMed](#)]
33. Miao, A.-J.; Luo, Z.; Chen, C.-S.; Chin, W.-C.; Santschi, P.H.; Quigg, A. Intracellular uptake: A possible mechanism for silver engineered nanoparticle toxicity to a freshwater alga *Ochromonas danica*. *PLoS ONE* **2010**, *5*, e15196. [[CrossRef](#)]
34. Siripattanakul-Ratpukdi, S.; Ploychankul, C.; Limpiyakorn, T.; Vangnai, A.S.; Rongsayamanont, C.; Khan, E. Mitigation of nitrification inhibition by silver nanoparticles using cell entrapment technique. *J. Nanopart. Res.* **2014**, *16*, 2218. [[CrossRef](#)]
35. Vance, M.E.; Kuiken, T.; Vejerano, E.P.; McGinnis, S.P.; Hochella, M.F., Jr.; Rejeski, D.; Hull, M.S. Nanotechnology in the real world: Redeveloping the nanomaterial consumer products inventory. *Beilstein J. Nanotechnol.* **2015**, *6*, 1769–1780. [[CrossRef](#)] [[PubMed](#)]
36. Ivask, A.; Kurvet, I.; Kasemets, K.; Blinova, I.; Aruoja, V.; Suppi, S.; Vija, H.; Käkinen, A.; Titma, T.; Heinlaan, M. Size-dependent toxicity of silver nanoparticles to bacteria, yeast, algae, crustaceans and mammalian cells in vitro. *PLoS ONE* **2014**, *9*, e102108. [[CrossRef](#)]
37. Fabrega, J.; Luoma, S.N.; Tyler, C.R.; Galloway, T.S.; Lead, J.R. Silver nanoparticles: Behaviour and effects in the aquatic environment. *Environ. Int.* **2011**, *37*, 517–531. [[CrossRef](#)]
38. Baker, T.J.; Tyler, C.R.; Galloway, T.S. Impacts of metal and metal oxide nanoparticles on marine organisms. *Environ. Pollut.* **2014**, *186*, 257–271. [[CrossRef](#)]
39. Klitzke, S.; Metreveli, G.; Peters, A.; Schaumann, G.E.; Lang, F. The fate of silver nanoparticles in soil solution—Sorption of solutes and aggregation. *Sci. Total Environ.* **2015**, *535*, 54–60. [[CrossRef](#)]
40. Nam, D.-H.; Lee, B.-C.; Eom, I.-C.; Kim, P.; Yeo, M.-K. Uptake and bioaccumulation of titanium- and silver-nanoparticles in aquatic ecosystems. *Mol. Cell. Toxicol.* **2014**, *10*, 9–17. [[CrossRef](#)]
41. Howe, P.D.; Dobson, S. *Silver and Silver Compounds: Environmental Aspects*; World Health Organization: Geneva, Switzerland, 2002.
42. Galazzi, R.M.; Júnior, C.A.L.; de Lima, T.B.; Gozzo, F.C.; Arruda, M.A.Z. Evaluation of some effects on plant metabolism through proteins and enzymes in transgenic and non-transgenic soybeans after cultivation with silver nanoparticles. *J. Proteom.* **2019**, *191*, 88–106. [[CrossRef](#)] [[PubMed](#)]
43. Benn, T.M.; Westerhoff, P. Nanoparticle silver released into water from commercially available sock fabrics. *Environ. Sci. Technol.* **2008**, *42*, 4133–4139. [[CrossRef](#)] [[PubMed](#)]
44. Lazim, Z.M.; Salmiati; Samaluddin, A.R.; Salim, M.R.; Arman, N.Z. Toxicity of Silver Nanoparticles and Their Removal Applying Phytoremediation System to Water Environment: An Overview. *J. Environ. Treat. Tech.* **2020**, *8*, 978–984.
45. Handy, R.D.; Shaw, B.J. Toxic effects of nanoparticles and nanomaterials: Implications for public health, risk assessment and the public perception of nanotechnology. *Health Risk Soc.* **2007**, *9*, 125–144. [[CrossRef](#)]
46. Owen, R.; Handy, R. *Formulating the Problems for Environmental Risk Assessment of Nanomaterials*; ACS Publications: Washington, DC, USA, 2007; pp. 5582–5588.
47. Pérez-de-Luque, A.; Rubiales, D. Nanotechnology for parasitic plant control. *Pest Manag. Sci. Former. Pestic. Sci.* **2009**, *65*, 540–545. [[CrossRef](#)] [[PubMed](#)]
48. Saharan, V. Advances in nanobiotechnology for agriculture. In *Current Topics in Biotechnology & Microbiology*; Dhingra, H.K., Nath Jha, P., Bajpai, P., Eds.; Lap Lambert Academic Publishing: Dudweiler Landstr, Germany, 2011; pp. 156–167.
49. Fernandes, J.P.; Mucha, A.P.; Francisco, T.; Gomes, C.R.; Almeida, C.M.R. Silver nanoparticles uptake by salt marsh plants—Implications for phytoremediation processes and effects in microbial community dynamics. *Mar. Pollut. Bull.* **2017**, *119*, 176–183. [[CrossRef](#)] [[PubMed](#)]
50. Shafer, M.M.; Overdier, J.T.; Armstrong, D.E. Removal, partitioning, and fate of silver and other metals in wastewater treatment plants and effluent-receiving streams. *Environ. Toxicol. Chem. Int. J.* **1998**, *17*, 630–641. [[CrossRef](#)]

51. Hanks, N.A.; Caruso, J.A.; Zhang, P. Assessing *Pistia stratiotes* for phytoremediation of silver nanoparticles and Ag (I) contaminated waters. *J. Environ. Manag.* **2015**, *164*, 41–45. [[CrossRef](#)] [[PubMed](#)]
52. Purcell, T.W.; Peters, J.J. Historical impacts of environmental regulation of silver. *Environ. Toxicol. Chem. Int. J.* **1999**, *18*, 3–8. [[CrossRef](#)]
53. U.S. Environmental Protection Agency (EPA). National Primary Drinking Water Regulation Table, EPA 816-F-09-0004. 2009. Available online: <https://www.epa.gov/ground-water-and-drinking-water/national-primary-drinking-water-regulation-table> (accessed on 5 August 2021).
54. World Health Organization (WHO). *Guidelines for Drinking-Water Quality*, 4th ed.; World Health Organization (WHO): Geneva, Switzerland, 2011; p. 415. Available online: http://apps.who.int/iris/bitstream/handle/10665/44584/9789241548151_eng.pdf;jsessionid=7602427D0558C27BC51742431A74F67E?sequence=1 (accessed on 5 August 2021).
55. Varner, K.; El-Badawy, A.; Feldhake, D.; Venkatapathy, R. *State-of-the-Science Review: Everything Nanosilver and More*; EPA/600/R-10/084:2010; US Environmental Protection Agency: Washington, DC, USA, 2010.
56. Bernas, L.; Winkelmann, K.; Palmer, A. Phytoremediation of silver species by waterweed (*Egeria densa*). *Chemist* **2017**, *90*, 7–13.
57. Valenti, L.E.; Giacomelli, C.E. Stability of silver nanoparticles: Agglomeration and oxidation in biological relevant conditions. *J. Nanopart. Res.* **2017**, *19*, 156. [[CrossRef](#)]
58. Moreno-Garrido, I.; Pérez, S.; Blasco, J. Toxicity of silver and gold nanoparticles on marine microalgae. *Mar. Environ. Res.* **2015**, *111*, 60–73. [[CrossRef](#)]
59. Lapresta-Fernández, A.; Fernández, A.; Blasco, J. Nanoecotoxicity effects of engineered silver and gold nanoparticles in aquatic organisms. *TrAC Trends Anal. Chem.* **2012**, *32*, 40–59. [[CrossRef](#)]
60. Navarro, E.; Baun, A.; Behra, R.; Hartmann, N.B.; Filser, J.; Miao, A.-J.; Quigg, A.; Santschi, P.H.; Sigg, L. Environmental behavior and ecotoxicity of engineered nanoparticles to algae, plants, and fungi. *Ecotoxicology* **2008**, *17*, 372–386. [[CrossRef](#)] [[PubMed](#)]
61. Ratte, H.T. Bioaccumulation and toxicity of silver compounds: A review. *Environ. Toxicol. Chem. Int. J.* **1999**, *18*, 89–108. [[CrossRef](#)]
62. Geisler-Lee, J.; Brooks, M.; Gerfen, J.R.; Wang, Q.; Fotis, C.; Sparer, A.; Ma, X.; Berg, R.H.; Geisler, M. Reproductive toxicity and life history study of silver nanoparticle effect, uptake and transport in *Arabidopsis thaliana*. *Nanomaterials* **2014**, *4*, 301–318. [[CrossRef](#)]
63. Çekiç, F.Ö.; Ekinçi, S.; İnal, M.S.; Özakça, D. Silver nanoparticles induced genotoxicity and oxidative stress in tomato plants. *Turk. J. Biol.* **2017**, *41*, 700–707. [[CrossRef](#)]
64. Cox, A.; Venkatachalam, P.; Sahi, S.; Sharma, N. Reprint of: Silver and titanium dioxide nanoparticle toxicity in plants: A review of current research. *Plant Physiol. Biochem.* **2017**, *110*, 33–49. [[CrossRef](#)]
65. Gupta, I.R.; Anderson, A.J.; Rai, M. Toxicity of fungal-generated silver nanoparticles to soil-inhabiting *Pseudomonas putida* KT2440, a rhizospheric bacterium responsible for plant protection and bioremediation. *J. Hazard. Mater.* **2015**, *286*, 48–54. [[CrossRef](#)]
66. Yan, A.; Chen, Z. Impacts of silver nanoparticles on plants: A focus on the phytotoxicity and underlying mechanism. *Int. J. Mol. Sci.* **2019**, *20*, 1003. [[CrossRef](#)] [[PubMed](#)]
67. Abbas, Q.; Yousaf, B.; Ullah, H.; Ali, M.U.; Zia-ur-Rehman, M.; Rizwan, M.; Rinklebe, J. Biochar-induced immobilization and transformation of silver-nanoparticles affect growth, intracellular-radicles generation and nutrients assimilation by reducing oxidative stress in maize. *J. Hazard. Mater.* **2020**, *390*, 121976. [[CrossRef](#)]
68. Qian, H.; Peng, X.; Han, X.; Ren, J.; Sun, L.; Fu, Z. Comparison of the toxicity of silver nanoparticles and silver ions on the growth of terrestrial plant model *Arabidopsis thaliana*. *J. Environ. Sci.* **2013**, *25*, 1947–1956. [[CrossRef](#)]
69. Nair, P.M.G.; Chung, I.M. Physiological and molecular level effects of silver nanoparticles exposure in rice (*Oryza sativa* L.) seedlings. *Chemosphere* **2014**, *112*, 105–113. [[CrossRef](#)]
70. Das, P.; Barua, S.; Sarkar, S.; Chatterjee, S.K.; Mukherjee, S.; Goswami, L.; Das, S.; Bhattacharya, S.; Karak, N.; Bhattacharya, S.S. Mechanism of toxicity and transformation of silver nanoparticles: Inclusive assessment in earthworm-microbe-soil-plant system. *Geoderma* **2018**, *314*, 73–84. [[CrossRef](#)]
71. Kaveh, R.; Li, Y.-S.; Ranjbar, S.; Tehrani, R.; Brueck, C.L.; Van Aken, B. Changes in *Arabidopsis thaliana* gene expression in response to silver nanoparticles and silver ions. *Environ. Sci. Technol.* **2013**, *47*, 10637–10644. [[CrossRef](#)]
72. Mirzajani, F.; Askari, H.; Hamzelou, S.; Schober, Y.; Römpf, A.; Ghassempour, A.; Spengler, B. Proteomics study of silver nanoparticles toxicity on *Oryza sativa* L. *Ecotoxicol. Environ. Saf.* **2014**, *108*, 335–339. [[CrossRef](#)] [[PubMed](#)]
73. Peharec Štefanić, P.; Jarnević, M.; Cvjetko, P.; Biba, R.; Šikić, S.; Tkalec, M.; Cindrić, M.; Letofsky-Papst, I.; Balen, B. Comparative proteomic study of phytotoxic effects of silver nanoparticles and silver ions on tobacco plants. *Environ. Sci. Pollut. Res.* **2019**, *26*, 22529–22550. [[CrossRef](#)]
74. Homaei, M.B.; Ehsanpour, A.A. Silver nanoparticles and silver ions: Oxidative stress responses and toxicity in potato (*Solanum tuberosum* L.) grown in vitro. *Hortic. Environ. Biotechnol.* **2016**, *57*, 544–553. [[CrossRef](#)]
75. Zou, X.; Li, P.; Huang, Q.; Zhang, H. The different response mechanisms of *Wolffia globosa*: Light-induced silver nanoparticle toxicity. *Aquat. Toxicol.* **2016**, *176*, 97–105. [[CrossRef](#)]
76. Song, U.; Jun, H.; Waldman, B.; Roh, J.; Kim, Y.; Yi, J.; Lee, E.J. Functional analyses of nanoparticle toxicity: A comparative study of the effects of TiO₂ and Ag on tomatoes (*Lycopersicon esculentum*). *Ecotoxicol. Environ. Saf.* **2013**, *93*, 60–67. [[CrossRef](#)]
77. Sun, J.; Wang, L.; Li, S.; Yin, L.; Huang, J.; Chen, C. Toxicity of silver nanoparticles to *Arabidopsis*: Inhibition of root gravitropism by interfering with auxin pathway. *Environ. Toxicol. Chem.* **2017**, *36*, 2773–2780. [[CrossRef](#)]

78. Wang, J.; Koo, Y.; Alexander, A.; Yang, Y.; Westerhof, S.; Zhang, Q.; Schnoor, J.L.; Colvin, V.L.; Braam, J.; Alvarez, P.J. Phytostimulation of poplars and Arabidopsis exposed to silver nanoparticles and Ag⁺ at sublethal concentrations. *Environ. Sci. Technol.* **2013**, *47*, 5442–5449. [[CrossRef](#)] [[PubMed](#)]
79. Ke, M.; Li, Y.; Qu, Q.; Ye, Y.; Peijnenburg, W.; Zhang, Z.; Xu, N.; Lu, T.; Sun, L.; Qian, H. Offspring toxicity of silver nanoparticles to Arabidopsis thaliana flowering and floral development. *J. Hazard. Mater.* **2020**, *386*, 121975. [[CrossRef](#)]
80. Minogiannis, P.; Valenti, M.; Kati, V.; Kalantzi, O.-I.; Biskos, G. Toxicity of pure silver nanoparticles produced by spark ablation on the aquatic plant Lemna minor. *J. Aerosol Sci.* **2019**, *128*, 17–21. [[CrossRef](#)]
81. Mylona, Z.; Panteris, E.; Moustakas, M.; Kevrekidis, T.; Malea, P. Physiological, structural and ultrastructural impacts of silver nanoparticles on the seagrass *Cymodocea nodosa*. *Chemosphere* **2020**, *248*, 126066. [[CrossRef](#)]
82. Yilmaz, M.; Yilmaz, A.; Karaman, A.; Aysin, F.; Aksakal, O. Monitoring chemically and green-synthesized silver nanoparticles in maize seedlings via surface-enhanced Raman spectroscopy (SERS) and their phytotoxicity evaluation. *Talanta* **2021**, *225*, 121952. [[CrossRef](#)] [[PubMed](#)]
83. Falco, W.F.; Scherer, M.D.; Oliveira, S.L.; Wender, H.; Colbeck, I.; Lawson, T.; Caires, A.R. Phytotoxicity of silver nanoparticles on Vicia faba: Evaluation of particle size effects on photosynthetic performance and leaf gas exchange. *Sci. Total Environ.* **2020**, *701*, 134816. [[CrossRef](#)]
84. Ma, C.; White, J.C.; Dhankher, O.P.; Xing, B. Metal-based nanotoxicity and detoxification pathways in higher plants. *Environ. Sci. Technol.* **2015**, *49*, 7109–7122. [[CrossRef](#)] [[PubMed](#)]
85. Apel, K.; Hirt, H. Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* **2004**, *55*, 373–399. [[CrossRef](#)]
86. Yadav, S.; Kushwaha, H.R.; Kumar, K.; Verma, P.K. Comparative structural modeling of a monothiol GRX from chickpea: Insight in iron–sulfur cluster assembly. *Int. J. Biol. Macromol.* **2012**, *51*, 266–273. [[CrossRef](#)]
87. Kumari, P.; Gupta, A.; Yadav, S. Thioredoxins as olecular Players in Plants, Pests, and Pathogens. In *Plant-Pest Interactions: From Molecular Mechanisms to Chemical Ecology*; Springer: Berlin/Heidelberg, Germany, 2021; pp. 107–125. Available online: https://doi.org/10.1007/978-981-15-2467-7_6 (accessed on 20 July 2021).
88. Abdel-Aziz, H.M.; Rizwan, M. Chemically synthesized silver nanoparticles induced physio-chemical and chloroplast ultrastructural changes in broad bean seedlings. *Chemosphere* **2019**, *235*, 1066–1072. [[CrossRef](#)]
89. Mahakham, W.; Sarmah, A.K.; Maensiri, S.; Theerakulpisut, P. Nanopriming technology for enhancing germination and starch metabolism of aged rice seeds using phytosynthesized silver nanoparticles. *Sci. Rep.* **2017**, *7*, 1–21. [[CrossRef](#)] [[PubMed](#)]
90. Wojtyła, Ł.; Lechowska, K.; Kubala, S.; Garnczarska, M. Molecular processes induced in primed seeds—increasing the potential to stabilize crop yields under drought conditions. *J. Plant Physiol.* **2016**, *203*, 116–126. [[CrossRef](#)] [[PubMed](#)]
91. Clément, L.; Hurel, C.; Marmier, N. Toxicity of TiO₂ nanoparticles to cladocerans, algae, rotifers and plants—effects of size and crystalline structure. *Chemosphere* **2013**, *90*, 1083–1090. [[CrossRef](#)] [[PubMed](#)]
92. Kruszcza, D.; Sawikowska, A.; Selvakesavan, R.K.; Krajewski, P.; Kachlicki, P.; Franklin, G. Silver nanoparticles affect phenolic and phytoalexin composition of Arabidopsis thaliana. *Sci. Total Environ.* **2020**, *716*, 135361. [[CrossRef](#)] [[PubMed](#)]
93. Khan, I.; Raza, M.A.; Awan, S.A.; Shah, G.A.; Rizwan, M.; Ali, B.; Tariq, R.; Hassan, M.J.; Alyemeni, M.N.; Brestic, M. Amelioration of salt induced toxicity in pearl millet by seed priming with silver nanoparticles (AgNPs): The oxidative damage, antioxidant enzymes and ions uptake are major determinants of salt tolerant capacity. *Plant Physiol. Biochem.* **2020**, *156*, 221–232. [[CrossRef](#)]
94. Park, S.; Park, H.H.; Ko, Y.-S.; Lee, S.J.; Le, T.S.; Woo, K.; Ko, G. Disinfection of various bacterial pathogens using novel silver nanoparticle-decorated magnetic hybrid colloids. *Sci. Total Environ.* **2017**, *609*, 289–296. [[CrossRef](#)]
95. Uddin, I.; Ahmad, K.; Khan, A.A.; Kazmi, M.A. Synthesis of silver nanoparticles using *Matricaria recutita* (Babunah) plant extract and its study as mercury ions sensor. *Sens. Bio-Sens. Res.* **2017**, *16*, 62–67. [[CrossRef](#)]
96. Samarajeewa, A.; Velicogna, J.; Princz, J.; Subasinghe, R.; Scroggins, R.; Beaudette, L. Effect of silver nano-particles on soil microbial growth, activity and community diversity in a sandy loam soil. *Environ. Pollut.* **2017**, *220*, 504–513. [[CrossRef](#)]
97. Grün, A.-L.; Straskraba, S.; Schulz, S.; Schloter, M.; Emmerling, C. Long-term effects of environmentally relevant concentrations of silver nanoparticles on microbial biomass, enzyme activity, and functional genes involved in the nitrogen cycle of loamy soil. *J. Environ. Sci.* **2018**, *69*, 12–22. [[CrossRef](#)]
98. Zhang, Z.; Gao, P.; Li, M.; Cheng, J.; Liu, W.; Feng, Y. Influence of Silver nanoparticles on nutrient removal and microbial communities in SBR process after long-term exposure. *Sci. Total Environ.* **2016**, *569*, 234–243. [[CrossRef](#)]
99. Shin, Y.-J.; Kwak, J.I.; An, Y.-J. Evidence for the inhibitory effects of silver nanoparticles on the activities of soil exoenzymes. *Chemosphere* **2012**, *88*, 524–529. [[CrossRef](#)] [[PubMed](#)]
100. Abdulsada, Z.; Kibbee, R.; Örmeci, B.; DeRosa, M.; Princz, J. Impact of anaerobically digested silver and copper oxide nanoparticles in biosolids on soil characteristics and bacterial community. *Chemosphere* **2021**, *263*, 128173. [[CrossRef](#)]
101. Zhai, Y.; Hunting, E.R.; Wouters, M.; Peijnenburg, W.J.; Vijver, M.G. Silver nanoparticles, ions, and shape governing soil microbial functional diversity: Nano shapes micro. *Front. Microbiol.* **2016**, *7*, 1123. [[CrossRef](#)]
102. Jain, A.; Kumar, S.; Seena, S. Can low concentrations of metal oxide and Ag loaded metal oxide nanoparticles pose a risk to stream plant litter microbial decomposers? *Sci. Total Environ.* **2019**, *653*, 930–937. [[CrossRef](#)]
103. Tripathi, D.K.; Tripathi, A.; Singh, S.; Singh, Y.; Vishwakarma, K.; Yadav, G.; Sharma, S.; Singh, V.K.; Mishra, R.K.; Upadhyay, R. Uptake, accumulation and toxicity of silver nanoparticle in autotrophic plants, and heterotrophic microbes: A concentric review. *Front. Microbiol.* **2017**, *8*, 7. [[CrossRef](#)]

104. Bondarenko, O.M.; Sihtmäe, M.; Kuzmičiova, J.; Ragelienė, L.; Kahru, A.; Daugelavičius, R. Plasma membrane is the target of rapid antibacterial action of silver nanoparticles in *Escherichia coli* and *Pseudomonas aeruginosa*. *Int. J. Nanomed.* **2018**, *13*, 6779. [[CrossRef](#)] [[PubMed](#)]
105. Dong, Y.; Zhu, H.; Shen, Y.; Zhang, W.; Zhang, L. Antibacterial activity of silver nanoparticles of different particle size against *Vibrio Natriegens*. *PLoS ONE* **2019**, *14*, e0222322. [[CrossRef](#)] [[PubMed](#)]
106. Judy, J.D.; Kirby, J.K.; Creamer, C.; McLaughlin, M.J.; Fiebiger, C.; Wright, C.; Cavagnaro, T.R.; Bertsch, P.M. Effects of silver sulfide nanomaterials on mycorrhizal colonization of tomato plants and soil microbial communities in biosolid-amended soil. *Environ. Pollut.* **2015**, *206*, 256–263. [[CrossRef](#)]
107. Sosnowska, M.E.; Jankiewicz, U.; Kutwin, M.; Chwalibog, A.; Gałazka, A. Influence of salts and metal nanoparticles on the activity and thermal stability of a recombinant chitinase from *Stenotrophomonas maltophilia* N4. *Enzym. Microb. Technol.* **2018**, *116*, 6–15. [[CrossRef](#)]
108. Batista, D.; Pascoal, C.; Cássio, F. How do physicochemical properties influence the toxicity of silver nanoparticles on freshwater decomposers of plant litter in streams? *Ecotoxicol. Environ. Saf.* **2017**, *140*, 148–155. [[CrossRef](#)] [[PubMed](#)]
109. Choi, O.; Hu, Z. Size dependent and reactive oxygen species related nanosilver toxicity to nitrifying bacteria. *Environ. Sci. Technol.* **2008**, *42*, 4583–4588. [[CrossRef](#)] [[PubMed](#)]
110. Beddow, J.; Stolpe, B.; Cole, P.; Lead, J.R.; Sapp, M.; Lyons, B.P.; Colbeck, I.; Whitby, C. Effects of engineered silver nanoparticles on the growth and activity of ecologically important microbes. *Environ. Microbiol. Rep.* **2014**, *6*, 448–458. [[CrossRef](#)] [[PubMed](#)]
111. Michels, C.; Yang, Y.; Moreira Soares, H.; Alvarez, P.J. Silver nanoparticles temporarily retard NO₂– production without significantly affecting N₂O release by *Nitrosomonas europaea*. *Environ. Toxicol. Chem.* **2015**, *34*, 2231–2235. [[CrossRef](#)] [[PubMed](#)]
112. Barker, L.; Giska, J.; Radniecki, T.; Semprini, L. Effects of short- and long-term exposure of silver nanoparticles and silver ions to *Nitrosomonas europaea* biofilms and planktonic cells. *Chemosphere* **2018**, *206*, 606–614. [[CrossRef](#)]
113. Grün, A.-L.; Emmerling, C. Long-term effects of environmentally relevant concentrations of silver nanoparticles on major soil bacterial phyla of a loamy soil. *Environ. Sci. Eur.* **2018**, *30*, 1–13. [[CrossRef](#)] [[PubMed](#)]
114. Kumar, N.; Palmer, G.R.; Shah, V.; Walker, V.K. The effect of silver nanoparticles on seasonal change in arctic tundra bacterial and fungal assemblages. *PLoS ONE* **2014**, *9*, e99953. [[CrossRef](#)] [[PubMed](#)]
115. Schlich, K.; Beule, L.; Hund-Rinke, K. Single versus repeated applications of CuO and Ag nanomaterials and their effect on soil microflora. *Environ. Pollut.* **2016**, *215*, 322–330. [[CrossRef](#)]
116. Zhang, L.; Wu, L.; Si, Y.; Shu, K. Size-dependent cytotoxicity of silver nanoparticles to *Azotobacter vinelandii*: Growth inhibition, cell injury, oxidative stress and internalization. *PLoS ONE* **2018**, *13*, e0209020. [[CrossRef](#)]
117. Yuan, Z.; Li, J.; Cui, L.; Xu, B.; Zhang, H.; Yu, C.-P. Interaction of silver nanoparticles with pure nitrifying bacteria. *Chemosphere* **2013**, *90*, 1404–1411. [[CrossRef](#)]
118. Huang, J.; Chong, C.; Runqing, L.; Wenzhu, G. Effects of silver nanoparticles on soil ammonia-oxidizing microorganisms under temperatures of 25 and 5 °C. *Pedosphere* **2018**, *28*, 607–616. [[CrossRef](#)]
119. He, S.; Feng, Y.; Ni, J.; Sun, Y.; Xue, L.; Feng, Y.; Yu, Y.; Lin, X.; Yang, L. Different responses of soil microbial metabolic activity to silver and iron oxide nanoparticles. *Chemosphere* **2016**, *147*, 195–202. [[CrossRef](#)]
120. Feng, Y.; Cui, X.; He, S.; Dong, G.; Chen, M.; Wang, J.; Lin, X. The role of metal nanoparticles in influencing arbuscular mycorrhizal fungi effects on plant growth. *Environ. Sci. Technol.* **2013**, *47*, 9496–9504. [[CrossRef](#)]
121. Abd-Alla, M.H.; Nafady, N.A.; Khalaf, D.M. Assessment of silver nanoparticles contamination on faba bean-Rhizobium leguminosarum bv. viciae-Glomus aggregatum symbiosis: Implications for induction of autophagy process in root nodule. *Agric. Ecosyst. Environ.* **2016**, *218*, 163–177. [[CrossRef](#)]
122. McGee, C.; Storey, S.; Clipson, N.; Doyle, E. Soil microbial community responses to contamination with silver, aluminium oxide and silicon dioxide nanoparticles. *Ecotoxicology* **2017**, *26*, 449–458. [[CrossRef](#)] [[PubMed](#)]
123. Shah, V.; Collins, D.; Walker, V.K.; Shah, S. The impact of engineered cobalt, iron, nickel and silver nanoparticles on soil bacterial diversity under field conditions. *Environ. Res. Lett.* **2014**, *9*, 024001. [[CrossRef](#)]
124. Kumar, N.; Shah, V.; Walker, V.K. Influence of a nanoparticle mixture on an arctic soil community. *Environ. Toxicol. Chem.* **2012**, *31*, 131–135. [[CrossRef](#)]
125. Colman, B.P.; Arnaout, C.L.; Anciaux, S.; Gunsch, C.K.; Hochella Jr, M.F.; Kim, B.; Lowry, G.V.; McGill, B.M.; Reinsch, B.C.; Richardson, C.J. Low concentrations of silver nanoparticles in biosolids cause adverse ecosystem responses under realistic field scenario. *PLoS ONE* **2013**, *8*, e57189. [[CrossRef](#)] [[PubMed](#)]
126. Peyrot, C.; Wilkinson, K.J.; Desrosiers, M.; Sauvé, S. Effects of silver nanoparticles on soil enzyme activities with and without added organic matter. *Environ. Toxicol. Chem.* **2014**, *33*, 115–125. [[CrossRef](#)] [[PubMed](#)]
127. Chen, M.; Xu, P.; Zeng, G.; Yang, C.; Huang, D.; Zhang, J. Bioremediation of soils contaminated with polycyclic aromatic hydrocarbons, petroleum, pesticides, chlorophenols and heavy metals by composting: Applications, microbes and future research needs. *Biotechnol. Adv.* **2015**, *33*, 745–755. [[CrossRef](#)]
128. Song, B.; Zeng, G.; Gong, J.; Liang, J.; Xu, P.; Liu, Z.; Zhang, Y.; Zhang, C.; Cheng, M.; Liu, Y. Evaluation methods for assessing effectiveness of in situ remediation of soil and sediment contaminated with organic pollutants and heavy metals. *Environ. Int.* **2017**, *105*, 43–55. [[CrossRef](#)]
129. Kumari, P.; Rastogi, A.; Shukla, A.; Srivastava, S.; Yadav, S. Prospects of genetic engineering utilizing potential genes for regulating arsenic accumulation in plants. *Chemosphere* **2018**, *211*, 397–406. [[CrossRef](#)]

130. Monica, R.C.; Cremonini, R. Nanoparticles and higher plants. *Caryologia* **2009**, *62*, 161–165. [CrossRef]
131. Jin, Y.; Liu, W.; Li, X.-L.; Shen, S.-G.; Liang, S.-X.; Liu, C.; Shan, L. Nano-hydroxyapatite immobilized lead and enhanced plant growth of ryegrass in a contaminated soil. *Ecol. Eng.* **2016**, *95*, 25–29. [CrossRef]
132. Fulekar, M. *Bioremediation Technology: Recent Advances*; Springer Science & Business Media: Berlin/Heidelberg, Germany, 2012.
133. Evangelou, M.W.; Papazoglou, E.G.; Robinson, B.H.; Schulin, R. Phytomanagement: Phytoremediation and the Production of Biomass for Economic Revenue on Contaminated ILand. In *Phytoremediation*; Springer: Berlin/Heidelberg, Germany, 2015; pp. 115–132. Available online: https://doi.org/10.1007/978-3-319-10395-2_9 (accessed on 22 April 2020).
134. Khan, N.; Bano, A. Role of plant growth promoting rhizobacteria and Ag-nano particle in the bioremediation of heavy metals and maize growth under municipal wastewater irrigation. *Int. J. Phytoremediat.* **2016**, *18*, 211–221. [CrossRef] [PubMed]
135. Padmapriya, S.; Murugan, N.; Ragavendran, C.; Thangabalu, R.; Natarajan, D. Phytoremediation potential of some agricultural plants on heavy metal contaminated mine waste soils, salem district, tamilnadu. *Int. J. Phytoremediat.* **2016**, *18*, 288–294. [CrossRef] [PubMed]
136. Yan, A.; Wang, Y.; Tan, S.N.; Mohd Yusof, M.L.; Ghosh, S.; Chen, Z. Phytoremediation: A promising approach for revegetation of heavy metal-polluted land. *Front. Plant Sci.* **2020**, *11*, 359. [CrossRef]
137. Bennicelli, R.; Stępniewska, Z.; Banach, A.; Szajnoch, K.; Ostrowski, J. The ability of *Azolla caroliniana* to remove heavy metals (Hg(II), Cr(III), Cr(VI)) from municipal waste water. *Chemosphere* **2004**, *55*, 141–146. [CrossRef] [PubMed]
138. Elmachli, S.; Chefetz, B.; Tel-Or, E.; Vidal, L.; Canals, A.; Gedanken, A. Removal of silver and lead ions from water wastes using *Azolla filiculoides*, an aquatic plant, which adsorbs and reduces the ions into the corresponding metallic nanoparticles under microwave radiation in 5 min. *Water Air Soil Pollut.* **2011**, *218*, 365–370. [CrossRef]
139. Macek, T.; Macková, M.; Káš, J. Exploitation of plants for the removal of organics in environmental remediation. *Biotechnol. Adv.* **2000**, *18*, 23–34. [CrossRef]
140. Glick, B.R. Phytoremediation: Synergistic use of plants and bacteria to clean up the environment. *Biotechnol. Adv.* **2003**, *21*, 383–393. [CrossRef]
141. Gong, X.; Huang, D.; Liu, Y.; Zeng, G.; Wang, R.; Wei, J.; Huang, C.; Xu, P.; Wan, J.; Zhang, C. Pyrolysis and reutilization of plant residues after phytoremediation of heavy metals contaminated sediments: For heavy metals stabilization and dye adsorption. *Bioresour. Technol.* **2018**, *253*, 64–71. [CrossRef] [PubMed]
142. Dubchak, S.; Bondar, O. Bioremediation and Phytoremediation: Best Approach for Rehabilitation of Soils for Future Use. In *Remediation Measures for Radioactively Contaminated Areas*; Gupta, D., Voronina, A., Eds.; Springer: Berlin/Heidelberg, Germany, 2019; pp. 201–221. Available online: https://doi.org/10.1007/978-3-319-73398-2_9 (accessed on 20 April 2020).
143. Sharma, P.; Pandey, S. Status of phytoremediation in world scenario. *Int. J. Environ. Bioremediat. Biodegrad.* **2014**, *2*, 178–191.
144. Huang, Z.; Zeng, Z.; Chen, A.; Zeng, G.; Xiao, R.; Xu, P.; He, K.; Song, Z.; Hu, L.; Peng, M. Differential behaviors of silver nanoparticles and silver ions towards cysteine: Bioremediation and toxicity to *Phanerochaete chrysosporium*. *Chemosphere* **2018**, *203*, 199–208. [CrossRef] [PubMed]
145. Romeh, A.A.A. Green silver nanoparticles for enhancing the phytoremediation of soil and water contaminated by fipronil and degradation products. *Water Air Soil Pollut.* **2018**, *229*, 1–13. [CrossRef]
146. Saraswathi, V.S.; Kamarudheen, N.; BhaskaraRao, K.; Santhakumar, K. Phytoremediation of dyes using *Lagerstroemia speciosa* mediated silver nanoparticles and its biofilm activity against clinical strains *Pseudomonas aeruginosa*. *J. Photochem. Photobiol. B Biol.* **2017**, *168*, 107–116. [CrossRef]
147. Abbas, S.; Nasreen, S.; Haroon, A.; Ashraf, M.A. Synthesis of silver and copper nanoparticles from plants and application as adsorbents for naphthalene decontamination. *Saudi J. Biol. Sci.* **2020**, *27*, 1016–1023. [CrossRef]
148. Omer, A.M. Energy, environment and sustainable development. *Renew. Sustain. Energy Rev.* **2008**, *12*, 2265–2300. [CrossRef]
149. Shrivastava, P. The role of corporations in achieving ecological sustainability. *Acad. Manag. Rev.* **1995**, *20*, 936–960. [CrossRef]
150. Wigger, K.A.; Shepherd, D.A. We're All in the Same Boat: A Collective Model of Preserving and Accessing Nature-Based Opportunities. *Entrep. Theory Pract.* **2020**, *44*, 587–617. [CrossRef]
151. Laguir, I.; Stekelorum, R.; El Baz, J. Going green? Investigating the relationships between proactive environmental strategy, GSCM practices and performances of third-party logistics providers (TPLs). *Prod. Plan. Control* **2020**, 1–14. [CrossRef]
152. Reeves, R.D.; Baker, A.J.; Jaffré, T.; Erskine, P.D.; Echevarria, G.; van der Ent, A. A global database for plants that hyperaccumulate metal and metalloid trace elements. *New Phytol.* **2018**, *218*, 407–411. [CrossRef] [PubMed]
153. Milner, M.J.; Kochian, L.V. Investigating heavy-metal hyperaccumulation using *Thlaspi caerulescens* as a model system. *Ann. Bot.* **2008**, *102*, 3–13. [CrossRef]
154. Frérot, H.; Faucon, M.P.; Willems, G.; Godé, C.; Courseaux, A.; Darracq, A.; Verbruggen, N.; Saumitou-Laprade, P. Genetic architecture of zinc hyperaccumulation in *Arabidopsis halleri*: The essential role of QTL × environment interactions. *New Phytol.* **2010**, *187*, 355–367. [CrossRef]
155. Caille, N.; Zhao, F.; McGrath, S. Comparison of root absorption, translocation and tolerance of arsenic in the hyperaccumulator *Pteris vittata* and the nonhyperaccumulator *Pteris tremula*. *New Phytol.* **2005**, *165*, 755–761. [CrossRef] [PubMed]
156. Krämer, U.; Cotter-Howells, J.D.; Charnock, J.M.; Baker, A.J.; Smith, J.A.C. Free histidine as a metal chelator in plants that accumulate nickel. *Nature* **1996**, *379*, 635–638. [CrossRef]
157. Krivoruchko, A.; Kuyukina, M.; Ivshina, I. Advanced Rhodococcus biocatalysts for environmental biotechnologies. *Catalysts* **2019**, *9*, 236. [CrossRef]

158. Ma, X.; Geiser-Lee, J.; Deng, Y.; Kolmakov, A. Interactions between engineered nanoparticles (ENPs) and plants: Phytotoxicity, uptake and accumulation. *Sci. Total Environ.* **2010**, *408*, 3053–3061. [[CrossRef](#)]
159. Rai, P.K.; Kim, K.-H.; Lee, S.S.; Lee, J.-H. Molecular mechanisms in phytoremediation of environmental contaminants and prospects of engineered transgenic plants/microbes. *Sci. Total Environ.* **2020**, *705*, 135858. [[CrossRef](#)] [[PubMed](#)]
160. Pilon-Smits, E. Phytoremediation. *Annu. Rev. Plant Biol.* **2005**, *56*, 15–39. [[CrossRef](#)] [[PubMed](#)]
161. Cota-Ruiz, K.; Delgado-Rios, M.; Martínez-Martínez, A.; Núñez-Gastelum, J.A.; Peralta-Videa, J.R.; Gardea-Torresdey, J.L. Current findings on terrestrial plants—Engineered nanomaterial interactions: Are plants capable of phytoremediating nanomaterials from soil? *Curr. Opin. Environ. Sci. Health* **2018**, *6*, 9–15. [[CrossRef](#)]
162. Rico, C.M.; Majumdar, S.; Duarte-Gardea, M.; Peralta-Videa, J.R.; Gardea-Torresdey, J.L. Interaction of nanoparticles with edible plants and their possible implications in the food chain. *J. Agric. Food Chem.* **2011**, *59*, 3485–3498. [[CrossRef](#)] [[PubMed](#)]
163. Yang, Q.; Xu, W.; Liu, G.; Song, M.; Tan, Z.; Mao, Y.; Yin, Y.; Cai, Y.; Liu, J.; Jiang, G. Transformation and uptake of silver nanoparticles and silver ions in rice plant (*Oryza sativa* L.): The effect of iron plaque and dissolved iron. *Environ. Sci. Nano* **2020**, *7*, 599–609. [[CrossRef](#)]
164. Oukarroum, A.; Barhoumi, L.; Pirastru, L.; Dewez, D. Silver nanoparticle toxicity effect on growth and cellular viability of the aquatic plant *Lemna gibba*. *Environ. Toxicol. Chem.* **2013**, *32*, 902–907. [[CrossRef](#)]
165. Prabakaran, K.; Li, J.; Anandkumar, A.; Leng, Z.; Zou, C.B.; Du, D. Managing environmental contamination through phytoremediation by invasive plants: A review. *Ecol. Eng.* **2019**, *138*, 28–37. [[CrossRef](#)]
166. Hashem, A.; Tabassum, B.; Abd_Allah, E.F. *Bacillus subtilis*: A plant-growth promoting rhizobacterium that also impacts biotic stress. *Saudi J. Biol. Sci.* **2019**, *26*, 1291–1297. [[CrossRef](#)]
167. Kaegi, R.; Voegelin, A.; Ort, C.; Sinnet, B.; Thalmann, B.; Krismer, J.; Hagendorfer, H.; Elumelu, M.; Mueller, E. Fate and transformation of silver nanoparticles in urban wastewater systems. *Water Res.* **2013**, *47*, 3866–3877. [[CrossRef](#)] [[PubMed](#)]