

The role of combination therapy with corticosteroids and long-acting β_2 -agonists in the prevention of exacerbations in COPD

Mario Cazzola¹
Nicola A Hanania²

¹Antonio Cardarelli Hospital, Department of Respiratory Medicine, Unit of Pneumology and Allergology, Naples, Italy; ²Baylor College of Medicine, Division of Pulmonary and Critical Care Medicine, Houston, TX, USA

Abstract: Acute exacerbations of COPD can complicate the course of the disease in patients with severe airway obstruction. Reduction of exacerbations is an important clinical outcome in evaluating new therapies in COPD. Combination therapies with long-acting β -agonists and inhaled corticosteroids have now been approved for use. Three 1-year randomized clinical trials, which studied the effect of combining a long-acting β_2 -agonist with an inhaled corticosteroid in COPD, documented that exacerbation frequency was lower with therapy than placebo. Combination therapy had a similar effect to its monocomponents in the trial evaluating salmeterol/fluticasone combination. However, when patients with more severe COPD were studied using a combination of budesonide and formoterol, a clear improvement was seen in the overall exacerbation rates compared with the use of a long-acting β_2 -agonist alone.

Keywords: COPD, exacerbation, inhaled corticosteroids, inhaled long-acting β_2 -agonists, budesonide, fluticasone, formoterol, salmeterol, combination therapy

Introduction

Acute exacerbations of COPD are characterized by a change in cough, sputum production, and respiratory symptoms from baseline that cause the patient to seek medical attention and involve increasing the dose and/or frequency of existing bronchodilator therapy. Systemic corticosteroids are also recommended in patients with more advanced COPD that may require hospitalization. Exacerbations may be mild, necessitating only outpatient therapy such as antibiotics and/or systemic steroids but often may be severe enough to require hospitalization and admission to the intensive care unit. Therefore, prevention of exacerbations is one of the most important goals of chronic obstructive pulmonary disease (COPD) management (Global Initiative for Chronic Obstructive Lung Disease 2005) because of the impact on patients' daily activities and health-related quality of life (HRQoL), with these effects lasting for several weeks or months (Haughney and Gruffydd-Jones 2004). Frequent exacerbations also lead to a poorer recovery in terms of improvement in health status (Haughney and Gruffydd-Jones 2004). Furthermore, the frequency of acute exacerbation has a significant impact on the accelerated decline of lung function (Donaldson et al 2002) and also on healthcare costs (Haughney and Gruffydd-Jones 2004). Consequently, strategies for preventing COPD exacerbations may have an important effect on the natural course, cost of management, and mortality in COPD (Martinez and Anzueto 2005).

In recent years, several large-scale clinical trials in COPD have analyzed the exacerbation-preventing aspects of different treatments (Niewoehner 2004; Solèr 2005). These include inhaled bronchodilators, inhaled corticosteroids (ICSs), and a limited number of antimicrobial strategies. However, most antimicrobial strategies

Correspondence: Mario Cazzola
Via del Parco Margherita 24, 80121
Napoli, Italy
Tel +39 081 7473334
Fax +39 081 404188
Email mcazzola@ubisoft.it

have been used therapeutically in the form of antibiotic treatment during an established exacerbation, apart from vaccinations against influenza and pneumococci, which can protect from these infections which are particularly dangerous for COPD patients (Cazzola et al 2005).

Effect of long-acting β_2 -agonists and inhaled corticosteroid in preventing COPD exacerbations

Long-acting β_2 -agonists (LABAs) have both bronchodilator and non-bronchodilator properties, whereas ICSs are potent anti-inflammatory agents (Cazzola and Dahl 2004). It is, therefore, obvious that LABAs and ICSs treat different aspects of COPD (Cazzola and Dahl 2004). LABAs are central in the symptomatic control of COPD (American Thoracic Society/European Respiratory Society 2005), whereas conflicting results have been reported with ICSs (Barnes 2000; Calverley 2000). This may reflect the fact that the inflammation in COPD is poorly suppressed by corticosteroids, with small reduction in inflammatory cells, cytokines, or proteases in induced sputum even with oral corticosteroids (Barnes 2004), although there are some subgroups of patients (eg, those with eosinophilia) who respond very well (Chanez et al 1997).

Nonetheless, findings from several clinical studies have shown that treatment with ICSs improve expiratory airflow and reduce symptoms and the rate of exacerbations associated with COPD (Alsaeedi et al 2002), and, moreover, reduce all-cause mortality in COPD (Sin et al 2005). ICSs in relatively high doses (eg, budesonide 800 μ g/day or fluticasone propionate 1 mg/day) reduce exacerbations by 20%–30% and improve the health status of COPD patients by a similar amount compared with placebo (Man and Sin 2005). In particular, a meta-analysis of six trials, which evaluated the long-term effects (≥ 6 months) of ICSs in stable COPD and provided information about total exacerbations, demonstrated that ICSs reduce exacerbation rates in COPD by nearly one third relative to placebo (relative risk [RR], 0.70; 95% CI, 0.58–0.84) (Alsaeedi et al 2002). In the ISOLDE (Inhaled Steroid in Obstructive Lung Disease in Europe) study, the major benefit of therapy was seen in those whose lung function was less than 50% of predicted; however, the number of patients having at least one exacerbation per year was reduced with ICSs irrespective of baseline lung function (Jones et al 2003). A convincing demonstration of the effectiveness of these agents in COPD patients has come from the COPE study (an investigation

of COPD in the Department of Pulmonology, Enschede, the Netherlands) (van der Valk et al 2002). This study showed that the discontinuation of therapy with ICSs was associated with a more rapid onset and a higher risk of recurrence of exacerbations, and with a significant deterioration in aspects of HRQoL, in the majority of patients. However, 40% of subjects in the COPE study experienced no untoward effect from the withdrawal of ICSs. This finding indicates that there is an urgent need to identify which subgroups of COPD patients respond well to prolonged ICSs therapy. In any case, a recent article that has reviewed the diverse approaches to data analysis used in the randomized trials of ICSs in the treatment of COPD and their meta-analysis, has concluded that the reports that ICSs reduce COPD exacerbation rates are the result of improper statistical analysis techniques (Suissa 2006). The only two studies that used the correct statistical approach found insignificant effects with these drugs. It is not a surprise, therefore, that the evaluation of a cohort of newly treated COPD patients formed from the Administrative Databases of Saskatchewan Health documented that ICSs do not seem to be beneficial in preventing the risk of a first exacerbation of COPD (de Melo et al 2004). Interestingly, a recent paper has documented that immortal time bias cannot account for the risk reduction associated with ICSs exposure in observational studies (Kiri et al 2005). This finding is strengthened by the observed relationship between increased regularity of ICSs prescriptions and reductions in event rates.

Although LABAs have been shown to reduce exacerbation rates of COPD, the magnitude of the effect appears modest at best. Published trials reveal that salmeterol is the only LABA with a significant effect on the rate of exacerbations. In a 1-year study comparing salmeterol with placebo, salmeterol significantly reduced the mean number of exacerbations per patient by 20% and the mean number of exacerbations requiring oral corticosteroids per patient by 29% (Calverley et al 2003a). Recently, it has been documented that the addition of salmeterol to existing treatment in patients with a history of COPD exacerbations and poor reversibility of airflow obstruction reduces exacerbations in patients who comply with treatment (Stockley et al 2006). On the contrary, formoterol, in twice daily doses, failed to show such an effect in two other trials (Calverley et al 2003b; Szafranski et al 2003), although a subsequent study has documented that it can influence exacerbation surrogates, for example “bad days” (Dahl et al 2002). Salmeterol has been shown in several model systems to reduce inflammatory cell activation,

to reduce formation of edema and to enhance edema clearance, and to mitigate epithelial damage induced by bacteria (Rennard 2004). Whether these effects have any clinical importance remains to be determined, but they may account for its ability to reduce frequency of exacerbations of COPD. Formoterol may have similar non-bronchodilator effects; however, studies are needed to document these effects.

Pharmacological rationale for combining long-acting β_2 -agonists and inhaled corticosteroid in COPD

Interaction at the molecular level between ICSs and LABAs

Combining two therapies that possess different modes of action could be expected to have a greater benefit in the management of COPD. There is an exciting possibility that the potential benefit in combining ICSs and LABAs might be due to a synergistic interaction. However, the basic molecular mechanism of such an interaction has still to be fully identified. Recent *in vitro* and *in vivo* evidence suggest a mechanistic interaction at the molecular level between ICSs and LABAs. Corticosteroids have been shown to up-regulate the β_2 -adrenoceptor in the human airways, which in turn may provide more receptors for β_2 -agonists to activate (Mak et al 1995; Baraniuk et al 1997). At the same time, LABAs may potentiate the molecular mechanism of corticosteroid actions by increasing the nuclear translocation of glucocorticoid receptors (GRs) and thus causing an additive or sometimes synergistic suppression of inflammatory mediator release (Eickelberg et al 1999; Korn et al 2001; Pang and Knox 2000, 2001). Similar interactions between ICSs and LABAs *in vivo* are a likely explanation for the effects seen in the study of Barnes and coworkers (2006) documenting that the combination of inhaled salmeterol and fluticasone propionate significantly reduced the absolute numbers of biopsy ($CD45^+$) leukocytes, $CD8^+$ cells, and $CD4^+$ cells together with decreases in cells expressing genes for the proinflammatory mediators IFN- γ and TNF- α . In the complementary airway compartment sampled by induced sputum, combination treatment significantly reduced sputum differential neutrophils and total eosinophils. Furthermore, *in vitro* studies suggest that the addition of a LABA to an ICS may counter some of the potential negative effects of the corticosteroid (Tse et al

2003) or alternatively, some of those elicited by LABA (Seeto et al 2003).

Increasing compliance

In addition to the complementary interaction of LABAs and ICSs at the molecular level, the delivery of these medications together in a single device may also help simplify treatment regimens and improve patient adherence (Cazzola and Dahl 2004). It is widely recognized that adherence to treatment of chronic obstructive airways diseases declines as the regime becomes more complicated, either by increasing the number of medications and/or the number of daily doses (Coutts et al 1992; Chapman et al 2000). As a result, improved compliance would be expected to occur through the use of a single combination inhaler simply because of the reduction in the number of medications and actuations required with such a regime. In particular, it would be expected that the use of combination therapy should improve compliance with ICS therapy as its use is linked with LABA therapy, which patients are more likely to take because of the greater symptomatic benefit.

Enhanced respiratory cytoprotection against viral and bacterial infection

Because of the above effects, it is not unexpected that long-term use of ICSs with LABAs may be associated with a reduction in COPD exacerbations (Cazzola and Dahl 2004). However, the capacity of both ICSs and LABAs to reduce the total number of bacteria adhering to the respiratory mucosa in a concentration-dependent manner without altering the bacterial tropism for mucosa, and to preserve ciliated cells, is important in this context (Dowling et al 1999). ICSs and LABAs, when administered together at low concentrations, exhibited a synergistic effect with respect to the preservation of ciliated cells, showing a trend toward reduced damage and a significant preservation of the number of ciliated cells compared with either agent alone at the same concentrations. This result may have clinical significance as it is thought that ciliated cells are the most sensitive to damage by bacterial infection (Tsang et al 1994). It is well-known that airway colonization and chronic infection contribute to progressive pulmonary damage in COPD patients via the action of proinflammatory substances in what is known as the "vicious circle theory" (Wilson 1991). Recently, the synergistic effects of salmeterol and fluticasone in human rhinovirus-induced proinflammatory cytokine production have also been documented (Edwards et al 2006).

Rhinoviruses are implicated in many acute exacerbations of COPD, perhaps by inducing proinflammatory cytokines (Seemungal et al 2001).

Effect of combination therapy in preventing COPD exacerbations

The efficacy of combination therapy with ICSs and LABAs delivered via a single inhaler to COPD patients has repeatedly been documented (Table 1). In particular, the impact on acute exacerbations of long-term treatments with combined therapy is shown in Figures 1 and 2, and Table 2. It is very important to also mention that the definition of exacerbation and the patient population enrolled in these studies are not uniform (Table 3), and these facts have to be taken in consideration when analyzing their results.

In a multicenter, randomized, placebo-controlled study, Hanania et al (2003) compared the efficacy and safety of fluticasone propionate (250 µg) and salmeterol (50 µg), when administered twice daily together in a single device for 6 months, with that of placebo and the individual agents alone in patients with COPD. This study was unable to observe significant differences among treatment groups in the number of exacerbations or the time to first exacerbation. However, this study was of short duration and was designed and powered to evaluate the treatment effect on FEV₁, which was the primary measure of efficacy, rather than to evaluate the rates of exacerbations.

In a study with a similar design but evaluating the higher dose formulation of this combination therapy, fluticasone propionate (500 µg) with salmeterol (50 µg) twice daily given for 6 months (Mahler et al 2002), there were no statistically significant differences between treatment groups in time to exacerbation. This study also demonstrated no difference in the number of participants who exacerbated once (placebo: 8.8%; salmeterol: 5.6%; fluticasone: 10.1%; fluticasone/salmeterol combination [FSC]: 8.5%), but no *p* values were presented. Also this study was of short duration and was designed and powered to evaluate the treatment effect on FEV₁, which was the primary measure of efficacy, rather than to evaluate the rates of exacerbations.

The TRISTAN study (Calverley et al 2003a), a randomized controlled trial over 1 year of combination treatment with salmeterol (50 µg) and fluticasone (500 µg) administered together in a single device twice daily versus each of the components alone and versus placebo, enrolled patients with a baseline FEV₁ before bronchodilation that

was 25%–70% of that predicted. In this study, the rate of exacerbations requiring medical intervention, defined as a worsening of COPD symptoms that required treatment with antibiotics or oral corticosteroids, or both, fell by 25% in the combination group and by 20% and 19% in the salmeterol and fluticasone groups, respectively, compared with placebo. In particular, there was a significant difference in the mean exacerbation rate per participant per year in favor of the combination therapy when compared with placebo (FSC: 0.97 vs placebo: 1.30, *p*<0.0001), but, interestingly, differences among the three active treatment arms were small and not statistically significant. The treatment effect was more pronounced in patients with severe disease (ie, a baseline FEV₁ <50% of predicted), who showed a 30% reduction with combination therapy compared with placebo, compared with a 10% reduction in patients who had a baseline FEV₁ greater than 50% of predicted. However, this beneficial effect was also seen for exacerbations treated with oral corticosteroids, which have conventionally been thought to be more severe episodes. The combination therapy was associated with an approximate 39% reduction in the number of this type of exacerbations compared with placebo (0.46 vs 0.76 exacerbations per year, *p*<0.0001) but, again, the differences between FSC and salmeterol or fluticasone were not significant. It is worth mentioning that the TRISTAN was longer, but still based on FEV₁. Hence, the ability to observe exacerbation differences may have been helped by study duration.

In a study using a design similar to that of the TRISTAN study but including patients with an FEV₁ below 50% predicted, Szafranski et al (2003) compared the effects of two inhalations twice daily of either budesonide/formoterol combination (BFC) 160/4.5 µg (delivered dose) in a single inhaler, budesonide 200 µg (metered dose), formoterol 4.5 µg, or placebo for 12 months. Data from this study demonstrate a significant reduction in the number of severe exacerbations by 24% in the BFC group vs placebo, 23% vs formoterol, and 11% vs budesonide. This reflects a mean reduction in the mean annual exacerbation rates of 0.42 (95% CI 0.14–0.70), 1.84–1.42 exacerbations per year when combination therapy was used. Compared with placebo, both BFC and budesonide significantly reduced the number of oral steroid courses used in association with exacerbations (31% and 29%, respectively, compared with 3% for formoterol vs placebo). In addition, BFC significantly reduced the number of oral steroid courses compared with formoterol (–28%). Mild exacerbations were also significantly reduced on BFC when

compared with placebo (−62%) and budesonide (−35%), but not with formoterol (−15%).

In another study investigating BFC, Calverley et al (2003b) studied patients who initially received formoterol alone (9 µg delivered dose twice daily) and a short course of oral prednisolone (30 mg once daily) for 2 weeks before randomization. The patients were subsequently randomized to twice daily inhaled BFC 320/9 µg (delivered dose), budesonide 400 µg (metered dose), formoterol 9 µg, or placebo for 12 months, BFC prolonged time to first exacerbation requiring medical intervention, defined as a need for antibiotics and/or oral corticosteroids and/or hospitalization due to respiratory symptoms, compared with all other treatments. Hazard rate analysis showed that the risk of having an exacerbation while being treated with BFC was reduced by 22.7%, 29.5%, and 28.5% vs budesonide, formoterol, and placebo, respectively. The exacerbation rate with BFC was reduced compared with placebo (23.6%) and formoterol (25.5%) but not with budesonide alone (13.6%). Neither budesonide nor formoterol affected either measure of exacerbation compared with placebo. Furthermore, BFC prolonged the time to first course of oral corticosteroids after randomization; risk reductions

were 32.7% and 33.8% vs budesonide and formoterol, respectively, and 42.3% vs placebo. BFC also reduced the rate of oral corticosteroid courses by 28.2%, 30.5%, and 44.7% vs budesonide, formoterol, and placebo, respectively; budesonide alone reduced the number of oral corticosteroid courses compared with placebo but formoterol did not.

Considerations on the role of the combination therapy in preventing COPD exacerbations

Data of three large studies (Calverley et al 2003a, 2003b; Szafranski et al 2003) that have examined the effects of a combination of LABA and ICS on the risk of developing exacerbation in COPD have shown that the patients receiving combination therapy were less likely to report an exacerbation than those receiving the LABA alone. Nevertheless, a recent Cochrane review (Nannini et al 2004) concluded that there were conflicting results when the different combination therapies were compared with the single components alone. In particular, the results of the studies showed that BFC and FSC were effective and reduced the frequency of exacerbations compared

Table 1 Long-term effect of combination therapy in COPD

Study	Treatment (µg bid) ^a	N	Duration (weeks)	Baseline FEV ₁ (% pred)	Primary outcome(s)
Fluticasone/salmeterol (FSC)					
Hanania et al 2003	FSC 250/50	178	24	41%	Changes in FEV ₁
	FP 250	183		42%	
	SAL 50	177		42%	
	PLA	185		42%	
Mahler et al 2002	FSC 500/50	165	24	41%	Changes in FEV ₁
	FP 500	168		41%	
	SAL 50	160		40%	
	PLA	181		41%	
Calverley et al 2003a (TRISTAN study)	FSC 500/50	358	52	45%	Changes in FEV ₁
	FP 500	374		45%	
	SAL 50	372		44%	
	PLA	361		44.2	
Budesonide/formoterol (BFC)					
Szafranski et al 2003	BFC 400/12	208	52	36%	Changes in FEV ₁
	BUD 400	198		37%	
	FORM 12	201		36%	
	PLA	205		36%	
Calverley et al 2003b	BFC 400/12	254	52	36%	Changes in FEV ₁
	BUD 400	257		36%	
	FORM 12	255		36%	
	PLA	256		36%	Time to first exacerbation

^a All doses are reported as metered doses.

Abbreviations: BFC, budesonide/formoterol combination; BUD, budesonide; FORM, formoterol; FP, fluticasone propionate; FSC, fluticasone/salmeterol combination; PLA, placebo; SAL, salmeterol.




Figure 1 Total severe exacerbation rate.

Notes: PLA, placebo; LABA, salmeterol 50 µg bid or formoterol 12 µg bid; ICS, fluticasone propionate 500 µg bid or budesonide 400 µg bid; LABA/ICS, salmeterol 50 µg/fluticasone 500 µg bid or formoterol 12 µg/budesonide 400 µg bid. * difference between combination treatment and single compound treatment and/or PLA statistically significant ($p < 0.05$); § difference between single compound treatment and PLA statistically significant ($p < 0.05$). Data from Calverley et al 2003a (TRISTAN); Calverley et al 2003b; Szafranski et al 2003.

with dummy medication to a level of three quarters of the previous rates, with one exacerbation prevented every 2–4 years in the participants in the clinical trials. When the combination treatment was compared with one of the component drugs given as single treatments, BFC was better than its component LABA at preventing the frequency of exacerbations, but FSC did not show a significant advantage over LABA. Recently, two different reviews (Halpin 2005; Cazzola 2006), which have evaluated the applicability and clinical relevance of number needed to treat (NNT) analysis for determining the effectiveness of combination therapies against COPD exacerbations, have calculated that for every 100 patients treated with BFC for 1 year, between 42 and 47 exacerbations requiring medical intervention would be prevented vs LABA therapy alone, whereas for every 100 patients treated with FSC for 1 year, only 7 exacerbations would be prevented vs LABA alone (Table 2). It should be noted, however, that the NNT value relating to FSC should be interpreted with caution as there was no significant difference between FSC and salmeterol alone ($p = 0.345$ for treatment ratios for combination vs salmeterol therapy). This means that the hypothetical 95% CI would include infinity and that an NNT of infinity should be more appropriately applied to the difference between the two treatments, which indicates no true net clinical benefit of FSC over salmeterol alone.

In any case, in a recent study investigating the effect withdrawal of fluticasone from patients with severe COPD ($FEV_1 < 50\%$ predicted) using FSC (COSMIC study) for

3 months, Wouters et al (2005) demonstrated an acute and sustained decrease in FEV_1 , increase in symptoms, and an increase in mild exacerbations in patients treated with salmeterol alone compared with those who continued treatment with FSC over the one year of the study. These results emphasize the key role of ICSs in managing COPD.

The conflicting findings between the results of the different studies may be accounted for by dissimilar study designs. In particular, definitions of exacerbation that have been used in the examined studies have been rather different (Table 3). For example, in Mahler's study (Mahler et al 2002), exacerbation was defined by treatment (mild = increased use of salbutamol; moderate = use of either oral antibiotics and/or corticosteroids; severe = hospitalization). In the Szafranski's study (Szafranski et al 2003), the definition of mild exacerbation was a day with ≥ 4 inhalations of reliever medication above the mean run-in use, whereas that of severe exacerbation was a requirement for oral steroids and/or antibiotics and/or hospitalization due to respiratory symptoms. In the TRISTAN study (Calverley et al 2003a), exacerbations were defined a priori as a worsening of COPD symptoms that required treatment with antibiotics or oral corticosteroids, or both. Episodes that required corticosteroid treatment or hospital admission were noted separately. Pauwels et al (2004) correctly highlighted that considerable heterogeneity in the aetiology and manifestation of COPD exacerbations makes identification and quantification of defining symptoms extremely difficult.

However, with the emergence of a number of drugs designed or developed specifically to treat COPD and associated exacerbations, the absence of a uniform definition is a significant issue. In fact, the choice of definition can significantly affect study outcomes, with varying criteria likely to result in different levels of demonstrated treatment success. Furthermore, emerging data confirm that exacerbation is a term not easily understood by patients, who prefer to use simpler words such as “crisis” (Haughney and Gruffydd-Jones 2004).

It is also interesting to stress that there was some difference in the patient populations enrolled in the studies described above. Patients enrolled in the TRISTAN study (Calverley et al 2003a) had to have suffered from at least one episode of acute COPD symptom exacerbation per year in the previous 3 years, and at least one exacerbation in the year immediately before trial entry that required treatment with oral corticosteroids, antibiotics, or both. In the Szafranski study (Szafranski et al 2003), patients had to have suffered from at least one severe (use of oral corticosteroids and/or antibiotics and/or hospitalization due to respiratory symptoms) COPD exacerbation within 2 ± 12 months before the first clinic visit. Moreover, the mean pre-trial FEV₁ was 45% predicted in the TRISTAN study (Calverley et al 2003a) and 36% predicted in the Szafranski’s study (Szafranski et al 2003). Moreover, the mean total exacerbation rate per patient per year under placebo was 1.30 in the TRISTAN study (Calverley et al 2003a) and 1.9 in the Szafranski study (Szafranski et al 2003). This differences indicate that

the Szafranski’s study (Szafranski et al 2003) probably enrolled patients with more severe disease or, at least, with a higher possibility of suffering from acute exacerbation. In fact, also in the Calverley study (Calverley et al 2003b), the mean pre-trial FEV₁ was 36% predicted and the mean total exacerbation rate per patient per year under placebo was 1.80. As correctly highlighted by Calverley et al (2003b), the more severe disease in the patients enrolled in BFC studies is the likely explanation of the greater number of episodes that they observed compared with other studies, a difference that increased the power of the study to detect an effect of treatment. It is also important to highlight that the authors of the three studies did not report the number of exacerbations in the year before the enrolment in the studies. This is a real lack of information because, as recently documented by Donaldson et al (2003), annual exacerbation frequency remains constant over time.

As Rabe (2003) stated in his editorial comment on the Calverley paper (Calverley et al 2003b), the data of the combination therapy trials in COPD seem to indicate that patients experiencing more severe exacerbations benefit the most from combination therapy, whereas, if the definition of a (mild) exacerbation for clinical trials also includes aggravation of symptoms, such as in the paper by Szafranski et al (2003), the relative effect of a long-acting bronchodilator might be more pronounced. This clearly highlights the importance of definitions of exacerbations for clinical trials and calls for studies comparing the effect of maximal bronchodilation with, for example, the combination of LABAs




Figure 2 Rate of exacerbations requiring oral corticosteroids.

Notes: PLA, placebo; LABA, salmeterol 50 µg bid or formoterol 12 µg bid; ICS, fluticasone propionate 500 µg bid or budesonide 400 µg bid; LABA/ICS, salmeterol 50 µg/ fluticasone 500 µg bid or formoterol 12 µg/budesonide 400 µg bid. * difference between combination treatment and single compound treatment and/or PLA statistically significant ($p < 0.05$); § difference between single compound treatment and PLA statistically significant ($p < 0.05$). Data from Calverley et al 2003a (TRISTAN); Calverley et al 2003b; Szafranski et al 2003.

Table 2 Absolute and relative risk reductions for exacerbations of COPD requiring medical intervention with number needed to treat in three randomized and double-blind multinational trials

Treatment	12 month frequency of exacerbation (mean rate/patient/year)		Absolute risk reduction	Relative risk reduction (combination vs LABA)	NTT	Reference
	combination therapy	LABA only				
Fluticasone/salmeterol (TRISTAN, n = 1456)	0.97	1.04	0.07	0.067 (6.7%)	14	Calverley et al 2003a
Budesonide/formoterol (n = 812)	1.42	1.84	0.42	0.23 (23%)	2.4	Szafranski et al 2003
Budesonide/formoterol (n = 1002)	1.38	1.85	0.47	0.25 (25%)	2.1	Calverley et al 2003b

Data from Cazzola 2006.

Abbreviations: LABA, long-acting β_2 -agonist; NTT, number needed to treat.

with long-acting anticholinergics (LAMAs) in COPD with mild and severe exacerbations as an outcome. In effect, some studies are documenting the clinical value of combining a LABA with a LAMA in COPD (Cazzola et al 2004; van Noord et al 2005, 2006), although the information of the impact of such a type of combination on exacerbations is still lacking.

Borrowing Jones' opinion (Jones 2004), we believe that there are a number of possible mechanisms to account for the apparent additive effect of inhaled combination therapy with ICSs and LABAs in preventing exacerbations and to explain the discrepancies observed between the studies. It is

likely that: (1) the individual drugs were not at the top of their dose–response curve for this outcome; (2) they have different, and additive, mechanisms of action, which can also be linked to the specific compound; or (3) the different agents were preventing different types of exacerbation. Jones (2004) has highlighted that an operational definition based upon a worsening of symptoms and the treatment required (moderate exacerbations are those that can be treated with oral corticosteroids and/or antibiotics, severe exacerbations are those requiring hospital admission) does not identify the type of exacerbation. This emphasizes again the importance for the need of a uniform definition of COPD exacerbation for clinical trials in order to reach more solid conclusions.

Whatever the case may be, it must be emphasized that a treatment regimen with administration of an ICS and LABA in a single inhaler prolongs the time to a first COPD exacerbation, compared with treatment with the single components in patients with an FEV₁ <50% predicted (Calverley et al 2003b). Intriguingly, this finding has been observed with BFC (Calverley et al 2003b), but not with FSC (Calverley et al 2003a). The prolonged time to first exacerbation may delay the deterioration of the disease and help maintain health status and patient well-being. It is an important outcome because it indicates that following aggressive treatment of an acute exacerbation with an oral corticosteroid, therapy with an ICS/LABA combination may better control the disease than treatment with bronchodilator alone. In particular, as highlighted by Calverley (2004), combining treatments is certainly something to be considered in patients who have reported exacerbations on a regular basis but have not yet received ICS or LABA therapy in combination, although there are concerns about

Table 3 Definitions of COPD exacerbation used in combination therapy studies

Study	Definition
Hanania et al 2003	Defined by treatment, with moderate exacerbations requiring treatment with antibiotics and/or corticosteroids, and severe exacerbations requiring hospitalization)
Mahler et al 2002	Defined by treatment (mild = increased use of salbutamol; moderate = use of either oral antibiotics and/or corticosteroids; severe = hospitalization)
Calverley et al 2003a (TRISTAN study)	Worsening of COPD symptoms that required treatment with antibiotics, oral corticosteroids, or both
Szafranski et al 2003	Mild: a day with ≥ 4 inhalations of reliever medication above the mean run-in use; severe: requirement for oral steroids and/or antibiotics and/or hospitalization due to respiratory symptoms
Calverley et al 2003b	Event requiring medical intervention (oral antibiotics and/or corticosteroids or hospitalization) due to respiratory symptoms

the increased risk of side-effects and cost of using LABA/ICS therapy in COPD.

Disclosures

Dr Cazzola has received financial support for research and attending meetings and fees for speaking and consulting from Abbott, Altana, AstraZeneca, Boehringer Ingelheim, Chiesi Farmaceutici, GSK, Meda, Menarini Farmaceutici, Novartis, Pfizer, and Valeas. Dr Hanania has received research grant support from GSK, Boehringer Ingelheim, AstraZeneca, Dey, Sepracor, and Altana and has served on the advisory board for GSK and Dey. No funding source had any role in the preparation of this paper.

References

- Alsaedi A, Sin DD, McAlister FA. 2002. The effects of inhaled corticosteroids in chronic obstructive pulmonary disease: a systematic review of randomized placebo-controlled trials. *Am J Med*, 113:59-65.
- American Thoracic Society/European Respiratory Society. 2005. Standards for the diagnosis and management of patients with COPD. Accessed 22 January 2006. URL: <http://www.thoracic.org>.
- Baraniuk, JN, Ali, M, Brody, D, et al. 1997. Glucocorticosteroids induce β_2 -adrenergic receptor function in human nasal mucosa. *Am J Respir Crit Care Med*, 155:704-10.
- Barnes PJ. 2000. Inhaled corticosteroids are not beneficial in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 161:342-4.
- Barnes PJ. 2004. Corticosteroid resistance in airway disease. *Proc Am Thorac Soc*, 1:264-8.
- Barnes NC, Qiu YS, Pavord ID, et al. 2006. Antiinflammatory effects of salmeterol/fluticasone propionate in chronic obstructive lung disease. *Am J Respir Crit Care Med*, 173:736-43.
- Calverley PM. 2000. Inhaled corticosteroids are beneficial in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 161:341-2.
- Calverley PM. 2004. Reducing the frequency and severity of exacerbations of chronic obstructive pulmonary disease. *Proc Am Thorac Soc*, 1:121-4.
- Calverley PM, Boonsawat W, Cseke Z, et al. 2003b. Maintenance therapy with budesonide and formoterol in chronic obstructive pulmonary disease. *Eur Respir J*, 22: 912-9.
- Calverley P, Pauwels R, Vestbo J, et al. 2003a. TRIal of Inhaled STeroids ANd long-acting β_2 agonists study group. Combined salmeterol and fluticasone in the treatment of chronic obstructive pulmonary disease: a randomised controlled trial. *Lancet*, 361: 449-56.
- Cazzola M. 2006. Application of number needed to treat (NNT) as a measure of treatment effect in respiratory medicine. *Treat Respir Med*, 5:79-84.
- Cazzola M, Dahl R. 2004. Inhaled combination therapy with long acting β_2 -agonists and corticosteroids in stable COPD. *Chest*, 126:220-37.
- Cazzola M, Di Marco F, Santus P, et al. 2004. The pharmacodynamic effects of single inhaled doses of formoterol, tiotropium and their combination in patients with COPD. *Pulm Pharmacol Ther*, 17:35-9.
- Cazzola M, Salzillo A, De Giglio C, et al. 2005. Treatment of acute exacerbation of severe-to-very severe COPD with azithromycin in patients vaccinated against *Streptococcus pneumoniae*. *Respir Med*, 99:663-9.
- Chaney P, Vignola AM, O'Shaughnessy T, et al. 1997. Corticosteroid reversibility in COPD is related to features of asthma. *Am J Respir Crit Care Med*, 155:1529-34.
- Chapman KR, Walker L, Cluley S, et al. 2000. Improving patient compliance with asthma therapy. *Respir Med*, 94:2-9.
- Coutts JAP, Gibson NA, Paton JY. 1992. Measuring compliance with inhaled medication in asthma. *Arch Dis Child*, 67:332-3.
- Dahl R, Greefhorst LA, Nowak D, et al. 2002. Inhaled formoterol dry powder versus ipratropium bromide in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 164:778-84.
- de Melo MN, Ernst P, Suissa S. 2004. Inhaled corticosteroids and the risk of a first exacerbation in COPD patients. *Eur Respir J*, 23:692-7.
- Donaldson GC, Seemungal TA, Bhowmik A, et al. 2002. Relationship between exacerbation frequency and lung function decline in chronic obstructive pulmonary disease. *Thorax*, 57:847-52.
- Donaldson GC, Seemungal TA, Patel IS, et al. 2003. Longitudinal changes in the nature, severity and frequency of COPD exacerbations. *Eur Respir J*, 22: 931-6.
- Dowling RB, Johnson M, Cole PJ, et al. 1999. Effect of fluticasone propionate and salmeterol on *Pseudomonas aeruginosa* infection of the respiratory mucosa *in vitro*. *Eur Respir J*, 14: 363-9.
- Edwards MR, Johnson M, Johnston SL. 2006. Combination therapy: synergistic suppression of virus induced chemokines in airway epithelial cells. *Am J Respir Cell Mol Biol*, Jan 19 [Epub ahead of print].
- Eickelberg O, Roth M, Lox R, et al. 1999. Ligand-independent activation of the glucocorticoid receptor by β_2 -adrenergic receptor agonists in primary human lung fibroblasts and vascular smooth muscle cells. *J Biol Chem*, 274:1005-10.
- Global Initiative for Chronic Obstructive Lung Disease. 2005. Global Strategy for the Diagnosis Management and Prevention of Chronic Obstructive Pulmonary Disease. NHBLI/WHO Workshop Report. Accessed 22 January 2006. URL: <http://www.goldcopd.com>.
- Halpin DM. 2005. Evaluating the effectiveness of combination therapy to prevent COPD exacerbations: the value of NNT analysis. *Int J Clin Pract*, 59:1187-94.
- Hanania NA, Darken P, Horstman D, et al. 2003. The efficacy and safety of fluticasone propionate (250 μ g)/salmeterol (50 μ g) combined in the Diskus inhaler for the treatment of COPD. *Chest*, 124:834-43.
- Haughney J, Gruffydd-Jones K. 2004. Patient-centred outcomes in primary care management of COPD - what do recent clinical trial data tell us? *Primary Care Resp J*, 13:185-97.
- Jones PW. 2004. Clinical effects of inhaled corticosteroids in chronic obstructive pulmonary disease. *Proc Am Thorac Soc*, 1:167-70.
- Jones PW, Willits LR, Burge PS, et al. 2003. Disease severity and the effect of fluticasone propionate on chronic obstructive pulmonary disease exacerbations. *Eur Respir J*, 21:68-73.
- Kiri VA, Pride NB, Soriano JB, et al. 2005. Inhaled corticosteroids in chronic obstructive pulmonary disease: results from two observational designs free of immortal time bias. *Am J Respir Crit Care Med*, 172:460-4.
- Korn SH, Jerre A, Brattsand R. 2001. Effects of formoterol and budesonide on GM-CSF and IL-8 secretion by triggered human bronchial epithelial cells. *Eur Respir J*, 17:1070-7.
- Mahler DA, Wire P, Horstman D, et al. 2002. Effectiveness of fluticasone propionate and salmeterol combination delivered via the Diskus device in the treatment of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 166:1084-91.
- Mak, JC, Nishikawa, M, Shirasaki, H, et al. 1995. Protective effects of a glucocorticoid on downregulation of pulmonary β_2 -adrenergic receptors *in vivo*. *J Clin Invest*, 96:99-106.
- Man SF, Sin DD. 2005. Inhaled corticosteroids in chronic obstructive pulmonary disease: is there a clinical benefit? *Drugs*, 65:579-91.
- Martinez FJ, Anzueto A. 2005. Appropriate outpatient treatment of acute bacterial exacerbations of chronic bronchitis. *Am J Med*, 118(Suppl 7A):S39-44.
- Nannini L, Cates CJ, Lasserson TJ, et al. 2004. Combined corticosteroid and long-acting β -agonist in one inhaler for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*, (3):CD003794.
- Niewoehn 2004. Interventions to prevent chronic obstructive pulmonary disease exacerbations. *Am J Med*, 117(Suppl 12A):S41-8.

- Pang L, Knox AJ. 2000. Synergistic inhibition by β_2 -agonists and corticosteroids on tumor necrosis factor- α -induced interleukin-8 release from cultured human airway smooth-muscle cells. *Am J Respir Cell Mol Biol*, 23:79-85.
- Pang L, Knox AJ. 2001. Regulation of TNF- β -induced eotaxin release from cultured human airway smooth muscle cells by β_2 -agonists and corticosteroids. *FASEB J*, 15:261-9.
- Pauwels R, Calverley P, Buist AS, et al. 2004. COPD exacerbations: the importance of a standard definition. *Respir Med*, 98:99-107.
- Rabe K. 2003. Combination therapy for chronic obstructive pulmonary disease: one size fits all? *Eur Respir J*, 22:874-5.
- Rennard SI. 2004. Treatment of stable chronic obstructive pulmonary disease. *Lancet*, 364:791-802.
- Seemungal T, Harper-Owen R, Bhowmik A, et al. 2001. Respiratory viruses, symptoms, and inflammatory markers in acute exacerbations and stable chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 164:1618-23.
- Seeto LJ, Burgess JK, Johnson PR, et al. 2003. Effect of fluticasone and salmeterol on human alveolar macrophage cytokine production in patients with chronic obstructive pulmonary disease (COPD) [abstract]. *Am J Respir Crit Care Med*, 167:A318.
- Sin DD, Wu L, Anderson JA, Anthonisen NR, et al. 2005. Inhaled corticosteroids and mortality in chronic obstructive pulmonary disease. *Thorax*, 60:992-7.
- Solèr M. 2005. Modulation of airway inflammation to prevent exacerbations of COPD. *Eur Respir Rev*, 14:78-82.
- Stockley RA, Chopra N, Rice L. 2006. Addition of salmeterol to existing treatment in patients with COPD: a 12 month study. *Thorax*, 61:122-8.
- Suissa S. 2006. Statistical treatment of exacerbations in therapeutic trials of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 173:842-6.
- Szafranski W, Cukier A, Ramirez A, et al. 2003. Efficacy and safety of budesonide/formoterol in the management of chronic obstructive pulmonary disease. *Eur Respir J*, 21:74-81.
- Tsang KW, Rutman A, Tanaka E, et al. 1994. Interaction of *Pseudomonas aeruginosa* with human respiratory mucosa *in vitro*. *Eur Respir J*, 7:1746-53.
- Tse R, Marroquin BA, Dorscheid DR, et al. 2003. β -adrenergic agonists inhibit corticosteroid-induced apoptosis of airway epithelial cells. *Am J Physiol Lung Cell Mol Physiol*, 285:L393-404.
- van der Valk P, Monninkhof E, van der Palen J, et al. 2002. Effect of discontinuation of inhaled corticosteroids in patients with chronic obstructive pulmonary disease: the COPE study. *Am J Respir Crit Care Med*, 166:1358-63.
- van Noord JA, Aumann JL, Janssens E, et al. 2005. Comparison of tiotropium once daily, formoterol twice daily and both combined once daily in patients with COPD. *Eur Respir J*, 26:214-22.
- van Noord JA, Aumann JL, Janssens E, et al. 2006. Effects of tiotropium with and without formoterol on airflow obstruction and resting hyperinflation in patients with COPD. *Chest*, 129:509-17.
- Wilson R. 1991. Infections of the airways. *Curr Opin Infect Dis*, 4:166-77.
- Wouters EFM, Postma DS, Fokkens B, et al. 2005. Withdrawal of fluticasone propionate from combined salmeterol/fluticasone treatment in patients with COPD causes an immediate and sustained disease deterioration: a randomised controlled trial. *Thorax*, 60:480-7.