

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Methyl (*E*)-2-cyano-3-(6-nitro-1,3-benzodioxol-5-yl)acrylateM. Bakthadoss,^{a,b} A. Devaraj,^a D. Lakshmanan^c and S. Murugavel^{d*}

^aDepartment of Organic Chemistry, University of Madras, Maraimalai Campus, Chennai 600 025, India, ^bDepartment of Chemistry, Pondicherry University, Puducherry 605 014, India, ^cDepartment of Physics, C. Abdul Hakeem College of Engineering & Technology, Melvisharam, Vellore 632 509, India, and ^dDepartment of Physics, Thanthai Periyar Government Institute of Technology, Vellore 632 002, India

Correspondence e-mail: smurugavel27@gmail.com


Received 4 October 2012; accepted 16 October 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.046; wR factor = 0.129; data-to-parameter ratio = 19.6.

In the title compound, $\text{C}_{12}\text{H}_8\text{N}_2\text{O}_6$, the 1,3-benzodioxole ring system is essentially planar [maximum deviation = 0.036 (2) Å] and the nitro group is oriented at a dihedral angle of 15.4 (1)° with respect to its mean plane. In the crystal, molecules are linked into $C(8)$ [101] chains by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, and weak aromatic $\pi-\pi$ stacking [centroid-centroid distance = 3.887 (1) Å] also occurs.

Related literature

For a related structure and background references, see: Karthikeyan *et al.* (2011); Loghmani-Khouzani *et al.* (2009).



Experimental

Crystal data

 $\text{C}_{12}\text{H}_8\text{N}_2\text{O}_6$ $M_r = 276.20$

Monoclinic, $P2_1/n$
 $a = 10.8191$ (9) Å
 $b = 7.3220$ (6) Å
 $c = 15.4133$ (13) Å
 $\beta = 91.691$ (2)°
 $V = 1220.47$ (18) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.12$ mm⁻¹
 $T = 293$ K
 $0.23 \times 0.22 \times 0.17$ mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.972$, $T_{\max} = 0.979$

14190 measured reflections
 3567 independent reflections
 2286 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.129$
 $S = 1.03$
 3567 reflections

182 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.20$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.18$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C3}-\text{H3B}\cdots\text{O4}^i$	0.97	2.51	3.247 (2)	132

Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT and XPREP (Bruker, 2004); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia (1997); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

The authors thank Dr Babu Vargheese, SAIF, IIT, Madras, India, for his help with the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6968).

References

- Bruker (2004). APEX2, SAINT and XPREP. Bruker AXS Inc., Madison, Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Karthikeyan, S., Sethusankar, K., Devaraj, A. & Bakthadoss, M. (2011). *Acta Cryst.* **E67**, o3469.
 Loghmani-Khouzani, H., Abdul Rahman, N., Robinson, W. T., Yaeghoobi, M. & Kia, R. (2009). *Acta Cryst.* **E65**, o2545.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

* Additional correspondence author, e-mail: bhakthadoss@yahoo.com.

supplementary materials

Acta Cryst. (2012). E68, o3184 [doi:10.1107/S1600536812043164]

Methyl (*E*)-2-cyano-3-(6-nitro-1,3-benzodioxol-5-yl)acrylate**M. Bakthadoss, A. Devaraj, D. Lakshmanan and S. Murugavel****Comment**

As part of our ongoing studies of benzodioxoles with possible biological activities (Karthikeyan *et al.*, 2011), the crystal structure of the title compound has been determined and the results are presented here.

Fig. 1. shows a displacement ellipsoid plot of (I), with the atom numbering scheme. The benzodioxole ring system is essentially planar [maximum deviation = 0.036 (2) Å for the C3 atom] and is oriented at a dihedral angle of 15.4 (1)° with respect to the nitro group. The sum of bond angles around N2 (360°) indicates that N2 is in sp^2 hybridization. The carbonitrile side chain (C9–C12–N1) is almost linear, with the angle around central carbon atom being 179.6 (2)°. The geometric parameters of the title molecule agrees well with those reported for similar structures (Karthikeyan *et al.*, 2011, Lohmani-Khouzani *et al.*, 2009).

The crystal packing features C—H···O hydrogen bonds. Atom C3 in the molecule at x, y, z directs its C—H bonds towards atom O4 at $1/2+x, 1/2-y, 1/2+z$, forming a C(8) chain along [101] (Fig. 2). Weak aromatic π — π interactions between the benzene rings of neighbouring molecules, with Cg — Cg^v distance of 3.887 (1) Å [Fig. 3; Cg is the centroid of the C1/C2/C4/C5/C6/C7 benzene ring, Symmetry code as in Fig. 3] are also observed.

Experimental


To a solution of methyl 2-cyanoacetate (0.1 g, 1 mmol) in dichloromethane (5 ml), pyrrolidine (0.073 g, 1 mmol) was added and stirred well for 10 minutes. To this solution 6-nitrobenzo [d][1,3]dioxole-5-carbaldehyde (0.2 g, 1 mmol) was added and allowed to stir well for 12 h. After the completion of the reaction as evidenced by *TLC*, the reaction mixture was poured into 2 *N* HCl solution (10 ml) and extracted using 20 ml of dichloromethane. The organic layer thus obtained was concentrated under reduced pressure. Column purification (silica gel, mesh size: 60–120) of the crude mixture using 10% ethyl acetate in hexanes successfully provided the desired title compound in 92% yield (0.25 g).

Refinement


All the H atoms were positioned geometrically with C–H = 0.93–0.97 Å and constrained to ride on their parent atom, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms and $1.2U_{\text{eq}}(\text{C})$ for other H atoms.

Computing details


Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia (1997)); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).


Figure 1

The molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level.


Figure 2

Part of the crystal structure of (I) showing C—H...O hydrogen bonds (dotted lines), with the formation of C(8) chains along [101]. [Symmetry codes: (i) $1/2+x, 1/2-y, 1/2+z$; (ii) $1+x, y, 1+z$; (iii) $3/2+x, 1/2-y, 3/2+z$; (iv) $2+x, y, 2+z$].


Figure 3

A view of π — π (green dotted lines) interactions in the crystal structure of the title compound. Cg denotes the centroid of the C1/C2/C4/C5/C6/C7 benzene ring [Symmetry code:(v) $I-x, I-y, -z$].

Methyl (*E*)-2-cyano-3-(6-nitro-1,3-benzodioxol-5-yl)acrylate

Crystal data

$C_{12}H_8N_2O_6$
 $M_r = 276.20$
 Monoclinic, $P2_1/n$
 Hall symbol: $-P\ 2_1n$
 $a = 10.8191(9)\ \text{\AA}$
 $b = 7.3220(6)\ \text{\AA}$
 $c = 15.4133(13)\ \text{\AA}$
 $\beta = 91.691(2)^\circ$
 $V = 1220.47(18)\ \text{\AA}^3$
 $Z = 4$

$F(000) = 568$
 $D_x = 1.503\ \text{Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$
 Cell parameters from 3589 reflections
 $\theta = 2.3\text{--}30.1^\circ$
 $\mu = 0.12\ \text{mm}^{-1}$
 $T = 293\ \text{K}$
 Block, colourless
 $0.23 \times 0.22 \times 0.17\ \text{mm}$

Data collection

Bruker APEXII CCD
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: $10.0\ \text{pixels mm}^{-1}$
 ω scans

Absorption correction: multi-scan
 (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.972, T_{\max} = 0.979$
 14190 measured reflections
 3567 independent reflections
 2286 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$
 $\theta_{\text{max}} = 30.1^\circ$, $\theta_{\text{min}} = 2.3^\circ$
 $h = -15 \rightarrow 14$

$k = -6 \rightarrow 10$
 $l = -21 \rightarrow 19$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.129$
 $S = 1.03$
 3567 reflections
 182 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0569P)^2 + 0.1956P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.20 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.18 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.34825 (14)	0.0705 (2)	0.14286 (9)	0.0415 (3)
H1	0.3194	0.0163	0.1929	0.050*
C2	0.44834 (14)	0.1836 (2)	0.14611 (9)	0.0408 (3)
C3	0.60882 (16)	0.3581 (3)	0.18676 (11)	0.0560 (4)
H3A	0.5974	0.4774	0.2126	0.067*
H3B	0.6912	0.3149	0.2026	0.067*
C4	0.49334 (13)	0.26561 (19)	0.07304 (10)	0.0395 (3)
C5	0.43953 (14)	0.2399 (2)	-0.00694 (9)	0.0406 (3)
H5	0.4697	0.2955	-0.0563	0.049*
C6	0.33678 (13)	0.12542 (19)	-0.01065 (8)	0.0372 (3)
C7	0.28993 (13)	0.03817 (19)	0.06181 (9)	0.0375 (3)
C8	0.18022 (14)	-0.0796 (2)	0.05853 (9)	0.0428 (3)
H8	0.1115	-0.0384	0.0266	0.051*
C9	0.17142 (13)	-0.2415 (2)	0.09786 (9)	0.0414 (3)
C10	0.05270 (16)	-0.3443 (2)	0.09545 (10)	0.0481 (4)
C11	-0.0446 (2)	-0.6111 (3)	0.14488 (14)	0.0865 (8)
H11A	-0.0937	-0.5637	0.1906	0.130*
H11B	-0.0215	-0.7349	0.1579	0.130*
H11C	-0.0917	-0.6077	0.0911	0.130*
C12	0.27569 (16)	-0.3259 (2)	0.14154 (11)	0.0484 (4)
N1	0.35768 (15)	-0.3935 (2)	0.17628 (12)	0.0710 (5)
N2	0.28260 (12)	0.09294 (19)	-0.09677 (8)	0.0472 (3)
O1	0.51784 (12)	0.23313 (16)	0.21683 (7)	0.0566 (3)

O2	0.59385 (11)	0.36932 (17)	0.09442 (7)	0.0549 (3)
O3	0.21209 (14)	-0.03492 (18)	-0.10801 (8)	0.0666 (4)
O4	0.31029 (14)	0.1961 (2)	-0.15503 (8)	0.0796 (5)
O5	-0.03967 (12)	-0.2902 (2)	0.06023 (9)	0.0704 (4)
O6	0.06541 (12)	-0.50122 (16)	0.13725 (8)	0.0589 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0503 (9)	0.0418 (8)	0.0322 (6)	-0.0033 (7)	0.0004 (6)	0.0053 (6)
C2	0.0479 (8)	0.0384 (8)	0.0357 (7)	0.0000 (6)	-0.0082 (6)	0.0024 (6)
C3	0.0527 (10)	0.0612 (10)	0.0530 (9)	-0.0109 (8)	-0.0154 (8)	0.0049 (8)
C4	0.0381 (7)	0.0362 (7)	0.0440 (7)	-0.0009 (6)	-0.0040 (6)	0.0060 (6)
C5	0.0440 (8)	0.0405 (7)	0.0374 (7)	0.0013 (6)	0.0008 (6)	0.0083 (6)
C6	0.0411 (7)	0.0374 (7)	0.0327 (6)	0.0029 (6)	-0.0050 (6)	0.0023 (6)
C7	0.0390 (7)	0.0346 (7)	0.0389 (7)	0.0005 (6)	-0.0001 (6)	0.0027 (6)
C8	0.0398 (8)	0.0483 (8)	0.0403 (7)	-0.0002 (7)	-0.0014 (6)	-0.0014 (6)
C9	0.0381 (7)	0.0464 (8)	0.0400 (7)	-0.0044 (6)	0.0054 (6)	-0.0008 (6)
C10	0.0468 (9)	0.0586 (10)	0.0395 (7)	-0.0135 (8)	0.0091 (7)	-0.0080 (7)
C11	0.0961 (16)	0.0936 (16)	0.0707 (13)	-0.0623 (14)	0.0181 (11)	-0.0093 (12)
C12	0.0437 (9)	0.0436 (8)	0.0584 (9)	-0.0069 (7)	0.0090 (7)	0.0084 (7)
N1	0.0541 (9)	0.0636 (10)	0.0951 (13)	0.0040 (8)	0.0009 (9)	0.0224 (9)
N2	0.0523 (8)	0.0526 (8)	0.0364 (6)	-0.0010 (7)	-0.0047 (6)	0.0011 (6)
O1	0.0685 (8)	0.0590 (7)	0.0412 (6)	-0.0174 (6)	-0.0175 (5)	0.0084 (5)
O2	0.0505 (7)	0.0605 (7)	0.0529 (6)	-0.0165 (6)	-0.0125 (5)	0.0109 (6)
O3	0.0882 (9)	0.0585 (8)	0.0520 (7)	-0.0219 (7)	-0.0152 (6)	-0.0052 (6)
O4	0.0919 (10)	0.1096 (12)	0.0367 (6)	-0.0364 (9)	-0.0120 (6)	0.0199 (7)
O5	0.0437 (7)	0.0963 (10)	0.0708 (8)	-0.0157 (7)	-0.0048 (6)	0.0024 (7)
O6	0.0640 (8)	0.0560 (7)	0.0575 (7)	-0.0254 (6)	0.0143 (6)	-0.0048 (6)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.363 (2)	C7—C8	1.467 (2)
C1—C7	1.403 (2)	C8—C9	1.336 (2)
C1—H1	0.9300	C8—H8	0.9300
C2—O1	1.3553 (17)	C9—C12	1.436 (2)
C2—C4	1.378 (2)	C9—C10	1.488 (2)
C3—O2	1.430 (2)	C10—O5	1.191 (2)
C3—O1	1.431 (2)	C10—O6	1.323 (2)
C3—H3A	0.9700	C11—O6	1.444 (2)
C3—H3B	0.9700	C11—H11A	0.9600
C4—O2	1.3589 (18)	C11—H11B	0.9600
C4—C5	1.361 (2)	C11—H11C	0.9600
C5—C6	1.392 (2)	C12—N1	1.136 (2)
C5—H5	0.9300	N2—O3	1.2169 (17)
C6—C7	1.3949 (19)	N2—O4	1.2177 (17)
C6—N2	1.4547 (18)		
C2—C1—C7	118.11 (13)	C1—C7—C8	118.17 (13)
C2—C1—H1	120.9	C9—C8—C7	125.06 (14)

C7—C1—H1	120.9	C9—C8—H8	117.5
O1—C2—C1	127.85 (13)	C7—C8—H8	117.5
O1—C2—C4	109.86 (13)	C8—C9—C12	121.93 (14)
C1—C2—C4	122.29 (13)	C8—C9—C10	120.68 (14)
O2—C3—O1	107.47 (12)	C12—C9—C10	117.36 (14)
O2—C3—H3A	110.2	O5—C10—O6	125.79 (15)
O1—C3—H3A	110.2	O5—C10—C9	123.73 (16)
O2—C3—H3B	110.2	O6—C10—C9	110.48 (15)
O1—C3—H3B	110.2	O6—C11—H11A	109.5
H3A—C3—H3B	108.5	O6—C11—H11B	109.5
O2—C4—C5	127.95 (13)	H11A—C11—H11B	109.5
O2—C4—C2	110.18 (13)	O6—C11—H11C	109.5
C5—C4—C2	121.88 (13)	H11A—C11—H11C	109.5
C4—C5—C6	116.13 (13)	H11B—C11—H11C	109.5
C4—C5—H5	121.9	N1—C12—C9	179.57 (18)
C6—C5—H5	121.9	O3—N2—O4	122.63 (13)
C5—C6—C7	123.46 (13)	O3—N2—C6	119.32 (13)
C5—C6—N2	115.83 (12)	O4—N2—C6	118.05 (13)
C7—C6—N2	120.64 (13)	C2—O1—C3	106.34 (11)
C6—C7—C1	118.12 (13)	C4—O2—C3	105.99 (12)
C6—C7—C8	123.67 (13)	C10—O6—C11	116.71 (16)
C7—C1—C2—O1	179.97 (15)	C7—C8—C9—C12	-6.3 (2)
C7—C1—C2—C4	-0.1 (2)	C7—C8—C9—C10	175.68 (13)
O1—C2—C4—O2	0.51 (18)	C8—C9—C10—O5	-0.7 (2)
C1—C2—C4—O2	-179.41 (14)	C12—C9—C10—O5	-178.88 (16)
O1—C2—C4—C5	-179.43 (13)	C8—C9—C10—O6	179.10 (13)
C1—C2—C4—C5	0.7 (2)	C12—C9—C10—O6	0.96 (19)
O2—C4—C5—C6	179.86 (14)	C5—C6—N2—O3	163.45 (14)
C2—C4—C5—C6	-0.2 (2)	C7—C6—N2—O3	-13.6 (2)
C4—C5—C6—C7	-0.7 (2)	C5—C6—N2—O4	-16.5 (2)
C4—C5—C6—N2	-177.74 (13)	C7—C6—N2—O4	166.40 (15)
C5—C6—C7—C1	1.2 (2)	C1—C2—O1—C3	-178.04 (16)
N2—C6—C7—C1	178.11 (13)	C4—C2—O1—C3	2.05 (18)
C5—C6—C7—C8	178.73 (13)	O2—C3—O1—C2	-3.74 (18)
N2—C6—C7—C8	-4.4 (2)	C5—C4—O2—C3	177.10 (16)
C2—C1—C7—C6	-0.8 (2)	C2—C4—O2—C3	-2.83 (17)
C2—C1—C7—C8	-178.41 (13)	O1—C3—O2—C4	4.01 (18)
C6—C7—C8—C9	135.22 (16)	O5—C10—O6—C11	-3.8 (2)
C1—C7—C8—C9	-47.3 (2)	C9—C10—O6—C11	176.34 (14)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3B \cdots O4 ⁱ	0.97	2.51	3.247 (2)	132

Symmetry code: (i) $x+1/2, -y+1/2, z+1/2$.