

Site-specific receptor methylation of FrzCD in *Myxococcus xanthus* is controlled by a tetra-trico peptide repeat (TPR) containing regulatory domain of the FrzF methyltransferase

■ **OnlineOpen:** This article is available free online at www.blackwell-synergy.com

Anslley E. Scott,¹ Eric Simon,² Samuel K. Park,¹
Philip Andrews² and David R. Zusman^{1*}

¹Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA.

²Department of Biological Chemistry, University of Michigan, Ann Arbor, MI, USA

Summary

Myxococcus xanthus is a gliding bacterium with a complex life cycle that includes swarming, predation and fruiting body formation. Directed movements in *M. xanthus* are regulated by the Frz chemosensory system, which controls cell reversals. The Frz pathway requires the activity of FrzCD, a cytoplasmic methyl-accepting chemotaxis protein, and FrzF, a methyltransferase (CheR) containing an additional domain with three tetra trico-peptide repeats (TPRs). To investigate the role of the TPRs in FrzCD methylation, we used full-length FrzF and FrzF lacking its TPRs (FrzF^{CheR}) to methylate FrzCD *in vitro*. FrzF methylated FrzCD on a single residue, E182, while FrzF^{CheR} methylated FrzCD on three residues, E168, E175 and E182, indicating that the TPRs regulate site-specific methylation. E168 and E182 were predicted consensus methylation sites, but E175 is methylated on an HE pair. To determine the roles of these sites *in vivo*, we substituted each methylatable glutamate with either an aspartate or an alanine residue and determined the impact of the point mutants on single cell reversals, swarming and fruiting body formation. Single, double and triple methylation site mutants revealed that each site played a unique role in *M. xanthus* behaviour and that the pattern of receptor methylation determined receptor activity. This work also shows that methylation can both activate and inactivate the receptor.

Accepted 28 May, 2008. *For correspondence. E-mail zusman@berkeley.edu; Tel. (+1) 510 642 2293; Fax (+1) 510 643 6334. Re-use of this article is permitted in accordance with the Creative Commons Deed, Attribution 2.5, which does not permit commercial exploitation.

Introduction

Myxococcus xanthus is a Gram-negative bacterium that has a complex life cycle that involves vegetative swarming, predation and fruiting body formation (Reichenbach, 1999; Shimkets, 1999; Kaiser, 2003; 2004; Berleman *et al.*, 2006). These behaviours require motility on solid surfaces: S-motility powered by Type IV pili (Wu and Kaiser, 1995) moves cells in groups and A-motility powered by unidentified motor proteins and putative adhesion complexes moves single cells (Mignot, 2007). To achieve directed movements, *M. xanthus* cells periodically reverse so that the leading pole becomes the lagging pole. The frequency of cell reversals is controlled by the *frz* chemosensory pathway and is important for directed cell movements (Blackhart and Zusman, 1985). *frz* mutants ($\Delta frzA$, $\Delta frzB$, $\Delta frzCD$, $\Delta frzE$, $\Delta frzF$) rarely reverse and are therefore defective in swarming and fruiting body formation, forming frizzy aggregates instead of fruiting bodies on starvation media (Zusman, 1982). In contrast, some constitutively signalling *frz* mutants hyper-reverse, forming very compact colonies with little cell spreading (Blackhart and Zusman, 1985; Bustamante *et al.*, 2004).

Frz proteins are homologous to bacterial chemotaxis proteins (McBride *et al.*, 1989; Zusman *et al.*, 2007). FrzCD, a cytoplasmic methyl-accepting chemotaxis protein (MCP) homologue, lacks the transmembrane and periplasmic domains common to most MCPs and has in its place a unique N-terminal domain. In contrast, the C-terminal domain of FrzCD is similar to other MCPs and contains potential methylation sites (Astling *et al.*, 2006). *In vitro* analysis has shown that FrzCD interacts with FrzE, a histidine kinase (CheA)-response regulator fusion protein by means of two CheW-like proteins, FrzA and FrzB (Astling, 2003). When stimulated, FrzE autophosphorylates and transfers a phosphoryl group to the dual response regulator FrzZ, triggering cell reversals for both the A- and S-motility systems (Inclán *et al.*, 2007).

It is not known how FrzCD receives signals as it lacks the usual signal-binding domain common to most MCPs. Furthermore, *frzCD* N-terminal domain deletion mutants show only minor defects in behaviour (Bustamante *et al.*, 2004). Previously, it was hypothesized that a signal input

to the Frz pathway may involve differential methylation of particular sites on the receptor (Igoshin *et al.*, 2004; Astling *et al.*, 2006). Astling *et al.* (2006) identified several putative methylation sites based on sequence in comparison with known *Escherichia coli* methylation sites. They systematically mutated these sites, substituting each EE, QQ, QE and EQ pair with an alanine pair (AA). This work suggested that FrzCD receptor activity could be turned on or off depending on the site that was methylated.

Based on this work, we hypothesized that differential methylation of FrzCD may be mediated by FrzF, a methyltransferase (CheR) homologue (McCleary *et al.*, 1990) that contains an additional domain with three tetra-trico-peptide repeats (TPRs) (Shiomi *et al.*, 2002). In other organisms, TPRs have been shown to mediate protein-protein interactions (Blatch and Lasse, 1999). A Basic Local Alignment Search Tool (BLAST) (Altschul *et al.*, 1990) analysis of all sequenced bacterial genomes revealed that dozens of bacterial species possess putative methyltransferases with one or more TPRs, including several within the α -proteobacteria, β -proteobacteria, δ -proteobacteria and high GC-rich Gram-positive bacteria. In *M. xanthus*, CheR4 and CheR6 are each predicted to contain one TPR (Scott, 2008). However, to our knowledge, no function has yet been attributed to TPRs in methyltransferases.

To investigate the role of the TPR containing domain in FrzCD methylation, we used full-length FrzF and FrzF lacking the TPRs (FrzF^{CheR}) to methylate FrzCD *in vitro*. We found that indeed the TPRs of FrzF negatively regulate FrzCD methylation. We used mass spectrometry to identify the methylated sites and site-directed mutagenesis to determine the function of these sites *in vivo*. This work showed that each FrzCD methylation site played specific roles in cell motility and behaviour.

Results

Methylation of FrzCD *in vitro* using purified FrzF and FrzF^{CheR}

Previous work showed that methylation of FrzCD is required for swarming and fruiting body formation in *M. xanthus* and that methylation is mediated by FrzF (McBride *et al.*, 1989; McCleary *et al.*, 1990). FrzF is a complex methyltransferase that contains an N-terminal domain with 31% sequence identity (83 of 271 amino acids) to the methyltransferase (CheR) of *E. coli* and a C-terminal domain with three TPRs (Fig. 1A). Because of its high homology to CheR, the N-terminal domain of FrzF was assumed to have methyltransferase activity, but the




Fig. 1. Methylation of FrzCD by FrzF and FrzF^{CheR} *in vitro*.

A. Cartoons show the domain organization of the wild-type FrzF protein and the FrzF^{CheR} protein, which lacks FrzF amino acids 450–580. TPR indicates a tetra-trico peptide repeat. Numbers refer to the amino acid position in full-length FrzF.

B. His-tagged FrzCD was expressed and purified from *E. coli* and incubated *in vitro* with purified FrzF or FrzF^{CheR} and S-adenosyl methionine (SAM) for 4 h at 32°C. Following the reaction, FrzCD was analysed by SDS-PAGE and Western immunoblotting using purified α -FrzCD antibodies. The white arrowhead shows the mobility of unmethylated FrzCD; the black arrowheads indicate methylated FrzCD. The position of the molecular weight markers is indicated on the left.

function of the C-terminal domain was unknown. To determine the activities of full-length FrzF and FrzF^{CheR} (FrzF lacking its TPR domains), we cloned His-tagged *frzCD*, *frzF* and *frzF^{CheR}* in expression vectors and purified the respective proteins from *E. coli*. FrzCD purified from *E. coli* was unmethylated (Fig. 1B, lane 3), indicating that the *E. coli* methyltransferase does not methylate FrzCD. To methylate FrzCD *in vitro*, we incubated FrzCD and the methyl donor S-adenosyl methionine (SAM) with either FrzF or FrzF^{CheR}. The reactions were monitored by Western immunoblot analysis using purified anti-FrzCD antibodies as methylated FrzCD migrates faster than unmethylated FrzCD in this SDS-PAGE system (McCleary *et al.*, 1990). We found that both FrzF and FrzF^{CheR} were able to methylate FrzCD in the presence of SAM (Fig. 1B), but that they produced different FrzCD methylation patterns. FrzF^{CheR} produced a faster migrating band of FrzCD (Fig. 1 lane 5) than FrzF (Fig. 1 lane 4). Neither FrzF nor FrzF^{CheR} were able to methylate FrzCD without SAM (data not shown). As methylated receptors migrate faster than unmethylated receptors in this gel system, we hypothesized that the faster migrating band observed with the FrzF^{CheR} sample represented a more methylated species of FrzCD.

Identifying the methylated residues of FrzCD

As the mobility of FrzCD on polyacrylamide gels does not give us quantitative data on methylation patterns, we analysed the *in vitro* methylated FrzCD by mass spectrometry. We prepared methylated FrzCD samples as described above by incubating purified FrzCD, SAM and either FrzF or FrzF^{CheR}; as a control, we also prepared non-methylated FrzCD in the same way except that we omitted FrzF. To generate peptides of the optimal size for mass spectrometry, we digested the methylated and unmethylated FrzCD samples with trypsin, chymotrypsin or GluC proteases. The combined tandem mass spectrometry (MS/MS) results accounted for 92.3% total sequence coverage including 100% sequence coverage of the C-terminus of FrzCD (amino acids 136–437), predicted to contain all putative methylation sites (Fig. S1).

A comparison between spectra of tryptic FrzCD peptides methylated by FrzF and by FrzF^{CheR} revealed that FrzF methylated FrzCD on a single glutamate residue and that FrzF^{CheR} methylated FrzCD on three residues (based on accurate precursor mass measurement and MS/MS data) within the peptide L148-K184. This peptide sequence includes five glutamate residues, all representing possible sites of methylation. However, while fragmentation was sufficient for the identification of the peptide, as well as the presence of methylation, it was not adequate to obtain the sequence coverage necessary to ascertain the specific sites of methylation (data not shown). To

address this, FrzCD was digested with the chymotrypsin and GluC proteases. Chymotrypsin allowed us to determine that FrzCD was methylated by FrzF on E182 (Table 1, Fig. S2). Specifically, the b-ion with the value of 1372.3 indicates that there is a methylated glutamate on the N-terminal side of residue L183 and the b-ion with the value of 1301.1 indicates that no residue is methylated to the N-terminal side of E182. Thus, the only residue that can be methylated on this peptide is E182. However, we found that the peptides were not always methylated. For instance, two of the nine ¹⁵⁷AASTQHETSSTEQAAA IHETTATMEEL₁₈₃ peptides analysed showed methylation on site E182 (Table 1, Fig. S2, and data not shown). No methylation was seen on sites E168 or E175 on any of the nine peptides. MASCOT scores for these nine peptides were all greater than 24 indicating that these results are highly significant.

We found that FrzF^{CheR} methylated FrzCD on residues E175 and E182 by using a chymotrypsin digest (Table 2, Fig. S3A). Specifically, b-ion 849.7 showed that the methylated glutamate was on the C-terminal side of residue I173 and b-ion 1040.6 showed that the methylated glutamate was on the N-terminal side of T176. Thus, E175 must be methylated as it is the only glutamate between residues I173 and T176. b-ion 1315.2 showed that another methylated glutamate was located on the

Table 1. Ions from MS/MS spectrum of a chymotryptic fragment^a of FrzCD show FrzF methylates site E182.

AA ^b	Theoretical ^c b-ions ^d	Observed ^e b-ions	Theoretical y-ions ^f	Observed y-ions
I173	842.9 (2+)	843.0 (2+)		
H174				
E175				
T176	1026.5 (2+)	1026.3 (2+)		
T177			824.4	824.0
A178	1112.5 (2+)	1112.5 (2+)	723.3	723.1
T179	1163.0 (2+)	1163.0 (2+)	652.3	652.1
M180	1236.6 (2+)	1236.9 (2+)	551.2	551.1
E181	1301.1 (2+)	1301.0 (2+)		
E182#	1372.6 (2+)	1372.3 (2+)		
L183				

a. No ions were observed for residues ¹⁵⁷AASTQHETSSTEQA AA172 of the chymotryptic peptide ¹⁵⁷AASTQHETSSTEQAAAIHETTATMEEL₁₈₃ (Fig. S2), so these residues were removed from the left column for simplicity.

b. Amino acids are indicated by their one letter code; numbers represent their position in FrzCD. The # symbol represents an amino acid that is methylated.

c. Theoretical ions are calculated by dividing the predicted mass of an ion by the ion's charge, $m/z = (M + nH^+)/n$.

d. A b-ion is an N-terminal charged fragment generated after ion activation causes a peptide bond to break.

e. Observed ions were found by digesting FrzCD with chymotrypsin and using tandem MS/MS.

f. A y-ion is a C-terminal charged fragment generated after ion activation causes a peptide bond to break.

All ions are 1+ charged unless otherwise indicated in parentheses.




Fig. 2. FrzCD methylation point mutants are stably expressed. An immunoblot of vegetative cell extracts from the methylation point mutants probed with the α -FrzCD antibody. Thirty micrograms of total protein from whole cell extracts were loaded per lane. The white arrowhead shows the mobility of un-methylated FrzCD; the black arrowhead indicates methylated FrzCD. DZ4717 (*6His::frzCD*) is the positive control strain used in this study. All point mutations were made in the DZ4717 background.

the non-essential *crtB* locus of the *M. xanthus* Δ *frzCD* strain. We expressed *frzCD* under control of the *tet* promoter by growing cells in the presence of anhydrotetracycline (Mignot *et al.*, 2007). As a positive control we made strain DZ4717, which contains the non-mutated *frzCD* gene in the *crtB* locus under the control of the *tet* promoter. We found that DZ4717 displayed phenotypes that were comparable to that of the wild-type strain, DZ2 (Fig. S5). To ensure that all strains expressed the FrzCD

protein, we examined each strain by immunoblot analysis using the anti-FrzCD antibody. Figure 2 shows that FrzCD was expressed at the same level in all strains and that any changes in phenotype were not due to altered expression.

We then determined the effects of mutating the methylation sites on *M. xanthus* fruiting body formation (Fig. 3) and swarming (Fig. 4), phenotypes that require a functioning Frz pathway. For reference, we also examined the phenotypes of the hypo-reversing Δ *frzCD* mutant and the




Fig. 3. Effect of FrzCD E to A and E to D methylation site mutations on aggregation and fruiting body formation. FrzCD methylation site glutamates (E) 168, 175 and 182 were changed to alanine (A) or aspartate (D) residues by site directed mutagenesis as described in *Experimental procedures*. Cells were spotted at 4×10^9 cells ml^{-1} on CF fruiting agar and incubated for 4 days at 32°C. Reference strains (DZ4717 (*6His::frzCD*), Δ *frzCD*, and *frzCD* Δ ₆₋₁₈₃) are shown in A–C. FrzCD E to A mutants are shown in the middle column (D, F, H, J and L) and FrzCD E to D mutants are shown in the right column (E, G, I, K and M). The small corner inset in each picture is a 7 \times magnification of a portion of the original.




Fig. 4. Effect of FrzCD E to A and E to D methylation site mutations on vegetative swarming. Reference strains [DZ4717 (*6His::frzCD*), $\Delta frzCD$ and $frzCD^{\Delta 6-182}$] are shown at the top (striped bars), FrzCD E to A mutants are shown in the left panel (dark grey), and FrzCD E to D mutants are shown in the right panel (light grey). The horizontal axis represents the relative distance swarmed by each strain compared with wild type. Error bars represent the standard deviation of the mean. Five microlitres of 4×10^9 cells ml^{-1} were spotted on CYE nutrient plates containing 0.4% agar and swarm expansion was measured after 3 days incubation at 32°C. Data shown are from two independent experiments with a total of seven measurements per strain.

hyper-reversing $frzCD^{\Delta 6-183}$ mutant. As shown previously (Bustamante *et al.*, 2004), the $\Delta frzCD$ control strain exhibited reduced swarming on rich media, formed tangled aggregates instead of fruiting bodies on starvation media, and reversed infrequently. In contrast, the $frzCD^{\Delta 6-183}$ control strain did not swarm on rich media, was unable to form aggregates on starvation media, and reversed much more frequently as single cells than wild type (Figs 3 and 4; Table 4). The methylation point mutants displayed varied phenotypes: some behaved like wild type, whereas others displayed $\Delta frzCD$, $frzCD^{\Delta 6-183}$ or intermediate phenotypes.

- (i) The most severe single mutant phenotypes were observed when we eliminated methylation at E182. The $frzCD^{E182D}$ mutant, similar to $\Delta frzCD$, showed reduced swarming (Fig. 4), formed tangled aggregates instead of fruiting bodies (Fig. 3I), and had a severely reduced single cell reversal frequency when compared with DZ4717 (Table 4). The $frzCD^{E182A}$ mutant displayed an intermediate swarming phenotype between

DZ4717 and the $frzCD^{E182D}$ mutant (Fig. 4), was able to form fruiting bodies (Fig. 3H), but still showed a reduced single cell reversal frequency (Table 4). Thus, FrzCD methylation site 182 is important for both social behaviours and single cell reversals.

- (ii) Single mutations in E175 or E168 resulted in less severe defects than single mutations in site E182. E to A or E to D mutations at site 175 allowed cells to swarm like DZ4717 (Fig. 4) and to form fruiting bodies (Figs 3F,G). Yet both of these mutants showed a decrease in reversal frequency when compared with DZ4717 (Table 4). The $frzCD^{E168D}$ mutant displayed no obvious defects. While the $frzCD^{E168A}$ mutant displayed a 30% decrease in swarming (Fig. 4), it showed normal single cell and fruiting body behaviour. Thus, site E175 is required for single cell reversal frequency, but neither site E168 nor E175 must be methylated for proper fruiting body formation.

In *E. coli*, methylation site mutants display additive phenotypes; defects increase in severity as additional methylation sites are mutated (Shapiro *et al.*, 1995). To test if methylation site mutations resulted in additive defects in *M. xanthus*, we constructed double and triple methylation site point mutants. We expected the double and triple mutants to display more severe defects than the single mutants. However, we found that the $frzCD^{E168D,E175D}$ and $frzCD^{E168D,E175D,E182D}$ mutants displayed DZ4717 phenotypes under all conditions tested (Fig. 3K and M, Fig. 4, Table 4). This was surprising as the $frzCD^{E175D}$ and the $frzCD^{E182D}$ mutants displayed reduced single cell reversal frequencies compared with DZ4717. These results suggest that the overall number of methylatable FrzCD residues is not crucial, but rather the pattern of FrzCD methylation is the determining factor for receptor function.


We were also surprised to find that the $frzCD^{E168A,E175A}$ and $frzCD^{E168A,E175A,E182A}$ mutants displayed severe defects under all conditions tested, as the single site

Table 4. Effect of methylation site mutations on single cell reversals.

Strain	Average reversals in 30 min (# cells)	Average reversals in 30 min E to D (# cells)
DZ4717 ^a	1.58 (84)	
$\Delta frzCD$	0.20 (59)	
$frzCD^{\Delta 6-183}$	12.14 (24)	
Site 168	1.67* (49)	1.57* (74)
Site 175	0.82 (44)	0.25 (58)
Site 182	0.58 (29)	0.22 (34)
Sites 168 + 175	6.25 (70)	1.34* (59)
Sites 168 + 175 + 182	4.63 (56)	1.18* (42)

a. DZ4717 has the same reversal frequency as wild type (see Fig. S5).

Asterisks indicate that reversals are statistically the same as DZ4717 (Student's *t*-test). Values that were statistically different from DZ4717 had *P*-values less than 0.005 (Student's *t*-test).



notype seen in the *frzCD*^{E182D} strain may be due to loss of methylation at other nearby methylation sites. In *E. coli*, replacing a methylatable E or Q residue with a D residue has been shown to prevent methylation at nearby sites (Shapiro *et al.*, 1995). This appears to hold true for the *frzCD*^{E182D} mutant. We do not see any FrzCD methylation in *M. xanthus* in the *frzCD*^{E182D} mutant (Fig. 2). As the *frzCD*^{E182D} mutant displayed a more severe phenotype than either the *frzCD*^{E168D,E175D,E182D} or *frzCD*^{E168D,E175D} mutants, we concluded that the *frzCD*^{E182D} mutant phenotype was not due to loss of methylation at sites E168 or E175. It is likely that the *frzCD*^{E182D} mutant lacks methylation at unidentified sites. We believe that additional FrzCD methylation sites exist because the *frzCD*^{E168D,E175D,E182D} mutant still displays a methylated band of FrzCD (Fig. 2). We anticipate that several additional methylation sites may be uncovered when glutamine residues are converted to glutamates by a methyltransferase *in vivo* (Stock and Koshland, 1978). Additionally, some methylation sites may only be modified *in vivo* under certain physiological conditions.

Mutations in FrzCD site E175 resulted in less severe mutant phenotypes compared with mutations in FrzCD site E182. Strains containing an aspartate or an alanine at site 175 displayed a decrease in single cell reversal frequency, but showed no obvious defect in social swarming or fruiting body development. These results suggest that site E175 is critical for coordinating single cell behaviour, but is dispensable for groups of cells. This is interesting because it suggests that FrzCD is able to regulate single cell behaviour and social behaviour independently.

FrzCD site 168 appears to play a role in regulating social swarming. The *frzCD*^{E168A} mutant displayed a reduction of social swarming, but did not have an obvious defect in fruiting body formation and single cells reversed normally. The *frzCD*^{E168D} mutant did not display any mutant phenotypes. Thus, site E168 appears to be a minor regulator of FrzCD when methylated alone. However, when both FrzCD sites E168 and E175 were replaced with alanines, *M. xanthus* cells displayed profound affects; cells were unable to swarm, they were defective in fruiting body formation, and single cells hyper-reversed. Thus, simultaneous methylation at these two sites seems to activate FrzCD. It was interesting that the *frzCD*^{E168D,E175D} mutant did not display any mutant phenotypes. Thus, the mutation at site 168 can rescue the hypo-reversing phenotype of the *frzCD*^{E175D} mutant.

In sum, we found that the TPRs of FrzF inhibit its ability to methylate FrzCD *in vitro*. Additionally, each site of FrzCD methylation plays a unique role in regulating FrzCD activity. Third, FrzCD appears to be able to control single cell and social behaviours independently. Last, our results confirm previous work by Astling *et al.* (2006),

which suggested that FrzCD methylation could both turn on and off receptor activity and that the pattern of methylation determines the activity of FrzCD, not the quantity of methylation.

Based on our current knowledge of the Frz pathway, we have proposed a model that addresses a putative input into FrzCD, the regulation of FrzCD activity, and how FrzCD activity regulates downstream signalling and cell behaviour. We believe that one input into FrzCD can be the methylation of FrzCD by FrzF. FrzF may be controlled by its TPRs. As FrzF^{CheR} methylates FrzCD more than FrzF *in vitro*, it is possible that the TPRs prevent the methyltransferase domain from methylating specific residues of FrzCD (such as E168 and E175) by forming a physical barrier between the methyltransferase domain and certain binding sites on FrzCD. To relieve this inhibition, the TPRs may bind to another protein. This binding may place the TPRs in a conformation in which they no longer impede the methyltransferase domain, thus allowing FrzF to methylate additional FrzCD residues. We hypothesize that a FrzF TPR-binding partner is up-regulated or activated as cells proceed through development because FrzCD methylation has been shown to increase as *M. xanthus* cells develop (McBride and Zusman, 1993). Additionally, we propose that the TPR-binding partner is upregulated by AsgA, CsgA, FruA and DevT and downregulated by RodK because FrzCD methylation is reduced in *asgA*, *csgA*, *fruA* and *devT* mutants, and increased in *rodK* mutants (Sogaard-Andersen and Kaiser, 1996; Geng *et al.*, 1998; Boysen *et al.*, 2002; Rasmussen *et al.*, 2005).

The second part of our model explains how the activity of FrzCD can be controlled through methylation. We propose that methylation can both turn on and turn off FrzCD activity. For instance, if FrzCD is methylated on one site (E182) its activity is inhibited, whereas if it is methylated on additional residues (E168 and E175) its activity is activated. These hypotheses are based on our FrzCD methylation point mutant data where mutations in site E182 were similar to a deletion of FrzCD and mutations that mimicked methylation at both 168 and 175 resembled a 'hyperactive' *frzCD* mutant (*frzCD*^{Δ6-183}).

The final part of our model involves how FrzCD activity regulates downstream proteins. It has been proposed that the CheA homologue, FrzE, stimulates cellular reversals by autophosphorylating and transferring a phosphoryl group to the dual receiver domain containing protein, FrzZ (McCleary and Zusman, 1990; Inclán *et al.*, 2007). We propose that certain methylation sites lead to an active form of FrzCD and this stimulates FrzE kinase activity, which leads to an increase in cellular reversals. Conversely, methylation states of FrzCD that inhibit activity result in a reduction of FrzE kinase activity and a reduction in cellular reversals.

Experimental procedures

Bacterial strains and culture conditions

The strains used in this study are listed in Table 5. *M. xanthus* was grown in CYE medium, which contains 10 mM morpholinepropanesulphonic acid (MOPs, pH 7.6), 1% (w/v) Bacto Casitone (BD Biosciences), 0.5% Bacto yeast extract and 4 mM MgSO₄ (Campos *et al.*, 1978). For *M. xanthus* swarming phenotypes 5 µl of cells at 4 × 10⁹ cells ml⁻¹ were spotted on CYE plates containing 0.5% Bacto agar (BD Biosciences). For developmental phenotypes 5 µl of cells at 2 × 10⁹ cells ml⁻¹ were spotted on CF agar which contains 1.5% Bacto agar, 10 mM MOPs (pH 7.6), 0.015% Bacto Casitone, 8 mM MgSO₄, 1 mM KH₂PO₄, 0.02% NH₄SO₄, 0.2% sodium citrate and 0.1% pyruvate (Hagen *et al.*, 1978).

Reversal frequency analysis

Cells were grown to mid-log phase and 10 µl of culture were spotted on 1/2 CTT (Hodgkin and Kaiser, 1977) with 1.5% agar. The cells were covered immediately with an oxygen permeable membrane (Yellow Springs Instrument) and allowed to settle for 1 h. Cells were filmed using a Labophot 2 microscope and a micropublisher 3.3 RTV digital camera (Q imaging) and reversal frequencies were analysed by eye. Only moving cells that did not touch another cell during filming were analysed. A student's *t*-test (two-tailed, type 3) was used to determine if the reversal frequency of each strain differed from wild type. Strains with a *P*-value less than 0.005 were considered to have different reversal frequencies from wild type. For each strain, a minimum of five movies were taken on two separate days.

Protein purification from *E. coli*

To obtain soluble protein, *frzF* and *frzF^{CheR}* amplified by PCR from genomic DNA and inserted into pET28a (Novagen) at the EcoRI and HindIII sites. The plasmid was sequenced and analysed using the Sequencher program (Gene Codes) and transformed into the *E. coli* Tuner protein expression strain (Novagen). Cells were grown to mid-log phase at 37°C and expression was induced by adding 1 mM of IPTG and transferring cells to 18°C for 16 h. Cells were isolated by centrifugation at 9000 *g*. Cells were lysed by sonication (Branson sonifier 450) (3 × 1 min output 4) in 20 mM Tris pH 7.2, 20 mM imidazole, 10% glycerol, 0.1% CHAPS buffer, 500 mM NaCl lysis buffer supplemented with 2% mammalian protease inhibitor cocktail (Sigma). Insoluble material was removed by centrifugation (20 min, 17 000 *g*) and filtration through a 0.22 micron filter (Corning 115 ml filter system). Cell lysate was injected onto a 5 ml HisTrap HP Nickel column (GE Healthcare) via an AKTA FPLC (GE Healthcare). Protein was washed with 50 ml of lysis buffer, then 50 ml of lysis buffer with 60 mM, 100 mM and 150 mM imidazole. FrzF^{CheR} was eluted using lysis buffer with 250 mM imidazole. The protein was concentrated using 30 000 molecular weight cutoff Centriprep columns (Amicon).

In vitro methylation of FrzCD

Each FrzCD methylation reaction mixture contained 3.5 µM

FrzCD, 6 µM SAM, 7 µM FrzF or FrzF^{CheR}, 10 mM TrisHCl pH 7.0, 1% glycerol and 50 mM KCl. Reactions took place at 32°C for 4 h. FrzCD methylation was observed initially by SDS-PAGE analysis and then by Mass Spectrometry (see below). Each reaction was repeated in triplicate on four separate days.

Sample preparation for mass spectrometry

Methylation of receptors was performed as described above. Methylated and unmethylated species were separated by SDS-PAGE. Gels were stained with Coloidal Coomassie (Invitrogen) and bands corresponding to unmethylated and methylated FrzCD were excised and diced into ~1–2 mm³ pieces. The gel slices were destained twice by incubation with 50 µl of 100 mM ammonium bicarbonate and 50 µl of acetonitrile at 37°C for 10 min. The gel slices were then dehydrated by incubation with 50 µl of acetonitrile at 37°C for 5 min. FrzCD contains no cysteine residues, so reduction and alkylation were not performed. Gel slices were incubated overnight at 37°C in 50 µl of 100 mM ammonium bicarbonate containing 150 ng of trypsin (Promega), chymotrypsin (Roche) or GluC (Roche). Peptides were extracted from the gel slices by incubation with 30 µl of extraction solution (2% acetonitrile and 1% formic acid) at 37°C for 30 min. The solution was removed and a second extraction was performed with 12 µl of acetonitrile and 12 µl of extraction solution at 37°C for 30 min. The peptide extract was dried down and reconstituted in 25 µl of 0.1% trifluoroacetic acid (TFA).

Liquid chromatography/mass spectrometry

High-performance liquid chromatography grade water and acetonitrile (Optima) and formic acid (Acros Organics) were purchased from Fisher Scientific. TFA was purchased from Sigma-Aldrich. Peptides were desalted and concentrated on a reversed-phase cartridge (Zorbax C₁₈; 5 mm by 0.3 mm i.d.; 5 µm; Agilent) then loaded onto a reverse-phase column (ProteoPepTM C18, 5 cm by 50 µm i.d.; 300 Å; 5 µm; New Objective) using an UltimateTM nanoliquid chromatography system (Dionex/LC Packings) coupled to a ThermoFinnigan Orbitrap tandem mass spectrometer equipped with a nano-spray source (Michrom Bioresources). The column was equilibrated for 5 min in 94% solvent A (0.1% formic acid) and 6% solvent B (90% acetonitrile, 0.1% formic acid). Solvent B was increased linearly to 40% at 40 min, 60% at 45 min and 100% B at 45.1 min, where it was held for 3 min. It was then set back to its initial solvent composition (6% B), where it was held for the duration of the run (60 min). MS survey scans were performed in the orbitrap followed by subsequent MS/MS scans of the three most abundant ions fragmented in the linear ion trap, with a dynamic exclusion of 30 s. Data files (.dta) for MS/MS spectra were generated by Bioworks Browser 3.2 EFI software (Thermo Fisher Scientific), converted to MASCOT generic format (.mgf), and searched against a proteomic database for *M. xanthus*. Peptides identified by MASCOT, with or without methyl modifications, at a significance score above 95% confidence, were validated manually. No fixed modifications were used in the searches while methionine oxidation, asparagine and glutamine deamidation, and aspartate and glutamate methylation were

Table 5. Strains and plasmids used in this study.

Strain or plasmid	Relevant feature	Source
<i>M. xanthus</i>		
DZ2	Wild type	Laboratory collection
VB197b	<i>frzCD</i> ^{Δ6-183}	Bustamante <i>et al.</i> (2004)
DZ4480	Δ <i>frzCD</i>	Bustamante <i>et al.</i> (2004)
DZ4717	Δ <i>frzCD</i> , 6His:: <i>frzCD</i> integrated in <i>crtB</i>	This study
DZ4707	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E168A} integrated in <i>crtB</i>	This study
DZ4708	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E175A} integrated in <i>crtB</i>	This study
DZ4709	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E182A} integrated in <i>crtB</i>	This study
DZ4710	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E168D} integrated in <i>crtB</i>	This study
DZ4711	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E175D} integrated in <i>crtB</i>	This study
DZ4712	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E182D} integrated in <i>crtB</i>	This study
DZ4713	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E168A E175A} integrated in <i>crtB</i>	This study
DZ4714	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E168D E175D} integrated in <i>crtB</i>	This study
DZ4718	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E168A E175A E182A} integrated in <i>crtB</i>	This study
DZ4719	Δ <i>frzCD</i> 6His:: <i>frzCD</i> ^{E168D E175D 182D} integrated in <i>crtB</i>	This study
<i>E. coli</i>		
Top10	General cloning strain	Invitrogen
Tuner	Protein expression strain	Novagen
Plasmids		
pET28a	Expression plasmid	Novagen
pAS201	pET28a with 6His:: <i>frzF</i> ^{cheR}	This study
pCT2	<i>M. xanthus</i> genomic integration plasmid	Mignot <i>et al.</i> (2007b)
pAS210	pCT2 with 6His:: <i>FrzCD</i>	This study
pAS211	pCT2 with 6His:: <i>frzCD</i> ^{E168A}	This study
pAS212	pCT2 with 6His:: <i>frzCD</i> ^{E175A}	This study
pAS213	pCT2 with 6His:: <i>frzCD</i> ^{E182A}	This study
pAS214	pCT2 with 6His:: <i>frzCD</i> ^{E168D}	This study
pAS215	pCT2 with 6His:: <i>frzCD</i> ^{E175D}	This study
pAS216	pCT2 with 6His:: <i>frzCD</i> ^{E182D}	This study
pAS217	pCT2 with 6His:: <i>frzCD</i> ^{E168A E175A}	This study
pAS218	pCT2 with 6His:: <i>frzCD</i> ^{E168D E175D}	This study
pAS221	pCT2 with 6His:: <i>frzCD</i> ^{E168A E175A E182A}	This study
pAS222	pCT2 with 6His:: <i>frzCD</i> ^{E168D E175D E182D}	This study

used as variable modifications. Three replicates of each sample were analysed.

Construction of mutants

All strains and plasmids are listed in Table 5. Point mutations were made using PCR. *M. xanthus* genomic DNA from strain DZ2 was used as a template for PCR. Oligonucleotides were prepared by operon. Complementary forward and reverse primers were used to amplify the gene with platinum HiFi Taq (Invitrogen). The primers for the coding strand are listed below with the lower case indicating the altered codon:

For *frzCD*^{E168D} catgagacgtctccacggaccagggcggccatccacg, For *frzCD*^{E168A} catgagacgtctccacggcgcagggcggccatccacg, For *frzCD*^{E175D} caggcggcggccatccacgacgaccgccaccatggaggag, For *frzCD*^{E175A} caggcggcggccatccacgaccgaccgaccgaccatggaggag, For *frzCD*^{E182D} Caccgagacgaccgccaccatggaggacctaagcagcgcgtcgccgc, For *frzCD*^{E182A} CACC ATGGAGgcgctgaagcagc. For *frzCD*^{E168D E175D} we used pAS215 as template and primer catgagacgtctccacggaccagggcggccatccacg, For *frzCD*^{E168A E175A} we used pAS212 and primer catgagacgtctccacggcgcagggcggccatccacg, For *frzCD*^{E168D E175D E182D} we used pAS218 as template and primer caccgagacgaccgccaccatggaggacctaagcagcgcgtcgccgc, For *frzCD*^{E168A E175A E182A} we used pAS217 (E168A E174A) and primer CACCATGGAGgcgCTGAAGCACGC.

Once each half of the insert was constructed the two PCR products per mutant were used as template and the primers. GATATCCAGCTGCCCGAGGAGGACGATG and GGCCAGT GCCAAGCTTCATTACTAGTCG were used to construct the complete insert. The insert was then placed into the digested (SmaI and HindIII) pCT2 (Mignot *et al.*, 2007) plasmid via the In-Fusion reaction (Clonetech). The resulting plasmids were confirmed by DNA sequencing.

Plasmids containing the wild-type *frzCD* and the point mutations were inserted into the *crtB* locus of *M. xanthus* strain Δ*frzCD* and confirmed by PCR and sequencing. In all figures 'wild type' corresponds to 6His::*frzCD* in the *crtB* locus. To induce expression of *frzCD* and the point mutants, 50 μg ml⁻¹ of anhydrotetracycline HCl (Reidel-de Haen) was used in all media.

Immunoblot analysis of FrzCD

Mycococcus xanthus strains were grown to mid-exponential phase, concentrated by centrifugation, and resuspended in 1 × SDS loading buffer lacking coloured dye. Cells were lysed by 10 s of sonication (Branson sonifier 450) on ice. Protein concentration was determined using the BCA (bicinchoninic acid) method (reagents from Pierce). Cells were resuspended to the same concentration in 2 × SDS loading buffer. Thirty micrograms of total protein were loaded per lane on

10% Tris HCl ready gels (Bio-Rad). After electrophoresis, the gel was transferred to a nitrocellulose membrane. Blots were probed with anti-FrzCD antibody as described (McCleary *et al.*, 1990) and with the antirabbit alexa fluor 680 (Molecular Probes) secondary antibody. Blots were visualized using an infrared imaging system (LiCor) and results were analysed using Odyssey software. The data were confirmed by three independent experiments done in duplicate.

Acknowledgements

We would like to acknowledge the contributions of Vivian Trang who isolated several point mutant strains. We would like to thank Lori Kohlstaedt (UC Berkeley Mass Spectrometry Facility) for mass spectrometry help and advice. We would like to thank Zusman lab and the Andrews lab past and present specifically Yuki Inclán, John Merlie, David Astling, Emilia Mauriello and Tam Mignot for many helpful suggestions and discussions throughout this work. We would like to thank Emilia Mauriello and Sophie Laurent for critically reading this manuscript. This work was supported by grants from the National Institutes of Health (GM20509) to DRZ and from the NIH/NCRR National Resource for Proteomics and Pathways (P41-18627) to PCA. AES was supported by a graduate fellowship from the National Science Foundation.

References

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990) Basic Local Alignment Search Tool. *J Mol Biol* **215**: 403–410.
- Astling, D.P. (2003) *Novel regulatory mechanisms of a chemotaxis pathway in the gliding bacterium Myxococcus xanthus*. PhD Thesis. Berkeley, CA: University of California, pp. 60–93.
- Astling, D.P., Lee, J.Y., and Zusman, D.R. (2006) Differential effects of chemoreceptor methylation-domain mutations on swarming and development in the social bacterium *Myxococcus xanthus*. *Mol Microbiol* **59**: 45–55.
- Berleman, J.E., Chumley, T., Cheung, P., and Kirby, J.R. (2006) Rippling is a predatory behavior in *Myxococcus xanthus*. *J Bacteriol* **188**: 5888–5895.
- Blackhart, B.D., and Zusman, D.R. (1985) 'Fizzy' genes of *Myxococcus xanthus* are involved in control of frequency of reversal of gliding motility. *Proc Natl Acad Sci USA* **82**: 8767–8770.
- Blatch, G.L., and Lassel, M. (1999) The tetratricopeptide repeat: a structural motif mediating protein–protein interactions. *Bioessays* **21**: 932–939.
- Boysen, A., Ellehaug, E., Julien, B., and Sogaard-Andersen, L. (2002) The DevT protein stimulates synthesis of FruA, a signal transduction protein required for fruiting body morphogenesis in *Myxococcus xanthus*. *J Bacteriol* **184**: 1540–1546.
- Bustamante, V.H., Martinez-Flores, I., Vlamakis, H.C., and Zusman, D.R. (2004) Analysis of the Frz signal transduction system of *Myxococcus xanthus* shows the importance of the conserved C-terminal region of the cytoplasmic chemoreceptor FrzCD in sensing signals. *Mol Microbiol* **53**: 1501–1513.
- Campos, J.M., Geisselsoder, J., and Zusman, D.R. (1978) Isolation of bacteriophage MX4, a generalized transducing phage for *Myxococcus xanthus*. *J Mol Biol* **119**: 167–178.
- Geng, Y., Yang, Z., Downard, J., Zusman, D., and Shi, W. (1998) Methylation of FrzCD defines a discrete step in the developmental program of *Myxococcus xanthus*. *J Bacteriol* **180**: 5765–5768.
- Hagen, D.C., Bretscher, A.P., and Kaiser, D. (1978) Synergism between morphogenetic mutants of *Myxococcus xanthus*. *Dev Biol* **64**: 284–296.
- Hodgkin, J., and Kaiser, D. (1977) Cell-to-cell stimulation of movement in nonmotile mutants of *Myxococcus*. *Proc Natl Acad Sci USA* **74**: 2938–2942.
- Igoshin, O.A., Goldbeter, A., Kaiser, D., and Oster, G. (2004) A biochemical oscillator explains several aspects of *Myxococcus xanthus* behavior during development. *Proc Natl Acad Sci USA* **101**: 15760–15765.
- Inclán, Y.F., Vlamakis, H.C., and Zusman, D.R. (2007) FrzZ, a dual CheY-like response regulator, functions as an output for the Frz chemosensory pathway of *Myxococcus xanthus*. *Mol Microbiol* **65**: 90–102.
- Kaiser, D. (2003) Coupling cell movement to multicellular development in myxobacteria. *Nat Rev Microbiol* **1**: 45–54.
- Kaiser, D. (2004) Signaling in myxobacteria. *Annu Rev Microbiol* **58**: 75–98.
- Kehry, M.R., and Dahlquist, F.W. (1982) The methyl-accepting chemotaxis proteins of *Escherichia coli*. Identification of the multiple methylation sites on methyl-accepting chemotaxis protein I. *J Biol Chem* **257**: 10378–10386.
- McBride, M.J., and Zusman, D.R. (1993) FrzCD, a methyl-accepting taxis protein from *Myxococcus xanthus*, shows modulated methylation during fruiting body formation. *J Bacteriol* **175**: 4936–4940.
- McBride, M.J., Weinberg, R.A., and Zusman, D.R. (1989) 'Fizzy' aggregation genes of the gliding bacterium *Myxococcus xanthus* show sequence similarities to the chemotaxis genes of enteric bacteria. *Proc Natl Acad Sci USA* **86**: 424–428.
- McCleary, W.R., and Zusman, D.R. (1990) Purification and characterization of the *Myxococcus xanthus* FrzE protein shows that it has autophosphorylation activity. *J Bacteriol* **172**: 6661–6668.
- McCleary, W.R., McBride, M.J., and Zusman, D.R. (1990) Developmental sensory transduction in *Myxococcus xanthus* involves methylation and demethylation of FrzCD. *J Bacteriol* **172**: 4877–4887.
- Mignot, T. (2007) The elusive engine in *Myxococcus xanthus* gliding motility. *Cell Mol Life Sci* **64**: 2733–2745.
- Mignot, T., Merlie, J.P., Jr and Zusman, D.R. (2007) Two localization motifs mediate polar residence of FrzS during cell movement and reversals of *Myxococcus xanthus*. *Mol Microbiol* **65**: 363–372.
- Nowlin, D.M., Bollinger, J., and Hazelbauer, G.L. (1987) Sites of covalent modification in Trg, a sensory transducer of *Escherichia coli*. *J Biol Chem* **262**: 6039–6045.
- Nowlin, D.M., Bollinger, J., and Hazelbauer, G.L. (1988) Site-directed mutations altering methyl-accepting residues of a sensory transducer protein. *Proteins* **3**: 102–112.
- Park, C., Dutton, D.P., and Hazelbauer, G.L. (1990) Effects of glutamines and glutamates at sites of covalent modification

- of a methyl-accepting transducer. *J Bacteriol* **172**: 7179–7187.
- Perez, E., Zheng, H., and Stock, A.M. (2006) Identification of methylation sites in *Thermotoga maritima* chemotaxis receptors. *J Bacteriol* **188**: 4093–4100.
- Rasmussen, A.A., Porter, S.L., Armitage, J.P., and Sogaard-Andersen, L. (2005) Coupling of multicellular morphogenesis and cellular differentiation by an unusual hybrid histidine protein kinase in *Myxococcus xanthus*. *Mol Microbiol* **56**: 1358–1372.
- Reichenbach, H. (1999) The ecology of the myxobacteria. *Environ Microbiol* **1**: 15–21.
- Scott, A.E. (2008) Receptor methylation controls single cell and group behaviors in the social bacterium *Myxococcus xanthus*. PhD Thesis. Berkeley, CA: University of California, p. 62.
- Shapiro, M.J., and Koshland, D.E., Jr (1994) Mutagenic studies of the interaction between the aspartate receptor and methyltransferase from *Escherichia coli*. *J Biol Chem* **269**: 11054–11059.
- Shapiro, M.J., Chakrabarti, I., and Koshland, D.E., Jr (1995) Contributions made by individual methylation sites of the *Escherichia coli* aspartate receptor to chemotactic behavior. *Proc Natl Acad Sci USA* **92**: 1053–1056.
- Shimkets, L.J. (1999) Intercellular signaling during fruiting-body development of *Myxococcus xanthus*. *Annu Rev Microbiol* **53**: 525–549.
- Shiomi, D., Zhulin, I.B., Homma, M., and Kawagishi, I. (2002) Dual recognition of the bacterial chemoreceptor by chemotaxis-specific domains of the CheR methyltransferase. *J Biol Chem* **277**: 42325–42333.
- Sogaard-Andersen, L., and Kaiser, D. (1996) C factor, a cell-surface-associated intercellular signaling protein, stimulates the cytoplasmic Frz signal transduction system in *Myxococcus xanthus*. *Proc Natl Acad Sci USA* **93**: 2675–2679.
- Stock, J.B., and Koshland, D.E., Jr (1978) A protein methyl-esterase involved in bacterial sensing. *Proc Natl Acad Sci USA* **75**: 3659–3663.
- Terwilliger, T.C., and Koshland, D.E., Jr (1984) Sites of methyl esterification and deamination on the aspartate receptor involved in chemotaxis. *J Biol Chem* **259**: 7719–7725.
- Terwilliger, T.C., Bogonez, E., Wang, E.A., and Koshland, D.E., Jr (1983) Sites of methyl esterification on the aspartate receptor involved in bacterial chemotaxis. *J Biol Chem* **258**: 9608–9611.
- Terwilliger, T.C., Wang, J.Y., and Koshland, D.E., Jr (1986) Surface structure recognized for covalent modification of the aspartate receptor in chemotaxis. *Proc Natl Acad Sci USA* **83**: 6707–6710.
- Wu, S.S., and Kaiser, D. (1995) Genetic and functional evidence that Type IV pili are required for social gliding motility. *Mol Microbiol* **18**: 547–558.
- Zusman, D.R. (1982) 'Frizzy' mutants: a new class of aggregation-defective developmental mutants of *Myxococcus xanthus*. *J Bacteriol* **150**: 1430–1437.
- Zusman, D.R., Scott, A.E., Yang, Z., and Kirby, J.R. (2007) Chemosensory pathways, motility and development in *Myxococcus xanthus*. *Nat Rev Microbiol* **5**: 862–872.

Supplementary material

This material is available as part of the online article from: <http://www.blackwell-synergy.com/doi/abs/10.1111/j.1365-2958.2008.06323.x>
(This link will take you to the article abstract).

Please note: Blackwell Publishing is not responsible for the content or functionality of any supplementary materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.