







OPEN ACCESS

Remote ischaemic conditioning in necrotising enterocolitis: a phase I feasibility and safety study

Carlos Zozaya ^{1,2}, Niloofar Ganji ², Bo Li,² Maarten Janssen Lok,² Carol Lee,² Yuhki Koike,³ Estelle Gauda,¹ Martin Offringa ⁴, Simon Eaton ⁵, Prakeshkumar S Shah,⁶ Agostino Pierro^{2,7}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/archdischild-2022-324174>).

For numbered affiliations see end of article.

Correspondence to

Dr Agostino Pierro, Division of General and Thoracic Surgery, Department of Translational Medicine, Hospital for Sick Children Research Institute, Toronto, ON M5G 1X8, Canada; agostino.pierro@sickkids.ca

Received 21 March 2022
Accepted 30 June 2022
Published Online First
8 August 2022

ABSTRACT

Objective Remote ischaemic conditioning (RIC) improves the outcome of experimental necrotising enterocolitis (NEC) by preserving intestinal microcirculation. The feasibility and safety of RIC in preterm infants with NEC are unknown. The study aimed to assess the feasibility and safety of RIC in preterm infants with suspected or confirmed NEC.

Design Phase I non-randomised pilot study conducted in three steps: step A to determine the safe duration of limb ischaemia (up to 4 min); step B to assess the safety of 4 repeated cycles of ischaemia-reperfusion at the maximum tolerated duration of ischaemia determined in step A; step C to assess the safety of applying 4 cycles of ischaemia-reperfusion on two consecutive days.

Setting Level III neonatal intensive care unit, The Hospital for Sick Children (Toronto, Canada).

Patients Fifteen preterm infants born between 22 and 33 weeks gestational age.

Intervention Four cycles of ischaemia (varying duration) applied to the limb via a manual sphygmomanometer, followed by reperfusion (4 min) and rest (5 min), repeated on two consecutive days.

Outcomes The primary outcomes were (1) feasibility defined as RIC being performed as planned in the protocol, and (2) safety defined as perfusion returning to baseline within 4 min after cuff deflation.

Results Four cycles/day of limb ischaemia (4 min) followed by reperfusion (4 min) and a 5 min gap, repeated on two consecutive days was feasible and safe in all neonates with suspected or confirmed NEC.

Conclusions This study is pivotal for designing a future randomised controlled trial to assess the efficacy of RIC in preterm infants with NEC.

Trial registration number NCT03860701.

INTRODUCTION

Necrotising enterocolitis (NEC) is one of the most common and severe gastrointestinal complications of prematurity. NEC affects 5%–10% of extremely preterm infants and is associated with high mortality (30%–50%).¹ Survivors have increased comorbidity (ie, bronchopulmonary dysplasia, postnatal growth restriction) and neurodevelopment impairment rates compared with preterm infants without NEC.^{2,3}

Current management is based on broad-spectrum antibiotics, bowel decompression and rest, analgesia, blood product transfusions and cardiovascular and respiratory support as required.¹ In case of bowel perforation or clinical deterioration despite

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ It has been demonstrated in preclinical models of NEC that remote ischaemic conditioning (RIC) effectively counteracts the intestinal damage due to NEC and prolongs survival by improving intestinal microcirculation.
- ⇒ RIC has never been investigated in preterm human neonates, the population that is at the highest risk for developing NEC.

WHAT THIS STUDY ADDS

- ⇒ This study demonstrates that it is safe, in preterm neonates with bodyweight between 840 and 2490 g, to apply four cycles of RIC each consisting of limb ischaemia (4 min) followed by reperfusion (4 min) and a 5 min gap, repeated on two consecutive days without any local or systemic adverse effects.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ This research is pivotal for the design of a randomised controlled trial aimed to assess the efficacy of RIC in preterm infants with NEC.

the implementation of adequate medical treatment, surgery is indicated.⁴ This has been the standard of care over the last three decades as there have not been significant changes in the management of NEC. However, NEC morbidity and mortality still remain high.^{5–7} Therefore, new preventive strategies and therapeutic options are urgently needed.

Remote ischaemic conditioning (RIC) is a manoeuvre that involves application of brief cycles of non-lethal ischaemia and reperfusion to a remote organ/site that conveys protection to distant organs withstanding ischaemic injury. We recently demonstrated that RIC promotes recovery from experimental NEC by reducing intestinal injury, inflammation and enhancing survival.⁸ The mechanism of action of RIC in NEC is dependent on improving intestinal microvascular blood flow by the vasodilatory action of hydrogen sulfide and nitric oxide.⁸

RIC has been applied to adults,^{9–11} children^{12–15} and neonatal patients >36 weeks of gestational age¹⁶ for different diagnoses. However, it has not been attempted in preterm neonates including those who had NEC, mostly due to concerns regarding feasibility and safety in this vulnerable population. Thus, our objective was to assess the feasibility and



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Zozaya C, Ganji N, Li B, et al. *Arch Dis Child Fetal Neonatal Ed* 2023;**108**:F69–F76.

safety of RIC in preterm infants with suspected or confirmed NEC. We hypothesised that RIC (up to four cycles of 4 min of ischaemia repeated after 24 hours) would be feasible and safe in this vulnerable population.

METHODS

Design and setting

We conducted a phase I feasibility and safety trial in the neonatal intensive care unit of the Hospital for Sick Children, Toronto (trial flow chart according to Consolidated Standards of Reporting Trials guidelines in online supplemental material). The trial was registered at ClinicalTrials.gov (NCT03860701).

Participants

Patients with suspected or confirmed NEC (stages I–III as per the modified Bell's classification)¹⁷ were eligible for this study if they weighed ≥ 750 g at the time of recruitment and were < 36 weeks of gestational age at birth. Exclusion criteria were: (1) major congenital heart disease, (2) congenital diaphragmatic hernia, (3) gastroschisis, (4) antecedent of limb ischaemia/limb thrombotic events, (5) antecedent diagnosis of occlusive arterial or venous thrombosis and (6) haemodynamic instability (defined as two inotropes/vasopressors administration, or one inotrope/vasopressor and hydrocortisone administration or one inotrope/vasopressor with dose increased in previous 6 hours).

Intervention

Limb ischaemia-reperfusion cycles were applied by placing an appropriately sized blood pressure cuff around a limb, and inflating (until blood flow distal to the cuff was interrupted with no pulse detected by pulse oximetry) and deflating it using a manual sphygmomanometer (Tyco, Welch Allyn, Skaneateles Falls, New York, USA). Before and after each ischaemia-reperfusion cycle, cutaneous integrity and perfusion (colour, capillary refill time) were assessed. The presence of grasp reflex was ascertained immediately before, right after and again 24 hours after RIC. Urine output (mL/kg/hour) during the 24 hours before and 24 hours after RIC was also recorded. Pain was evaluated 30 s after each ischaemic phase and 6 hours after the last RIC cycle using the Premature Infant Pain Profile (PIPP score).^{18,19} According to the protocol, RIC was due to be stopped if there was no return of limb perfusion on clinical assessment and/or no return of oxygen saturation to baseline values $\pm 3\%$ within 4 min in the hand or foot of the limb receiving RIC. If perfusion did not return, the limb was warmed and elevated for 10 min, followed by nitroglycerine patch application and Doppler ultrasound if necessary.

To determine tolerability of the ischaemia period up to a maximum of 4 min, as well as feasibility of applying repeated cycles of ischaemia-reperfusion on two consecutive days, the study was conducted in three steps.

Step A

Four cycles of limb ischaemia-reperfusion in 1 day, increasing the ischaemia time by 1 min in each cycle starting at 1 min in the first cycle to a maximum of 4 min in the last cycle. Each ischaemic cycle was followed by reperfusion (4 min) and rest (5 min) before the next cycle.

Step B

Four cycles of ischaemia-reperfusion in 1 day, consisting of the longest safe duration of ischaemia determined in step A, followed by reperfusion (4 min) and rest (5 min).

Step C

Four cycles of ischaemia-reperfusion as in step B but repeated on two consecutive days (24 ± 2 hours apart).

STUDY OUTCOMES

The *primary outcomes* were the feasibility and safety of RIC. Feasibility was defined as the ability to apply the RIC manoeuvre as planned in the protocol (available on request). Safety was defined as the return of limb perfusion to baseline within 4 min after cuff deflation assessed by clinical assessment of skin colour, pulse oximetry and capillary refill time.

Secondary outcomes included:

1. Peripheral nerve injury indicated by loss of grasp reflex after RIC.
2. Cutaneous injury identified as a new-onset skin breakdown, bruising, ecchymosis or petechiae within 24 hours after RIC.
3. Pain assessed by measuring the PIPP score before RIC, 30 s after each RIC cycle and 6 hours after the last cycle. This validated score ranges from 0 to 21 with 0–6 indicating no pain, 6–12 mild-to-moderate pain and > 12 severe pain.¹⁹
4. Change in urine output (mL/kg/hour) measured over 24 hours before and after RIC.

The primary and secondary outcomes after each study step (A, B or C) were reviewed anonymously by an independent safety panel of experts including two neonatologists, one paediatric surgeon, one cardiologist and one nurse practitioner, not involved in the circle of care, trial design and execution. This panel determined whether it was appropriate (i) to proceed to the next study step, (ii) recruit more patients in the same step and (iii) stop the trial altogether.

Statistical methods

Data are presented as median (range) and n (%) as indicated. A two-way analysis of variance with main effects only and post hoc Tukey's multiple comparisons test was used to compare pre-RIC and post-RIC oxygen saturation, capillary refill time (s) and PIPP score. A paired non-parametric Wilcoxon test was used to compare pre-RIC and post-RIC urine output (mL/kg/hour). P value < 0.05 was considered statistically significant. GraphPad Prism 8 (GraphPad Software, San Diego, California, USA) was used for data analysis and presentation.

RESULTS

Fifteen neonates were recruited, five in each step of the study (tables 1 and 2). All neonates were preterm with a birth weight ranging from 490 to 2010 g. NEC was diagnosed between 7 and 58 days and RIC was performed after a median of 4 days (2–33) from NEC diagnosis. Table 1 reports patients' demographics and NEC Bell's stage at the time of RIC. Thirteen (87%) patients were mechanically ventilated at the time of RIC. Five (33%) received medical treatment and 10 (67%) required an operation for NEC.

The independent panel of experts confirmed the feasibility and safety after each study step, did not require recruitment of further patients and did not stop the trial at any step.

Primary outcomes

RIC was completed according to the protocol in all neonates (14 in the arm and 1 in the leg). Arterial flow and limb perfusion returned to baseline within 4 min in all patients, as confirmed by clinical assessment of limb, return of pulse waveform displayed by pulse-oximetry, return to peripheral oxygen saturation to pre-RIC value (figures 1A–3A and figure 3E) and capillary refill time

Table 1 Characteristics of the study population

Demographics	Step A (n=5)	Step B (n=5)	Step C (n=5)
Sex (male/female)	4/1	3/2	2/3
Gestational age (weeks+days)	27.9 (24.1–33.1)	24.1 (23.4–25)	25.3 (22.6–26.4)
Gestational age <28 weeks (n, %)	3 (60)	5 (100)	5 (100)
Birth weight (g)	1193 (711–2010)	610 (490–750)	700 (503–740)
Birth weight <1000 g (n, %)	1 (20)	5 (100)	5 (100)
Age at diagnosis (days)	31 (7–56)	24 (14–42)	17 (7–58)
Weight at RIC (g)	1770 (1143–2490)	951 (840–1064)	1435 (937–1597)
Postnatal age at RIC (days)	34 (9–59)	28 (26–44)	43 (19–65)
Invasive mechanical ventilation at the time of RIC	3 (60)	5 (100)	5 (100)
Inotropic support at the time of RIC	1 (20)	0	0
NEC Bell's stage at the time of RIC (n, %)			
I	1 (20)	1 (20)	0 (0)
II	1 (20)	1 (20)	1 (20)
III	3 (60)	3 (60)	4 (80)
NEC treatment (n, %)			
Medical	3 (60)	1 (20)	1 (20)
Surgical	2 (40)	4 (80)	4 (80)
Death (n)	0	1	1

Data are expressed as median (range) or number of patients (percentage), depending on the variable.
NEC, necrotising enterocolitis; RIC, remote ischaemic conditioning.

preservation (figures 1B–3B and 3F). Two enrolled patients died before discharge; however, none of the deaths were attributed to the RIC procedure. One patient died due to septic shock 4 days after RIC and the other died due to multiorgan failure 2 months after RIC.

Secondary outcomes

All patients had a normal grasp reflex both before and after RIC. No new-onset skin lesions (breakdown, bruising or ecchymosis)

were observed. PIPP scores obtained before and after RIC are reported in figures 1C–3C and 3G. There was no significant difference in PIPP score comparing its value before and 6 hours after RIC (table 2). Compared with the assessment before RIC, the PIPP score did not change from 'no pain' or 'mild-to-moderate' to 'severe' pain 6 hours after the last ischaemic cycle. The PIPP score in five neonates (33%) increased from 'no pain' to 'mild-to-moderate' pain, and in two neonates (13%) decreased from 'mild-to-moderate' pain to 'no pain'. No patient

Table 2 Safety outcomes of the RIC manoeuvre

Outcomes	Step A (n=5)	Step B (n=5)	Step C (n=5)	
			Day 1	Day 2
Oxygen saturation (%)*				
Pre-RIC	98 (97–99)	95 (94–98)	95 (92–100)	91 (91–93)
Post-RIC	96 (95–99)	96 (95–100)	100 (93–100)	94 (92–100)
Change (pre-post RIC)†	2 (–1 to 4)	–2 (–4 to 3)	–1 (–5 to 1)	–2 (–9 to 0)
Capillary refill time (s)*				
Pre-RIC	3 (1–3)	3 (2–3)	3 (2–4)	3 (2–3)
Post-RIC	3 (2–4)	3 (2–4)	3 (2–4)	3 (2–3)
Change (pre-post RIC)†	0 (–2 to 0)	0 (0 to 1)	0 (0 to 1)	0
PIPP score‡				
Pre-RIC	3 (2–7)	5 (2–9)	5 (3–11)	4 (4–7)
Post-RIC	5 (4–6)	7 (3–10)	6 (2–9)	7 (3–8)
Change (pre-post RIC)†	–1 (–4 to 2)	–1 (–7 to 1)	0 (–1 to 2)	–1 (–4 to 4)
Urine output (mL/kg/hour)				
24 hours pre-RIC	3.5 (0.6–8)	4.7 (3.2–6)	5.2 (3.4–14.7)	4.3 (3.4–5.4)
24 hours post-RIC	4.8 (3.1–6.4)	5 (3.7–6.5)	4.3 (3.7–5.4)	3.5 (2–4.8)
Change (pre-post RIC)†	1.9 (–3.7 to 5.8)	0.5 (0.2 to 1.2)	–0.1 (–10.4 to 0.2)	0.8 (–1 to 3.5)

Data are expressed as median (range).

*Oxygen saturation (%) and capillary refill time (s) were measured before the first cycle of ischaemia (pre-RIC) and during the 4 min reperfusion time following the last cycle of ischaemia (post-RIC).

†Change was calculated as pre-RIC minus post-RIC values.

‡PIPP score determined before RIC and 6 hours after the last cycle.

PIPP, Premature Infant Pain Profile; RIC, remote ischaemic conditioning.




Figure 1 Physiological variables recorded to assess the safety of remote ischaemic conditioning (RIC) in step A (n=5). (A) Peripheral oxygen (O₂) saturation and (B) capillary refill time (CRT in s) were assessed before and after every cycle of ischaemia. (C) Pain was assessed according to the Premature Infant Pain Profile (PIPP score) and measured before the first RIC cycle, 30 s after each cycle and 6 hours after the last cycle of ischaemia. (D) Urine output (mL/kg/hour) was measured 24 hours before and after RIC. Data are represented as median with range. A two-way analysis of variance with main effects only and post hoc Tukey's multiple comparisons test was used to compare pre-RIC and post-RIC oxygen saturation, CRT (s) and PIPP score. A paired non-parametric Wilcoxon test was used to compare pre-RIC and post-RIC urine output (mL/kg/hour). There was no significant difference between pre-RIC and post-RIC values of any of the listed parameters.

became oligo-anuric (defined as urine output <1 mL/kg/hour) after RIC (figures 1D–3D and 3H, table 2).

DISCUSSION

In this phase I feasibility and safety trial, we demonstrated that RIC consisting of four consecutive cycles of 4 min ischaemia followed by 4 min reperfusion on two consecutive days is feasible in preterm infants with NEC. No significant adverse events related to RIC were detected. Reperfusion occurred promptly after RIC in all patients without any impairment in vascular flow and no cutaneous, nor nerve injury was observed.

To our knowledge, RIC has not been evaluated in preterm infants with NEC. Eighty-seven per cent of enrolled neonates in this phase I trial were born before 28 weeks gestation and 73% were extremely low birth weight (<1000 g), representing the population at higher risk of NEC. In this study, we assessed the feasibility and safety of RIC in fragile preterm neonates with either suspected or defined medical and surgical NEC, suggesting that RIC is safe and can be applied even during acute illness.

Prematurity and enteral formula feeding are two of the most important risk factors for NEC.^{1 8 20 21} Additionally, ischaemia plays a role in both NEC pathogenesis and pathophysiology.^{20–22}

Physiologically, intestinal blood flow is known to increase above baseline to fulfil the increased intestinal oxygen demand after feeds.²⁰ However, postprandial hyperaemia is remarkably compromised in premature infants that later developed NEC,²³ and we have demonstrated that NEC in human preterm infants is associated with impairments in intestinal microcirculation.^{8 20} Furthermore, NEC risk is higher in infants with intrauterine growth restriction and circulatory impairment, respiratory distress syndrome and associated hypoxia requiring surfactant treatment or early onset sepsis.²⁴ In addition, persistent ductus arteriosus can lead to bowel hypoperfusion²⁵ and is also associated with NEC.²⁶ Previous studies on splanchnic circulation using near-infrared spectroscopy showed that regional oxygen saturation is lower, especially after feeds in babies who later developed NEC.^{22 27} Lastly, at the disease onset, local bowel inflammation is associated with impaired microcirculation which leads to bowel wall ischaemia and eventually transmural necrosis.²⁸ In fact, patched coagulative necrosis (a marker of ischaemia) is the main pathological finding in NEC patients, and hypoxia markers like hypoxia-inducible factor 1 α and glucose transporter 1 are elevated in NEC intestinal resection specimens.²⁹




Figure 2 Physiological variables recorded to assess the safety of remote ischaemic conditioning (RIC) in step B (n=5). (A) Peripheral oxygen (O₂) saturation and (B) capillary refill time (CRT in s) were assessed before and after every cycle of ischaemia. (C) Pain was assessed according to the Premature Infant Pain Profile (PIPP score) and measured before the first RIC cycle, 30s after each cycle and 6 hours after the last cycle of ischaemia. (D) Urine output (mL/kg/hour) was measured 24 hours before and after RIC. Data are represented as median with range. A two-way analysis of variance with main effects only and post hoc Tukey's multiple comparisons test was used to compare pre-RIC and post-RIC oxygen saturation, CRT (s) and PIPP score. A paired non-parametric Wilcoxon test was used to compare pre-RIC and post-RIC urine output (mL/kg/hour). There was no significant difference between pre-RIC and post-RIC values of any of the listed parameters.

Considering the role of ischaemia in NEC, modulating the immature intestinal microvasculature could prove to be a viable strategy to counteract the feeding-induced hypoxia and prevent the progression of NEC. For this reason, we evaluated RIC, a manoeuvre which can improve the blood supply to the intestine. RIC involves application of brief cycles of ischaemia and reperfusion to a limb to protect distant organs from sustained ischaemic damage—in this case the intestine. Prior to the current trial, experimental studies performed by our group demonstrated that RIC induced endothelium-dependent vasodilation, preserved bowel microcirculation and reduced gut injury and mortality due to NEC.⁸

RIC implies inflating a standard blood pressure cuff around a limb, aiming to interrupt distal arterial blood supply to produce transient skeletal muscle ischaemia. To assess safety of this procedure in the study population, potential complications/side effects were monitored including limb ischaemia, peripheral nerve damage, pain and change in urinary output.

First, limb ischaemia is a potential complication of peripheral or central arterial lines in neonates. Vascular obstruction is known to present as a pale and poorly perfused limb distal to the occluded point. We assumed RIC could have induced persistent

limb ischaemia after cuff deflation secondary to arterial thrombosis or vasospasm. To prove this was not the case, limb perfusion was continuously monitored during the procedure and cautiously assessed after RIC, both visually and with pulse oximetry. During RIC, the limb became dusky (venous congestion), and the pulse wave was lost due to total arterial compression as expected and desired. Once the blood pressure cuff was deflated, we observed rapid reperfusion, usually with evident hyperaemia in visual inspection. Capillary refill time was assessed before and after every ischaemia cycle to quantify limb perfusion and always recovered within 4 min.

Second, to assess peripheral nerve integrity, palmar or plantar grasp reflex (depending on the limb used) was assessed before and after RIC as it is a useful clinical marker of potential nerve ischaemic injury involving both sensorial and motor fibres.²⁹ The grasp reflex was preserved in all patients.

Third, RIC requires limb compression with pressure above systolic pressure affecting venous blood return and potentially leading to microhaemorrhage (petechiae, ecchymosis, bruising) and/or skin breakdown in this vulnerable patient population. However, none of the neonates developed cutaneous bleeding or changes in skin integrity after RIC.




Figure 3 Physiological variables recorded to assess the safety of remote ischaemic conditioning (RIC) in step C day 1 and 2 (n=5 in each day). (A, E) Peripheral oxygen (O₂) saturation and (B, F) capillary refill time (CRT in s) were assessed before and after every cycle of ischaemia. The day 2 CRT for all patients is reported, although lines are superimposed. (C, G) Pain was assessed according to the Premature Infant Pain Profile (PIPP score) and measured before the first RIC cycle, 30 s after each cycle and 6 hours after the last cycle of ischaemia. (D, H) Urine output (mL/kg/hour) was measured 24 hours before and after RIC. Data are represented as median with range. A two-way analysis of variance with main effects only and post hoc Tukey's multiple comparisons test was used to compare pre-RIC and post-RIC oxygen saturation, CRT (s) and PIPP score. A paired non-parametric Wilcoxon test was used to compare pre-RIC and post-RIC urine output (mL/kg/hour). There was no significant difference between pre-RIC and post-RIC values of any of the listed parameters.

Forth, pain has also been related to worsening neurodevelopment in preterm infants.³⁰ NEC is a cause of pain, making it challenging to separate pain due to NEC from that due to RIC. To add objectivity to our prospective evaluation, pain was assessed before and after RIC using the validated PIPP score.¹⁹ There was no development of persistent or severe pain after RIC, and in none of the neonates, the RIC manoeuvre was discontinued due to pain.

At last, in theory, muscle ischaemia could have led to muscle necrosis and rhabdomyolysis.³¹ The most severe potential complication of rhabdomyolysis is oliguric acute kidney injury secondary to renal tubules obstruction with myoglobin affecting >50% of them. We did not find any difference in urine output 24 hours before and after RIC and none of the neonates studied became oligo-anuric after RIC. Moreover, without significant renal impairment, rhabdomyolysis is a self-resolving transient condition.

Two of our patients died, none of them of causes directly related to RIC but due to the underlying disease. Indeed, the study's mortality was not higher than the mortality of NEC in preterm infants. Although there is no evidence of harm in this phase I preliminary trial, there may be beneficial or detrimental effects that can only become apparent in larger phase II–III trials. We acknowledge that the small sample size of the present study, although typical of phase I trials, did not allow us to rule out other potential complications. Specifically, the possibility that reactive oxygen metabolites can cause reperfusion injury in other organs and may worsen chronic lung disease, retinopathy of prematurity or other complications of prematurity, as well as potential effects of RIC on long-term neurological development will have to be assessed in future phase II and III trials.

RIC is simply performed by inflating and deflating a blood pressure cuff, similar to blood pressure measurement, and can be performed with minimal invasiveness and cost in preterm neonates with NEC, making it potentially applicable worldwide.

CONCLUSIONS

In this study, we demonstrated that RIC is feasible and safe in small and fragile preterm neonates with NEC. These novel findings are essential to embark on a multicentre randomised controlled trial to investigate the efficacy of RIC in reducing NEC progression, morbidity and mortality.

Author affiliations

¹Division of Neonatology, The Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada

²Translational Medicine, Hospital for Sick Children Research Institute, University of Toronto, Toronto, Ontario, Canada

³Departments of Gastrointestinal and Pediatric Surgery, Mie University, Tsu, Mie, Japan

⁴Child Health Evaluative Sciences, The Hospital for Sick Children, Toronto, Ontario, Canada

⁵Paediatric Surgery, Institute of Child Health, Great Ormond Street Hospital for Children, London, UK

⁶Department of Paediatrics, Mount Sinai Hospital, Toronto, Ontario, Canada

⁷Division of General and Thoracic Surgery, The Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada

Correction notice This article has been updated since it was published online. An Orcid ID has been added to one of the authors.

Acknowledgements We are very grateful to the members of the independent safety panel of experts who reviewed the results after every step of this study: Luc Mertens MD, PhD (Division of Cardiology, The Hospital for Sick Children, Toronto, Ontario, Canada); Yenge Diambomba MD, FRCP (Department of Pediatrics, Neonatal Intensive Care Unit, Mount Sinai Hospital, Toronto, Ontario, Canada); Eugene Ng, MD, FRCP, FAAP (Department of Newborn and Developmental Paediatrics, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada); David Wesson MD

(Pediatric Surgery Division, Texas Children's Hospital, Houston, Texas, USA) and Irina Pashynskyy MScN, NP-PhC (Division of General and Thoracic Surgery, Toronto, Ontario, Canada).

Contributors All authors approved the final manuscript as submitted and agree to be for all accountable aspects of the work. CZ was involved in the study design, performing the clinical procedures, acquisition of data, interpretation of data and writing of the paper. NG, MJL, BL, CL, YK, MO and EG were involved in interpreting the data and writing of the paper. SE and PSS were involved in the study's design, interpretation of data and paper writing. AP was involved in the conceptualisation of study hypothesis, design of the study, interpretation of data, writing of the paper and study supervision, and is responsible for the overall content as the guarantor.

Funding SE acknowledges support from the NIHR Biomedical Research Centre at Great Ormond Street, London, UK. NG was the recipient of Restracom Scholarship from the Hospital for Sick Children, Toronto, Ontario, Canada. AP was supported by the Canadian Institutes of Health Research (CIHR) Foundation Grant 353857. The other authors received no additional funding. The listed funding sources had no role in the design and conduct of the study.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study was approved by Institutional Research Ethics Board (REB) at the Hospital for Sick Children (REB: 1000061144). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Individual deidentified participant data will not be made available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Carlos Zozaya <http://orcid.org/0000-0003-0062-9258>

Niloofar Ganji <http://orcid.org/0000-0002-6705-2812>

Martin Offringa <http://orcid.org/0000-0002-4402-5299>

Simon Eaton <http://orcid.org/0000-0003-0892-9204>

REFERENCES

- 1 Neu J, Walker WA. Necrotizing enterocolitis. *N Engl J Med* 2011;364:255–64.
- 2 Rees CM, Pierro A, Eaton S. Neurodevelopmental outcomes of neonates with medically and surgically treated necrotizing enterocolitis. *Arch Dis Child Fetal Neonatal Ed* 2007;92:F193–8.
- 3 Zozaya C, Avila-Alvarez A, Arruza L, et al. The effect of morbidity and sex on postnatal growth of very preterm infants: a multicenter cohort study. *Neonatology* 2019;115:348–54.
- 4 Zozaya C, García González I, Avila-Alvarez A, et al. Incidence, treatment, and outcome trends of necrotizing enterocolitis in preterm infants: a multicenter cohort study. *Front Pediatr* 2020;8:188.
- 5 Shah PS, Dunn M, Aziz K, et al. Sustained quality improvement in outcomes of preterm neonates with a gestational age less than 29 weeks: results from the Evidence-based Practice for Improving Quality Phase 3¹. *Can J Physiol Pharmacol* 2019;97:213–21.
- 6 Stoll BJ, Hansen NI, Bell EF, et al. Trends in care practices, morbidity, and mortality of extremely preterm neonates, 1993–2012. *JAMA* 2015;314:1039–51.
- 7 Luig M, Lui K, NSW & ACT NICUS Group. Epidemiology of necrotizing enterocolitis--Part I: Changing regional trends in extremely preterm infants over 14 years. *J Paediatr Child Health* 2005;41:169–73.
- 8 Koike Y, Li B, Ganji N, et al. Remote ischemic conditioning counteracts the intestinal damage of necrotizing enterocolitis by improving intestinal microcirculation. *Nat Commun* 2020;11:1–15.
- 9 Eitel I, Stiermaier T, Rommel KP, et al. Cardioprotection by combined intrahospital remote ischaemic preconditioning and postconditioning in ST-elevation myocardial infarction: the randomized LIPSI conditioning trial. *Eur Heart J* 2015;36:3049–57.

- 10 Gaspar A, Lourenço AP, Pereira Miguel Álvares, *et al.* Randomized controlled trial of remote ischaemic conditioning in ST-elevation myocardial infarction as adjuvant to primary angioplasty (RIC-STEMI). *Basic Res Cardiol* 2018;113:14.
- 11 Bøtker HE, Kharbanda R, Schmidt MR, *et al.* Remote ischaemic conditioning before hospital admission, as a complement to angioplasty, and effect on myocardial salvage in patients with acute myocardial infarction: a randomised trial. *Lancet* 2010;375:727–34.
- 12 Guerra GG, Joffe AR, Seal R, *et al.* Pilot randomized controlled trial on early and late remote ischemic preconditioning prior to complex cardiac surgery in young infants. *Paediatr Anaesth* 2017;27:433–41.
- 13 Pedersen KR, Ravn HB, Povlsen JV, *et al.* Failure of remote ischemic preconditioning to reduce the risk of postoperative acute kidney injury in children undergoing operation for complex congenital heart disease: a randomized single-center study. *J Thorac Cardiovasc Surg* 2012;143:576–83.
- 14 McCrindle BW, Clarizia NA, Khaikin S, *et al.* Remote ischemic preconditioning in children undergoing cardiac surgery with cardiopulmonary bypass: a single-center double-blinded randomized trial. *J Am Heart Assoc* 2014;3. doi:10.1161/JAHA.114.000964. [Epub ahead of print: 28 Jul 2014].
- 15 Zhou W, Zeng D, Chen R, *et al.* Limb ischemic preconditioning reduces heart and lung injury after an open heart operation in infants. *Pediatr Cardiol* 2010;31:22–9.
- 16 Jones BO, Pepe S, Sheeran FL, *et al.* Remote ischemic preconditioning in cyanosed neonates undergoing cardiopulmonary bypass: a randomized controlled trial. *J Thorac Cardiovasc Surg* 2013;146:1334–40.
- 17 Walsh MC, Kliegman RM. Necrotizing enterocolitis: treatment based on staging criteria. *Pediatr Clin North Am* 1986;33:179–201.
- 18 Stevens B, Johnston C, Petryshen P, *et al.* Premature infant pain profile: development and initial validation. *Clin J Pain* 1996;12:13–22.
- 19 McNair C, Ballantyne M, Dionne K, *et al.* Postoperative pain assessment in the neonatal intensive care unit. *Arch Dis Child Fetal Neonatal Ed* 2004;89:F537–41.
- 20 Chen Y, Koike Y, Miyake H, *et al.* Formula feeding and systemic hypoxia synergistically induce intestinal hypoxia in experimental necrotizing enterocolitis. *Pediatr Surg Int* 2016;32:1115–9.
- 21 Chen Y, Koike Y, Chi L, *et al.* Formula feeding and immature gut microcirculation promote intestinal hypoxia, leading to necrotizing enterocolitis. *Dis Model Mech* 2019;12. doi:10.1242/dmm.040998. [Epub ahead of print: 09 12 2019].
- 22 Patel AK, Lazar DA, Burrin DG, *et al.* Abdominal near-infrared spectroscopy measurements are lower in preterm infants at risk for necrotizing enterocolitis. *Pediatr Crit Care Med* 2014;15:735–41.
- 23 Gamsu HR, Kempley ST. Enteral hypoxia/ischaemia and necrotizing enterocolitis. *Seminars in Neonatology* 1997;2:245–54.
- 24 Rose AT, Patel RM. A critical analysis of risk factors for necrotizing enterocolitis. *Semin Fetal Neonatal Med* 2018;23:374–9.
- 25 Ledo A, Aguar M, Núñez-Ramiro A, *et al.* Abdominal near-infrared spectroscopy detects low mesenteric perfusion early in preterm infants with hemodynamic significant ductus arteriosus. *Neonatology* 2017;112:238–45.
- 26 Dollberg S, Lusky A, Reichman B. Patent ductus arteriosus, indomethacin and necrotizing enterocolitis in very low birth weight infants: a population-based study. *J Pediatr Gastroenterol Nutr* 2005;40:184–8.
- 27 Kuik SJ, van Zoonen AGJF, Bos AF, *et al.* The effect of enteral bolus feeding on regional intestinal oxygen saturation in preterm infants is age-dependent: a longitudinal observational study. *BMC Pediatr* 2019;19:404.
- 28 Nowicki PT. Ischemia and necrotizing enterocolitis: where, when, and how. *Semin Pediatr Surg* 2005;14:152–8.
- 29 Salandy S, Rai R, Gutierrez S, *et al.* Neurological examination of the infant: a comprehensive review. *Clin Anat* 2019;32:770–7.
- 30 Vinall J, Grunau RE. Impact of repeated procedural pain-related stress in infants born very preterm. *Pediatr Res* 2014;75:584–7.
- 31 Vanholder R, Sever MS, Ereik E, *et al.* Rhabdomyolysis. *J Am Soc Nephrol* 2000;11:1553–61.