

# Combined Interhemispheric and Transsylvian Approach for Resection of Craniopharyngioma

Tomohiro Inoue<sup>1</sup> Hideaki Ono<sup>2</sup> Akira Tamura<sup>2</sup> Isamu Saito<sup>2</sup>

<sup>1</sup>Department of Neurosurgery, NTT Medical Center Tokyo, Tokyo, Japan

<sup>2</sup>Department of Neurosurgery, Fuji Brain Institute and Hospital, Fujinomiya Shi, Japan

Address for correspondence Tomohiro Inoue, MD, PhD, Department of Neurosurgery, NTT Medical Center Tokyo, 5-9-22 Higashi Gotanda Shinakawa-ku, Tokyo 141-0022, Japan (e-mail: t.inouenttsu@gmail.com).

J Neurol Surg B 2018;79(suppl S3):S251.

## Abstract

### Keywords

- ▶ interhemispheric approach
- ▶ lamina terminalis
- ▶ pterional approach
- ▶ craniopharyngioma

We present a 37-year-old male case of cystic suprasellar huge craniopharyngioma, who presented with significant memory disturbance due to obstructive hydrocephalus. Combined interhemispheric and pterional approach was chosen to resect huge suprasellar tumor. Interhemispheric trans-lamina terminalis approach was quite effective to resect third ventricular tumor, while pterional approach was useful to dissect tumor out of basilar perforators and stalk.

The link to the video can be found at: <https://youtu.be/BoYIPa96kdo>.

### Conflict of Interest

None.



[www.thieme.com/skullbasevideos](http://www.thieme.com/skullbasevideos)

[www.thieme.com/jnlsbvideos](http://www.thieme.com/jnlsbvideos)

received  
September 18, 2017  
accepted  
December 18, 2017  
published online  
February 20, 2018

DOI <https://doi.org/10.1055/s-0038-1625968>.  
ISSN 2193-6331.

© 2018 Georg Thieme Verlag KG  
Stuttgart · New York

License terms

