



Predicting adverse outcomes in oncological nephrectomy: let's take a look at the patient around the tumor

Pedro F. S. Freitas^{1^}, Rafael S. R. Vidal¹, Carlos H. Watanabe¹, Marco A. Arap²

¹Urologic Oncology Division, Hospital Sírio-Libanês, Brasília, DF, Brazil; ²Urologic Oncology Division, Hospital Sírio-Libanês, São Paulo, SP, Brazil

Correspondence to: Pedro F. S. Freitas, MD. Urologic Oncology Division, Hospital Sírio-Libanês Brasília. SGAS 613, s/n, Lote 94, Asa Sul. Brasília, DF, Brazil. Email: pedro.freitas05@gmail.com.

Comment on: Maurits JSF, Sedelaar JPM, Aben KKH, *et al.* The association of body composition with postoperative complications and length of hospital stay after radical or partial nephrectomy in patients with renal cell cancer: a multicenter population-based cohort study. *Transl Androl Urol* 2022;11:1667-79.

Keywords: Renal cell carcinoma (RCC); nephrectomy; obesity; body composition; skeletal muscle

Submitted Dec 03, 2022. Accepted for publication Dec 19, 2022. Published online Jan 09, 2023.

doi: 10.21037/tau-22-803

View this article at: <https://dx.doi.org/10.21037/tau-22-803>

The management of renal cell carcinoma (RCC) underwent a revolution in the last decades. From an exclusively surgical alternative—open radical nephrectomy, we have seen the rise of ablation and surveillance, which ultimately made their way into the guidelines (1). On the surgical front, there was a shift towards nephron-sparing and minimally invasive procedures, even for larger and more challenging cases (2). At the same time, we are operating on older and more frail patients, often presenting with locally advanced and metastatic disease. Finally, there was a 50% increase in the overall incidence of RCC between 1996 and 2016 in the United States (3).

Therefore, clinical decision-making for patients with RCC has grown in complexity. To refine conventional risk assessment comprising clinical tumor, node, metastasis (TNM) staging, performance status, and patient comorbidities, tumor scores like the RENAL or Padua were developed to estimate the difficulty and risks involved in tumor resection, while improving uniformity of studies on partial nephrectomy (4,5). Still, there remains a gap of unexplained variation in the occurrence of adverse outcomes, which are often missed by conventional evaluation, with potential implications for the management of RCC (6).

The study presented by Maurits *et al.* (7) on this issue of *Translational Andrology and Urology* addresses the equally important, and less investigated, determinant of nephrectomy outcomes: patient features. More specifically, they investigated whether body composition variables—skeletal muscle density (SMD), body mass index (BMI), and adipose tissue distribution—impact on postoperative morbidity of RCC patients.

Nutritional status is a key determinant of perioperative complications, as both obesity (8) and malnutrition (9) may negatively impact on surgical outcomes of cancer patients. The challenge, however, is to evaluate them in a standard, simple, and readily available way that predicts adverse outcomes, making it useful in clinical practice. Even though the picture is evident in the extremes of cachexia and morbid obesity (BMI >40 kg/m²), most RCC patients present within intermediate nutritional categories, whose evaluation is further complicated by the fact that obesity can hide sarcopenia (10).

The authors evaluated SMD and adipose tissue by the BMI and its distribution in the subcutaneous (SAT) and visceral compartments (VAT) in patients undergoing radical or partial nephrectomy for RCC. They sought to verify whether these CT-derived variables had any

[^] ORCID: 0000-0003-4497-3923.

impact on surgical outcomes, namely major postoperative complications (Clavien-Dindo 3 or higher) and prolonged length of hospital stay (LOHS), defined as ≥ 7 days.

This large, retrospective multicenter series found obesity significantly associated with major postoperative complications. On the other hand, CT-based body composition variables concerning skeletal muscle (SMD and skeletal muscle index) and adipose tissue (VAT, SAT) showed no significant association with major complications after radical or partial nephrectomy.

The impact of obesity on nephrectomy for RCC has been extensively studied, especially with the advent of laparoscopy. Initially regarded as a contraindication to this technique, subsequent series reported no significant increase in warm ischemia time, blood loss, positive margins, complications, and length of stay (LOS) for obese patients undergoing partial or radical nephrectomy (11-15). Indeed, overweight and class I obesity have been associated with *better* surgical and oncologic outcomes, resulting in the so-called 'obesity paradox' (10). Such paradox consists in that surgical complications (13) and cancer specific survival (16) do not differ from the general population despite obesity being a recognized risk factor for RCC and obese patients presenting with more comorbidities (14).

The reason for Maurits *et al.*'s (7) observation of increased major postoperative complications among obese patients may be found in their high rate of radical nephrectomy (78%), especially considering that 42% were performed for pT1 tumors, and in about a third the access was open. Although compatible with clinical practice at the study period [2008–2012], these features differ from the series that have reported oncological nephrectomy to be safe in obese patients based on nephron sparing and minimally invasive surgery (11-15).

BMI's inconsistency to accurately determine prognosis in oncological nephrectomy population has fueled research on alternative surrogates, such as the distribution of adipose tissue. The literature yields conflicting observations on this topic: while some studies report increased adverse outcomes (12,17), others (11,18) found no association with high visceral or perinephric fat on CT.

The lack of significant association between SAT, and especially visceral adipose tissue with complications in this (7) and previous studies (11,18) leaves the value of this marker to be determined. Accordingly, skeletal muscle and adipose tissue indexes failed to improve the accuracy of a model based on established predictors of major complications [age, stage, and American Society of

Anesthesiologists (ASA) score] as reported by Maurits *et al.* (7). In their series, 78% of patients undergoing radical nephrectomy harbored a pT1 tumor, suggesting that the utility of CT-based body composition analysis in the general RCC population RCC is limited by its heterogeneity and the low number of major complications in the study population.

On the other hand, the same authors (7) showed a high SMD at the level of L3 vertebrae on CT scan to be associated with a 38% reduction in the risk of prolonged hospital stay. Interestingly, the adipose tissue parameters (VAT, SAT, and BMI) did not significantly affect LOHS in the study population.

Skeletal muscle and its deficiency (sarcopenia) are promising prognostic markers, which have been associated with adverse surgical and clinical outcomes in other malignancies (19,20). They have emerged not only as indicators of nutritional status, but low skeletal muscle radiodensity has also been linked with increased postoperative systemic inflammation (21) and dose-limiting toxicity in systemic therapy for metastatic disease (6). The few studies investigating sarcopenia's role in RCC report increased rates of major complications in nephrectomy for locally advanced disease (22), and worse overall survival in cytoreductive nephrectomy (23). The present study (7) adds to the existing evidence by showing increased LOS in patients with low SMD. Interestingly, low SMD was associated with major complications in subgroups of patients undergoing radical (but not partial), and laparoscopic/robot-assisted (but not open) nephrectomy (7).

Finally, body composition assessment is plagued with disagreement on measurement criteria and reference values in the general and oncologic populations (6). CT-based SMD has been measured either as total psoas muscle area (22) or SMD index, which comprises all muscle at the level of L3 vertebrae (16,23). Whereas the former is simple and can be measured by non-radiologists, the latter is more accurate but requires specific software and a specialized reader. No consensus has been reached so far; to illustrate one of the many issues to be overcome before such tools be adopted into clinical practice.

Still, the present study (7) connects with previous research on locally advanced (22), and cytoreductive nephrectomy (23) to suggest a possible role for CT-based body composition analysis in subgroups with such an elevated risk that it becomes clinically significant. Patients with vena cava thrombus, metastatic disease, and those >80 years old are at particular risk of adverse outcomes, for

whom we often hesitate to propose surgery. Future studies on CT-based body composition variables should focus on these patients for whom nephrectomy yields a narrow risk-benefit ratio, preferably relying on prospectively collected imaging and multicenter collaboration to overcome their low frequency.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Translational Andrology and Urology*. The article did not undergo external peer review.

Conflicts of Interest: The authors have completed the ICMJE uniform disclosure form (available at <https://tau.amegroups.com/article/view/10.21037/tau-22-803/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Ljungberg B, Albiges L, Abu-Ghanem Y, et al. European Association of Urology Guidelines on Renal Cell Carcinoma: The 2022 Update. *Eur Urol* 2022;82:399-410.
2. Banegas MP, Harlan LC, Mann B, et al. Toward greater adoption of minimally invasive and nephron-sparing surgical techniques for renal cell cancer in the United States. *Urol Oncol* 2016;34:433.e9-433.e17.
3. Ries L, Melbert D, Krapcho M. SEER cancer statistics review 1975–2012. Bethesda, MD: NCI; 2014.
4. Kutikov A, Uzzo RG. The R.E.N.A.L. nephrometry score: a comprehensive standardized system for quantitating renal tumor size, location and depth. *J Urol* 2009;182:844-53.
5. Ficarra V, Novara G, Secco S, et al. Preoperative aspects and dimensions used for an anatomical (PADUA) classification of renal tumours in patients who are candidates for nephron-sparing surgery. *Eur Urol* 2009;56:786-93.
6. Vrieling A, Kampman E, Knijnenburg NC, et al. Body Composition in Relation to Clinical Outcomes in Renal Cell Cancer: A Systematic Review and Meta-analysis. *Eur Urol Focus* 2018;4:420-34.
7. Maurits JSE, Sedelaar JPM, Aben KKH, et al. The association of body composition with postoperative complications and length of hospital stay after radical or partial nephrectomy in patients with renal cell cancer: a multicenter population-based cohort study. *Transl Androl Urol* 2022;11:1667-79.
8. Gong EM, Orvieto MA, Lyon MB, et al. Analysis of impact of body mass index on outcomes of laparoscopic renal surgery. *Urology* 2007;69:38-43.
9. Morgan TM, Tang D, Stratton KL, et al. Preoperative nutritional status is an important predictor of survival in patients undergoing surgery for renal cell carcinoma. *Eur Urol* 2011;59:923-8.
10. Kott O, Golijanin B, Pereira JF, et al. The BMI Paradox and Robotic Assisted Partial Nephrectomy. *Front Surg* 2020;6:74.
11. Ioffe E, Hakimi AA, Oh SK, et al. Effect of visceral obesity on minimally invasive partial nephrectomy. *Urology* 2013;82:612-8.
12. Gorin MA, Mullins JK, Pierorazio PM, et al. Increased intra-abdominal fat predicts perioperative complications following minimally invasive partial nephrectomy. *Urology* 2013;81:1225-30.
13. Aboumarzouk OM, Stein RJ, Haber GP, et al. Laparoscopic partial nephrectomy in obese patients: a systematic review and meta-analysis. *BJU Int* 2012;110:1244-50.
14. Sperling CD, Xia L, Berger IB, et al. Obesity and 30-Day Outcomes Following Minimally Invasive Nephrectomy. *Urology* 2018;121:104-11.
15. Sharma V, Aggarwal A, McGuire BB, et al. Open vs Minimally Invasive Partial Nephrectomy: Assessing the Impact of BMI on Postoperative Outcomes in 3685 Cases from National Data. *J Endourol* 2015;29:561-7.
16. Psutka SP, Boorjian SA, Moynagh MR, et al. Decreased

- Skeletal Muscle Mass is Associated with an Increased Risk of Mortality after Radical Nephrectomy for Localized Renal Cell Cancer. *J Urol* 2016;195:270-6.
17. Hagiwara M, Miyajima A, Hasegawa M, et al. Visceral obesity is a strong predictor of perioperative outcome in patients undergoing laparoscopic radical nephrectomy. *BJU Int* 2012;110:E980-4.
 18. Darbas T, Forestier G, Leobon S, et al. Impact of Body Composition in Overweight and Obese Patients With Localised Renal Cell Carcinoma. *In Vivo* 2020;34:2873-81.
 19. Psutka SP, Carrasco A, Schmit GD, et al. Sarcopenia in patients with bladder cancer undergoing radical cystectomy: impact on cancer-specific and all-cause mortality. *Cancer* 2014;120:2910-8.
 20. Lieffers JR, Bathe OF, Fassbender K, et al. Sarcopenia is associated with postoperative infection and delayed recovery from colorectal cancer resection surgery. *Br J Cancer* 2012;107:931-6.
 21. Aubrey J, Esfandiari N, Baracos VE, et al. Measurement of skeletal muscle radiation attenuation and basis of its biological variation. *Acta Physiol (Oxf)* 2014;210:489-97.
 22. Peyton CC, Heavner MG, Rague JT, et al. Does Sarcopenia Impact Complications and Overall Survival in Patients Undergoing Radical Nephrectomy for Stage III and IV Kidney Cancer? *J Endourol* 2016;30:229-36.
 23. Sharma P, Zargar-Shoshtari K, Caracciolo JT, et al. Sarcopenia as a predictor of overall survival after cytoreductive nephrectomy for metastatic renal cell carcinoma. *Urol Oncol* 2015;33:339.e17-23.

Cite this article as: Freitas PFS, Vidal RSR, Watanabe CH, Arap MA. Predicting adverse outcomes in oncological nephrectomy: let's take a look at the patient around the tumor. *Transl Androl Urol* 2023;12(1):5-8. doi: 10.21037/tau-22-803