

Mitotic centromere-associated kinesin is a novel marker for prognosis and lymph node metastasis in colorectal cancer

K Ishikawa^{1,2}, Y Kamohara^{1,2}, F Tanaka², N Haraguchi^{1,2}, K Mimori^{1,2}, H Inoue^{1,2} and M Mori^{*1,2}

¹Department of Molecular and Surgical Oncology, Medical Institute of Bioregulation, Kyushu University, 4546 Tsurumibaru, Beppu 874-0838, Japan;

²Core Research for Evolutional Science and Technology (CREST), Japan Science and Technology Agency (JST), 4-1-8 Honcho Kawaguchi, Saitama, Japan

Mitotic centromere-associated kinesin (MCAK) is a microtubule depolymerase that is essential for proper kinetochore–microtubule attachment during spindle formation. Overexpression of MCAK has been correlated with aggressive forms of carcinoma, resulting in poor prognosis of colorectal cancer. The purpose of this study was to quantify MCAK expression in malignant and benign colorectal tissues and to determine if MCAK expression levels correlate with clinicopathologic factors and prognosis in colorectal cancer patients. Paired colorectal tissue samples from tumours and the corresponding normal tissues were obtained from 120 patients with colorectal cancer who underwent surgical resection. The real-time reverse transcriptase-PCR and immunohistochemistry were used to analyse mRNA and protein expression status with respect to various clinicopathological factors. MCAK expression was higher in colorectal cancer tissue ($P < 0.01$) than in corresponding normal tissue, and this elevated expression level was markedly associated with factors such as lymph node metastasis ($P = 0.0023$), venous invasion ($P = 0.019$), peritoneal dissemination ($P = 0.021$) and Dukes' classification ($P = 0.0023$). Patients with high MCAK mRNA expression also showed a far poorer survival rate than those with low MCAK mRNA expression ($P < 0.01$). Elevated MCAK expression was an independent predictor of overall survival and lymph node metastasis. These data suggest that MCAK expression may serve as a good marker of prognosis and lymph node metastasis in colorectal cancer.

British Journal of Cancer (2008) 98, 1824–1829. doi:10.1038/sj.bjc.6604379 www.bjcancer.com

Published online 27 May 2008

© 2008 Cancer Research UK

Keywords: MCAK; colorectal cancer; prognosis; lymph node metastasis

Colorectal cancer is one of the most prevalent cancers in the world, accounting for nearly 10% of all cancer cases (Parkin *et al*, 2001). In Japan, the incidence of colorectal cancer has significantly increased during the past 50 years, in concert with changes in eating habits, other lifestyle parameters and longevity (Kono, 1996). Colorectal cancer is now the second most important cause of death from neoplastic disease in Japan (Tsukuma *et al*, 2004). Identification of the genes responsible for the development and progression of colorectal cancer is critical if improvements are to be made in the diagnosis and treatment of the disease.

Chromosomal instability, a recognisable feature of human cancers, is caused by continuous chromosomal missegregation during mitosis (Tomonaga *et al*, 2005). A physical connection (i.e., a kinetochore) between the spindle microtubules (MTs) and centromeric DNA is essential for appropriate chromosomal segregation (Tomonaga *et al*, 2005). During spindle formation, mitotic centromere-associated kinesin (MCAK), a microtubule depolymerase, ensures proper kinetochore–MT attachment

(Holmfeldt *et al*, 2005). An attachment failure leads to missegregation of individual chromosomes.

MCAK, located at chromosome 1p34.1, is a member of the kinesin-13 subfamily that shares homology with other members of the kinesin superfamily (Wordeman and Mitchison, 1995; Vale and Fletterick, 1997; Lawrence *et al*, 2004) that play important roles in intracellular transport and cell division (Wittmann *et al*, 2001). Mitotic centromere-associated kinesin is present throughout the cell, but is especially concentrated at the centromeres, kinetochores and spindle poles (Gorbsky, 2004). Although other kinesins act to transport cargo, MCAK and other kinesin-13 members catalyse MT disassembly, an important factor in normal chromosome movement (Helenius *et al*, 2006). Mitotic centromere-associated kinesin destabilises MTs from either end, and this activity and localisation are under the regulation of mitotic kinases (Wordeman *et al*, 2007). Overexpression of wild-type MCAK induces destabilisation of MTs in both mitotic and interphase cells (Kline-Smith and Walczak, 2002).

We recently reported that elevated MCAK expression is associated with lymphatic invasion, lymph node metastasis and poor prognosis in gastric cancer (Nakamura *et al*, 2007). Although MCAK mRNA is also highly expressed in colorectal cancer tissues (Mori *et al*, 1993a), no clinicopathologic analysis of MCAK expression in human colorectal cancer has been reported. In this study, we explored MCAK gene expression in human colorectal cancer and evaluated possible associations

*Correspondence: Dr M Mori, Department of Molecular and Surgical Oncology, Medical Institute of Bioregulation, Kyushu University, 4546 Tsurumibaru, Beppu 874-0838, Japan;

E-mail: mmori@beppu.kyushu-u.ac.jp

Received 4 February 2008; revised 7 April 2008; accepted 7 April 2008; published online 27 May 2008

of gene expression with clinicopathological factors and prognosis.

MATERIALS AND METHODS

Cell lines and clinical tissue samples

Cell lines derived from human colorectal cancer (DLD1, COLO201, COLO205, COLO320DM, HT29, HCT15, WiDr, SW480, LS174T, CCK81, RCM1 and CaR1) were obtained from the Cell Resource Centre for Biomedical Research Institute of Development, Aging and Cancer (Tohoku University, Sendai, Japan) and maintained in RPMI-1640 containing 10% fetal bovine serum and antibiotics at 37°C in a 5% humidified CO₂ atmosphere.

One hundred and twenty patients (70 men, 50 women) with colorectal cancer who underwent surgery at the Medical Institute of Bioregulation at Kyushu University from 1994 to 1999 were included in this study. Primary colorectal cancer specimens and adjacent normal colorectal mucosa were obtained from patients after informed consent had been obtained in accordance with the institutional guidelines of the hospital. Immediately after resection, the necrotic and ulcerated portions of the tumours were removed and normal colonic mucosa was dissociated from muscle and connective tissue. All specimens were immediately frozen in liquid nitrogen and kept at -80°C until RNA extractions were performed. Every patient was definitively identified as having colorectal cancer based on the clinicopathologic findings. None of the patients received chemotherapy or radiotherapy prior to surgery. Clinicopathological factors were assessed according to the criteria of the Japanese Classification of Colorectal Carcinoma (Japanese Society for Cancer Colon and Rectum, 1998). All patients were closely followed after surgery at regular 1-month intervals for 2–72 months, with a mean follow-up period of 38.4 months. This study was conducted under the supervision of the ethical board of Kyushu University.

RNA preparation and reverse transcription

Total RNA was isolated using a modified acid guanidinium-phenol-chloroform procedure with DNase (Mimori *et al*, 1997). cDNA was synthesised from 2.5 µg of total RNA as described previously (Mori *et al*, 1993b).

MCAK gene amplification

A 237-bp MCAK fragment was amplified with the following primers: 5'-GATGGAAGCCTGCTAACG-3' (forward) and 5'-GAGCAGATTCCGCTTGTTC-3' (reverse). The forward primer is located in exon 8 and the reverse primer in exon 9. The amplification parameters were 1 min at 95°C, 1 min at 62°C and 1 min at 72°C for 28 cycles. An 8-µl aliquot of each reaction mixture was size-fractionated in a 2% agarose gel and visualised by ethidium bromide staining. To ensure that the RNA was not degraded, a PCR assay with primers specific for the glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) gene was performed for 1 min at 95°C, 1 min at 56°C and 1 min at 72°C for 28 cycles. The *GAPDH* primers 5'-TTGGTATCGTGAAGGACTCA-3' (forward) and 5'-TGTCATCATATTTGGCAGGTT-3' (reverse) produced a 270-bp amplicon (Ogawa *et al*, 2004). cDNA from the Human Universal Reference Total RNAs (Clontech, Palo Alto, CA, USA) was studied concurrently as a source of positive controls. Confirmation of reverse transcriptase-PCR (RT-PCR) products was performed by direct sequencing analysis.

Real-time quantitative reverse transcriptase-PCR

Real-time monitoring of the PCRs was performed using the LightCycler FastStart DNA Master SYBR Green I Kit (Roche Diagnostics, Tokyo, Japan). The amplification protocol consisted

of 35 cycles of denaturation at 95°C for 10 s, annealing at 64°C for 10 s and elongation at 72°C for 10 s. The products were then subjected to a temperature gradient from 68°C to 95°C at 0.1°C s⁻¹ with continuous fluorescence monitoring to produce a melting curve of the products. After proportional background adjustment, the fit point method was used to determine the cycle in which the log-linear signal was distinguished from the background, and that cycle number was used as a crossing-point value. The standard curve was produced by measuring the crossing point of each standard value (twofold serially diluted cDNAs from the Human Universal Reference Total RNAs) and plotting them against the logarithmic value of the concentration. The concentration of each sample was then calculated by setting their crossing points to the standard curve. The expression levels were normalised against *GAPDH* mRNA expression (Ogawa *et al*, 2004).

Immunohistochemistry

Colorectal cancer surgical specimens from formalin-fixed, paraffin-embedded tissues were used for MCAK immunohistochemistry. After deparaffinisation and blocking, the antigen-antibody reaction was incubated overnight at 4°C. ENVISION reagents (ENVISION + Dual Link/HRP; Dako Cytomation, Glostrup, Denmark) were applied to detect the signal from the antigen-antibody reaction. All sections were counterstained with haematoxylin. The primary anti-MCAK goat polyclonal antibody (ab5966; Abcam, Cambridge, UK) was used at a dilution of 1:500.

Statistical analysis

For continuous variables, the data were expressed as the mean ± s.d. The relationship between MCAK mRNA expression and the clinicopathological factors was analysed by the χ^2 test and Student's *t*-test. Kaplan-Meier survival curves were plotted and compared with the generalised log rank test. Prognostic factors were evaluated by univariate and multivariate analyses (Cox proportional hazard regression model). A logistic regression model was used to identify the independent predictors of lymph node metastasis. All tests were analysed by JMP software (SAS Institute, Cary, NC, USA), and the findings were considered statistically significant at *P* < 0.05.

RESULTS

MCAK mRNA expression in colorectal cancer cell lines and clinical tissue specimens

Figure 1A shows MCAK gene expression status in human colorectal cancer cell lines from RT-PCR analysis. Twelve of the 13 cell lines (92%) expressed the MCAK gene; COLO201 was the exception. We also performed RT-PCR analysis of MCAK in colorectal cancers and paired normal samples obtained from seven patients. In all seven cases, MCAK expression was higher in the cancer tissues than in paired normal tissue (Figure 1B). Quantitative real-time RT-PCR was performed using the Human Total RNA Master Panel (Clontech) to characterise MCAK mRNA expression. MCAK expression was extremely high in the testis. Lower expression was found in the digestive organs (colon and intestines), possibly a reflection of basic mitosis of the mucosa (data not shown). Moreover, quantitative real-time RT-PCR on 120 paired clinical samples showed that 91 of 120 cases (75.8%) exhibited higher levels of MCAK mRNA in tumours than in paired normal samples. The mean expression value of MCAK mRNA in tumour samples, 0.53 ± 0.037 (mean ± s.d., normalised by *GAPDH* gene expression), was significantly higher than the value, 0.32 ± 0.037, for the corresponding normal samples (*P* < 0.01; Student's *t*-test).




Figure 1 *MCAK* mRNA expression in colorectal cancer cell lines and clinical tissue specimens. **(A)** Reverse transcriptase-PCR analysis of *MCAK* expression in human colorectal cancer cell lines. Twelve of the 13 cell lines (92%) expressed the *MCAK* gene, whereas COLO201 did not. **(B)** Reverse transcriptase-PCR analysis of *MCAK* expression in clinical samples of colorectal cancer. *MCAK* expression was determined in the colorectal tumours (T) and paired normal (N) samples obtained from seven patients. In all seven cases, *MCAK* expression was higher in the tumours than in the normal tissues. *GAPDH* was used as a loading control. m = marker; NC = negative control; PC = positive control.




Figure 2 Immunohistochemical staining for *MCAK* in tumour specimens. **(A and B)** A representative positive stain for *MCAK* in tissue from a colorectal cancer patient. Positive staining is observed in the cytoplasm of cancer cells, but not in the stromal cells nor in normal epithelium. Solid line arrows indicate cancer cells that are positive for *MCAK* expression. **(C and D)** A representative negative stain for *MCAK* in a colorectal cancer specimen. Dotted line arrows indicate cancer cells that are negative for *MCAK* expression. Mitotic centromere-associated kinesin protein expression was evaluated in tumours and corresponding normal tissues from 15 representative colorectal cancer cases. The six tumours that were negative for protein expression (**C and D**) also exhibited lower mRNA expression levels, whereas seven of nine tumours with a positive immunohistochemical expression (**A and B**) displayed elevated mRNA expression levels. The expression of *MCAK* mRNA was thereby associated with protein expression. Each solid square in (**A**) and (**C**) was magnified $\times 100$ and shown in (**B**) and (**D**), respectively (original magnification; (**A**) and (**C**), $\times 40$; (**B**) and (**D**), $\times 100$).

Immunohistochemistry

A positive immunohistochemical staining pattern for *MCAK* in tissue from a colorectal cancer patient is shown in Figure 2A and B. Mitotic centromere-associated kinesin protein staining was observed in the cytoplasm of cancer cells, but not in the stromal cells nor in normal epithelium. Mitotic centromere-associated kinesin protein expression was examined in the tumours and corresponding normal tissues from 15 representative colorectal cancer cases. Of these 15 cases, seven exhibited a higher level of

MCAK mRNA expression in tumour tissues, whereas the remaining eight cases showed a lower level of *MCAK* mRNA in cancerous tissue. All sections were independently examined for protein expression and scored as positive or negative when $>10\%$ (positive) or $<10\%$ (negative) of carcinoma cells were stained in an examined area of a specimen. As a result, nine of the representative cases showed a positive reaction for *MCAK*, whereas six cases were negative. The data were similar to those obtained from *MCAK* mRNA expression analysis. The six negative tumours (Figure 2C and D) also showed lower mRNA expression levels,

while seven of the nine positive tumours (Figure 2A and B) showed higher mRNA expression levels. These data suggest that MCAK mRNA expression is associated with protein expression.

MCAK mRNA expression and clinicopathological characteristics

The experimental samples were divided into two expression groups. Patients who had less than the median expression ratio of tumour to normal tissue (T/N) were assigned to the low-expression group ($n=81$); the others were assigned to the high-expression group ($n=39$). Clinicopathological factors related to MCAK expression status are listed in Table 1. The incidence of three of these factors was positively correlated with increased expression (i.e., the high-expression group): (1) lymph node metastasis was significantly higher ($P=0.0023$) in the high-expression group (25 of 39, 64.1%) than in the low-expression group (28 of 81, 34.6%); (2) the incidence of venous invasion was significantly higher ($P=0.019$) in the high-expression group (13 of 39, 33.3%) than in the low-expression group (12 of 81, 14.8%); and (3) the incidence of peritoneal dissemination was significantly higher in the high-expression group (4 of 39, 10.3%) than in the low-expression group (1 of 81, 1.23%). Moreover, the incidence of advanced stage cancer (according to Dukes' classification) was significantly higher ($P=0.0023$) in the high-expression group (26 of 39, 66.7%) than in the low-expression group (30 of 81, 37.0%). The other clinicopathological factors were not correlated with MCAK overexpression.

Relationship between MCAK expression and prognosis

The 5-year survival rate was significantly lower in patients with elevated MCAK expression ($P<0.01$; Figure 3). Table 2 provides the univariate and multivariate analyses of factors related to patient prognosis. Univariate analysis showed that the following factors were significantly related to postoperative survival: tumour size ($P=0.009$), serosal invasion ($P<0.0001$), lymph node metastasis ($P<0.0001$), lymphatic invasion ($P<0.0001$) and MCAK mRNA expression ($P=0.019$). Multivariate regression analysis indicated that inclusion in the MCAK mRNA high-expression group (relative risk (RR), 1.42; 95% confidence interval (CI), 1.01–2.00; $P<0.05$) was an independent predictor of overall survival, as were serosal invasion (RR, 2.16; 95% CI, 1.52–3.12; $P<0.0001$) and lymph node metastasis (RR, 1.82; 95% CI, 1.26–2.76; $P=0.0012$).

Multivariate analysis for lymph node metastasis

Univariate and multivariate logistic regression analyses were performed on lymph node metastasis (Table 3). Univariate analysis revealed a significant relationship between lymph node metastasis and the following factors: histological grade ($P=0.04$), tumour size ($P=0.01$), depth of invasion ($P=0.02$), lymphatic invasion ($P<0.0001$), venous invasion ($P=0.003$) and MCAK mRNA expression ($P=0.003$). Multivariate regression analysis indicated that inclusion in the MCAK mRNA high-expression group (RR, 4.02; 95% CI, 1.53–11.3; $P<0.006$) was an independent predictor of lymph node metastasis in addition to lymphatic invasion (RR, 7.67; 95% CI, 3.01–21.1; $P<0.0001$).

DISCUSSION

In this study, we demonstrated that MCAK is expressed at higher levels in colorectal cancer cells than in the corresponding normal tissues. Mitotic centromere-associated kinesin overexpression causes a moderate increase in the frequency of multipolar spindles (Holmfeldt *et al*, 2004) and monopolar spindles (Kline-Smith and Walczak, 2002), which can contribute to the gain or loss of

Table 1 Clinicopathological factors and MCAK mRNA expression in 120 colorectal cancers

Factors	High expression (%) (n = 39)	Low expression (%) (n = 81)	P-value
Age (mean \pm s.d.)	65.3 \pm 12.3	66.9 \pm 11.3	0.49
Gender			
Male	22 (56.4)	48 (59.3)	0.76
Female	17 (43.6)	33 (40.7)	
Histological grade			
Well	15 (38.5)	29 (35.8)	0.78
Others (moderately, poorly and mucinous)	24 (61.5)	52 (64.2)	
Tumour size			
< 30 mm (small)	8 (20.5)	17 (21.0)	0.95
\geq 30 mm (large)	31 (79.5)	64 (79.0)	
Serosal invasion			
Absent	26 (66.7)	59 (72.8)	0.49
Present	13 (33.3)	22 (27.2)	
Lymph node metastasis			
Absent	14 (35.9)	53 (65.4)	0.0023*
Present	25 (64.1)	28 (34.6)	
Lymphatic invasion			
Absent	22 (56.4)	53 (65.4)	0.34
Present	17 (43.6)	28 (34.6)	
Venous invasion			
Absent	26 (66.7)	69 (85.2)	0.019*
Present	13 (33.3)	12 (14.8)	
Liver metastasis			
Absent	33 (84.6)	73 (90.1)	0.38
Present	6 (15.4)	8 (9.9)	
Peritoneal dissemination			
Absent	35 (89.7)	80 (98.8)	0.021*
Present	4 (10.3)	1 (1.2)	
Dukes' classification			
A and B	13 (33.3)	51 (63.0)	0.0023*
C and D	26 (66.7)	30 (37.0)	

s.d. = standard deviation. * $P<0.05$.

chromosomes in daughter cells. We reported that overexpression of MCAK leads to the increased migratory and proliferative ability of gastric cancer cells (Nakamura *et al*, 2007). Microtubules, key components of the cytoskeleton, play essential roles in mitosis, cell migration, and cell signalling and trafficking (Honore *et al*, 2005). Mitotic centromere-associated kinesin is present throughout the cell, and overexpression of MCAK may accelerate MT turnover, resulting in the gain of cancer cell motility. This result may partially explain the finding that gastric cancer patients with tumours that express high levels of MCAK had higher rates of lymphatic invasion and metastasis, and a poorer prognosis. Along the same line, downregulation of MCAK in a breast cancer study resulted in tumour growth suppression (Shimo *et al*, 2008). These findings suggest that overexpression of MCAK enhances the malignancy of cancer cells.

Clinicopathological analysis revealed that tumours with high MCAK expression were associated with lymph node metastasis, venous invasion, peritoneal dissemination and advanced Dukes'

stage. In our previous study on gastric cancer, *MCAK* gene expression in cancer tissues was significantly higher than expression in nonmalignant tissue, and elevated *MCAK* expression was significantly associated with lymphatic invasion and lymph node metastasis (Nakamura *et al*, 2007). These results suggest that *MCAK* plays a pivotal role in the progression of colorectal cancer and gastric cancer. Moreover, in both studies, patients with high *MCAK* expression had a poorer survival rate than those with low *MCAK* expression, and high *MCAK* expression was an independent




Figure 3 Kaplan–Meier survival curves of patients with colorectal cancer based on *MCAK* mRNA expression status. Patients in the high-expression group (bold line) showed significantly poorer prognosis than those in the low-expression group (dotted line; $P < 0.01$, log rank test).

prognostic factor for overall survival in the Cox proportional hazard regression model. To our knowledge, this is the first report on correlations between *MCAK* gene expression and clinicopathological factors in colorectal cancer. In the majority of colorectal cancer reports, gene expression is secondary to TNM or Dukes' classification as a prognostic marker. Because its expression is an independent prognostic factor, the expression profile of *MCAK* may contribute to the creation of a new clinical classification system.

Another aspect of this study is the relationship between *MCAK* expression and lymphatic extension. Interestingly, multivariate analysis for lymph node metastasis revealed that *MCAK* mRNA expression is a good predictor of lymph node metastasis. Lymph node metastases are the most important predictors of survival in non-stage IV colorectal cancer (Schumacher *et al*, 2007). Today, we have several options for curative treatment of early colorectal cancer, such as endoscopic mucosal resection (EMR), endoscopic submucosal dissection (ESD) and laparoscopic-assisted colectomy with regional lymphadenectomy. In limited cases, patients are eligible for such less invasive surgery. Patients suitable for those endoscopic treatments are selected by the preoperative diagnosis of lymph node metastasis, such as macroscopic type, tumour size, presence of an ulcer and histology of biopsy specimens. However, some patients are misdiagnosed and actually do have lymph node metastasis before the operation. By incorporating the genetic diagnosis of *MCAK* gene expression, preoperative selection for patients without lymph node metastasis may be possible. Furthermore, it has been reported that with SELEX, *MCAK* was detected in the peripheral blood of a colon cancer patient (Scanlan *et al*, 2002), suggesting that the blood level of *MCAK* can be used for the prediction of lymph node metastasis. The analysis of *MCAK* using biopsied specimens and blood

Table 2 Univariate and multivariate analysis for overall survival (Cox proportional hazards regression model)

Factors	Univariate analysis			Multivariate analysis		
	RR	95% CI	P-value	RR	95% CI	P-value
Age (<60/60 ≤)	0.82	0.58–1.20	0.29	—	—	—
Gender (male/female)	0.96	0.68–1.33	0.81	—	—	—
Histological grade (well/moderately, poorly and mucinous)	1.21	0.86–1.76	0.29	—	—	—
Tumour size (<30 mm/≥30 mm)	1.93	1.16–3.94	0.009	1.29	0.74–2.70	0.4
Serosal invasion (absent/present)	2.40	1.72–3.38	<0.0001	2.16	1.52–3.12	<0.0001
Lymph node metastasis (absent/present)	2.25	1.58–3.38	<0.0001	1.82	1.26–2.76	0.0012
Lymphatic invasion (absent/present)	2.30	1.64–3.31	<0.0001	—	—	—
Venous invasion (absent/present)	1.43	0.99–2.00	0.055	—	—	—
<i>MCAK</i> mRNA expression (T/N <3/T/N >3)	1.49	1.07–2.07	0.019	1.42	1.01–2.00	0.048

CI = confidence interval; RR = relative risk.

Table 3 Univariate and multivariate analysis for lymph node metastasis (logistic regression model)

Factors	Univariate analysis			Multivariate analysis		
	RR	95% CI	P-value	RR	95% CI	P-value
Age (<60/60 ≤)	0.44	0.18–1.108	0.08	—	—	—
Gender (male/female)	0.86	0.41–1.79	0.69	—	—	—
Histological grade (well/moderately, poorly and mucinous)	2.26	1.05–5.02	0.04	2.21	0.84–6.09	0.11
Tumour size (<30 mm/≥30 mm)	4.08	1.51–13.1	0.01	2.52	0.72–10.1	0.17
Tumour stage (T1/T2–T4)	4.40	1.34–19.9	0.02	1.25	0.26–7.09	0.79
Lymphatic invasion (absent/present)	9.11	3.98–22.3	<0.0001	7.67	3.01–21.1	<0.0001
Venous invasion (absent/present)	4.41	1.56–8.25	0.003	2.01	0.66–6.48	0.23
<i>MCAK</i> mRNA expression (T/N <3/T/N >3)	3.38	1.54–7.67	0.003	4.02	1.53–11.3	0.006

CI = confidence interval; RR = relative risk.

samples could provide a more accurate evaluation, which might translate to minimally invasive treatments for early colorectal cancer.

In conclusion, the expression of MCAK in colorectal cancer was elevated relative to levels in normal colorectal tissue, and this overexpression was associated with lymph node metastasis, venous invasion, peritoneal dissemination and poor prognosis. In particular, MCAK mRNA overexpression was associated with prognosis and lymph node metastasis in a multivariate analysis. Therefore, MCAK could be a useful predictive marker of lymph node metastasis, which might permit minimally invasive and curative treatments combining EMR and ESD for early colorectal cancer.

REFERENCES

- Gorbsky GJ (2004) Mitosis: MCAK under the aura of Aurora B. *Curr Biol* **14**: R346–R348
- Helenius J, Brouhard G, Kalaidzidis Y, Diez S, Howard J (2006) The depolymerizing kinesin MCAK uses lattice diffusion to rapidly target microtubule ends. *Nature* **441**: 115–119
- Holmfeldt P, Stenmark S, Gullberg M (2004) Differential functional interplay of TOGp/XMAP215 and the Kin I kinesin MCAK during interphase and mitosis. *EMBO J* **23**: 627–637
- Holmfeldt P, Zhang X, Stenmark S, Walczak CE, Gullberg M (2005) CaMKIIγ-mediated inactivation of the Kin I kinesin MCAK is essential for bipolar spindle formation. *EMBO J* **24**: 1256–1266
- Honore S, Pasquier E, Braguer D (2005) Understanding microtubule dynamics for improved cancer therapy. *Cell Mol Life Sci* **62**: 3039–3056
- Kline-Smith SL, Walczak CE (2002) The microtubule-destabilizing kinesin XKCM1 regulates microtubule dynamic instability in cells. *Mol Biol Cell* **13**: 2718–2731
- Kono S (1996) Dietary factors for gastrointestinal cancers: a worldwide overview. *Gann Monogr Cancer Res* **44**: 29–39
- Lawrence CJ, Dawe RK, Christie KR, Cleveland DW, Dawson SC, Endow SA, Goldstein LS, Goodson HV, Hirokawa N, Howard J, Malmberg RL, McIntosh JR, Miki H, Mitchison TJ, Okada Y, Reddy AS, Saxton WM, Schliwa M, Scholey JM, Vale RD, Walczak CE, Wordeman L (2004) A standardized kinesin nomenclature. *J Cell Biol* **167**: 19–22
- Mimori K, Mori M, Shiraishi T, Fujie T, Baba K, Haraguchi M, Abe R, Ueo H, Akiyoshi T (1997) Clinical significance of tissue inhibitor of metalloproteinase expression in gastric carcinoma. *Br J Cancer* **76**: 531–536
- Mori M, Barnard GF, Staniunas RJ, Jessup JM, Steele Jr GD, Chen LB (1993a) Prothymosin-α mRNA expression correlates with that of c-myc in human colon cancer. *Oncogene* **8**: 2821–2826
- Mori M, Staniunas RJ, Barnard GF, Jessup Jr JM, Steele GD, Chen LB (1993b) The significance of carbonic anhydrase expression in human colorectal cancer. *Gastroenterology* **105**: 820–826
- Nakamura Y, Tanaka F, Haraguchi N, Mimori K, Matsumoto T, Inoue H, Yanaga K, Mori M (2007) Clinicopathological and biological significance of mitotic centromere-associated kinesin overexpression in human gastric cancer. *Br J Cancer* **97**: 543–549

ACKNOWLEDGEMENTS

We thank Dr Y Nakamura, Ms T Shimooka, Ms K Ogata, Ms M Kasagi and Ms Y Nakagawa for their technical assistance and advice. This work was supported in part by the following grants and foundations: CREST, Japan Science and Technology Agency; Japan Society for the Promotion of Science Grant-in-Aid for Scientific Research, grants 17109013, 17591411, 17591413, 18390367, 18590333, 18659384, 18790964, 19890336 and 19591509; The Ministry of Education, Culture, Sports, Science and Technology Grant-in-Aid for Scientific Research on Priority Areas, Grant 18015039; Third Term Comprehensive Ten-Year Strategy for Cancer Control, Grant 16271201.

- Ogawa K, Utsunomiya T, Mimori K, Tanaka Y, Tanaka F, Inoue H, Murayama S, Mori M (2004) Clinical significance of elongation factor-1 delta mRNA expression in oesophageal carcinoma. *Br J Cancer* **91**: 282–286
- Parkin DM, Bray F, Ferlay J, Pisani P (2001) Estimating the world cancer burden: Globocan 2000. *Int J Cancer* **94**: 153–156
- Japanese Society for Cancer Colon and Rectum (1998) *General Rules for Clinical and Pathological Studies on Cancer of the Colon, Rectum and Anus* 6th edn. Tokyo: Kanehara
- Scanlan MJ, Welt S, Gordon CM, Chen YT, Gure AO, Stockert E, Jungbluth AA, Ritter G, Jager D, Jager E, Knuth A, Old LJ (2002) Cancer-related serological recognition of human colon cancer: identification of potential diagnostic and immunotherapeutic targets. *Cancer Res* **62**: 4041–4047
- Schumacher P, Dineen S, Barnett Jr C, Fleming J, Anthony T (2007) The metastatic lymph node ratio predicts survival in colon cancer. *Am J Surg* **194**: 827–831; discussion 831–822
- Shimo A, Tanikawa C, Nishidate T, Lin ML, Matsuda K, Park JH, Ueki T, Ohta T, Hirata K, Fukuda M, Nakamura Y, Katagiri T (2008) Involvement of kinesin family member 2C/mitotic centromere-associated kinesin overexpression in mammary carcinogenesis. *Cancer Sci* **99**: 62–70
- Tomonaga T, Matsushita K, Ishibashi M, Nezu M, Shimada H, Ochiai T, Yoda K, Nomura F (2005) Centromere protein H is up-regulated in primary human colorectal cancer and its overexpression induces aneuploidy. *Cancer Res* **65**: 4683–4689
- Tsukuma H, Ajiki W, Oshima A (2004) [Cancer incidence in Japan]. *Gan To Kagaku Ryoho* **31**: 840–846
- Vale RD, Fletterick RJ (1997) The design plan of kinesin motors. *Annu Rev Cell Dev Biol* **13**: 745–777
- Wittmann T, Hyman A, Desai A (2001) The spindle: a dynamic assembly of microtubules and motors. *Nat Cell Biol* **3**: E28–E34
- Wordeman L, Mitchison TJ (1995) Identification and partial characterization of mitotic centromere-associated kinesin, a kinesin-related protein that associates with centromeres during mitosis. *J Cell Biol* **128**: 95–104
- Wordeman L, Wagenbach M, von Dassow G (2007) MCAK facilitates chromosome movement by promoting kinetochore microtubule turnover. *J Cell Biol* **179**: 869–879