




ELSEVIER

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Parasite Epidemiology and Control

journal homepage: www.elsevier.com/locate/parepi

Seroprevalence canine survey for selected vector-borne pathogens and its relationship with poverty in metropolitan Pereira, Colombia, 2020[☆]

D. Katterine Bonilla-Aldana^{a,b,c,q}, Erwin J. Gutiérrez-Grajales^c,
 J. Paola Martínez-Arboleda^{a,b}, María Angelica Reina-Mora^{a,b},
 Adrián E. Trejos-Mendoza^{a,b}, Soffia Pérez-Vargas^{a,b}, Lorenzo Valencia-Mejía^{a,b},
 Luisa F. Marín-Arboleda^{a,b}, Daniela Osorio-Navia^{a,b}, Mariana Chacón-Peña^{a,b},
 Luz Victoria González-Colonia^d, Jaime A. Cardona-Ospina^{b,e,f,q},
 Erika Vanessa Jiménez-Posada^f, Andrés Díaz^g, Jean Carlos Salazar^g,
 Manuel Sierra^h, Fausto Muñoz-Lara^{i,p}, Lysien I. Zambrano^h,
 Eduardo Ramírez-Vallejo^j, Juan Camilo Álvarez^k,
 Ingrid Lorena Jaramillo-Delgado^{c,k}, Samuel Pecho-Silva^{l,m},
 Alberto Paniz-Mondolfiⁿ, Álvaro A. Faccini-Martínez^{c,o,q},
 Alfonso J. Rodríguez-Morales^{b,c,e,f,l,q,*}

^a Semillero de Investigación en Zoonosis (SIZOO), Fundación Universitaria Autónoma de las Américas, Pereira, Risaralda, Colombia

^b Institución Universitaria Visión de las Américas, Pereira, Risaralda, Colombia

^c Red Colombiana de Enfermedades Transmitidas por Garrapatas en Pequeños Animales (RECEPA) – Colombian Network of Tick-Borne Diseases in Small Animals (RECEPA), Pereira, Risaralda, Colombia

^d San Lucas Centro Veterinario y Diagnóstico, Pereira, Risaralda, Colombia

^e Grupo de Investigación Biomedicina, Faculty of Medicine, Fundación Universitaria Autónoma de las Américas, Pereira, Risaralda, Colombia

^f Grupo de Investigación en Infecciones Emergentes y Medicina Tropical, Instituto para la Investigación en Ciencias Biomédicas, SCI-HELP, Pereira, Risaralda, Colombia

^g Vitalcare, Armenia, Quindío, Colombia

^h Unit of Scientific Research, School of Medical, Faculty of Medical Sciences, Universidad Nacional Autónoma de Honduras (UNAH), Tegucigalpa, Honduras

ⁱ Department of Internal Medicine, Faculty of Medical Sciences, Universidad Nacional Autónoma de Honduras (UNAH), Tegucigalpa, Honduras

^j IPS Cardiológica Eduardo Ramírez, Pereira, Risaralda, Colombia

^k Grupo de Investigación One-Health, Departamento de Investigación de Enfermedades Infecciosas en Animales, Centro de Diagnóstico Especializado Testmol, Medellín, Antioquia, Colombia

^l Universidad Científica del Sur, Lima, Peru

^m Hospital Nacional Edgardo Rebagliati Martins, Lima, Peru

ⁿ Laboratory of Medical Microbiology, Department of Pathology, Molecular and Cell-based Medicine, The Mount Sinai Hospital-Icahn School of Medicine at Mount Sinai, New York, USA

^o Department of Pathology, University of Texas Medical Branch, Galveston, TX, USA

^p Department of Internal Medicine, Hospital Escuela, Tegucigalpa, Honduras

^q Committee of Tropical Medicine, Zoonoses and Travel Medicine, Asociación Colombiana de Infectología, Bogotá, Colombia

[☆] This study was presented in part at the American Society for Microbiology (ASM) & the Federation of European Microbiological Societies (FEMS) World Microbe Forum, June 20–24, 2021 (iPoster #6498).

* Corresponding author at: Institución Universitaria Visión de las Américas, Pereira, Risaralda, Colombia.

E-mail addresses: arodriguezmo@cientifica.edu.pe, alfonso.rodriguez@uam.edu.co (A.J. Rodríguez-Morales).

<https://doi.org/10.1016/j.parepi.2022.e00249>

Received 1 November 2021; Received in revised form 1 March 2022; Accepted 29 March 2022

Available online 1 April 2022

2405-6731/© 2022 The Authors. Published by Elsevier Ltd on behalf of World Federation of Parasitologists. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

ARTICLE INFO

Keywords:

Anaplasma phagocytophilum
Anaplasma platys
Ehrlichia canis
Ehrlichia ewingii
Dirofilaria immitis
 Tick-borne diseases
 Hemothropic pathogens
 Canine
 Zoonotic
 Colombia

ABSTRACT

Background: Tick-borne diseases (TBD) and dirofilariosis are currently not under surveillance in most Latin American countries. In addition, there is a significant lack of studies describing the current situation in most endemic areas, including Colombia. Therefore, seroprevalence studies are crucial for understanding the epidemiology of these vector-borne diseases.

Methods: A serosurvey for TBD and dirofilariosis among 100 dogs was carried out in the municipality of Pereira, located in the Coffee-Triangle region, Colombia. Samples were tested using a rapid assay test system (SNAP® 4Dx®); based on an enzyme immunoassay technique, screening for antibodies to *Anaplasma phagocytophilum/platys* (sensitivity 99.1%), *Borrelia burgdorferi* s.l. (98.8%), and *Ehrlichia canis/ewingii* (96.2%) by using specific antigens and checking for *Dirofilaria immitis* antigen based on specific antibodies (99.2%). Bivariate analyses were performed on Stata®14, significant $p < 0.05$.

Findings: Global seroprevalence to the selected vector-borne pathogens was 74% (95%CI 65–83%). The highest seroprevalence was found for *E. canis/ewingii* (74%), followed by *A. phagocytophilum/platys* (16%). Seropositivity for *Borrelia* spp. and *Dirofilaria* spp. was 0%. All *Anaplasma* spp. seropositive dogs showed co-detection of *Ehrlichia* spp. (16%). Seroprevalence was significantly higher among dogs from families of lower socioeconomic status/level (I, 86%, followed by level II (74%), and III (36%) ($p = 0.001$). All dogs exhibiting anorexia (12%) were invariably seropositive (100%) ($p = 0.029$). Seroprevalence was higher among those showing mucocutaneous paleness (95%) compared to those without paleness (68%) ($p = 0.013$) (OR = 9.3; 95%CI 1.18–72.9). There was high variability in seroprevalence through the studied areas,




Fig. 1. Relative location of Pereira and Dosquebradas municipalities, in the Risaralda department, Colombia, Latin America.

ranging from 0% (La Libertad Park) up to Combia, Cesar Nader, Las Brisas and Saturno localities (100%) ($p = 0.033$).

Interpretation: Given the high seroprevalence obtained in an area with documented ticks, there is a potential risk of zoonotic transmission to humans. Further seroprevalence studies in humans are needed to assess the prevalence of infections. Poverty is highly associated with these tick-borne pathogens in Pereira, as shown in the present study.

1. Introduction

Tick-borne diseases (TBD) are an essential group of infectious diseases that may affect both animals and humans, particularly in the tropical and subtropical regions of the world. Among these diseases is Lyme borreliosis, spotted fever group rickettsioses, Colorado tick fever, monocytic ehrlichiosis, tularemia, granulocytic anaplasmosis, among others (Alkishe et al., 2021; Karim et al., 2021; Negi et al., 2021).

Among the most vulnerable populations for these vector-borne diseases (Bonilla-Aldana et al., 2021) are dogs (Bonilla-Aldana et al., 2020b; Rodriguez-Morales et al., 2018a; Rodriguez-Morales et al., 2018b). Multiple studies worldwide have assessed the prevalence of such diseases in canine populations (Petruccioli et al., 2020). However, many countries still lack detailed studies, with many aspects concerning their epidemiology that remain yet to be clarified, including many societal aspects (Bayles and Allan, 2014; Peretti-Watel et al., 2019; Walker, 2011).

Unfortunately, TBD (such as ehrlichiosis, anaplasmosis), and dirofilariosis, are not reportable and currently not under surveillance in most Latin American countries (Baneth et al., 2020; Hernandez-Velasco et al., 2020; Mendes-de-Almeida et al., 2021; Polo et al., 2015; Robles et al., 2018; Wisely and Glass, 2019). In addition, there is a significant gap of knowledge about their circulation across endemic areas in certain countries, such as Colombia (Jaimes-Duenez et al., 2018; Miranda and Mattar, 2015; Pesapane et al., 2019;




Fig. 2. Urban areas (communes) of Pereira and the study areas in the metropolitan area (Pereira and Dosquebradas municipalities).

Santodomingo et al., 2019; Vargas-Hernandez et al., 2016). That is why seroprevalence studies remain crucial better to understand the diverse epidemiological aspects of these vector-borne diseases.

Herein, we present a prospective study aimed to evaluate the seroprevalence of *A. phagocytophilum/platys*, *Borrelia burgdorferi* sensu lato (s.l.), *E. canis/ewingii*, and *Dirofilaria immitis* infection and scrutinize the main associated factors related to the epidemiology of tick-borne diseases among dogs inhabiting the municipality of Pereira, Risaralda department, Colombia, in 2020.

2. Materials and methods

Pereira is the principal city of the Coffee-Triangle region, which includes three departments (first administrative territory level) and 53 municipalities (second administrative territory level). Pereira is the capital of Risaralda (967,780 habitants in 2020), a department surrounded by six other western departments (Antioquia, Caldas, Tolima, Quindio, Valle del Cauca and Chocó) (Rodríguez-Morales et al., 2015). Pereira's landscape embraces both urban and rural areas. The first consists of 20 communities (the city) and the second by 12 corregimientos (sub-municipalities) (both tertiary administrative territory level) (Fig. 1). As per the National Statistics Department (DANE, www.dane.gov.co), the total population for 2020 reached 477,027 inhabitants. The metropolitan area includes Dosquebradas and La Virginia municipalities (Fig. 1), with 709,338 inhabitants for 2020. Rio Otún, Centro, San Joaquín, Del Café, Boston, El Oso, Consoá and Cuba, are the most populated areas of the Pereira municipality, making up 51% of its population (Fig. 2). The municipality of Pereira extends over an area of 702 Km² (4°48'51"N 75°41'40"W). The climate is tropical, with an annual median temperature of 18.8 °C (median minimum of 15.8 °C, median maximum of 26.3 °C).

Based on the municipality dog's census, we calculated a minimum sample of 98 dogs to be assessed. Finally, 100 with infection suspicion were included distributed in four urban communes (out of 20) and four rural corregimientos (out of 12) of the municipality Pereira and its neighboring municipality of Dosquebradas (Fig. 1). Only one dog was included from those owners who voluntarily accepted to participate with their dogs, regardless of where co-living was in the same house.

Blood samples from each dog were individually collected from the radial vein into a sterile vacuum tube (Vacutainer, Becton, Dickinson and Company Franklin Lakes, NJ, USA) without anticoagulant in the morning time. In the evening, samples were centrifuged at 1300–1800 ×g for 20 min, followed by serum separation from the clot.

To determine selected-vector-borne pathogens exposure, a rapid enzyme-linked immunosorbent assay (ELISA) kit (SNAP® 4Dx® Plus Test Kit, IDEXX Laboratories, Inc., Westbrook, ME, USA) was used following the manufacturer's instructions (Petruccioli et al., 2020). This qualitative test allowed us to simultaneously detect the presence of circulating antibodies (IgG and IgM) against immunodominant proteins of *Ehrlichia canis/ewingii* (p30 and p30–1, sensitivity of 96.2%), *Anaplasma phagocytophilum/platys* (p44/MSP2, sensitivity of 99.1%), *Borrelia burgdorferi* s.l. (C6, the sensitivity of 98.8%), and *Dirofilaria immitis* antigens (principally produced by adult females) based on specific antibodies (sensitivity of 99.2%). The SNAP® 4Dx® Plus Test Kit showed a specificity of ~100% for the microorganisms mentioned above (Chandrashekar et al., 2010; Petruccioli et al., 2020; Stillman et al., 2014). 4Dx® Plus Test Kit has been validated in dogs (Chandrashekar et al., 2010; Petruccioli et al., 2020; Stillman et al., 2014).

Besides the general demographic, clinical and laboratory surveys, a questionnaire for social variables related to living conditions and households was also performed, including aspects such as: house vulnerability (defined according to five possible types of houses, from those with luxury and appropriate sanitary conditions to those without luxury and inappropriate sanitary conditions, last category considered as a vulnerable house); house location (in rural or urban areas); environmental elements close proximity to the house (e.g. small lakes, small rivers or wetlands); materials employed in the building of the walls (e.g. blocks, bricks, cement, wood, cardboard-tin) and the floors (e.g. cement, wood, soil); access to tap water; need of water collection and its keeping at appropriate receptacles; and disposal of sewage water and waste disposal (e.g. by an urban service) (Quintero et al., 2012), among other social aspects such as the socioeconomic level of the house which is determined by the National Statistics Department (DANE).

Statistical analysis was performed using Stata 14®IC (Stata Corp., College Station, Texas, USA). Chi-square tests were used to compare proportions of positivity related to categorical dependent variables and establish statistical significance. In addition, Association between seropositivity and independent variables such as social house conditions, dog living conditions, and the socioeconomic level was evaluated. For all the independent variables, chi-square (χ^2) and Fisher tests were used to assess associations and significance. In those significant values ($p < 0.05$), also the odds ratio with their 95% confidence interval (95%CI) was calculated.

The location of the houses hosting the dogs was georeferenced with their correspondent coordinates by the free mobile cell-GPS software application "Herramientas de GPS" v.3.1.0.5, developed by Virtual Maze® (www.virtualmaze.com). Seroprevalence was also presented by geographical information systems (GIS)-based maps. The georeferenced places were incorporated in the software Google Earth Pro®, and the layers of communes, corregimientos and municipalities, provided by the Open Data Pereira Geographical Information System (<https://mapas-pereira.opendata.arcgis.com/>), in .kml files, were included to locate the coordinate's points in the corresponding shapes. Microsoft Access® software was used to design the spatial database to import the GIS software's incidence rates by corregimientos and comunas at Pereira municipality. The Client GIS software open source was Kosmo Desktop 3.0 RC1® (SAIG S.L., Madrid, Spain). For access to geographic data required and sharing results with institutions, support was provided by the spatial data infrastructure for the department by the Regional Information System of the Coffee-Triangle ecoregion (SIR) as standardized and reported before (Bonilla-Aldana et al., 2020a; Idarraga-Bedoya et al., 2020; Rodríguez-Morales et al., 2017). The shapefiles of corregimientos and comunas (.shp) were linked to a database through spatial joined operation to produce digital maps of the seroprevalence for the study area. Data used for the GIS-based maps were derived from the geographical origin of the serosurvey at the different geographic levels (Idarraga-Bedoya et al., 2020).

3. Results

The mean age of the canine population was 4.2 years (\pm 2.89 years, range 0.21–12.56), 53% were female, and 47% were males (Table 1). The average age did not differ significantly by sex (males 4.34 ± 3.27 years; females 4.10 ± 2.59 years) ($p = 0.6345$). Other dog variables are presented in Table 1. From the total number of canines, 89% presented clinical alterations, of them 100% presented anaemia, 54% lethargy, 39% alopecia, 21% decay, 21% mucocutaneous paleness, among others.

The global seroprevalence to the selected vector-borne pathogens was 74% (95%CI 65–83%). The highest seroprevalence was found for *E. canis/ewingii* (74%), followed by *A. phagocytophilum/platys* (16%). Seropositive for *Borrelia* spp. and *Dirofilaria* spp. was 0%. All the *Anaplasma* spp. seropositive dogs were also positive for *Ehrlichia* spp. (16%).

Twenty-two per cent of the houses had rudimentary walls (adobe, cardboard, and palm), 84% rudimentary roofs (wood, zinc, and palms), 41% rudimentary floors (wood and earth), and 73% of the houses had only one bathroom. In those with only one bathroom, 100% had seropositive dogs, whilst in those houses with two or more bathrooms had 65% ($p = 0.025$). Although there was water supply (100%) in all the houses, 28% collected and store water (25% in tanks and 3% in plastic recipients). The water supply in 75% was by the public system, but the rest (25%) utilized water wells, rivers, water streams, and rainwater; 56% and the reported presence of rats in the house. Thirty-nine per cent of the dogs were allowed to be inside the house with the family; 10% of the dogs remained leashed in the house or the yard (all of them, 100%, were seropositive), whilst 90% were free to go anywhere (71% were seropositive) ($\chi^2 = 3.903$; $p = 0.048$); 82% of the dogs had access to wild areas. Regarding house waste, 81% was collected by the urban system, but the rest (29%) was burned. Illiteracy was reported in 2% of dog owners; 8% of dog owners were unemployed at the survey. Twelve per cent of owners lived in overcrowded houses (≥ 3 persons/room); in 70% of the owners, the number of persons per bathroom in their houses was ≥ 3 .

The seroprevalence at rural (79%) and urban areas (67%) was not significantly different ($p = 0.181$). However, the seroprevalence was significantly higher in dogs from families of the lower socioeconomic level (I [Low-low, the lowest], 86%), followed by level II (74%), and III (36%) ($\chi^2 = 14.162$; $p = 0.001$); no dogs from the highest strata (IV and V) were available for our study (Fig. 3). There was high variability in seroprevalence by studied areas, ranging from 0% (La Libertad Park) up to Combia, Cesar Nader, Las Brisas and

Table 1
Physical, clinical, and living conditions of the studied dogs.

Variable	Mean/n	SD/%	Minimum	Maximum
Age (years)	4.20	2.89	0.21	12.56
Sex				
Male	53	53.00		
Female	47	47.00		
Weight (kilograms)	13.99	7.76	4.00	40.00
Body condition				
1 (Very slim)	0	0.00		
2 (Slim)	21	21.00		
3 (Ideal)	50	50.00		
4 (Overweight)	29	29.00		
5 (Obese)	0	0.00		
Body temperature (°C)	38.84	0.96	30.50	40.00
Cardiac rate (beats/min)	94.80	9.86	50.00	115.00
Respiratory rate (breaths/min)	30.83	4.51	24.00	55.00
Capillary Refill Time (seconds)				
1	0	0.00		
2	60	60.00		
3	28	28.00		
4	10	10.00		
5	2	2.00		
Recent deworming (last month)				
Yes	16	16.00		
No	84	84.00		
Any vaccine (in the last year)				
Yes	45	45.00		
Rabies vaccine				
Yes	43	43.00		
No	55	55.00		
Number of other dogs in the house				
1	5	5.00		
2	13	13.00		
3	12	12.00		
>3	64	64.00		
Number of cats in the house				
0	32	32.00		
1	16	16.00		
2	9	9.00		
3	5	5.00		
>3	34	34.00		




Fig. 3. Seroprevalence by socioeconomic strata.

Saturno (100%) ($p = 0.033$). In the rural areas, Cerritos was the corregimiento with the highest seroprevalence (100%), followed by Combia Baja and Caimalito (Fig. 4), whilst in the urban area, Oriente and Villasantana were the communes with the highest seroprevalence (100%) (Fig. 4).

All the dogs with anorexia (12%) were seropositive (100%) ($\chi^2 = 4.791; p = 0.029$). Seroprevalence was higher in those with mucocutaneous paleness (95%) compared to those without it (68%) ($\chi^2 = 6.232; p = 0.013$) (OR = 9.3; 95%CI 1.18–72.9).

4. Discussion

Poverty has been linked as a determinant to multiple disease groups, especially infectious diseases (Bukhman et al., 2020; Hummel et al., 2021; The Lancet Child Adolescent Health, 2019). Among the infectious diseases linked to poverty, tropical diseases and vector-borne diseases, such as malaria (Bi and Tong, 2014; Ricci, 2012; Santos-Vega et al., 2016; Tusting et al., 2016) and dengue (Bavia et al., 2020; Bonifay et al., 2017; Mulligan et al., 2015), have been well documented. Housing conditions are related to the occurrence and persistence of tropical infectious diseases (Cardona-Arias, 2018; Ercumen et al., 2019; Pham-Duc et al., 2013; Quintero et al., 2012; Snyman et al., 2015). There is a global lack of studies exploring the links between tick-borne diseases, such as *Ehrlichia* spp. infection, the primary pathogen that elicited antibodies in our study, and poverty. Poverty is also related to pet care as well as to tick control in dogs. In this study, none of the owners provided any veterinary care to their dogs over the last previous six months.

A recent study in Argentina, assessing different infectious diseases among dogs from a rural area in the humid Chaco, found 7.9% of them prevalent for ehrlichiosis. They suggest that their findings likely reflect structural poverty, poor sanitation and lack of a safe water supply (Enriquez et al., 2019). In another study from Brazil, assessing carthorses from low-income owners, authors, found that 27.4% were positive for *Ehrlichia* exposure (Vieira et al., 2016). However, none of these studies linked the pathogen prevalence to social factors or poverty directly and statistically. In our study, we observed a social profile of the studied population, but especially a significantly higher seroprevalence among those in the poor socioeconomic level, which is at the same time related to multiple other potentially associated factors that may be suitable for tick-borne and other tropical diseases.

This study was performed in Colombia, where there is also a lack of seroprevalence studies about *Ehrlichia* spp. (less than 20 in PubMed database), 11 in canine populations, none in Pereira and Risaralda (Arroyave et al., 2020b; Bonilla-Aldana et al., 2021; Forero-Becerra et al., 2021; Hidalgo et al., 2009; McCown et al., 2014; Miranda and Mattar, 2015; Pesapane et al., 2019; Vargas-Hernandez et al., 2012). Given the high seroprevalence obtained in an area with ticks' infestation (data not shown), the potential risk for zoonotic transmission to humans remains latent, deserving immediate seroprevalence studies.

Albeit their importance, emerging infectious diseases such as those caused by *Ehrlichia* and *Anaplasma* spp. are still neglected in many aspects that need to be understood, including clinical and epidemiological topics. Tick-borne diseases may lead to severe clinical consequences in animals and humans, leading to potentially fatal outcomes (Hardalo et al., 1995; Jahangir et al., 1998; Paddock et al., 1993; Tsiodras et al., 2017). Even for some species, there is still questioning whether these may cause, in addition, to dogs, infection in humans, such as is the case of *A. platys* (Arraga-Alvarado et al., 2014). Thus, more studies assessing their circulation in different hosts are needed to understand the ecoepidemiological associated factors, including the social ones, such as poverty (Godfrey and Randolph,




Fig. 4. Seroprevalence in the municipalities of Pereira and Dosquebradas, metropolitan area, rural and urban areas of the capital of Risaralda.

2011; Randolph, 2010; Sumilo et al., 2008). For example, in Eastern Europe, background socioeconomic conditions determine susceptibility to risk of tick-borne encephalitis; some authors suggested that increased unemployment during the economic recession of 2009 triggered a sudden increase in risk (Godfrey and Randolph, 2011). Additionally, regarding ticks, an ongoing assessment (unpublished data) has identified the presence of *Rhipicephalus sanguineus* sensu lato, in the area, then, this is expectable according to the results of the current study. *R. sanguineus* sensu lato has a wide distribution in Colombia according to recent studies (Páez-Triana et al., 2021). In a recent study in Medellín, Colombia, with detection of *E. canis* and *A. platys* among dogs, *R. sanguineus* s.l. was the main tick identified (Arroyave et al., 2020a). Contrary, the presence of anthropophilic *Ixodes* ticks in Colombia is probably very low (Guglielmono et al., 2021), and even, only recently (2021), it was published the first report in the country of a species of the *Ixodes* genus parasitizing a human (Quintero et al., 2021). Then, as expected, the seroprevalence and circulation of *B. burgdorferi* s.l., is rare in

Colombia, although a new putative taxon within the genus *Borrelia* has been detected from certain wild animals (e.g. bats) (Muñoz-Leal et al., 2021). Consistently, a previous study (2014) in Medellín, Barranquilla and Cartagena, also found 0% samples positive for *B. burgdorferi* s.l. in dogs (McCown et al., 2014). In the case of *Dirofilaria*, its prevalence in dogs has been found consistently low in different areas, e.g. 0% in Medellín, 2% in Barranquilla, 3% in Cartagena (McCown et al., 2014). In a recent study, its prevalence among dogs of Bucaramanga was 0.5% (by immunochromatography test kit) and 6.5% (by blood smears and modified Knott's test) (Muñoz et al., 2020).

Although we should acknowledge as a limitation that this study only assessed 100 dogs, preventing the possibility to run multivariate analyses, clearly evaluated multiple relevant social variables in addition to the clinical conditions of the animals, and the seroprevalence by a widely used (Kotwa et al., 2020; Mendes-de-Almeida et al., 2021; Petruccelli et al., 2020; Selim et al., 2021) serological diagnostic test for selected vector-borne pathogens in dogs, in different areas of the municipality, for the first also mapping, using geographical information system (GIS), the location of the assessed population, but particularly the distribution of the seropositive populations. Indeed, the use of GIS for mapping infectious diseases is highly valuable and previously used in our setting for human and animal diseases, such as malaria, Zika, neosporosis, among others (Bonilla-Aldana et al., 2020a; Idarraga-Bedoya et al., 2020; Rodríguez-Morales et al., 2015; Rodríguez-Morales et al., 2017). Its use, as shown here, would be helpful also in ehrlichiosis. These maps clearly showed the high seroprevalence in rural and poor communes in the urban area. The individual assessed conditions suggested that poverty is highly associated with these tick-borne pathogens in Pereira, as observed in this study.

Although *Ehrlichia chaffeensis*, *E. ewingii* and *A. phagocytophilum* are recognized as zoonotic agents, the role *E. canis* and *A. platys* as human pathogens is controversial (Ismail and McBride, 2017; Rar et al., 2021). Some reports have identified *E. canis* and *A. platys* in human samples (Arraga-Alvarado et al., 2014; Bouza-Mora et al., 2017; Perez et al., 2006). In the case of *A. phagocytophilum* this is the etiological agent of the human anaplasmosis (Price et al., 2022; Taank et al., 2020). In the case of *A. platys*, some reports suggest the possible human infection (Arraga-Alvarado et al., 2014). Then, there is a need for surveillance and study of human-dog interaction, assessing the potential for infections also in humans.

No previous studies on the epidemiology of tick-borne diseases and dirofilariasis in the city have been published so far. The current study was a cross-sectional, seroprevalence assessment, but further studies are expected.

5. Conclusions

Given the high seroprevalence obtained in an area with documented ticks, there is a potential risk of zoonotic transmission to humans. Further seroprevalence studies in humans are needed to assess the prevalence of infections. Poverty is highly associated with these tick-borne pathogens in Pereira, as shown in the present study.

Ethics approval and consent to participate

All persons gave their informed consent before their inclusion in the study. Furthermore, according to standard protocols and guidelines from the Animal Ethics Committee (CICUA) at the Fundación Universitaria Autónoma de las Américas, Colombia, animal procedures were performed according to standard protocols and guidelines. Therefore, the CICUA approved and endorsed this study (Acta No. 27 de 2019).

Consent for publication

All persons gave their informed consent before their inclusion in the study.

Availability of data and materials

Available upon reasonable request.

Funding

This study has been funded by Fundación Universitaria Autónoma de las Américas (UAM) (Schools of Veterinary Medicine and Zootechnics, and Medicine, Pereira), Universidad Nacional Autónoma de Honduras (Faculty of Medical Sciences), Sci-Help, Vitalcare, IPS Cardiológica Eduardo Ramírez and Centro de Diagnóstico Especializado Testmol, RECEPA, the family Trejos-Mendoza, and the Dirección de Investigación Científica, Humanística y Tecnológica (DICIHT) (2-05-01-01). The current article processing charges (publication fees) were funded by the Facultad de Ciencias Médicas (FCM) (2-03-01-01), Universidad Nacional Autónoma de Honduras (UNAH), Tegucigalpa, MDC, Honduras, Central America (granted to Dr. Zambrano).

Authors' contributions

DKBA conceived the investigation and supervised the field and laboratory work and data analyses (Conceptualization, Methodology, Project administration).

JPMA, MARM, AETM, SPV, LVM, LFMA, DON, MCP and EG, performed field and laboratory work (Data curation, Investigation); LVGC and AJRM revised and analyzed the clinical records of the patients and supervised laboratory work (Data curation,

Investigation). SPV, AJRM and DKBA prepared the database with laboratory results and clinical data (Formal Analysis). AJRM and DKBA analyzed the data and wrote the first draft of the manuscript (Writing – original draft). All authors revised and approved the final version (Writing – review & editing).

Declaration of Competing Interest

None.

Acknowledgements

Semillero de Investigación en Zoonosis of the UAM. Also, to Diana Marcela Rojas Gallardo and Amparo Juliet Vanegas Quiceno, from the Laboratories of UAM. This study was previously presented in part at the American Society for Microbiology (ASM) & the Federation of European Microbiological Societies (FEMS) World Microbe Forum, June 20-24, 2021 (iPoster #6498). This study was part of the undergraduate thesis for Veterinary Medicine and Zootechnics (DVM) of JP Martínez-Arboleda, MA Reina-Mora, AE Trejos-Mendoza, S Pérez-Vargas, L Valencia-Mejía, LF Marín-Arboleda, D Osorio-Navia, and M Chacón-Peña, under the supervision of DK Bonilla-Aldana.

References

- Alkhishe, A., Raghavan, R.K., Peterson, A.T., 2021. Likely geographic distributional shifts among medically important tick species and tick-associated diseases under climate change in North America: a review. *Insects* 12.
- Arraga-Alvarado, C.M., Qurollo, B.A., Parra, O.C., Berrueta, M.A., Hegarty, B.C., Breitschwerdt, E.B., 2014. Case report: molecular evidence of *Anaplasma platys* infection in two women from Venezuela. *Am. J. Trop. Med. Hyg.* 91, 1161–1165.
- Arroyave, E., Cornwell, E.R., McBride, J.W., Díaz, C.A., Labruna, M.B., Rodas, J.D., 2020a. Detection of tick-borne rickettsial pathogens in naturally infected dogs and dog-associated ticks in Medellín, Colombia. *Rev. Bras. Parasitol. Vet.* 29, e005320.
- Arroyave, E., Rodas-Gonzalez, J.D., Zhang, X., Labruna, M.B., Gonzalez, M.S., Fernandez-Silva, J.A., McBride, J.W., 2020b. *Ehrlichia canis* TRP36 diversity in naturally infected-dogs from an urban area of Colombia. *Ticks Tick Borne Dis.* 11, 101367.
- Baneth, G., Nachum-Biala, Y., Birkenheuer, A.J., Schreag, M.E., Prince, H., Florin-Christensen, M., Schnitger, L., Aroch, I., 2020. A new piroplasmid species infecting dogs: morphological and molecular characterization and pathogen of *Babesia negevi* n. sp. *Parasit. Vectors* 13, 130.
- Bavia, L., Melanda, F.N., de Arruda, T.B., Mosimann, A.L.P., Silveira, G.F., Aoki, M.N., Kuczera, D., Sarzi, M.L., Junior, W.L.C., Conchon-Costa, I., Pavanelli, W.R., Duarte Dos Santos, C.N., Barreto, R.C., Bordignon, J., 2020. Epidemiological study on dengue in southern Brazil under the perspective of climate and poverty. *Sci. Rep.* 10, 2127.
- Bayles, B.R., Allan, B.F., 2014. Social-ecological factors determine spatial variation in human incidence of tick-borne ehrlichiosis. *Epidemiol. Infect.* 142, 1911–1924.
- Bi, Y., Tong, S., 2014. Poverty and malaria in the Yunnan province, China. *Infect. Dis. Poverty* 3, 32.
- Bonifay, T., Douine, M., Bonnefoy, C., Hurpeau, B., Nacher, M., Djossou, F., Epelboin, L., 2017. Poverty and Arbovirus outbreaks: when Chikungunya virus hits more precarious populations than dengue virus in French Guiana. *Open Forum. Infect. Dis.* 4, ofx247.
- Bonilla-Aldana, D.K., Bonilla-Aldana, J.L., Garcia-Bustos, J.J., Lozada, C.O., Rodriguez-Morales, A.J., 2020a. Geographical trends of chikungunya and Zika in the Colombian Amazonian gateway department, Caqueta, 2015-2018 - implications for public health and travel medicine. *Travel Med. Infect. Dis.* 35, 101481.
- Bonilla-Aldana, D.K., Pomares-Cantillo, L.H., Beltran-Sanchez, C.A., Bettin-Martinez, A.C., Campo-Urbina, M.L., Rodriguez-Morales, A.J., Perez-Doria, A., 2020b. Molecular detection of *Anaplasma* spp. in domestic dogs from urban areas of Soledad, Atlantico, Colombia. *Infez. Med.* 28, 373–383.
- Bonilla-Aldana, D.K., Quintero-Rada, K., Montoya-Posada, J.P., Soler-Tovar, D., Barato, P., Arteaga-Livias, K., Zambrano, L.I., Faccini-Martínez, Á.A., Rodriguez-Morales, A.J., 2021. Bovine Ehrlichiosis prevalence: a systematic review and Meta-analysis of molecular studies. *World's Vet. J.* 11, 01–15.
- Bouza-Mora, L., Dolz, G., Solórzano-Morales, A., Romero-Zuñiga, J.J., Salazar-Sánchez, L., Labruna, M.B., Aguiar, D.M., 2017. Novel genotype of *Ehrlichia canis* detected in samples of human blood bank donors in Costa Rica. *Ticks Tick Borne Dis.* 8, 36–40.
- Bukhman, G., Mocumbi, A.O., Atun, R., Becker, A.E., Bhutta, Z., Binagwaho, A., Clinton, C., Coates, M.M., Dain, K., Ezzati, M., Gottlieb, G., Gupta, I., Gupta, N., Hyder, A.A., Jain, Y., Kruk, M.E., Makani, J., Marx, A., Miranda, J.J., Norheim, O.F., Nugent, R., Roy, N., Stefan, C., Wallis, L., Mayosi, B., Lancet, N.P.C.S.G., 2020. The Lancet NCDI poverty commission: bridging a gap in universal health coverage for the poorest billion. *Lancet* 396, 991–1044.
- Cardona-Arias, J.A., 2018. Social determinants of intestinal parasitism, malnutrition, and anemia: systematic review. *Rev. Panam. Salud Publica* 41, e143.
- Chandrashekar, R., Mainville, C.A., Beall, M.J., O'Connor, T., Eberts, M.D., Alleman, A.R., Gaunt, S.D., Breitschwerdt, E.B., 2010. Performance of a commercially available in-clinic ELISA for the detection of antibodies against *Anaplasma phagocytophilum*, *Ehrlichia canis*, and *Borrelia burgdorferi* and *Dirofilaria immitis* antigen in dogs. *Am. J. Vet. Res.* 71, 1443–1450.
- Enriquez, G.F., Macchiaverna, N.P., Argibay, H.D., Lopez Arias, L., Farber, M., Gurtler, R.E., Cardinal, M.V., Garbossa, G., 2019. Polyparasitism and zoonotic parasites in dogs from a rural area of the Argentine Chaco. *Vet. Parasitol. Reg. Stud. Rep.* 16, 100287.
- Ercumen, A., Benjamin-Chung, J., Arnold, B.F., Lin, A., Hubbard, A.E., Stewart, C., Rahman, Z., Parvez, S.M., Unicomb, L., Rahman, M., Haque, R., Colford Jr., J.M., Luby, S.P., 2019. Effects of water, sanitation, handwashing and nutritional interventions on soil-transmitted helminth infections in young children: a cluster-randomized controlled trial in rural Bangladesh. *PLoS Negl. Trop. Dis.* 13, e0007323.
- Forero-Becerra, E., Patel, J., Martínez-Díaz, H.C., Betancourt-Ruiz, P., Benavides, E., Duran, S., Olaya, M.L., Bolanos, E., Hidalgo, M., McBride, J.W., 2021. Seroprevalence and genotypic analysis of *Ehrlichia canis* infection in dogs and humans in Cauca, Colombia. *Am. J. Trop. Med. Hyg.* 104 (5), 1771–1776. <https://doi.org/10.4269/ajtmh.20-0965>.
- Godfrey, E.R., Randolph, S.E., 2011. Economic downturn results in tick-borne disease upsurge. *Parasit. Vectors* 4, 35.
- Guglielmo, A.A., Nava, S., Robbins, R.G., 2021. Animal and Human Parasitism, and Lists of Hosts of Neotropical Ixodidae, Neotropical Hard Ticks (Acari: Ixodida: Ixodidae): A Critical Analysis of their Taxonomy, Distribution, and Host Relationships. Springer International Publishing, Cham, pp. 297–369.
- Hardalo, C.J., Quagliarello, V., Dumler, J.S., 1995. Human granulocytic ehrlichiosis in Connecticut: report of a fatal case. *Clin. Infect. Dis.* 21, 910–914.
- Hernandez-Velasco, A., Sanchez-Montes, S., Romero-Salas, D., Cruz-Romero, A., Jimenez-Hernandez, J.A., Becker, I., Aguilar-Dominguez, M., de Leon, A.P., 2020. First record of natural infection with *Anaplasma marginale* in sucking lice infesting the water buffalo (*Bubalus bubalis*) in Mexico. *Parasitol. Res.* 119, 3853–3856.
- Hidalgo, M., Vesga, J.F., Lizarazo, D., Valbuena, G., 2009. A survey of antibodies against *Rickettsia rickettsii* and Ehrlichia chafeensis in domestic animals from a rural area of Colombia. *Am. J. Trop. Med. Hyg.* 80, 1029–1030.
- Hummel, C., Knaul, F.M., Touchton, M., Guachalla, V.X.V., Nelson-Nunez, J., Boulding, C., 2021. Poverty, precarious work, and the COVID-19 pandemic: lessons from Bolivia. *Lancet Glob. Health* 9, e579–e581.
- Idarraga-Bedoya, S.E., Alvarez-Chica, J., Bonilla-Aldana, D.K., Moore, D.P., Rodriguez-Morales, A.J., 2020. Seroprevalence of Neospora caninum infection in cattle from Pereira, Colombia (small star, filled). *Vet. Parasitol. Reg. Stud. Rep.* 22, 100469.
- Ismail, N., McBride, J.W., 2017. Tick-borne emerging infections: Ehrlichiosis and Anaplasmosis. *Clin. Lab. Med.* 37, 317–340.
- Jahangir, A., Kolbert, C., Edwards, W., Mitchell, P., Dumler, J.S., Persing, D.H., 1998. Fatal pancarditis associated with human granulocytic Ehrlichiosis in a 44-year-old man. *Clin. Infect. Dis.* 27, 1424–1427.

- Jaimes-Duenez, J., Triana-Chavez, O., Mejia-Jaramillo, A.M., 2018. Genetic, host and environmental factors associated with a high prevalence of *Anaplasma marginale*. *Ticks Tick Borne Dis.* 9, 1286–1295.
- Karim, S., Kumar, D., Budachetri, K., 2021. Recent advances in understanding tick and rickettsiae interactions. *Parasite Immunol.* 43, e12830.
- Kotwa, J.D., Jardine, C.M., Pearl, D.L., Berke, O., Mercer, N.J., Peregrine, A.S., 2020. Evaluation of the SNAP(R) 4Dx(R) plus test for the detection of *Dirofilaria immitis* antigen and characterization of exposure to tick-borne pathogens in wild canids in southern Ontario. *Vet. Parasitol.* 283, 109176.
- McCown, M.E., Monteroso, V.H., Cardona, W., 2014. Surveillance for *Ehrlichia canis*, *Anaplasma phagocytophilum*, *Borrelia burgdorferi*, and *Dirofilaria immitis* in dogs from three cities in Colombia. *J. Spec. Oper. Med.* 14, 86–90.
- Mendes-de-Almeida, F., Alves, L.C., do Amaral Fernandes, P., de Menezes Leivas, R., Labarthe, N., 2021. Infection with *Dirofilaria immitis* and Other Infections in Cats and Dogs from Rio de Janeiro. The Need for Prophylactic Enforcement. *Acta Parasitol. Brazil.*
- Miranda, J., Mattar, S., 2015. Molecular detection of *Anaplasma* sp. and *Ehrlichia* sp. in ticks collected in domestic animals, Colombia. *Trop. Biomed.* 32, 726–735.
- Mulligan, K., Dixon, J., Sinn, C.L., Elliott, S.J., 2015. Is dengue a disease of poverty? A systematic review. *Pathog. Glob. Health* 109, 10–18.
- Muñoz, A.A.F., Martínez, A.R., Pinilla, J.C., 2020. Prevalence of *Dirofilaria immitis* in shelter dogs in Bucaramanga metropolitan area, Colombia. *Veterin. Parasitol. Region. Stud. Rep.* 22, 100489.
- Muñoz-Leal, S., Faccini-Martínez, Á.A., Pérez-Torres, J., Chala-Quintero, S.M., Herrera-Sepúlveda, M.T., Cuervo, C., Labruna, M.B., 2021. Novel *Borrelia* genotypes in bats from the Macaregua cave, Colombia. *Zoonoses Public Health* 68, 12–18.
- Negi, T., Kandari, L.S., Arunachalam, K., 2021. Update on prevalence and distribution pattern of tick-borne diseases among humans in India: a review. *Parasitol. Res.* 120, 1523–1539.
- Paddock, C.D., Suchard, D.P., Grumbach, K.L., Hadley, W.K., Kerschmann, R.L., Abbey, N.W., Dawson, J.E., Anderson, B.E., Sims, K.G., Dumler, J.S., et al., 1993. Brief report: fatal seronegative ehrlichiosis in a patient with HIV infection. *N. Engl. J. Med.* 329, 1164–1167.
- Páez-Triana, L., Muñoz, M., Herrera, G., Moreno-Pérez, D.A., Tafur-Gómez, G.A., Montenegro, D., Patarroyo, M.A., Paniz-Mondolfi, A., Ramírez, J.D., 2021. Genetic diversity and population structure of *Rhipicephalus sanguineus* sensu lato across different regions of Colombia. *Parasit. Vectors* 14, 424.
- Peretti-Watel, P., Ward, J., Lutaud, R., Seror, V., 2019. Lyme disease: insight from social sciences. *Med. Mal. Infect.* 49, 133–139.
- Perez, M., Bodor, M., Zhang, C., Xiong, Q., Rikihisa, Y., 2006. Human infection with *Ehrlichia canis* accompanied by clinical signs in Venezuela. *Ann. N. Y. Acad. Sci.* 1078, 110–117.
- Pesapane, R., Foley, J., Thomas, R., Castro, L.R., 2019. Molecular detection and characterization of *Anaplasma platys* and *Ehrlichia canis* in dogs from northern Colombia. *Vet. Microbiol.* 233, 184–189.
- Petruccioli, A., Ferrara, G., Iovane, G., Schettini, R., Ciarcia, R., Caputo, V., Pompameo, M., Pagnini, U., Montagnaro, S., 2020. Seroprevalence of *Ehrlichia* spp., *Anaplasma* spp., *Borrelia burgdorferi* sensu lato, and *Dirofilaria immitis* in stray dogs, from 2016 to 2019, in southern Italy. *Animals (Basel)* 11.
- Pham-Duc, P., Nguyen-Viet, H., Hattendorf, J., Zinsstag, J., Phung-Dac, C., Zurbrugg, C., Odermatt, P., 2013. *Ascaris lumbricoides* and *Trichuris trichiura* infections associated with wastewater and human excreta use in agriculture in Vietnam. *Parasitol. Int.* 62, 172–180.
- Polo, G., Labruna, M.B., Ferreira, F., 2015. Satellite Hyperspectral imagery to support tick-borne infectious diseases surveillance. *PLoS One* 10, e0143736.
- Price, K.J., Ayres, B.N., Maes, S.E., Witmier, B.J., Chapman, H.A., Coder, B.L., Boyer, C.N., Eisen, R.J., Nicholson, W.L., 2022. First detection of human pathogenic variant of *Anaplasma phagocytophilum* in field-collected *Haemaphysalis longicornis*, Pennsylvania, USA. *Zoonoses Public Health* 69, 143–148.
- Quintero, K., Duran, C., Duri, D., Medina, F., Garcia, J., Hidalgo, G., Nakal, S., Echeverría-Ortega, M., Albano, C., Incani, R.N., Cortez, J., Jimenez, S., Diaz, M., Maldonado, C., Matute, F., Rodríguez-Morales, A.J., 2012. Household social determinants of ascariasis and trichuriasis in north Central Venezuela. *Int. Health* 4, 103–110.
- Quintero, J.C., Félix, M.L., Venzal, J.M., Nava, S., 2021. *Ixodes tropicalis* (Acari: Ixodidae) infesting a human and molecular detection of *Rickettsia bellii*, Colombia. *Biomedica* 41, 347–352.
- Randolph, S.E., 2010. To what extent has climate change contributed to the recent epidemiology of tick-borne diseases? *Vet. Parasitol.* 167, 92–94.
- Rar, V., Tkachev, S., Tikunova, N., 2021. Genetic diversity of *Anaplasma* bacteria: twenty years later. *Infect. Genet. Evol.* 91, 104833.
- Ricci, F., 2012. Social implications of malaria and their relationships with poverty. *Mediterr. J. Hematol. Infect. Dis.* 4, e2012048.
- Robles, A., Fong, J., Cervantes, J., 2018. *Borrelia* infection in Latin America. *Rev. Investig. Clin.* 70, 158–163.
- Rodríguez-Morales, A.J., Orrego-Acevedo, C.A., Zambrano-Munoz, Y., Garcia-Folleco, F.J., Herrera-Giraldo, A.C., Lozada-Riascos, C.O., 2015. Mapping malaria in municipalities of the coffee triangle region of Colombia using geographic information systems (GIS). *J. Infect. Public Health* 8, 603–611.
- Rodríguez-Morales, A.J., Ruiz, P., Tabares, J., Ossa, C.A., Yepes-Echeverry, M.C., Ramirez-Jaramillo, V., Galindo-Marquez, M.L., Garcia-Loaiza, C.J., Sabogal-Roman, J.A., Parra-Valencia, E., Lagos-Grisales, G.J., Lozada-Riascos, C.O., de Pijper, C.A., Grobusch, M.P., 2017. Mapping the ecoinfectology of Zika virus infection in urban and rural areas of Pereira, Risaralda, Colombia, 2015–2016: implications for public health and travel medicine. *Travel Med. Infect. Dis.* 18, 57–66.
- Rodríguez-Morales, A.J., Bonilla-Aldana, D.K., Escalera-Antezana, J.P., Alvarado-Arnez, L.E., 2018a. Research on Babesia: a bibliometric assessment of a neglected tick-borne parasite. *F1000Res* 7, 1987.
- Rodríguez-Morales, A.J., Bonilla-Aldana, D.K., Idarraga-Bedoya, S.E., Garcia-Bustos, J.J., Cardona-Ospina, J.A., Faccini-Martínez, A.A., 2018b. Epidemiology of zoonotic tick-borne diseases in Latin America: are we just seeing the tip of the iceberg? *F1000Res* 7, 1988.
- Santodomingo, A., Sierra-Orozco, K., Cotes-Perdomo, A., Castro, L.R., 2019. Molecular detection of *Rickettsia* spp., *Anaplasma platys* and *Theileria equi* in ticks collected from horses in Tayrona National Park, Colombia. *Exp. Appl. Acarol.* 77, 411–423.
- Santos-Vega, M., Bouma, M.J., Kohli, V., Pascual, M., 2016. Population density, climate variables and poverty synergistically structure spatial risk in urban malaria in India. *PLoS Negl. Trop. Dis.* 10, e0005155.
- Selim, A., Alanazi, A.D., Sazmand, A., Otranto, D., 2021. Seroprevalence and associated risk factors for vector-borne pathogens in dogs from Egypt. *Parasit. Vectors* 14, 175.
- Snyman, K., Mwangwa, F., Bigira, V., Kapisi, J., Clark, T.D., Osterbauer, B., Greenhouse, B., Sturrock, H., Gosling, R., Liu, J., Dorsey, G., 2015. Poor housing construction associated with increased malaria incidence in a cohort of young Ugandan children. *Am. J. Trop. Med. Hyg.* 92, 1207–1213.
- Stillman, B.A., Monn, M., Liu, J., Thatcher, B., Foster, P., Andrews, B., Little, S., Eberts, M., Breitschwerdt, E.B., Beall, M.J., Chandrashekar, R., 2014. Performance of a commercially available in-clinic ELISA for detection of antibodies against *Anaplasma phagocytophilum*, *Anaplasma platys*, *Borrelia burgdorferi*, *Ehrlichia canis*, and *Ehrlichia ewingii* and *Dirofilaria immitis* antigen in dogs. *J. Am. Vet. Med. Assoc.* 245, 80–86.
- Sumilo, D., Bormane, A., Asokliene, L., Vasilenko, V., Golovljova, I., Avsic-Zupanc, T., Hubalek, Z., Randolph, S.E., 2008. Socio-economic factors in the differential upsurge of tick-borne encephalitis in central and Eastern Europe. *Rev. Med. Virol.* 18, 81–95.
- Taank, V., Ramasamy, E., Sultana, H., Neelakanta, G., 2020. An efficient microinjection method to generate human anaplasmosis agent *Anaplasma phagocytophilum*-infected ticks. *Sci. Rep.* 10, 15994.
- The Lancet Child Adolescent Health, 2019. Tackling the multidimensionality of child poverty. *Lancet Child Adolesc. Health* 3 (4), 199. [https://doi.org/10.1016/S2352-4642\(19\)30067-7](https://doi.org/10.1016/S2352-4642(19)30067-7).
- Tsiodras, S., Spanakis, N., Spanakos, G., Pervanidou, D., Georgakopoulou, T., Campos, E., Petra, T., Kanellopoulos, P., Georgiadis, G., Antalis, E., Kontos, V., Giannopoulos, L.A., Tselentis, Y., Papa, A., Tsakris, A., Saroglou, G., 2017. Fatal human anaplasmosis associated with macrophage activation syndrome in Greece and the public health response. *J. Infect. Public Health* 10, 819–823.
- Tusting, L.S., Rek, J., Arinaitwe, E., Staedke, S.G., Kanya, M.R., Cano, J., Bottomley, C., Johnston, D., Dorsey, G., Lindsay, S.W., Lines, J., 2016. Why is malaria associated with poverty? Findings from a cohort study in rural Uganda. *Infect. Dis. Poverty* 5, 78.
- Vargas-Hernandez, G., Andre, M.R., Faria, J.L., Munhoz, T.D., Hernandez-Rodriguez, M., Machado, R.Z., Tinucci-Costa, M., 2012. Molecular and serological detection of *Ehrlichia canis* and *Babesia vogeli* in dogs in Colombia. *Vet. Parasitol.* 186, 254–260.
- Vargas-Hernandez, G., Andre, M.R., Cendales, D.M., Sousa, K.C., Goncalves, L.R., Rondelli, M.C., Machado, R.Z., Tinucci-Costa, M., 2016. Molecular detection of *Anaplasma* species in dogs in Colombia. *Rev. Bras. Parasitol. Vet.* 25, 459–464.

- Vieira, T.S., Vieira, R.F., Krawczak, F.S., Soares, H.S., Guimaraes, A.M., Barros-Filho, I.R., Marcondes, M., Labruna, M.B., Biondo, A.W., Vidotto, O., 2016. Ehrlichia sp. infection in carthorses of low-income owners, southern Brazil. *Comp. Immunol. Microbiol. Infect. Dis.* 48, 1–5.
- Walker, A.R., 2011. Eradication and control of livestock ticks: biological, economic and social perspectives. *Parasitology* 138, 945–959.
- Wisely, S.M., Glass, G.E., 2019. Advancing the science of tick and tick-borne disease surveillance in the United States. *Insects* 10.