

catena-Poly[[diaquazinc(II)]- μ -L-cysteinato(2-)- κ^4 S:S,N,O-[di- μ -sulfido-bis[oxidomolybdate(V)](Mo—Mo)]- μ -L-cysteinato(2-)- κ^4 S,N,O:S]

Takashi Shibahara,* Shinobu Ogasahara and Genta Sakane

Department of Chemistry, Okayama University of Science, Ridai-cho, Okayama 700-0005, Japan

Correspondence e-mail: shiba@chem.ous.ac.jp


Received 6 March 2008; accepted 21 March 2008

Key indicators: single-crystal X-ray study; $T = 93$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.020; wR factor = 0.049; data-to-parameter ratio = 21.8.

The title compound, $[Mo_2Zn(C_3H_5NO_2S)_2O_2S_2(H_2O)_2]$, forms a one-dimensional chain. The cysteine S atom of the dinuclear molybdenum complex anion coordinates to the zinc ion, which has a tetrahedral environment by the additional coordination of two water molecules. The one-dimensional chains are connected to each other by hydrogen bonds. The Zn—S(cysteine) distances [2.3599 (6) and 2.3072 (6) Å] are close to the value in ZnS (2.35 Å). The distances and angles within the complex are very close to those reported for the sodium and potassium di- μ -sulfide species.

Related literature

For related literature, see: Brown & Jeffreys (1973); Hong *et al.* (1983); Kay & Mitchell (1970); Knox & Prout (1969); Shibahara *et al.* (1987); Lee *et al.* (1989); Liu & Williams (1981); Xing *et al.* (1998).



Experimental

Crystal data

| | |
|---|-----------------------------------|
| $[Mo_2Zn(C_3H_5NO_2S)_2O_2S_2(H_2O)_2]$ | $V = 843.23$ (16) Å ³ |
| $M_r = 627.69$ | $Z = 2$ |
| Monoclinic, $P2_1$ | $Mo K\alpha$ radiation |
| $a = 8.6881$ (11) Å | $\mu = 3.40$ mm ⁻¹ |
| $b = 10.3529$ (8) Å | $T = 93.1$ K |
| $c = 9.8686$ (11) Å | $0.35 \times 0.30 \times 0.10$ mm |
| $\beta = 108.2022$ (14)° | |

Data collection

| | |
|---|--|
| Rigaku Mercury diffractometer | 9357 measured reflections |
| Absorption correction: multi-scan (Jacobson, 1998) | 4556 independent reflections |
| $T_{\min} = 0.382$, $T_{\max} = 0.727$ | 4549 reflections with $F^2 > 2\sigma(F^2)$ |
| | $R_{\text{int}} = 0.019$ |

Refinement

| | |
|---------------------------------|--|
| $R[F^2 > 2\sigma(F^2)] = 0.019$ | H atoms treated by a mixture of independent and constrained refinement |
| $wR(F^2) = 0.049$ | $\Delta\rho_{\text{max}} = 0.47$ e Å ⁻³ |
| $S = 1.02$ | $\Delta\rho_{\text{min}} = -0.94$ e Å ⁻³ |
| 4556 reflections | Absolute structure: Flack (1983), with 2010 Friedel pairs |
| 209 parameters | Flack parameter: 0.002 (7) |
| 15 restraints | |

Table 1
Selected geometric parameters (Å, °).

| | | | |
|------------------------|------------|-------------------------|-------------|
| Mo1—S1 | 2.3201 (6) | Mo2—S4 | 2.5428 (6) |
| Mo1—S2 | 2.3378 (6) | Zn1—O8 | 2.0052 (17) |
| Mo1—S3 | 2.5572 (6) | Zn1—O7 | 2.0275 (19) |
| Mo1—Mo2 | 2.8354 (3) | Zn1—S4 ⁱ | 2.3072 (6) |
| Mo2—S2 | 2.3276 (6) | Zn1—S3 | 2.3599 (6) |
| Mo2—S1 | 2.3368 (6) | | |
| O8—Zn1—O7 | 96.70 (7) | O8—Zn1—S3 | 93.73 (5) |
| O8—Zn1—S4 ⁱ | 129.42 (5) | O7—Zn1—S3 | 104.55 (6) |
| O7—Zn1—S4 ⁱ | 107.94 (6) | S4 ⁱ —Zn1—S3 | 120.25 (2) |

Symmetry code: (i) $x - 1, y, z$.

Table 2
Hydrogen-bond geometry (Å, °).

| $D-H \cdots A$ | $D-H$ | $H \cdots A$ | $D \cdots A$ | $D-H \cdots A$ |
|----------------------------------|-------|--------------|--------------|----------------|
| O7—H11 \cdots O3 ⁱⁱ | 0.84 | 2.03 | 2.832 (2) | 161 |
| O8—H13 \cdots O5 ⁱⁱ | 0.84 | 1.77 | 2.604 (2) | 171 |
| O8—H14 \cdots O4 ⁱ | 0.84 | 2.00 | 2.789 (2) | 158 |

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x, y + \frac{1}{2}, -z + 2$.

Data collection: *CrystalClear* (Rigaku, 1999); cell refinement: *CrystalClear*; data reduction: *CrystalStructure* (Rigaku, 2007); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *CrystalStructure*; software used to prepare material for publication: *CrystalStructure*.

This work was partly supported by a Special Grant for Cooperative Research administered by the Japan Private School Promotion Foundation.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WK2080).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Brown, D. H. & Jeffreys, J. A. D. (1973). *J. Chem. Soc. Dalton Trans.* pp. 732–735.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Hong, M.-C., Shi, J., Huang, Z. & Liu, H. (1983). *Jiegou Huaxue*, **2**, 153–156.
- Jacobson, R. (1998). Private communication to Rigaku Corporation, Tokyo, Japan.
- Kay, A. & Mitchell, P. C. H. (1970). *J. Chem. Soc. A*, pp. 2421–2428.
- Knox, J. R. & Prout, C. K. (1969). *Acta Cryst. B* **25**, 1857–1866.
- Lee, M. S., Gippert, G. P., Soman, K. V., Case, D. A. & Wright, P. E. (1989). *Science*, **245**, 635–637.
- Liu, H. & Williams, G. J. B. (1981). *Acta Cryst. B* **37**, 2065–2067.
- Rigaku (1999). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2007). *CrystalStructure*. Rigaku Americas Corporation, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shibahara, T., Kuroya, H., Matsumoto, K. & Ooi, S. (1987). *Bull. Chem. Soc. Jpn.* **60**, 2277–2279.
- Xing, Y.-H., Xu, J.-Q., Li, D.-M., Sun, H.-R., Bu, W.-M., Ye, L., Yang, G.-D. & Fan, Y.-G. (1998). *Synth. React. Inorg. Met. Org. Chem.* **28**, 1493–1503.

supplementary materials

Acta Cryst. (2008). E64, m605-m606 [doi:10.1107/S1600536808007757]

catena-Poly[[diaquazinc(II)]- μ -L-cysteinato(2-)- κ^4 S:S,N,O-[di- μ -sulfido-bis[oxidomolybdate(V)](Mo-Mo)]- μ -L-cysteinato(2-)- κ^4 S,N,O:S]

T. Shibahara, S. Ogasahara and G. Sakane

Comment

Molybdenum and L-cysteine are important components of many enzymes. X-ray structures of sulfur/oxygen-bridged dinuclear molybdenum complexes with L-cysteine ligands: $\text{Na}_2[\text{Mo}_2(\mu\text{-S})_2\text{O}_2(\text{cys})_2]\cdot 4\text{H}_2\text{O}$ (Brown & Jeffreys, 1973; Hong *et al.*, 1983); $\text{K}_2[\text{Mo}_2\text{S}_2\text{O}_2(\text{cys})_2]\cdot 4\text{H}_2\text{O}\cdot \text{CH}_3\text{OH}$ (Xing *et al.*, 1998), $\text{Ca}[\text{Mo}_2(\mu\text{-S})(\mu\text{-O})\text{O}_2(\text{cys})_2]\cdot 3\text{H}_2\text{O}$ (Shibahara *et al.*, 1987), and $\text{Na}_2[\text{Mo}_2\text{O}_4(\text{cys})_2]\cdot 5\text{H}_2\text{O}$ (Knox & Prout, 1969; Kay & Mitchell, 1970; Liu & Williams, 1981), have been reported, where alkaline or alkaline earth metals are counter cations, and the existence of metal-oxygen (cysteine oxygen) bonds has been reported. Seeking another crystal structure type, we used Zn^{2+} ion, as the counter ion. The present structural study of the complex compound $\text{Zn}[\text{Mo}_2\text{O}_2\text{S}_2(\text{cys})_2]\cdot 2\text{H}_2\text{O}$ (I) reveals the existence of Zn—S(cysteine sulfur) bonds, which result in polymerization; this type of Zn—S bond has been found in zinc finger proteins. The asymmetric unit of I is shown in Fig. 1 and a view of part of a one-dimensional polymeric chain of I is shown in Fig. 2. The zinc ion bridges the molybdenum complex anions: the coordination of the cysteine sulfur in the complex anion to the zinc ion results in the formation of one dimensional chains, where the zinc forms a tetrahedral structure by the additional coordination of two water molecules. The one dimensional chains are connected to each other by hydrogen bonds. Intra-chain hydrogen bonds also exist. The dimensions of the molybdenum complex and of the zinc tetrahedron are listed in Table 1, and the hydrogen bonds are listed in Table 2. The Zn—S(cysteine) distances (2.3599 (6), 2.3072 (6) Å) are close to that in ZnS (2.35 Å). The distances and angles within the complex are very close to those reported in the sodium and potassium salts in the di- μ -sulfide species.

Experimental

The title compound was prepared by the addition of ZnCl_2 to a diluted aqueous solution of $\text{Na}_2[\text{Mo}_2\text{O}_2\text{S}_2(\text{cys})_2]\cdot 4\text{H}_2\text{O}$. A crystal suitable for single-crystal X-ray diffraction was selected directly from the prepared sample.

Refinement

H atoms bonded to C, N, and O (H_2O) atoms were located in a difference map and refined with distance restraints of C—H = 0.99 (1), N—H = 0.92 (1), and O—H, 0.84 (1) Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N}, \text{O})$. The absolute structure was confirmed by the value of Flack parameter (0.003 (7)).

Figures




Fig. 1. The asymmetric unit of I with atom labels and 50% probability displacement ellipsoids for non-H atoms.

supplementary materials




Fig. 2. A view of part of a one-dimensional polymeric chain with hydrogen bonds (dashed lines).

catena-Poly[[diaquazinc(II)]- μ -L-cysteinato(2-)- κ^4 S,N,O-[di- μ -sulfido- bis[oxidomolybdate(V)](Mo—Mo)]- μ -L-cysteinato(2-)- κ^4 S,N,O:S]

Crystal data

| | |
|---|---|
| [Mo ₂ Zn(C ₃ H ₅ NO ₂ S) ₂ O ₂ S ₂ (H ₂ O) ₂] | $F_{000} = 612.00$ |
| $M_r = 627.69$ | $D_x = 2.472 \text{ Mg m}^{-3}$ |
| Monoclinic, $P2_1$ | Mo $K\alpha$ radiation |
| Hall symbol: P 2yb | $\lambda = 0.71070 \text{ \AA}$ |
| $a = 8.6881 (11) \text{ \AA}$ | Cell parameters from 3004 reflections |
| $b = 10.3529 (8) \text{ \AA}$ | $\theta = 5.5\text{--}30.0^\circ$ |
| $c = 9.8686 (11) \text{ \AA}$ | $\mu = 3.41 \text{ mm}^{-1}$ |
| $\beta = 108.2022 (14)^\circ$ | $T = 93.1 \text{ K}$ |
| $V = 843.23 (16) \text{ \AA}^3$ | Platelet, orange |
| $Z = 2$ | $0.35 \times 0.30 \times 0.10 \text{ mm}$ |

Data collection

| | |
|---|--|
| Rigaku Mercury diffractometer | 4549 reflections with $F^2 > 2\sigma(F^2)$ |
| Detector resolution: 14.63 pixels mm^{-1} | $R_{\text{int}} = 0.019$ |
| ω scans | $\theta_{\text{max}} = 30.0^\circ$ |
| Absorption correction: multi-scan (Jacobson, 1998) | $h = -12 \rightarrow 11$ |
| $T_{\text{min}} = 0.382$, $T_{\text{max}} = 0.727$ | $k = -14 \rightarrow 14$ |
| 9357 measured reflections | $l = -13 \rightarrow 13$ |
| 4556 independent reflections | |

Refinement

| | |
|--|---|
| Refinement on F^2 | $w = 1/[\sigma^2(F_o^2) + (0.0303P)^2 + 0.5406P]$ |
| | where $P = (F_o^2 + 2F_c^2)/3$ |
| $R[F^2 > 2\sigma(F^2)] = 0.019$ | $(\Delta/\sigma)_{\text{max}} = 0.001$ |
| $wR(F^2) = 0.049$ | $\Delta\rho_{\text{max}} = 0.47 \text{ e \AA}^{-3}$ |
| $S = 1.02$ | $\Delta\rho_{\text{min}} = -0.94 \text{ e \AA}^{-3}$ |
| 4556 reflections | Extinction correction: none |
| 209 parameters | Absolute structure: Flack (1983), with 2010 Friedel pairs |
| 15 restraints | Flack parameter: 0.002 (7) |
| H atoms treated by a mixture of independent and constrained refinement | |

*Special details***Geometry.** ENTER SPECIAL DETAILS OF THE MOLECULAR GEOMETRY

Refinement. Refinement was performed using all reflections. The weighted R-factor (wR) and goodness of fit (S) are based on F^2 . R-factor (gt) are based on F. The threshold expression of $F^2 > 2.0 \sigma > F^2$ is used only for calculating R-factor (gt).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|----------------|---------------|---------------|----------------------------------|
| Mo1 | -0.056910 (19) | 0.591229 (16) | 0.808774 (16) | 0.00615 (4) |
| Mo2 | 0.113206 (19) | 0.533750 (19) | 0.613333 (16) | 0.00647 (4) |
| Zn1 | -0.46728 (3) | 0.59442 (3) | 0.84366 (3) | 0.01008 (4) |
| S1 | -0.07812 (7) | 0.40574 (6) | 0.67221 (6) | 0.00917 (8) |
| S2 | 0.16158 (6) | 0.70395 (5) | 0.77534 (6) | 0.00880 (8) |
| S3 | -0.22984 (6) | 0.46976 (5) | 0.93439 (5) | 0.00813 (8) |
| S4 | 0.39318 (6) | 0.59322 (6) | 0.60271 (5) | 0.01009 (8) |
| O1 | -0.2167 (2) | 0.68209 (17) | 0.71583 (17) | 0.0104 (2) |
| O2 | 0.00101 (19) | 0.60344 (18) | 0.45909 (16) | 0.0111 (2) |
| O3 | 0.2203 (2) | 0.45050 (18) | 1.21945 (17) | 0.0125 (3) |
| O4 | 0.13324 (19) | 0.49461 (16) | 0.98677 (16) | 0.0089 (2) |
| O5 | 0.4690 (2) | 0.2383 (2) | 0.7999 (2) | 0.0217 (3) |
| O6 | 0.2865 (2) | 0.39663 (18) | 0.77426 (17) | 0.0125 (3) |
| O7 | -0.3954 (2) | 0.77830 (18) | 0.9006 (2) | 0.0182 (3) |
| O8 | -0.54704 (19) | 0.55572 (18) | 1.00931 (17) | 0.0136 (3) |
| N1 | -0.0163 (2) | 0.70357 (19) | 1.01408 (19) | 0.0087 (3) |
| N2 | 0.1744 (2) | 0.3658 (2) | 0.4962 (2) | 0.0102 (3) |
| C1 | 0.0084 (2) | 0.6095 (2) | 1.1320 (2) | 0.0092 (3) |
| C2 | 0.1322 (2) | 0.5110 (2) | 1.1161 (2) | 0.0083 (3) |
| C3 | -0.1525 (2) | 0.5421 (2) | 1.1147 (2) | 0.0099 (3) |
| C4 | 0.3455 (2) | 0.3284 (2) | 0.5675 (2) | 0.0112 (3) |
| C5 | 0.3694 (2) | 0.3165 (2) | 0.7269 (2) | 0.0116 (4) |
| C6 | 0.4534 (2) | 0.4378 (2) | 0.5447 (2) | 0.0130 (4) |
| H1 | 0.0683 | 0.7596 | 1.0256 | 0.010* |
| H2 | -0.1027 | 0.7551 | 1.0124 | 0.010* |
| H3 | 0.1090 | 0.2957 | 0.4919 | 0.012* |
| H4 | 0.1624 | 0.3890 | 0.4039 | 0.012* |
| H5 | -0.2299 | 0.6042 | 1.1334 | 0.012* |
| H6 | -0.1366 | 0.4727 | 1.1869 | 0.012* |
| H7 | 0.0460 | 0.6482 | 1.2291 | 0.011* |
| H8 | 0.4318 | 0.4416 | 0.4403 | 0.016* |
| H9 | 0.5706 | 0.4243 | 0.5928 | 0.016* |
| H10 | 0.3821 | 0.2458 | 0.5359 | 0.013* |
| H11 | -0.3638 | 0.8290 | 0.8487 | 0.022* |
| H12 | -0.4303 | 0.8035 | 0.9659 | 0.022* |
| H13 | -0.5293 | 0.6197 | 1.0646 | 0.016* |
| H14 | -0.6424 | 0.5281 | 0.9816 | 0.016* |

supplementary materials

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|--------------|--------------|--------------|---------------|--------------|---------------|
| Mo1 | 0.00760 (7) | 0.00633 (8) | 0.00567 (7) | 0.00053 (6) | 0.00374 (5) | 0.00047 (6) |
| Mo2 | 0.00756 (7) | 0.00729 (8) | 0.00565 (7) | 0.00067 (6) | 0.00364 (5) | 0.00022 (6) |
| Zn1 | 0.01030 (10) | 0.01118 (12) | 0.00928 (10) | 0.00082 (10) | 0.00382 (8) | -0.00105 (9) |
| S1 | 0.0114 (2) | 0.0085 (2) | 0.0091 (2) | -0.00148 (18) | 0.00549 (17) | -0.00119 (17) |
| S2 | 0.0109 (2) | 0.0077 (2) | 0.0098 (2) | -0.00074 (18) | 0.00601 (17) | -0.00107 (17) |
| S3 | 0.0087 (2) | 0.0085 (2) | 0.0086 (2) | -0.00003 (18) | 0.00480 (16) | -0.00014 (17) |
| S4 | 0.0088 (2) | 0.0127 (2) | 0.0098 (2) | -0.0003 (2) | 0.00436 (16) | 0.00152 (19) |
| O1 | 0.0111 (6) | 0.0117 (8) | 0.0102 (6) | 0.0024 (5) | 0.0060 (5) | 0.0015 (5) |
| O2 | 0.0113 (6) | 0.0135 (8) | 0.0097 (6) | 0.0019 (6) | 0.0048 (5) | 0.0014 (6) |
| O3 | 0.0152 (7) | 0.0123 (8) | 0.0101 (6) | 0.0030 (6) | 0.0039 (5) | 0.0014 (6) |
| O4 | 0.0102 (6) | 0.0099 (7) | 0.0074 (6) | 0.0011 (5) | 0.0041 (5) | 0.0002 (5) |
| O5 | 0.0216 (8) | 0.0185 (9) | 0.0221 (8) | 0.0064 (7) | 0.0027 (7) | 0.0083 (7) |
| O6 | 0.0166 (7) | 0.0129 (8) | 0.0087 (6) | 0.0010 (6) | 0.0049 (5) | -0.0004 (5) |
| O7 | 0.0217 (9) | 0.0131 (8) | 0.0250 (8) | -0.0031 (7) | 0.0146 (7) | -0.0054 (7) |
| O8 | 0.0096 (6) | 0.0217 (10) | 0.0103 (6) | -0.0025 (5) | 0.0045 (5) | -0.0045 (6) |
| N1 | 0.0111 (7) | 0.0069 (8) | 0.0096 (7) | 0.0002 (6) | 0.0053 (6) | 0.0004 (6) |
| N2 | 0.0096 (7) | 0.0125 (9) | 0.0084 (7) | 0.0010 (6) | 0.0026 (6) | -0.0032 (6) |
| C1 | 0.0131 (8) | 0.0086 (10) | 0.0067 (7) | 0.0002 (7) | 0.0043 (6) | -0.0005 (7) |
| C2 | 0.0099 (8) | 0.0060 (10) | 0.0107 (8) | -0.0019 (6) | 0.0058 (6) | -0.0008 (6) |
| C3 | 0.0118 (8) | 0.0109 (9) | 0.0088 (7) | -0.0005 (7) | 0.0060 (6) | 0.0001 (7) |
| C4 | 0.0114 (8) | 0.0130 (10) | 0.0088 (8) | 0.0038 (7) | 0.0024 (6) | -0.0032 (7) |
| C5 | 0.0126 (9) | 0.0095 (10) | 0.0124 (9) | -0.0007 (7) | 0.0034 (7) | 0.0003 (7) |
| C6 | 0.0117 (9) | 0.0186 (11) | 0.0099 (8) | 0.0019 (7) | 0.0052 (7) | -0.0024 (8) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|---------------------|-------------|--------|-----------|
| Mo1—O1 | 1.6905 (17) | O5—C5 | 1.238 (3) |
| Mo1—O4 | 2.2366 (16) | O6—C5 | 1.279 (3) |
| Mo1—N1 | 2.2662 (19) | O7—H11 | 0.837 |
| Mo1—S1 | 2.3201 (6) | O7—H12 | 0.835 |
| Mo1—S2 | 2.3378 (6) | O8—H13 | 0.841 |
| Mo1—S3 | 2.5572 (6) | O8—H14 | 0.838 |
| Mo1—Mo2 | 2.8354 (3) | N1—C1 | 1.481 (3) |
| Mo2—O2 | 1.6914 (16) | N1—H1 | 0.915 |
| Mo2—N2 | 2.2419 (19) | N1—H2 | 0.917 |
| Mo2—O6 | 2.3044 (18) | N2—C4 | 1.484 (3) |
| Mo2—S2 | 2.3276 (6) | N2—H3 | 0.914 |
| Mo2—S1 | 2.3368 (6) | N2—H4 | 0.915 |
| Mo2—S4 | 2.5428 (6) | C1—C3 | 1.523 (3) |
| Zn1—O8 | 2.0052 (17) | C1—C2 | 1.526 (3) |
| Zn1—O7 | 2.0275 (19) | C1—H7 | 0.995 |
| Zn1—S4 ⁱ | 2.3072 (6) | C3—H5 | 0.989 |
| Zn1—S3 | 2.3599 (6) | C3—H6 | 0.991 |
| S3—C3 | 1.852 (2) | C4—C5 | 1.527 (3) |

| | | | |
|------------------------|--------------|---------------------------|-------------|
| S4—C6 | 1.838 (3) | C4—C6 | 1.531 (4) |
| S4—Zn1 ⁱⁱ | 2.3072 (6) | C4—H10 | 0.996 |
| O3—C2 | 1.237 (3) | C6—H8 | 0.989 |
| O4—C2 | 1.290 (2) | C6—H9 | 0.989 |
| O1—Mo1—O4 | 162.73 (7) | C6—S4—Zn1 ⁱⁱ | 102.50 (7) |
| O1—Mo1—N1 | 94.11 (7) | C6—S4—Mo2 | 99.79 (8) |
| O4—Mo1—N1 | 69.93 (6) | Zn1 ⁱⁱ —S4—Mo2 | 99.22 (2) |
| O1—Mo1—S1 | 104.04 (6) | C2—O4—Mo1 | 119.10 (14) |
| O4—Mo1—S1 | 89.50 (4) | C5—O6—Mo2 | 117.99 (14) |
| N1—Mo1—S1 | 154.97 (5) | Zn1—O7—H11 | 123.4 |
| O1—Mo1—S2 | 102.18 (6) | Zn1—O7—H12 | 110.6 |
| O4—Mo1—S2 | 84.32 (4) | H11—O7—H12 | 122.9 |
| N1—Mo1—S2 | 87.91 (5) | Zn1—O8—H13 | 109.3 |
| S1—Mo1—S2 | 104.59 (2) | Zn1—O8—H14 | 111.1 |
| O1—Mo1—S3 | 91.69 (6) | H13—O8—H14 | 116.3 |
| O4—Mo1—S3 | 78.52 (4) | C1—N1—Mo1 | 107.99 (13) |
| N1—Mo1—S3 | 76.70 (5) | C1—N1—H1 | 113.9 |
| S1—Mo1—S3 | 85.55 (2) | Mo1—N1—H1 | 109.3 |
| S2—Mo1—S3 | 160.022 (19) | C1—N1—H2 | 108.3 |
| O1—Mo1—Mo2 | 105.18 (6) | Mo1—N1—H2 | 112.6 |
| O4—Mo1—Mo2 | 91.51 (4) | H1—N1—H2 | 104.8 |
| N1—Mo1—Mo2 | 138.35 (5) | C4—N2—Mo2 | 108.76 (13) |
| S1—Mo1—Mo2 | 52.760 (15) | C4—N2—H3 | 109.3 |
| S2—Mo1—Mo2 | 52.411 (14) | Mo2—N2—H3 | 113.3 |
| S3—Mo1—Mo2 | 137.441 (15) | C4—N2—H4 | 109.0 |
| O2—Mo2—N2 | 91.69 (8) | Mo2—N2—H4 | 109.8 |
| O2—Mo2—O6 | 162.14 (7) | H3—N2—H4 | 106.6 |
| N2—Mo2—O6 | 70.64 (6) | N1—C1—C3 | 108.11 (17) |
| O2—Mo2—S2 | 103.00 (6) | N1—C1—C2 | 107.01 (16) |
| N2—Mo2—S2 | 156.97 (5) | C3—C1—C2 | 109.49 (18) |
| O6—Mo2—S2 | 93.10 (5) | N1—C1—H7 | 114.5 |
| O2—Mo2—S1 | 102.18 (6) | C3—C1—H7 | 107.9 |
| N2—Mo2—S1 | 89.44 (5) | C2—C1—H7 | 109.8 |
| O6—Mo2—S1 | 80.89 (5) | O3—C2—O4 | 123.7 (2) |
| S2—Mo2—S1 | 104.38 (2) | O3—C2—C1 | 121.88 (18) |
| O2—Mo2—S4 | 98.59 (6) | O4—C2—C1 | 114.38 (18) |
| N2—Mo2—S4 | 77.44 (5) | C1—C3—S3 | 109.75 (13) |
| O6—Mo2—S4 | 75.53 (5) | C1—C3—H5 | 109.6 |
| S2—Mo2—S4 | 82.83 (2) | S3—C3—H5 | 112.2 |
| S1—Mo2—S4 | 155.71 (2) | C1—C3—H6 | 109.6 |
| O2—Mo2—Mo1 | 104.26 (5) | S3—C3—H6 | 109.0 |
| N2—Mo2—Mo1 | 140.49 (5) | H5—C3—H6 | 106.7 |
| O6—Mo2—Mo1 | 91.59 (4) | N2—C4—C5 | 107.66 (18) |
| S2—Mo2—Mo1 | 52.737 (15) | N2—C4—C6 | 107.82 (19) |
| S1—Mo2—Mo1 | 52.226 (15) | C5—C4—C6 | 108.60 (18) |
| S4—Mo2—Mo1 | 133.300 (15) | N2—C4—H10 | 116.5 |
| O8—Zn1—O7 | 96.70 (7) | C5—C4—H10 | 107.4 |
| O8—Zn1—S4 ⁱ | 129.42 (5) | C6—C4—H10 | 108.7 |

supplementary materials

| | | | |
|-------------------------|-------------|----------|-------------|
| O7—Zn1—S4 ⁱ | 107.94 (6) | O5—C5—O6 | 125.6 (2) |
| O8—Zn1—S3 | 93.73 (5) | O5—C5—C4 | 119.9 (2) |
| O7—Zn1—S3 | 104.55 (6) | O6—C5—C4 | 114.4 (2) |
| S4 ⁱ —Zn1—S3 | 120.25 (2) | C4—C6—S4 | 110.80 (15) |
| Mo1—S1—Mo2 | 75.014 (19) | C4—C6—H8 | 104.8 |
| Mo2—S2—Mo1 | 74.853 (18) | S4—C6—H8 | 108.1 |
| C3—S3—Zn1 | 98.94 (7) | C4—C6—H9 | 114.2 |
| C3—S3—Mo1 | 100.08 (7) | S4—C6—H9 | 109.0 |
| Zn1—S3—Mo1 | 97.09 (2) | H8—C6—H9 | 109.7 |


Symmetry codes: (i) $x-1, y, z$; (ii) $x+1, y, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|----------------------------|--------------|--------------------|-------------|----------------------|
| O7—H11···O3 ⁱⁱⁱ | 0.84 | 2.03 | 2.832 (2) | 161 |
| O7—H11···O4 ⁱⁱⁱ | 0.84 | 2.75 | 3.139 (2) | 110 |
| O7—H12···O5 ⁱⁱⁱ | 0.84 | 2.53 | 3.239 (3) | 144 |
| O7—H12···O6 ⁱⁱⁱ | 0.84 | 2.66 | 3.285 (2) | 133 |
| O7—H12···O8 ^{iv} | 0.84 | 2.64 | 3.093 (2) | 116 |
| O8—H13···O5 ⁱⁱⁱ | 0.84 | 1.77 | 2.604 (2) | 171 |
| O8—H14···O4 ⁱ | 0.84 | 2.00 | 2.789 (2) | 158 |
| O8—H14···O6 ⁱ | 0.84 | 2.37 | 2.844 (2) | 116 |
| N1—H2···O4 ⁱⁱⁱ | 0.92 | 2.49 | 3.179 (2) | 132 |
| N1—H2···O7 | 0.92 | 2.45 | 3.224 (2) | 143 |
| N2—H3···O2 ^v | 0.91 | 2.32 | 3.212 (2) | 164 |
| N2—H4···O1 ^v | 0.92 | 2.56 | 2.934 (2) | 105 |
| N2—H4···O3 ^{vi} | 0.92 | 2.13 | 3.011 (2) | 161 |

Symmetry codes: (iii) $-x, y+1/2, -z+2$; (iv) $-x-1, y+1/2, -z+2$; (i) $x-1, y, z$; (v) $-x, y-1/2, -z+1$; (vi) $x, y, z-1$.

Fig. 1



supplementary materials

Fig. 2

