

POSTER PRESENTATION

Open Access

Dynamics of self-sustained activity in random networks with strong synapses

Håkon Enger^{1*}, Tom Tetzlaff¹, Birgit Kriener¹, Marc-Oliver Gewaltig^{2,3}, Gaute T Einevoll¹

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

An understanding of short-term memory requires models of neural networks which are able to sustain activity in the absence of external input for several seconds. About half a century ago, [1] predicted the existence of self-sustained activity in neural networks with strong synapses. Despite this finding, most previous studies on the dynamics of neural networks are restricted to weak synapses, i.e. to a regime where the diffusion approximation is applicable. Recently, it has been shown in simulations that self-sustained activity emerges in networks of integrate-and-fire (IaF) neurons with strong synapses modeled as currents [2] (rather than as conductances, see [3]). Above a critical synaptic weight, the lifetime of

self-sustained activity increases rapidly (Fig. 1C). We present a stochastic model of the dynamics of a balanced random network of IaF neurons with current-based synapses in the strong-synapse regime. Based on the network's firing-rate transfer [4], we show that the firing-rate dynamics becomes bistable if the synapses are sufficiently strong: in addition to the quiescent state, a second stable fixed point at moderate firing rates is created (sketched in Fig. 1A). Firing-rate fluctuations can destabilize this fixed point, thereby limiting the lifetime of self-sustained activity (sketched in Fig. 1B). The magnitude of these fluctuations is mainly determined by the amount of spike-train correlations [5]. Our model




Figure 1 A. Phase space of the firing-rate dynamics. Dependence of the rate change dr/dt on the rate r (sketch). **B.** Firing-rate distribution in the self-sustained state (sketch). **C.** Dependence of the lifetime of self-sustained activity (color coded) on the level g of inhibition and the synaptic weight J in a random network of 100000 excitatory and 25000 inhibitory IaF neurons with 1% connectivity (simulation results).

* Correspondence: hakon.enger@umb.no

¹Dept. of Mathematical Sciences and Technology, Norwegian University of Life Sciences, NO-1432 Ås, Norway

Full list of author information is available at the end of the article

explains the existence and the lifetime of self-sustained activity, and how the lifetime depends on the network size, the connectivity, the level of inhibition and the synapse strength. The results of our model are confirmed by network simulations.

Acknowledgements

We acknowledge support by the NOTUR and eScience programs of the Research Council of Norway. All network simulations were carried out with the neural simulation tool NEST (see <http://www.nest-initiative.org>).

Author details

¹Dept. of Mathematical Sciences and Technology, Norwegian University of Life Sciences, NO-1432 Ås, Norway. ²Honda Research Institute Europe GmbH, D-63073 Offenbach/Main, Germany. ³Bernstein Center for Computational Neuroscience, D-79104 Freiburg, Germany.

Published: 18 July 2011

References

1. Griffiths JS: **On the stability of brain-like structures.** *Biophys J* 1963, **3**:299-308.
2. Gewaltig M: **Self-sustained activity in networks of integrate and fire neurons without external noise.** *Front. Comput. Neurosci. Conference Abstract: Bernstein Conference on Computational Neuroscience 2009*, doi: 10.3389/conf.neuro.10.2009.14.051.
3. Kuhn A, Aertsen A, Rotter S: **Neuronal integration of synaptic input in the fluctuation-driven regime.** *J Neurosci* 2004, **24**(10):2345-2356.
4. Siegert AJF: **On the first passage time probability problem.** *Phys Rev* 1951, **81**:617-623.
5. Kriener B, Tetzlaff T, Aertsen A, Diesmann M, Rotter S: **Correlations and population dynamics in cortical networks.** *Neural Comput* 2008, **20**:2185-2226.

doi:10.1186/1471-2202-12-S1-P89

Cite this article as: Enger *et al.*: Dynamics of self-sustained activity in random networks with strong synapses. *BMC Neuroscience* 2011 **12**(Suppl 1):P89.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

