

Paradoxical effects of ZIM3, a CRISPRi effector, on human induced pluripotent stem-cell-derived cardiomyocyte electrophysiology

Julie L. Han , Yuli W. Heinson^a, Maria R. Pozo , Weizhen Li  and Emilia Entcheva ^{a,*}

^aDepartment of Biomedical Engineering, The George Washington University, Washington, DC 20052, USA

*To whom correspondence should be addressed: Email: entcheva@gwu.edu

Edited By: Dennis Discher

Abstract

We show that zinc finger imprinted 3 (Zim3), when used as Zim3-KRAB-dCas9 effector in interference CRISPR, without any guide RNAs, paradoxically up-regulates key cardiac ion channel genes in human-induced pluripotent stem-cell-derived cardiomyocytes (iPSC-CMs), responsible for healthy resting membrane potential, repolarization of the action potential, and electrical transmission of signals. These were found to yield expected functional enhancements consistent with a more mature iPSC-CM phenotype, with potentially desirable properties.

Zinc finger imprinted 3 (Zim3 or ZNF657/264) is a protein-coding gene in humans, found at low levels in testis, brain, skeletal muscle, and skin (1). As other ZNF-KRAB (Krueppel-associated box) domains, which bind to DNA and through complexing with KRAB-associated protein 1 (KAP1), affect chromatin organization (2), Zim3-KRAB may be involved in the regulation of transcription at multiple genomic loci. In the human heart, the expression of Zim3/ZNF657/ZNF264 has not been examined in detail. We analyzed RNAseq data in human adult heart (left ventricle [LV]) from the Gene Tissue Expression (GTEx) database (3) and in human-induced pluripotent stem-cell-derived cardiomyocytes (iPSC-CMs; Fig. 1). The Zim3 gene identifier was only expressed in a very small fraction of the samples, at negligible levels, both in the LV and in our iPSC-CMs. ZNF657 was not expressed in either group of samples. The ZNF264 gene was most consistently expressed—in all adult heart samples and in all iPSC-CM samples but also at very low levels. For comparison, we also show the gene expression for KCNH2, encoding for the main repolarizing K⁺ channel in cardiomyocytes. To quantitatively compare the two datasets, we normalized the transcripts per million (TPM) by the respective geometric mean (GM) across the transcriptome in each sample and displayed log₁₀ of the ratios (4). The adult human heart (LV) has a higher ZNF264 expression than the human iPSC-CMs; however, both have very low expression overall.

Recently, Zim3-KRAB has been identified as a potent effector (repressor) in combination with dCas9 for interference CRISPR (CRISPRi) gene regulation studies (5). Indeed, using Zim3-KRAB-dCas9 adenoviral delivery in combination with suitable gRNAs in postdifferentiated iPSC-CMs, we have shown robust suppression of the expression of KCNH2 with concomitant functional




Fig. 1. Endogenous gene expression of Zim3/ZNF657/ZNF264 in human adult heart (LV) from GTEx and in human iPSC-CMs, along a well-expressed cardiac gene, KCNH2. The Zim3 and ZNF657 gene identifiers were either not expressed or expressed in only a small subset of samples at negligible levels. For ZNF264 and for KCNH2, we show log₁₀ of normalized TPM, log₁₀(TPM/GM).

consequence (6). While performing control experiments, we observed unexpected effects of simply expressing Zim3-KRAB-dCas9 alone on cardiac electrophysiology, without the addition of any gRNAs or with the use of scrambled/control gRNA.

Competing Interest: E.E. and J.L.H. have filed a provisional patent.

Received: June 2, 2023. **Accepted:** January 12, 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of National Academy of Sciences. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.




Fig. 2. Effects of dCas9/Zim3 expression in human iPSC-CMs: RNA, protein, and electrophysiological consequences. A) Experimental workflow with all-optical electrophysiology (7) and molecular characterization. B) Expression of mCherry tagged constructs (mCherry, MeCP2-KRAB-dCa9-mCherry, and Zim3-KRAB-dCa9-mCherry) in human iPSC-CMs. Scale bar 10 μm . C) Functional results from all-optical electrophysiology: changes in APD and calcium transient duration under 0.5 Hz pacing ($n=9-34$ biologically independent samples). D) Relative changes in spontaneous frequency (delta) with respect to the mean for the dCas9/Zim3 group for each run ($n=14-40$ biologically independent samples per group). E) Changes in conduction shown in example activation maps and as normalized conduction velocity (CV_{norm}) at 0.75 and 1.0 Hz pacing ($n=2-9$ biologically independent samples). Data shown as mean \pm SEM. F) Molecular analysis (qPCR) in pristine samples (top) and after all-optical electrophysiology (bottom)—KCNH2, GJA1, and KCHJ2 ($n=4-16$ biologically independent samples, $n=3$ technical replicates). G) Western blot (Wes by ProteinSimple) of Cx43 post all-optical electrophysiology ($n=6-13$ biologically independent samples). Statistical comparisons were done using one-way ANOVA with post hoc Tukey correction. In all cases, $**P < 0.01$. Biorender was used.

Namely, we found Zim3-KRAB-dCas9 expression to significantly up-regulate the mRNA expression of key cardiac ion channels: KCNJ2, encoding for the Kir2.1 protein—inward rectifier K⁺ channel, contributing to the maintenance of healthy negative resting membrane potential; KCNH2, encoding for the rapid delayed rectifier that controls action potential duration (APD); and GJA1, encoding for Cx43—the main gap-junctional protein in ventricular cardiomyocytes; for Cx43 this was also confirmed at the protein level. Using all-optical electrophysiology (Fig. 2A and B), we corroborated physiologically relevant functional effects (Fig. 2C–E), expected for up-regulated KCNH2 (APD shortening in paced conditions; Fig. 2C and F); up-regulated KCNJ2 (significantly reduced spontaneous firing; Fig. 2D and F);

and up-regulated GJA1 (conduction velocity increase; Fig. 2E–G). In combination, these electrophysiological changes are perceived as desirable and a signature of more mature iPSC-CMs.

These gene expression changes and functional effects cannot be attributed to the expression of dCas9, as we have shown that regular KRAB-dCas9 (using the KRAB domain of KOX1/ZNF10) without gRNAs does not alter these genes (6). The fluorescent reporter, mCherry, also does not have such effects. Included here earlier-generation effector construct for CRISPRi—MeCP2-KRAB-dCas9 (a non-KRAB effector, MeCP2, combined with KOX1/ZNF10) (8) caused similar, but much milder changes when delivered in the same type of viral vector—adenovirus with a spleen

focus-forming virus (SFFV) promoter and an mCherry reporter, at the same multiplicity of infection (MOI) 1,000, as Zim3-KRAB-dCas9. Interestingly, MeCP2 has been shown previously to act not only as a transcriptional repressor but also as an activator for many genes (9). No such studies have been conducted with Zim3/ZNF657/ZNF264. It is possible that some but not all KRAB domains or transcription effectors, alone, or when linked to dCas9 can exert the effects seen here. We cannot exclude the SFFV promoter as a possible culprit—it has rarely been used with cardiomyocytes before and is shared by these two constructs (not used in the original KRAB-dCas9 delivery in Ref. (6)). Overall, we are unaware of any single perturbation acting as a potent agonist for all three genes. These findings should be taken into consideration when performing CRISPRi studies in iPSC-CMs. Zim3 also needs to be investigated further as a potential therapeutic tool for iPSC-CMs maturation for their use in drug screening, disease modeling, and regenerative applications; furthermore, it would be interesting to explore if Zim3 may exert similar effects in vivo in the adult human heart as a therapeutic tool.

Supplementary Material

[Supplementary material](#) is available at *PNAS Nexus* online.

Funding

The authors declare no funding.

Author Contributions

J.L.H. and E.E. designed the experiments. Y.W.H. built some of the imaging tools needed. J.L.H. and Y.W.H. performed most experiments and analyzed the data. M.R.P. and W.L. conducted and analyzed some of the functional experiments. E.E. and J.L.H. wrote the article.

Preprints

This manuscript was posted as a preprint: doi.org/10.1101/2023.06.10.544468.

Data Availability

All data are included in the presented figure.

References

- Kim J, Bergmann A, Wehri E, Lu X, Stubbs L. 2001. Imprinting and evolution of two Kruppel-type zinc-finger genes, ZIM3 and ZNF264, located in the PEG3/USP29 IMPRINTED DDomain. *Genomics*. 77:91–98.
- Stoll GA, Pandiloski N, Douse CH, Modis Y. 2022. Structure and functional mapping of the KRAB-KAP1 repressor complex. *EMBO J*. 41:e111179.
- GTEX Consortium. 2020. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science*. 369:1318–1330.
- Pressler MP, Horvath A, Entcheva E. 2022. Sex-dependent transcription of cardiac electrophysiology and links to acetylation modifiers based on the GTEx database. *Front Cardiovasc Med*. 9:941890.
- Alerasool N, Segal D, Lee H, Taipale M. 2020. An efficient KRAB domain for CRISPRi applications in human cells. *Nat Methods*. 17:1093–1096.
- Han JL, Heinson YW, Chua CJ, Liu W, Entcheva E. 2023. CRISPRi gene modulation and all-optical electrophysiology in post-differentiated human iPSC-cardiomyocytes. *Commun Biol*. 6:1236.
- Heinson YW, Han JL, Entcheva E. 2023. OptoDyCE-plate as an affordable high throughput imager for all optical cardiac electrophysiology. *J Mol Cell Cardiol Plus*. 6:100054.
- Yeo NC, et al. 2018. An enhanced CRISPR repressor for targeted mammalian gene regulation. *Nat Methods*. 15:611–616.
- Chahrour M, et al. 2008. MeCP2, a key contributor to neurological disease, activates and represses transcription. *Science*. 320:1224–1229.