

## Laboratory Investigation

# Effects of Tumor Necrosis Factor Alpha Blocker Adalimumab in Experimental Spinal Cord Injury

Alp Özgün Börcek, M.D.,<sup>1</sup> Soner Çivi, M.D.,<sup>2</sup> Özgür Öcal, M.D.,<sup>1</sup> Özlem Gülbahar, M.D.<sup>3</sup>

Department of Neurosurgery,<sup>1</sup> Gazi University Faculty of Medicine, Ankara, Turkey

Department of Neurosurgery,<sup>2</sup> Medicana Private Hospital, Konya, Turkey

Department of Biochemistry,<sup>3</sup> Gazi University Faculty of Medicine, Ankara, Turkey

**Objective :** Tumor necrosis factor alpha (TNF- $\alpha$ ) have proven effects in pathogenesis of neuroinflammation after spinal cord injury (SCI). Current study is designed to evaluate the effects of an anti-TNF- $\alpha$  agent, adalimumab, on spinal cord clip compression injury in rats.

**Methods :** Thirty two male adult Wistar rats were divided into four groups (sham, trauma, infliximab, and adalimumab groups) and SCI was introduced using an aneurysm clip. Animals in treatment groups received 5 mg/kg subcutaneous adalimumab and infliximab right after the trauma. Malondialdehyde (MDA) levels were studied in traumatized spinal cord tissues 72 hours after the injury as a marker of lipid peroxidation.

**Results :** Animals that received anti-TNF- $\alpha$  agents are found to have significantly decreased MDA levels. MDA levels were significantly different between the trauma and infliximab groups ( $p < 0.01$ ) and trauma and adalimumab groups ( $p = 0.022$ ). There was no significant difference in neurological evaluation of the rats using Tarlov scale.

**Conclusion :** These results suggest that, like infliximab, adalimumab has favorable effects on lipid peroxidation induced by spinal cord trauma in rats.

**Key Words :** Adalimumab · Spinal cord injury · Tumor necrosis factor alpha.

## INTRODUCTION

Spinal cord injury (SCI) with its devastating burden both for the victim and the health care system continues to be a great problem without significant improvement in management. The annual incidence of SCI is estimated as 10–50 cases per million<sup>37,40</sup> and there is no established treatment modality on clinical base except for methylprednisolone even though the effect is controversial<sup>7-12</sup>.

Present basic science experiment is designed to demonstrate the effects of a potential pharmacological agent—adalimumab (Humira, Abbott, Abbott Park, IL, USA)—on SCI in rats. Previously we have demonstrated the effects of a tumor necrosis factor-alpha (TNF- $\alpha$ ) inhibitor infliximab's (Remicade, Merck&Co, Whitehouse Station, NJ, USA) effects on different experimental models of neuronal injury<sup>24,31,32</sup>. Adalimumab, human recombinant monoclonal IgG1 antibody specific for cytokine TNF- $\alpha$ , is the third TNF- $\alpha$  inhibitor approved for inflammatory diseases such as ulcerative colitis after infliximab and etanercept<sup>14</sup>.

As emerged from the experimental knowledge, injury to neu-

ral structures results from two different mechanisms. The first is the primary insult that is resulting from the mechanical event itself and the other is the secondary insult that is resulting mainly from the inflammatory response of the organism's to the primary insult. Neutrophils are blamed for orchestration of secondary injury by releasing various types of enzymes, reactive oxygen species and proinflammatory factors. These factors play important roles in glio-neuronal cell damage<sup>1,2,19,25-27,30</sup>.

## MATERIALS AND METHODS

All experiments were approved by our Institutional Review Board and performed in accordance with the local guidelines to minimize animal discomfort.

Thirty two male adult Wistar rats weighing between 250 and 300 g were used in this study. Animals were kept under constant laboratory conditions. After spinal cord trauma, manual emptying of the bladders was performed for the injured animals.

The experimental methodology used in this study was described in detail elsewhere<sup>32</sup>. Briefly, after the period of habituation, rats

• Received : March 10, 2014 • Revised : August 4, 2014 • Accepted : August 24, 2014

• Address for reprints : Alp Özgün Börcek, M.D.

Department of Neurosurgery, Gazi University Faculty of Medicine, Beyin ve Sinir Cerrahisi AD, Ankara 06500, Turkey

Tel : +90-312 202 5110, Fax : +90-312 212 90 04, E-mail : alpborecek@gmail.com

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

were randomly divided into four groups. Induction of anesthesia was performed with proper administration of ketamine hydrochloride (50 mg/kg, Ketalar, Pfizer, Istanbul, Turkey) and xylazine (10 mg/kg, Rompun, Bayer, Istanbul, Turkey). Their midbacks were prepared in a sterile fashion. Using surgical microscope, a midline incision was performed over T5 and T12 vertebrae. A total laminectomy was performed over the segments of T7 and T10 with utmost care preventing SCI. The dura was left intact. Using an aneurysm clip (70 g closing force, Yasargil FE 721, Aesculap, Istanbul, Turkey) compression was applied to the spinal cord at the midpoint of the laminectomy site. The clip was removed after 1 minute of application and the wound is irrigated with normal saline before closure of the layers. Rats in the control group did not receive clip compression procedure. No further treatment was applied to the animals in the trauma group after clip compression. Animals in treatment groups received 5 mg/kg subcutaneous adalimumab and infliximab respectively right after wound closure.

After 72 hours, rats were killed with overdose pentobarbital. Trauma site being at the epicenter, 2 cm length spinal cord segments were removed en block. Tissue samples were immediately stored in a freezer for malondialdehyde (MDA) assay.

### Neurological evaluation

Preoperative and postoperative neurological assessments were performed every day until the end of the study by an observer who is blinded to the groups. The Tarlov scoring system was used<sup>35)</sup>. According to this system 5 tier scale was used to evaluate the neurological status: score 0, spastic paraplegia and no movement of the lower limbs; score 1, spastic paraplegia and slight movement of the lower limbs; score 2, good movement of the lower limbs, but inability to stand; score 3, able to stand, but unable to walk normally; score 4, complete recovery and normal gait/hopping.

### Determination of lipid peroxidation in traumatized spinal cord tissue

The level of lipid peroxides in traumatized spinal cord tissue were measured as thiobarbituric acid-reactive material and determined using the method of Mihara and Uchiyama<sup>34)</sup>. MDA has been identified as the product of lipid peroxidation that reacts with thiobarbituric acid to give a red species absorbing at 535 nm. The assay procedure for lipid peroxide in spinal cord tissue was set up as follows. Tissues were homogenized in 10 volumes (wt/vol) of cold 1.5% of KCl. One half a milliliter (0.5 mL) of homogenate was mixed with 3 mL of 1% of H<sub>3</sub>PO<sub>4</sub> (Carlo Erba Reagents, Cat No: 30406, Val de Reuil, France) and 1 mL of 0.6% of thiobarbituric acid (Merck & Co., Cat No: M.108180.0025, White House Station, NJ, USA). The mixture was then heated in boiling water for 60 minutes. After cooling, the color was extracted into 4 mL n-butanol (Sigma-Aldrich, Cat No: 24124, St. Louis, MO, USA), and the absorbance was recorded at 535 and 520 nm with a spectrophotometer (U-2900, Hitachi High Tech, Tokyo, Japan). Using tetramethoxypropane as the standard, tissue lipid peroxide

levels were calculated as nanomoles per gram of wet tissue<sup>32)</sup>.

### Statistical analyzes

Commercially available software package SPSS (SPSS Inc., Released 2007, SPSS for windows, version 16.0., Chicago, IL, USA) was used to compare MDA levels between groups. Results are expressed as mean values and probability level less than 0.05 was accepted as significant. Nonparametric tests of Kruskal-Wallis and Mann-Whitney U were used to determine differences between groups regarding mean MDA levels.

## RESULTS

### Lipid peroxidation


The mean MDA level of the control group was 23.87±8.96 (range: 12.00–35.00), trauma group was 79.12±24.19 (range: 47.00–123.00), infliximab group was 11.25±1.90 (range: 8.00–14.00) and adalimumab group was 22.75±8.10 (range: 11.00–31.00). MDA levels were statistically significantly different between groups,  $\chi^2(3)=23.063$ ,  $p<0.01$ . Pairwise comparisons were performed using Dunn's procedure with a Bonferroni correction for multiple comparisons. MDA levels were significantly different between the trauma and infliximab groups ( $p<0.01$ ) and trauma and adalimumab groups ( $p=0.022$ ). Boxplot in Fig. 1 demonstrate mean MDA levels of groups.

### Neurological evaluation


Line chart in Fig. 2 demonstrates mean Tarlov score changes throughout the experiment. There was no significant difference between trauma and treatment groups regarding Tarlov score.

## DISCUSSION

Traumatic injury to the spinal cord itself creates an injury site within traumatic tissue. This is the primary injury and apart from



**Fig. 1.** Box plot demonstrating mean MDA levels. MDA levels were significantly different between the trauma and infliximab groups (theta,  $p<0.01$ ) and trauma and adalimumab groups (asterisk,  $p=0.022$ ) (central line for each group indicates median, with the box showing upper and lower quartiles and the whiskers showing the range). MDA: malondialdehyde.



**Fig. 2.** Line graph demonstrating progress in mean Tarlov scores in groups.

preventive measures aimed to prevent accidents, there is virtually nothing that can be done in terms of treatment for now. On the other hand, secondary injury resulting from the changes in the trauma milieu is the main target of research<sup>33</sup>. Secondary injury results from cascades of ischemia, edema, excitatory amino acid influx and oxidative injury<sup>4,16</sup>.

Central nervous system is considered as a relatively “immune free” environment in normal circumstances. After an insult, this changes<sup>6,16</sup> and it was demonstrated that reducing or blocking inflammatory response may have beneficial effects<sup>6,10,21,31</sup>.

Oxidative stress occurring after trauma injures nucleic acids, proteins, carbohydrates and lipids<sup>17</sup>. Since biological membranes constitute of lipids, lipid peroxidation targets them<sup>17,22</sup>. The lipid rich environment of nervous tissue makes it susceptible to such injuries. MDA is a three carbon, low molecular weight aldehyde which is considered as an important lipid peroxidation marker. Levels of MDA increases in several diseases<sup>22,31</sup> and SCI<sup>20,24,32</sup>.

TNF- $\alpha$  which is a proinflammatory cytokine produced by monocytes, macrophages and T lymphocytes was proven to play important roles in various diseases such as rheumatoid arthritis, psoriatic arthritis, Crohn’s disease, psoriasis and ankylosing spondylitis<sup>13,29</sup>. Blockage of TNF- $\alpha$  causes immune system suppression<sup>18</sup> and this effect is used to treat aforementioned diseases<sup>13,14,18,29,36</sup>. Besides its roles in such diseases TNF- $\alpha$  has demonstrated roles in neural tissue damage and neuroinflammation<sup>5,23,38,39</sup>. Harrington et al.<sup>28</sup> have demonstrated an increase in TNF- $\alpha$  levels in cerebrospinal fluid in the first hour of SCI. Wang et al.<sup>39</sup> had similar findings in traumatized spinal cord

tissue. Authors of these studies concluded that TNF- $\alpha$  has important roles in pathophysiology of traumatic SCI.

Etanercept, infliximab, adalimumab, golimumab, and certolizumab are the anti-TNF- $\alpha$  agents that are authorized for treatment of diseases mediated by TNF- $\alpha$ <sup>15</sup>. Only infliximab<sup>32</sup> and etanercept<sup>3</sup> has been evaluated in SCI and their favorable effects have been demonstrated. This is the first study evaluating the effects of adalimumab on SCI. The study is built on the hypothesis that TNF- $\alpha$  blockage by adalimumab may have beneficial effects in SCI as demonstrated in previous studies of different agents targeting TNF- $\alpha$ . Results here have showed that adalimumab has favorable effects on lipid peroxidation in spinal cord injured rats. The decrease in MDA levels was significant when compared to controls; however, there were no significant difference between adalimumab and infliximab. This suggests, these two agents have similar effects on lipid peroxidation levels in experimental SCI. On the other hand we could not demonstrate any favorable effect in neurological status. This may be the result of 72 hours of evaluation. This study is primarily designed to demonstrate biochemical effects of the agents on SCI and longer periods may reveal differences.

## CONCLUSION

TNF- $\alpha$  blockers are already used for treatment of various diseases and their benefits and side effects are well studied. Although further experiments are required, it may be time to try these agents in clinical setting of SCI.

## References

- Balentine JD : Pathology of experimental spinal cord trauma. I. The necrotic lesion as a function of vascular injury. *Lab Invest* 39 : 236-253, 1978
- Balentine JD : Pathology of experimental spinal cord trauma. II. Ultrastructure of axons and myelin. *Lab Invest* 39 : 254-266, 1978
- Bayrakli F, Balaban H, Ozum U, Duger C, Topaktas S, Kars HZ : Etanercept treatment enhances clinical and neuroelectrophysiological recovery in partial spinal cord injury. *Eur Spine J* 21 : 2588-2593, 2012
- Beattie MS, Hermann GE, Rogers RC, Bresnahan JC : Cell death in models of spinal cord injury. *Prog Brain Res* 137 : 37-47, 2002
- Bethea JR, Nagashima H, Acosta MC, Briceno C, Gomez F, Marcillo AE, et al. : Systemically administered interleukin-10 reduces tumor necrosis factor-alpha production and significantly improves functional recovery following traumatic spinal cord injury in rats. *J Neurotrauma* 16 : 851-863, 1999
- Blight AR : Macrophages and inflammatory damage in spinal cord injury. *J Neurotrauma* 9 Suppl 1 : S83-S91, 1992
- Bracken MB : Methylprednisolone and acute spinal cord injury : an update of the randomized evidence. *Spine (Phila Pa 1976)* 26 (24 Suppl) : S47-S54, 2001
- Bracken MB : Methylprednisolone in the management of acute spinal cord injuries. *Med J Aust* 153 : 368, 1990
- Bracken MB : Steroids for acute spinal cord injury. *Cochrane Database Syst Rev* 1 : CD001046, 2012
- Bracken MB, Collins WF, Freeman DF, Shepard MJ, Wagner FW, Silten RM, et al. : Efficacy of methylprednisolone in acute spinal cord injury. *JAMA* 251 : 45-52, 1984

11. Bracken MB, Shepard MJ, Collins WF, Holford TR, Young W, Baskin DS, et al. : A randomized, controlled trial of methylprednisolone or naltrexone in the treatment of acute spinal-cord injury. Results of the Second National Acute Spinal Cord Injury Study. *N Engl J Med* 322 : 1405-1411, 1990
12. Bracken MB, Shepard MJ, Hellenbrand KG, Collins WF, Leo LS, Freeman DF, et al. : Methylprednisolone and neurological function 1 year after spinal cord injury. Results of the National Acute Spinal Cord Injury Study. *J Neurosurg* 63 : 704-713, 1985
13. Burmester GR, Mease P, Dijkmans BA, Gordon K, Lovell D, Panaccione R, et al. : Adalimumab safety and mortality rates from global clinical trials of six immune-mediated inflammatory diseases. *Ann Rheum Dis* 68 : 1863-1869, 2009
14. Burness CB, Keating GM : Adalimumab : a review of its use in the treatment of patients with ulcerative colitis. *BioDrugs* 27 : 247-262, 2013
15. Caminero A, Comabella M, Montalban X : Tumor necrosis factor alpha (TNF- $\alpha$ ), anti-TNF- $\alpha$  and demyelination revisited : an ongoing story. *J Neuroimmunol* 234 : 1-6, 2011
16. Carlson SL, Parrish ME, Springer JE, Doty K, Dossett L : Acute inflammatory response in spinal cord following impact injury. *Exp Neurol* 151 : 77-88, 1998
17. Catalá A : Lipid peroxidation of membrane phospholipids generates hydroxy-alkenals and oxidized phospholipids active in physiological and/or pathological conditions. *Chem Phys Lipids* 157 : 1-11, 2009
18. Diak P, Siegel J, La Grenade L, Choi L, Lemery S, McMahon A : Tumor necrosis factor alpha blockers and malignancy in children : forty-eight cases reported to the Food and Drug Administration. *Arthritis Rheum* 62 : 2517-2524, 2010
19. Ducker TB, Kindt GW, Kempf LG : Pathological findings in acute experimental spinal cord trauma. *J Neurosurg* 35 : 700-708, 1971
20. Emmez H, Börcek AÖ, Kaymaz M, Kaymaz F, Durdağ E, Civi S, et al. : Neuroprotective effects of gabapentin in experimental spinal cord injury. *World Neurosurg* 73 : 729-734, 2010
21. Fleming JC, Norenberg MD, Ramsay DA, Dekaban GA, Marcillo AE, Saenz AD, et al. : The cellular inflammatory response in human spinal cords after injury. *Brain* 129 (Pt 12) : 3249-3269, 2006
22. Grotto D, Maria LS, Valentini J, Paniz C, Schmitt G, Garcia SC, et al. : Importance of the lipid peroxidation biomarkers and methodological aspects for malondialdehyde quantification. *Quim Nova* 32 : 169-174, 2009
23. Guadagno J, Xu X, Karajigikar M, Brown A, Cregan SP : Microglia-derived TNF $\alpha$  induces apoptosis in neural precursor cells via transcriptional activation of the Bcl-2 family member Puma. *Cell Death Dis* 4 : e538, 2013
24. Guven C, Borcek AO, Cemil B, Kurt G, Yildirim Z, Ucanus NL, et al. : Neuroprotective effects of infliximab in experimental spinal cord ischemic injury. *J Clin Neurosci* 17 : 1563-1567, 2010
25. Hall ED : Inhibition of lipid peroxidation in CNS trauma. *J Neurotrauma* 8 Suppl 1 : S31-S40; discussion S41, 1991
26. Hall ED, Braughler JM : Free radicals in CNS injury. *Res Publ Assoc Res Nerv Ment Dis* 71 : 81-105, 1993
27. Hamada Y, Ikata T, Katoh S, Nakauchi K, Niwa M, Kawai Y, et al. : Involvement of an intercellular adhesion molecule 1-dependent pathway in the pathogenesis of secondary changes after spinal cord injury in rats. *J Neurochem* 66 : 1525-1531, 1996
28. Harrington JF, Messier AA, Levine A, Szymdynger-Chodobska J, Chodobski A : Shedding of tumor necrosis factor type 1 receptor after experimental spinal cord injury. *J Neurotrauma* 22 : 919-928, 2005
29. Kaymakcalan Z, Sakorafas P, Bose S, Scesney S, Xiong L, Hanzatian DK, et al. : Comparisons of affinities, avidities, and complement activation of adalimumab, infliximab, and etanercept in binding to soluble and membrane tumor necrosis factor. *Clin Immunol* 131 : 308-316, 2009
30. Klebanoff SJ, Vadas MA, Harlan JM, Sparks LH, Gamble JR, Agosti JM, et al. : Stimulation of neutrophils by tumor necrosis factor. *J Immunol* 136 : 4220-4225, 1986
31. Kurt G, Cemil B, Borcek AO, Borcek P, Akyurek N, Sepici A, et al. : Infliximab administration reduces neuronal apoptosis on the optic pathways in a rabbit hydrocephalus model : a preliminary report. *Br J Neurosurg* 24 : 275-279, 2010
32. Kurt G, Ergün E, Cemil B, Börcek AO, Börcek P, Gülbahar O, et al. : Neuroprotective effects of infliximab in experimental spinal cord injury. *Surg Neurol* 71 : 332-336; discussion 336, 2009
33. Kwon BK, Okon E, Hillyer J, Mann C, Baptiste D, Weaver LC, et al. : A systematic review of non-invasive pharmacologic neuroprotective treatments for acute spinal cord injury. *J Neurotrauma* 28 : 1545-1588, 2011
34. Mihara M, Uchiyama M : Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. *Anal Biochem* 86 : 271-278, 1978
35. Papakostas JC, Matsagas MI, Toumpoulis IK, Malamou-Mitsi VD, Pappa LS, Gkrepi C, et al. : Evolution of spinal cord injury in a porcine model of prolonged aortic occlusion. *J Surg Res* 133 : 159-166, 2006
36. Reinisch W, Sandborn WJ, Hommes DW, D'Haens G, Hanauer S, Schreiber S, et al. : Adalimumab for induction of clinical remission in moderately to severely active ulcerative colitis : results of a randomised controlled trial. *Gut* 60 : 780-787, 2011
37. Sekhon LH, Fehlings MG : Epidemiology, demographics, and pathophysiology of acute spinal cord injury. *Spine (Phila Pa 1976)* 26 (24 Suppl) : S2-S12, 2001
38. Tang X, Wang Y, Zhou S, Qian T, Gu X : Signaling pathways regulating dose-dependent dual effects of TNF- $\alpha$  on primary cultured Schwann cells. *Mol Cell Biochem* 378 : 237-246, 2013
39. Wang CX, Nuttin B, Heremans H, Dom R, Gybels J : Production of tumor necrosis factor in spinal cord following traumatic injury in rats. *J Neuroimmunol* 69 : 151-156, 1996
40. Yu SH, Cho DC, Kim KT, Nam KH, Cho HJ, Sung JK : The neuroprotective effect of treatment of valproic Acid in acute spinal cord injury. *J Korean Neurosurg Soc* 51 : 191-198, 2012