

RESEARCH ARTICLE

Open Access

# Bovine herpesvirus 4 glycoprotein B is indispensable for lytic replication and irreplaceable by VSVg

Valentina Franceschi, Antonio Capocéfalo, Sandro Cvirani and Gaetano Donofrio\*

## Abstract

**Background:** Bovine herpesvirus 4 (BoHV-4) is a gammaherpesvirus, belonging to Rhadinovirus genus, with no clear association with disease. However, there is increasing evidence of its secondary pathogenic role in cases of post-partum metritis in cattle. BoHV-4 Open Reading Frame 8 (ORF8) codifies for glycoprotein B (gB) that shows a heterodimeric structure, composed of two subunits and covalently linked by disulfide bonds and responsible for host cell adhesion through binding to heparan sulfates associated with cellular proteoglycans. Here we describe the generation of several tagged soluble forms of gB ectodomain, in order to test their ability to neutralize BoHV-4 infection.

**Results:** The results show, however, that none of these soluble forms are able to block viral infectivity. To better understand the role of gB during BoHV-4 lytic replication, a recombinant BoHV-4 was generated by homologous recombination from a BoHV-4 cloned as a Bacterial artificial chromosome (BAC) (pBAC-BoHV-4-A), in which most of the BoHV-4 gB ORF was substituted by the insertion of a DNA stuffer selectable cassette. The resulting recombinant BoHV-4 genome (pBAC-BoHV-4- $\Delta$ gB-KanaGalK) was completely unable to reconstitute infectious replicating viral particles (Infectious Replicating Viral Particles, IRVPs) and to replicate when transfected in permissive cell lines in comparison to its revertant clone (pBAC-BoHV-4- $\Delta$ gB-Rev) or pBAC-BoHV-4-A parental clone.

**Conclusion:** This demonstrates that the BoHV-4 replicating cycle is dependent on gB. Moreover, when gB was deleted from a recombinant BoHV-4 genome delivering an heterologous glycoprotein, Vesicular Stomatitis Virus Glycoprotein (VSVg), VSVg was unable to complement gB. This study provides direct evidence that gB is necessary for BoHV-4 lytic replication.

**Keywords:** Bovine herpesvirus 4, Glycoprotein B, Homologous recombination, Gene deletion, Vesicular stomatitis virus glycoprotein

## Background

The Herpesvirus envelope contains a variable number of glycoproteins involved in virus attachment, penetration, budding and spreading among infected cells. Some of these proteins are extremely conserved in functions and sequences, while others are typical of a peculiar virus genus or species.

The entry model mechanism of Human Herpes Virus 1 (HHV-1) is based firstly on the interaction between glycoproteins gC and gB and the host cell heparan

sulfates associated with glycosaminoglycans (GAGs) and secondly, by gD-mediated penetration that is also responsible for the fusion via gB, gH and gL [1]. Five glycoproteins are conserved in all herpesviruses, gB, gH, gL, gM and gN, and numerous studies have been carried out in order to characterize them [2].

The gB gene is one of the most conserved glycoproteins among the herpesvirus family. Homologues of gB have in fact been identified in every human and animal herpesvirus studied [3]. The hypothesis that gB function may be conserved in all herpesviruses is corroborated by its secondary structure, that shows the consistent conservation of cysteine and proline residues and the carboxy-terminal portion of the proteins [3].

\* Correspondence: gaetano.donofrio@unipr.it  
Dipartimento di Scienze Medico Veterinarie, Università di Parma, via del Taglio 10, 43126, Parma, Italy

Herpesvirus 4 (BoHV-4) belongs to the gammaherpesvirus, Rhadinovirus genus, and has been isolated worldwide both in healthy animals and in animals with a range of diseases varying from ocular discharge, conjunctivitis, dermatitis, respiratory diseases and abortion [4]. The pathogenic role of BoHV-4 is still unclear, even if there is increasing evidence of a secondary pathogenic role in bovine post-partum metritis [5,6].

Ten glycoproteins have been identified in BoHV-4 [7] and gB was demonstrated to be involved in cell host contact through heparan sulfate interactions [8]. It is not known if this interaction is sufficient to induce virus penetration and fusion or if a cellular receptor is specifically required, as for example occurs for Human Herpesvirus 5 (HHV-5) [9]. Structurally, gB is a heterodimeric protein, composed of two subunits linked by disulfide bonds. The protein is derived from a precursor that is first glycosylated, then trimmed and cleaved to acquire the mature gB form, thanks to a putative cleavage site which is present nearly in the middle of the sequence [3,10]. Similarly to Human Herpesvirus 4 (HHV-4), Murid Herpesvirus 4 (MuHV-4), Human Herpesvirus 8 (HHV-8) and HHV-5 [10], gB is one of the major components of the BoHV-4 virion. However the direct role of BoHV-4 gB for the progression of the virus into the lytic cycle has yet to be demonstrated. In the present study, the generation of several tagged soluble forms of BoHV-4 gB ectodomain was described, and their inability to block viral infectivity was also assessed. The cloning of different isolates of BoHV-4 genome as Bacterial artificial chromosomes (BAC) [11-13] allowed the generation of a gB-deleted mutant BoHV-4, which provided further evidence that BoHV-4 gB has an indispensable role in BoHV-4 lytic replication; moreover VSVg was unable to complement BoHV-4 gB deletion, underlying the primary role of gB during BoHV-4 life cycle.

## Methods

### DNA preparation and PCR

Virus infected cells were lysed overnight in Proteinase K Buffer containing 10 mM Tris-HCl, pH 7.5, 1 mM EDTA, 0.5% sodium dodecyl sulfate (SDS) and 100 µg of proteinase K per ml at 37°C. Nucleic acids were extracted by treatment with phenol-chloroform and precipitated with ethanol. Treatment with 100 µg/ml of RNase A (Sigma) was performed for 1 h, after which the DNA was extracted with phenol and precipitated with ethanol again. The samples were kept at -20°C. One microgram of DNA sample was amplified over 35 cycles, each cycle consisting of denaturation at 94°C for 1 min, primer annealing at 55°C for 1 min, and chain elongation with 1 U of Pfu DNA polymerase (Fermentas) at 72°C for 2 min. PCR amplification was carried out in a final volume of 50 µl of 10 mM Tris-hydrochloride pH 8.3 containing 0.2

mM deoxynucleotide triphosphates, 3 mM MgCl<sub>2</sub>, 50 mM KCl, 5% DMSO and 0.25 µM of each primer. The primers used for the amplification of BoHV-4 ORF8, coding for gB protein, are Sgfl-gB-sense and MluI-gB-antisense (see Table 1). The so generated 2185 bp amplicon was then checked in 1% agarose gel and visualized after ethidium bromide staining in 1× TAE buffer (40 mM Tris-acetate, 1 mM EDTA) and used to subclone gB in an expression cassette. The specificity of the PCR product was determined by sequencing.

### Plasmids

To generate pgB<sub>Fc</sub>, BoHV-4 gB ORF (from 1–2175 nt) was amplified by PCR using Sgfl-gB-sense and MluI-gB-antisense primer (see Table 1), cut with Sgfl/MluI and inserted in pCMV6-AC-Fc (Origene) cut with the same enzymes. gB/Fc ORF was cut from pgB/Fc firstly with BamHI, blunt-ended, consequently cut with PmeI and then sub-cloned in pWPI (Addgene; <http://www.addgene.org/12254/>) cut with PmeI. Thus pWPI/gB<sub>Fc</sub> was obtained.

The chimeric ORF (1510 bp) coding for gB<sub>truncated</sub>-gD<sub>106</sub> was obtained by gene synthesis (Eurofins MWG Operon) where an NheI and a BamHI restriction sites were included to the 5' and 3' end respectively. This gB<sub>truncated</sub>-gD<sub>106</sub> ORF was excised from the vector (pBluescript) by NheI/BamHI digestion and inserted in pEGFP-C1 (Clontech) depleted of Green Fluorescent Protein (GFP) after cutting with the same enzymes and pCMV-gB<sub>truncated</sub>-gD<sub>106</sub> was obtained. Murine Fc was then amplified by PCR from pCMV6-AC-Fc vector with Sall-Fc-sense and BamHI-Fc-antisense primers (see Table 1) and used to substitute the gD<sub>106</sub> tag in pCMV-gB<sub>truncated</sub>-gD<sub>106</sub>.

pCMV-gB<sub>truncated</sub>-gD<sub>106</sub> was then cut with NdeI and Sall and the gB<sub>truncated</sub> was exchanged with the full length 2614 bp gB ectodomain, excised from pgB/Fc, cut with NdeI and XhoI. Sall and XhoI are compatible enzymes and the generated ends are in frame. The so generated pCMVgBgD<sub>106</sub> was then cut with BamHI, blunt-ended and the fusion peptide gBgD<sub>106</sub> was inserted in pWPI opened with PmeI and pWPI/gBgD<sub>106</sub> was generated.

pBluescript/gB<sub>truncated</sub>-gD<sub>106</sub> was cut with NheI and BamHI, the fragment was blunt-ended and inserted in PmeI cut pWPI to generate pWPI/gB<sub>truncated</sub>-gD<sub>106</sub>.

pCMV-gB<sub>truncated</sub>-gD<sub>106</sub> was cut with Sall/BamHI to excise gD<sub>106</sub> that was replaced with the 752bp Sall/BamHI cut murine Fc amplicon for generating pCMV-gB<sub>truncated</sub>-Fc.

The gB<sub>truncated</sub>-Fc was at the end excised from pCMV-gB<sub>truncated</sub>-Fc, cut with NheI/BamHI, blunt-ended and inserted in PmeI cut pWPI to obtain pWPI/gB<sub>truncated</sub>-Fc.

**Table 1 List of primers used in this work**

Primer name	Primer sequence 5'-3'
Sgfl-gB-sense	5'-CCC CCC GCG ATC GCA TGT ATT ATA AGA CTA TCT TAT TCT TCG CT-3'
Mlul-gB-antisense	5'-CCC CCC ACG CGT AAG GTC TGC CAT CAT TTC AGA GAG ATC TTT-3'
Sall-Fc-sense	5'-CCC GTC GAC CGT ACG CGG CCG CTC GAG ATG CCC AGA-3'
BamHI-Fc-antisense	5'-CCC GGA TCC GCG GCC GGC CGT TTA AAC TCA TTT ACC CGG-3'
gB-left-sense	5'-CCG GAA TTC AGA ACC CAC ACT ATA GGG GAA ACG ACC TCA-3'
gB-left-antisense	5'-TCT CCA GAG GAA GAT GAT GCA CTG CAT ACT CTG-3'
gB-right-sense	5'-CCCC CTG CAG TCA CAT CCT AGA GGA ATT AAC-3'
gB-right-antisense	5'-AGA CTT GCA AGC TTC TGT GTG TAG TAA TTA-3'
Smal-Asel-Kana-sense	5'-AAC CCC CGG GAT TAA TCC GGA ATT GCC AGC TGG GG-3'
Smal-Mlul-Kana anti	5'-CCA ACC CGG GAC GCG TGA AAT TGT AAG CGT TAA TAA T-3'

pTZ-KanaGalK, was generated by sub-cloning the 2232 bp galactokinase prokaryotic expression cassette (GalK), along with the kanamycin resistant expression cassette (Kana), into the pTZ57R shuttle vector, cut with KpnI/PstI [14]. The targeting vector, pgB-KanaGalK-gB, was generated firstly by the insertion of the 1167bp left gB homology region amplicon (gB-left-sense and antisense, see Table 1) cut with EcoRI/KpnI, in pTZ-KanaGalK, cut with the same enzymes; in this intermediate construct, cut with PstI/HindIII, was consequently sub-cloned the 600bp right gB homology region amplicon (gB-right-sense and antisense; see Table 1), cut with the same enzymes.

#### Cells, cell culture electroporation and recombinant virus reconstitution

Bovine Embryo Kidney [(BEK) from M. Ferrari, Istituto Zooprofilattico Sperimentale, Brescia, Italy], MDBK (Madin Darby Bovine Kidney cells), BEK/cre [12], HEK (Human Embryo Kidney) and BEC (Bovine endometrial stromal cells) cell lines were cultured in Dulbecco's modified Eagle Minimal Essential Medium (DMEM) (Lonza) containing 10% fetal bovine serum (FBS), 2 mM l-glutamine, 100 IU/ml penicillin (SIGMA) and 100 µg/ml streptomycin (SIGMA).

Cells were sub cultured to a fresh culture vessel when growth reached 70 to 90% confluence (i.e., every 3 to 5 days) and were incubated at 37°C in a humidified atmosphere of 95% air–5% CO<sub>2</sub>.

Plasmid DNAs (5 µg) in 500 µl DMEM without serum were electroporated (Equibio apparatus, 270 V, 960 µF, 4-mm gap cuvettes) into BEK or BEK/cre cells from a confluent 25-cm<sup>2</sup> flask. Electroporated cells were then returned to the flask, fed the next day, and split 1:2 when they reached confluence at 2 days post electroporation. Cells were left to grow until Cytopathic Effect (CPE) appeared. Recombinant viruses were propagated by infecting confluent monolayers of MDBK cells at a multiplicity of infection (m.o.i.) of 0.5 50% tissue culture

infectious doses (TCID<sub>50</sub>) per cell and maintaining them in MEM with 10% FBS for 2 h. The medium was removed and replaced by fresh MEM containing 10% FBS. When approximately 90% of the cell monolayer exhibited CPE (~72 h post infection), the virus was prepared by freezing and thawing cells three times and pelleting virions through 30% sucrose, as described previously [12]. Virus pellets were resuspended in cold MEM without FBS. TCID<sub>50</sub> were determined on MDBK or BEK cells by limiting dilution.

#### Viruses

BoHV-4-A, BoHV-4-A-ΔTK-EGFP, BAC-BoHV-4-A, BAC-BoHV-4-AΔL1.7VSVG, BAC-BoHV-4-AΔL1.7VSVGΔgB-KanaGalK, and BAC-BoHV-4-A-ΔgB-KanaGalK were propagated by infecting confluent monolayers of MDBK or BEK cells at a m.o.i. of 0.5 and maintained in minimal essential medium (MEM) (SIGMA) with 2% FBS for 2 h. The medium was then removed and replaced by fresh MEM containing 10% FBS. When approximately 90% of the cell monolayer exhibited CPE (approximately 72 h post infection), the virus was prepared by freezing and thawing cells three times and pelleting the virions through 30% sucrose, as previously described [12]. Virus pellets were resuspended in cold MEM without FBS. TCID<sub>50</sub> were determined in MDBK cells by limiting dilution.

#### *In vitro* neutralization assay (NTA)

HEK cells were seeded in a 175 cm<sup>2</sup> flask and when reached the 85% of confluence, were transiently transfected with the plasmids carrying the gB tagged constructs.

In particular, 14 ml of DMEM without serum were incubated with 43,75 µl of LTX Lipofectamine (Invitrogen) and 17,5 µg of pWPI/gB<sub>FC</sub>, pWPI/gB<sub>truncatedFC</sub>, or pWPI/gB<sub>truncatedGD106</sub> at room temperature for at least 15 minutes. This transfection solution was then added carefully to the HEK cell monolayer and left to incubate 6 hours at 37°C with 5% CO<sub>2</sub>. The transfection solution was then removed and replaced by 21 ml of 1:1 DMEM/

F12 medium without serum, after a washing, to remove any traces of the transfection solution. After 96 hours the supernatant was collected, clarified at 3000 r.p.m. for 5 minutes and stored at  $-20^{\circ}\text{C}$ .

The expression of gB was assessed by Western Blotting (the amount of gB was estimated to be between 5 and 50  $\mu\text{g}/\text{ml}$ , depending on the preparation) and consequently the supernatant was used for the neutralization assay.

$10^5$  HEK cells/well were seeded in a 12 multiwells plate and left to attach to the well.

After few hours the cells were pre-incubated 30 minutes at  $37^{\circ}\text{C}$  with 1ml/well of the supernatant recovered from HEK transfected with pWPI/gB<sub>Fc</sub> or pWPI/gB<sub>truncated</sub>gD<sub>106</sub> or pWPI/gB<sub>truncated</sub>Fc with medium without serum. The supernatant was then removed and substituted with ten-fold BoHV-4-A- $\Delta\text{TK}$ -EGFP virus dilution, starting from 1 to  $10^{-3}$  m.o.i. The cells were then incubated overnight at  $37^{\circ}\text{C}$  in 5% humidified atmosphere.

This assay was also repeated in BESC cells, with the same protocol, but seeding  $3 \times 10^3$  cells/well in a 96 multiwell plate; the cells were then treated with the gB supernatant and infected with two-fold dilutions of the virus, starting from 10 m.o.i.

These two protocols were also tested using a contact time varying from 30 to 60 minutes and/or an incubation contact temperature of  $4^{\circ}\text{C}$  or  $37^{\circ}\text{C}$ .

The virus spreading was then monitored during the following days by fluorescence and contrast microscope analysis.

### Recombineering and selection

Recombineering was performed as previously described [15] with some modifications. Five hundred  $\mu\text{l}$  of a  $32^{\circ}\text{C}$  overnight culture of SW102 containing BAC-BoHV-4-A, and BAC-BoHV-4- $\Delta\text{L1.7}$ -VSVG were diluted in 25 ml Luria-Bertani (LB) medium with or without chloramphenicol (SIGMA) selection (12.5  $\mu\text{g}/\text{ml}$ ) in a 50 ml baffled conical flask and grown at  $32^{\circ}\text{C}$  in a shaking water bath to an  $\text{OD}_{600}$  of 0.6. Then, 12 ml were transferred to another baffled 50 ml conical flask and heat-shocked at  $42^{\circ}\text{C}$  for exactly 15 minutes in a shaking water bath. The remaining culture was left at  $32^{\circ}\text{C}$  as the un-induced control. After 15 minutes the two samples, induced and un-induced, were briefly cooled in ice/water bath slurry and then transferred to two 15 ml Falcon tubes and pelleted using 5000 r.p.m. (Eppendorf centrifuge) at  $0^{\circ}\text{C}$  for 5 min. The supernatant was poured off and the pellet was resuspended in 1 ml ice-cold ddH<sub>2</sub>O by gently swirling the tubes in ice/water bath slurry. Subsequently, 9 ml ice-cold ddH<sub>2</sub>O was added and the samples pelleted again. This step was repeated once more, the supernatant was removed and

the pellet (50  $\mu\text{l}$  each) was kept on ice until electroporated with gel-purified  $\sim 3.8$  kb fragment (gB-KanaGalK-gB) obtained by cutting pgB-KanaGalK-gB with EcoRI/HindIII (Fermentas). An aliquot of 25  $\mu\text{l}$  was used for each electroporation in a 0.1 cm cuvette at 25  $\mu\text{F}$ , 2.5 kV and 201  $\Omega$ . After electroporation, the bacteria were recovered in 1 ml LB (15 ml Falcon tube) for 1 hour in a  $32^{\circ}\text{C}$  shaking water bath. For the counter selection step (see below), the bacteria were recovered in 10 ml LB in a 50 ml baffled conical flask and incubated for 4.5 hours in a  $32^{\circ}\text{C}$  shaking water bath.

After the recovery period, bacteria were washed twice in sterile 1x M9 salts (6 g/l Na<sub>2</sub>HPO<sub>4</sub>, 3 g/l KH<sub>2</sub>PO<sub>4</sub>, 1 g/l NH<sub>4</sub>Cl, 0.5 g/l NaCl) (SIGMA) as follows: 1 ml culture was pelleted in an eppendorf tube at 13,200 r.p.m. for 15 seconds and the supernatant was removed with a pipette. The pellet was resuspended in 1 ml of 1x M9 salts, and pelleted again. This washing step was repeated once more. After the second wash, the supernatant was removed and the pellet was resuspended in 1 ml of 1x M9 salts before plating serial dilutions (100  $\mu\text{l}$  each of 1:10, 1:100 and 1:1000 dilutions) on M63 minimal medium plates [15 g/l agar (DIFCO, BD Biosciences), 0.2% D-galactose (SIGMA), 1 mg/l D-biotin (SIGMA), 45 mg/l L-leucine (SIGMA) and 50 mg/l kanamycin (SIGMA)]. Washing in M9 salts is necessary to remove any rich media from the bacteria prior to selection on minimal medium plates. Plates were incubated 3–5 days at  $32^{\circ}\text{C}$ . Several selected colonies were picked, streaked on McConkey agar indicator plates (DIFCO, BD Biosciences) containing 50  $\mu\text{g}/\text{ml}$  of kanamycin and incubated at  $32^{\circ}\text{C}$  for 3 days until red colonies appeared. Red colonies were grown overnight in 5 ml of LB containing 50  $\mu\text{g}/\text{ml}$  of kanamycin and BAC-BoHV-4- $\Delta\text{gB}$ -KanaGalK were purified and analysed through HindIII restriction enzyme digestion for gB-KanaGalK-gB fragment targeted integration into the BoHV-4-A gB locus.

SW102 bacteria containing BAC-BoHV-4- $\Delta\text{gB}$ -KanaGalK genome were also grown, heat induced as described above and electroporated with a  $\sim 2.9$ kb gel purified fragment (gB<sub>locus</sub>) obtained by amplifying the gB locus with the primers gB-left-sense and gB-right-antisense. For the counter selection step, the bacteria were recovered in 10 ml LB in a 50 ml baffled conical flask and incubated for 4.5 hours in a  $32^{\circ}\text{C}$  shaking water bath. Bacterial serial dilutions were plated on M63 minimal medium plates containing 15 g/l agar, 0.2% glycerol (SIGMA), 1 mg/l D-biotin, 45 mg/l L-leucine, 0.2% 2-deoxy-galactose (DOG, SIGMA) and 12.5  $\mu\text{g}/\text{ml}$  chloramphenicol. Plates were incubated for 3–5 days at  $32^{\circ}\text{C}$ . Several selected colonies were picked up, streaked on McConkey agar indicator plates (DIFCO, BD Biosciences) containing 12.5  $\mu\text{g}/\text{ml}$  of chloramphenicol and incubated at  $32^{\circ}\text{C}$  for 3 days until white colonies appeared. White colonies were grown



in duplicate for 5–8 hours in 1 ml of LB containing 50 µg/ml of kanamycin or LB containing 12.5 µg/ml of chloramphenicol. Only those colonies growing on chloramphenicol and not on kanamycin were kept and grown overnight in 5 ml of LB containing 12.5 µg/ml of chloramphenicol. BAC-BoHV-4-gB<sub>revertant</sub> was purified and analysed through HindIII restriction enzyme digestion for gB locus fragment targeted integration. Original detailed protocols for recombineering can also be found at the recombineering website [16].

### Restriction enzyme analysis and non isotopic Southern hybridization

Fifteen µl of DNA prepared from bacteria containing pBAC-BoHV-4-A and derivatives were restriction enzyme-digested with HindIII, separated by electrophoresis overnight in a 0.8% agarose gel, stained with ethidium bromide, capillary transferred to a positively charged nylon membrane (Roche), and cross-linked by UV irradiation by standard procedures. The membrane was pre-hybridized in 50 ml of hybridization solution (7% SDS, 0.5 M phosphate, pH 7.2, 1 mM EDTA) for 2 hours at 65°C in a rotating hybridization oven (Techna instruments). Probe preparation and digoxigenin non-isotopic labelling was performed by PCR.

Southern Blotting probe was designed spanning Kana region using the primer pair SmaI-Asel-Kana sense and SmaI-MluI-Kana anti (see Table 1).

PCR amplification was carried out in a final volume of 50 µl of 10 mM Tris-HCl, pH 8.3, containing 0.2 mM deoxynucleotide triphosphates, 0.02 mM alkaline labile digoxigenin-dUTP (Roche), 3 mM MgCl<sub>2</sub>, 50 mM KCl, and 0.25 µM of each primer over 35 cycles, each cycle consisting of denaturation at 94°C for 1 minute, primer annealing at 55°C for 1 minute, and chain elongation with 1 U of Taq polymerase (Boehringer Diagnostics) at 72°C for 1 minute. A parallel reaction omitting digoxigenin-dUTP was performed, because digoxigenin incorporation into the amplicon can be checked through the size shift of the amplicon by gel electrophoresis. Five µl of the probe were added to 100 µl of ddH<sub>2</sub>O into a screw-cap tube, denatured in boiling water for 5 minutes, and cooled down on ice for another 2 minutes. Denatured probe was added to 50 ml of pre-heated 65°C hybridization solution to the pre-hybridized membrane and hybridized overnight at 65°C in a rotating hybridization oven (Techna Instruments). Following hybridization, the membrane was washed twice for 30 minutes with 100 ml of washing solution I (0.5× SSC [1× SSC is 0.15 M NaCl plus 0.015 M sodium citrate] and 0.1% SDS) and twice for 30 minutes with 100 ml of washing solution II (40 mM phosphate, pH 7.2, 0.05% SDS) at 65°C. On a freshly washed dish, the membrane was incubated for 30 min at room temperature in 25 ml of blocking solution (100 mM maleic acid, pH

7.5, 150 mM NaCl, 1% blocking reagent [Roche] or 5% skim milk). Anti-digoxigenin Fab fragment (150 U/200 µl [Roche]), diluted 1:15,000 in 25 ml of blocking solution, was applied to the membrane for 30 minutes under gentle shaking at room temperature and washed twice for 15 minutes with 100 ml of washing solution (100 mM maleic acid, pH 7.5, 150 mM NaCl, 0.3% Tween 20). Detection was performed following equilibration of the membrane in detection buffer (100 mM Tris-HCl, pH 9.5, 1 mM EDTA) for 2 minutes at room temperature. Chemiluminescent substrate (CSPD, Roche) was added by scattering the drops over the surface of the membrane after placement of the membrane between two plastic sheets, and any bubbles present under the sheet were eliminated with a damp lab tissue to create a liquid seal around the membrane. Signal detection was obtained, exposing the membrane to X-ray film. The exposure time was adjusted with the intensity of the signal.

### Western immunoblotting

Cell lysates containing 50 µg of total protein were electrophoresed through 8 to 12% SDS-polyacrylamide gels and transferred to nylon membranes by electro blotting. Membranes were incubated with monoclonal anti-BoHV-1-gD antibody (clone 1B8F11; VRMD, Inc., Pullman, WA), probed with horseradish peroxidase-labelled anti-mouse immunoglobulin G (IgG) antibody (SIGMA), and visualized by enhanced chemiluminescence (Millipore). For detecting the Fc was used an anti-mouse IgG whole molecule (A9044, Sigma, Saint Louis, Missouri).


## Results and discussion

### Soluble gB fragments did not inhibit virus attachment and replication

Recent studies have demonstrated that HHV-4 infection can be prevented by treating cells with a library of peptides homologous to the ectodomain of the HHV-4 gB. [9]. To verify if BoHV-4 gB could act as a soluble ligand for a putative BoHV-4 host cell receptor and block BoHV-4 infection, BoHV-4 gB ectodomain was sub-cloned and expressed as a secreted form into the supernatant of transiently transfected cells. BoHV-4 gB has a length of 873 aa and corresponds to an ORF of 2622 bp (Figure 1A and B). Several studies have demonstrated that the mature form of gB is heterodimeric and is generated by the presence of an internal putative protease site (*RQKRS*) in the second half of its aminoacidic sequence that is responsible for the cleavage of the two subunits, that are then covalently linked by disulfide bonds (Figure 1B) [3,10].

The secreted forms of gB were obtained on the basis of its transmembrane domain and cytoplasmic tail exclusion and/or substitution. Firstly, the full gB ectodomain (720 aa) was fused to a murine Fc tag encoding the





**Figure 2 BoHV-4 gB chimeric peptide expression.** **A)** Western Immunoblotting of pWPI/gBFc transfected HEK293 cell lysate or supernatant, in presence or absence of a reducing agent, β-mercaptoethanol. The cell extract of pEGFP-C1 transfected HEK 293 was used as a negative control. As a positive control, the supernatant of cells transfected with a plasmid carrying only the murine Fc was used. **B)** Western Immunoblotting of pWPI/gBFc and pWPI/gB<sub>truncated</sub>Fc transfected HEK293 cell lysates with the presence of β-mercaptoethanol. **C)** Western Immunoblotting of pWPI/gBgD<sub>106</sub> and pWPI/gB<sub>truncated</sub>gD<sub>106</sub> transfected HEK293 cell lysates and supernatant. The control used was the same of the Figure 2A.

gB<sub>truncated</sub>gD<sub>106</sub> respectively, were generated replacing the Fc tag with the gD<sub>106</sub> soluble epitope tag of BoHV-1 in order to verify the influence of the tag peptide fused with gB in terms of secretion (Figure 2B and Additional file 3: Figure S3 and Additional file 4: Figure S4).

Surprisingly, only gB<sub>truncated</sub>gD<sub>106</sub> was efficiently secreted in transiently transfected HEK cells, as shown in Figure 2B and C. Although gBgD<sub>106</sub> was well expressed, its secretion levels were very low. It was outside the scope of the present work to determine the reasons of the inability of gBgD<sub>106</sub> to be efficiently secreted and gBgD<sub>106</sub> was eliminated from the study.


The supernatants containing gBFc, gB<sub>truncated</sub>Fc and gB<sub>truncated</sub>D<sub>106</sub> were used in an *in vitro* neutralization assay (NTA) to verify if one of these secreted forms could successfully prevent BoHV-4 attachment and penetration. The NTA was performed on different cell lines (HEK, BEK, MDBK and BESC), employing a recombinant BoHV-4 expressing GFP (BoHV-4EGFPΔTK) [18], at different temperatures (37 and 4°C), time of contact (30 and 60 minutes), number of seeded cells and amount of secreted peptides. BoHV-4 attachment and penetration was monitored at different times (24 and 48 hours) by GFP expression under fluorescence microscopy. No blocking activity was achieved with any of the experimental conditions tested (Figure 3 and data not shown). The secreted gB forms tested were apparently unable to prevent BoHV-4 attachment and penetration and this was in contrast with that reported for some Herpesvirus [19], but in agreement with others [20].

The inability of gBFc, gB<sub>truncated</sub>Fc and gB<sub>truncated</sub>gD<sub>106</sub> to prevent BoHV-4 attachment and penetration of the host cell gave rise to several hypothesis: *a)* the folding of the secreted peptide could be improper; *b)* the amount of the secreted peptide could be insufficient; *c)* viral glycoproteins different from gB could be responsible for mediating virus-cell interaction and internalization, subordinating gB to a secondary role.

**BoHV-4-ΔgB was unable to replicate in host cells**

Therefore, it was of interest to investigate the above hypotheses through the disruption of the gB gene in BoHV-4 genome. The BoHV-4 gB ORF was disrupted by site





**Figure 3 Secreted gB neutralization assay.** Representative fluorescence and phase contrast images (10X) of MDBK cells untreated **A)** or treated **B)** for 1 hour at 37°C with BoHV-4 gB enriched supernatant and infected with one M.O.I. of BoHV-4-EGFP $\Delta$ TK. Infection was monitored under a fluorescence microscopy at 24 and 48 hours after the infection. This experiment, along with others made under different condition, as indicated into the text, was repeated three times with identical results.

specific insertional mutagenesis mediated by heat inducible homologous recombination in a strain of BoHV-4 genome cloned as a Bacterial artificial chromosome (BAC), pBAC-BoHV-4-A [12].

A targeting cassette (gB-KanaGalK-gB), excised from pgB-KanaGalK-gB vector, containing the 2232 bp Kana/GalK DNA stuffer double selecting cassette [21], flanked by two BoHV-4 gB gene homology regions, was electroporated in SW102 *E.coli* containing pBAC-BoHV-4-A genome and pBAC-BoHV-4-A $\Delta$ gB-KanaGalK was generated (Figure 4A). In the pBAC-BoHV-4-A $\Delta$ gB-KanaGalK genome, most of the gB ORF has been replaced by Kana/GalK DNA stuffer double selection cassette (Figure 4A). All recombinant selected clones were shown to be authentic when analysed by PCR and sequencing (data not shown), HindIII restriction enzyme digestion and Southern hybridization (Figure 4B). In order to obtain the revertant genome, pBAC-BoHV-4-A-gB<sub>revertant</sub> a large DNA fragment obtained by PCR from the BoHV-4-A genome, was electroporated in SW102 *E.coli* containing the pBAC-BoHV-4-A $\Delta$ gB-KanaGalK genome. Following a double negative selection [12], several revertant clones were obtained as showed by PCR, sequencing (data not shown), HindIII restriction enzyme digestion and Southern hybridization (Additional file 5: Figure S5A and B). The selected clones' stability was assessed by serially passaging over 25 days and analysis by HindIII restriction enzyme digestion (Figure 4C and Additional file 5: Figure S5C). To reconstitute infectious viruses, pBAC-BoHV-4-A $\Delta$ gB-KanaGalK and pBAC-BoHV-4-A-gB<sub>revertant</sub> were electroporated into BEK or BEK/cre cells to excise the BAC cassette. Surprisingly, the viable virus was obtained only

from pBAC-BoHV-4-A-gB<sub>revertant</sub> but not from pBAC-BoHV-4-A $\Delta$ gB-KanaGalK, in both BEK and BEK/cre cells (Figure 5A). The deletion of gB rendered BoHV-4 unable to be reconstituted and productively replicated. BoHV-4-A-gB<sub>revertant</sub> growth characteristics were identical to the parental BoHV-4-A (Figure 5B).


#### VSVg could not complement BoHV-4 gB

Various attempts have been reported regarding gB complementation in several gammaherpesviruses using heterologous gB [20,22]. It was thus decided here to try to complement gB with VSVg, which has been previously shown to complement HSV-1 gB [20] and which is also widely used to pseudo-type lentiviral vectors [23].

BoHV-4 gB ORF was deleted as was done for pBAC-BoHV-4-A $\Delta$ gB-KanaGalK, in BoHV-4-A $\Delta$ L1.7-VSVG (Additional file 6: Figure S6A, B and C). BoHV-4-A $\Delta$ L1.7-VSVG is a replicating competent attenuated BoHV-4 expressing VSVg, where the VSVg expression cassette is inserted into BoHV-4 L1.7 gene locus (Capocefalo et al.; *submitted paper*). When pBAC-BoHV-4-A $\Delta$ L1.7-VSVG $\Delta$ gB-KanaGalK was electroporated into BEK or BEK/cre it was unable to reconstitute replicating viral particles (data not shown), as was observed for the deletion of gB from BoHV-4-A. This is a further demonstration that BoHV-4 gB is essential and cannot be complemented by a heterologous protein such as VSVg.

Virus entry into host cells normally requires one glycoprotein involved in cell binding and membrane fusion, whereas herpesviruses use several viral glycoproteins to enter cells. Herpesvirus entry relies on a complex multiprotein mechanism that is not yet well






**Figure 4 BoHV-4 gB disruption.** **A)** Overall strategy to delete a 1261 bp sequence from the ORF8 coding for gB, via heat inducible homologous recombination. The 2232 bp Kana-Galk selectable DNA stuffer, flanked by ORF8 homologous regions, was introduced between positions 11640 and 12901 of the BoHV-4-A strain cloned as a BAC. The expected ORF8 locus (A, bottom) has an increased size of the HindIII fragment (5280 instead of 4314 bp), generated by HindIII restriction enzyme digestion. The diagrams here presented are not to scale. **B)** HindIII Restriction profile and corresponding Southern Blotting of six representative targeted clones, compared to the untargeted control. Southern Blotting was performed with a probe spanning Kana sequence and confirmed the above data. **C)** Clonal stability of the pBAC-BoHV-4-A-ΔgBKanaGalk in *Escherichia coli* SW102 cells, passaged for 25 consecutive days and analyzed by HindIII digestion and agarose gel electrophoresis.

understood. Even if the role of each single protein potentially involved is widely studied, their action as a complex, determining virus adhesion and cell entry has not been fully clarified.

Gammaherpesviruses can infect different cell types in a cell-dependent manner, as is the case for HHV-4, which is able to infect B cells through the action of the gp350, gp42, gB, gH and gL proteins, whereas it can infect epithelial cells only with gB, gH and gL [24]. In this complicated panorama, the role of gB is usually conserved with respect to virus adhesion and fusion to the host cells. Indeed, a gB homologue has been identified in all Gammaherpesviruses [25].

In MuHV-4, gB and gH are required for cell fusion and the N-terminal half of gB is thought to be responsible for the fusion, containing a putative fusion loop. A change in pH drives the fusion and gB changes its state from a pre-fusion state to a post-fusion one. Specific mAbs directed against gB block membrane fusion, maintain gB in its post-fusion form, block the release of virions from late endosomes [26], thus neutralizing infection. Of the ten glycoproteins that have been described in BoHV-4 [7], gB is the most abundant and one of the major components of the virion. Similarly to the majority of other gammaherpesviruses, like Equine Herpesvirus 2 (EHV-2) and 5 (EHV-5) and MuHV-4,



**Figure 5 A) Viral reconstitution.** Representative fluorescence and phase contrast images (10X) of BoHV-4-A-ΔgB<sub>revertant</sub> virus reconstitution following pBAC-BoHV-4-A-ΔgB<sub>revertant</sub> electroporation in BEK or BEK/cre cells. pBAC-BoHV-4-A-ΔgB<sub>KanaGalK</sub> was unable to reconstitute IRVPs after electroporation in BEK or BEK/cre cells. **B)** Replication kinetics of BoHV-4-A-ΔgB<sub>revertant</sub> compared with BoHV-4-A and BoHV-4-A-ΔgB<sub>KanaGalK</sub>. The data presented are the means ± standard errors of triplicate measurements.

BoHV-4 gB is present in the virion in its cleaved form [10,27]. The BoHV-4 ORF8 coding for gB is 2622 bp in length and has a strict homology with HHV-4 gB, indicating that these sequence similarities could correspond to similar secondary and tertiary structures [3]. BoHV-4 gB transcript is unspliced and shows a typical TATA box sequence positioned 80 nucleotides upstream of the first ATG and a putative polyadenylation signal can be identified in 3' respect to the stop codon. BoHV-4 gB is a heterodimeric glycoprotein, called gp6-gp10-gp17, with an apparent molecular size of 150 kDa-120 kDa-51 kDa respectively. gp10-gp17 are linked by disulfide bonds and gp6 by non-covalent bonds and all these subunits derive from the proteolytical cleavage of the gB gene product [7]. BoHV-4 gB interaction with heparan-like molecules present on host cell GAGs was previously shown [8], demonstrating BoHV-4 gB contribution to virus entry and fusion, but its direct necessity has not yet been demonstrated.

## Conclusion

The present study may be considered the direct demonstration that gB is essential for BoHV-4 replication and its deletion is not compatible with virus survival. Moreover, despite the structural similarity between BoHV-4 gB and VSVg proteins, gB substitution with VSVg is not effective, emphasising the unique and specific role of gB in BoHV-4 life cycle.

## Additional files

**Additional file 1: Figure S1. A)** Chimeric peptide gB/Fc sequence and predicted amino-acid product. In black is highlighted the gB sequence, in grey the putative protease site, RQKRS, and in sky blue the Fc sequence. **B)** Representative contrast phase and fluorescence images of pWPV/gBFC transfected HEK cells at 24 hours after transfection.

**Additional file 2: Figure S2. A)** Chimeric peptide gB<sub>truncated</sub><sup>Fc</sup> sequence and predicted amino-acid product. In black is highlighted the

gB truncated sequence and in sky blue the Fc sequence.

**B)** Representative contrast phase and fluorescence images of pWPI/gB<sub>truncated</sub>Fc transfected HEK cells at 24 hours after transfection.

**Additional file 3: Figure S3. A)** Chimeric peptide gB/gD<sub>106</sub> sequence and predicted amino-acid product. In black is highlighted the gB sequence, in grey the putative protease site, *RQKRS*, in yellow the gD<sub>106</sub> sequence. **B)** Representative contrast phase and fluorescence images of pWPI/gB<sub>truncated</sub>gD<sub>106</sub> transfected HEK cells at 24 hours after transfection.

**Additional file 4: Figure S4. A)** Chimeric peptide gB<sub>truncated</sub>-gD<sub>106</sub> sequence and predicted amino-acid product. In black is highlighted the gB truncated sequence and in yellow the gD<sub>106</sub> tag sequence. **B)** Representative contrast phase and fluorescence images of pWPI/gB<sub>truncated</sub>-gD<sub>106</sub> transfected HEK cells at 24 hours after transfection.

**Additional file 5: Figure S5. A)** Overall strategy to reconstitute the ORF8 complete gB locus, via heat inducible homologous recombination. The 2911 bp gB locus, amplified by PCR (between positions 10576 and 13487 of BoHV-4-A genome), was introduced to replace the 2232 bp Kana-GalK selectable DNA stuffer, flanked by ORF8 homologous regions in the BoHV-4-A ΔgBKanaGalK strain cloned as a BAC. The expected ORF8 locus (A, bottom) has decreased size of the HindIII fragment (4314 instead of 5280 bp), generated by HindIII restriction enzyme digestion (diagram not on scale). **B)** HindIII restriction profile and corresponding Southern Blotting of five representative targeted clones, compared to the untargeted control. Southern Blotting was performed with a probe spanning Kana sequence and confirmed the above data. **C)** Clonal stability of the pBAC-BoHV-4-A-ΔgB<sub>revertant</sub> in *Escherichia coli* SW102 cells, passaged for 25 consecutive days and analyzed by HindIII digestion and agarose gel electrophoresis.

**Additional file 6: Figure S6.** Overall strategy to delete a 1261 bp sequence from the ORF8 codifying for gB, via heat inducible homologous recombination. The 2232 bp Kana-GalK selectable DNA stuffer, flanked by ORF8 homologous regions, was introduced between positions 11640 and 12901 of the BoHV-4-ADL1.7-eF-VSVG strain cloned as a BAC. The expected ORF8 locus (A, bottom) has an increased size of the HindIII fragment (5280 instead of 4314 bp), generated by HindIII restriction enzyme digestion (diagram not on scale). **B)** HindIII Restriction profile and corresponding Southern Blotting of six representative targeted clones, compared to the untargeted control. Southern Blotting was performed with a probe spanning Kana sequence and confirmed the above data. **C)** Clonal stability of the pBAC-BoHV-4-A-ΔgBKanaGalK in *Escherichia coli* SW102 cells, passaged for 30 consecutive days and analyzed by HindIII digestion and agarose gel electrophoresis.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contribution

VF performed the experiments and contributed to write the paper; AC contributed to perform the experiments; SC intellectually contributed; GD designed the study, interpret the data, performed the experiments and wrote the paper. All authors read and approved the final manuscript.

#### Acknowledgements

We would like to thank Dr. Giulio Grandi and Professor Laura Kramer for reading the paper and Italian Ministry of University and Scientific Research (Italian National Grant MIUR, PRIN 2008) for the financial support.

Received: 8 September 2012 Accepted: 7 November 2012

Published: 9 January 2013

#### References

1. Campadelli-Fiume G, Cocchi F, Menotti L, Lopez M: **The novel receptors that mediate the entry of herpes simplex viruses and animal alphaherpesviruses into cells.** *Rev Med Virol* 2000, **10**(5):305–319.
2. Lake CM, Molesworth SJ, Hutt-Fletcher LM: **The Epstein-Barr virus (EBV) gN homolog BLRF1 encodes a 15-kilodalton glycoprotein that cannot be authentically processed unless it is coexpressed with the EBV gM homolog BBRF3.** *J Virol* 1998, **72**(7):5559–5564.

3. Goltz M, Broll H, Mankertz A, Weigelt W, Ludwig H, Buhk HJ, Borchers K: **Glycoprotein B of bovine herpesvirus type 4: its phylogenetic relationship to gB equivalents of the herpesviruses.** *Virus Genes* 1994, **9**(1):53–59.
4. Thiry E, Bublot M, Dubuisson J, Van Bressems MF, Lequarre AS, Lomonte P, Vanderplasschen A, Pastoret PP: **Molecular biology of bovine herpesvirus type 4.** *Vet Microbiol* 1992, **33**(1–4):79–92.
5. Donofrio G, Cavirani S, van Santen V, Flammini CF: **Potential secondary pathogenic role for bovine herpesvirus 4.** *J Clin Microbiol* 2005, **43**(7):3421–3426.
6. Donofrio G, Herath S, Sartori C, Cavirani S, Flammini CF, Sheldon IM: **Bovine herpesvirus 4 is tropic for bovine endometrial cells and modulates endocrine function.** *Reproduction* 2007, **134**(1):183–197.
7. Dubuisson J, Boulanger D, Bublot M, Thiry E, Pastoret PP: **Proteins specified by bovine herpesvirus type 4: structural proteins of the virion and identification of two major glycoproteins by using monoclonal antibodies.** *J Gen Virol* 1989, **70**(Pt 7):1743–1753.
8. Vanderplasschen A, Bublot M, Dubuisson J, Pastoret PP, Thiry E: **Attachment of the gammaherpesvirus bovine herpesvirus 4 is mediated by the interaction of gp8 glycoprotein with heparinlike moieties on the cell surface.** *Virology* 1993, **196**(1):232–240.
9. Melnik LI, Garry RF, Morris CA: **Peptide inhibition of human cytomegalovirus infection.** *Virol J* 2011, **8**:76.
10. Lomonte P, Filee P, Lyaku JR, Bublot M, Pastoret PP, Thiry E: **Glycoprotein B of bovine herpesvirus 4 is a major component of the virion, unlike that of two other gammaherpesviruses, Epstein-Barr virus and murine gammaherpesvirus 68.** *J Virol* 1997, **71**(4):3332–3335.
11. Gillet L, Daix V, Donofrio G, Wagner M, Koszinowski UH, China B, Ackermann M, Markine-Goriaynoff N, Vanderplasschen A: **Development of bovine herpesvirus 4 as an expression vector using bacterial artificial chromosome cloning.** *J Gen Virol* 2005, **86**(Pt 4):907–917.
12. Donofrio G, Sartori C, Franceschi V, Capocéfalo A, Cavirani S, Taddei S, Flammini CF: **Double immunization strategy with a BoHV-4-vectorialized secreted chimeric peptide BVDV-E2/BoHV-1-gD.** *Vaccine* 2008, **26**(48):6031–6042.
13. Donofrio G, Franceschi V, Capocéfalo A, Cavirani S, Sheldon IM: **Isolation and characterization of bovine herpesvirus 4 (BoHV-4) from a cow affected by post partum metritis and cloning of the genome as a bacterial artificial chromosome.** *Reprod Biol Endocrinol* 2009, **7**:83.
14. Franceschi V, Capocéfalo A, Ravanetti L, Vanderplasschen A, Gillet L, Cavirani S, van Santen VL, Donofrio G: **Bovine herpesvirus 4 immediate early 2 (Rta) gene is an essential gene and is duplicated in bovine herpesvirus 4 isolate U.** *Vet Microbiol* 2011, **148**(2–4):219–231.
15. Warming S, Costantino N, Court DL, Jenkins NA, Copeland NG: **Simple and highly efficient BAC recombineering using galK selection.** *Nucleic Acids Res* 2005, **33**(4):e36.
16. *The recombineering website:* <http://web.ncicrf.gov/research/brb/recombineering/Information.aspx>.
17. Segura MF, Hanniford D, Menendez S, Reavie L, Zou X, Alvarez-Diaz S, Zakrzewski J, Blochin E, Rose A, Bogunovic D, et al: **Aberrant miR-182 expression promotes melanoma metastasis by repressing FOXO3 and microphthalmia-associated transcription factor.** *Proc Natl Acad Sci USA* 2009, **106**(6):1814–1819.
18. Donofrio G, Cavirani S, Taddei S, van Santen VL: **Potential of bovine herpesvirus 4 as a gene delivery vector.** *J Virol Methods* 2002, **101**(1–2):49–61.
19. Leoni V, Gianni T, Salvioli S, Campadelli-Fiume G: **Herpes simplex virus glycoproteins gH/gL and gB bind Toll-like receptor 2, and soluble gH/gL is sufficient to activate NF-kappaB.** *J Virol* 2012, **86**(12):6555–6562.
20. Lee SK, Compton T, Longnecker R: **Failure to complement infectivity of EBV and HSV-1 glycoprotein B (gB) deletion mutants with gBs from different human herpesvirus subfamilies.** *Virology* 1997, **237**(1):170–181.
21. Donofrio G, Sartori C, Ravanetti L, Cavirani S, Gillet L, Vanderplasschen A, Taddei S, Flammini CF: **Establishment of a bovine herpesvirus 4 based vector expressing a secreted form of the bovine viral diarrhoea virus structural glycoprotein E2 for immunization purposes.** *BMC Biotechnol* 2007, **7**:68.
22. Pertel PE, Spear PG, Longnecker R: **Human herpesvirus-8 glycoprotein B interacts with Epstein-Barr virus (EBV) glycoprotein 110 but fails to complement the infectivity of EBV mutants.** *Virology* 1998, **251**(2):402–413.



23. Tang J, Yang T, Ghosh HP, Geller AI: **Helper virus-free HSV-1 vectors packaged both in the presence of VSV G protein and in the absence of HSV-1 glycoprotein B support gene transfer into neurons in the rat striatum.** *J Neurovirol* 2001, **7**(6):548–555.
24. Borza CM, Hutt-Fletcher LM: **Alternate replication in B cells and epithelial cells switches tropism of Epstein-Barr virus.** *Nat Med* 2002, **8**(6):594–599.
25. Pereira L: **Function of glycoprotein B homologues of the family herpesviridae.** *Infect Agents Dis* 1994, **3**(1):9–28.
26. Glauser DL, Kratz AS, Gillet L, Stevenson PG: **A mechanistic basis for potent, glycoprotein B-directed gammaherpesvirus neutralization.** *J Gen Virol* 2011, **92**(Pt 9):2020–2033.
27. Holloway SA, Lindquister GJ, Studdert MJ, Drummer HE: **Identification, sequence analysis and characterisation of equine herpesvirus 5 glycoprotein B.** *Arch Virol* 1999, **144**(2):287–307.

doi:10.1186/1746-6148-9-6

**Cite this article as:** Franceschi et al.: Bovine herpesvirus 4 glycoprotein B is indispensable for lytic replication and irreplaceable by VSvg. *BMC Veterinary Research* 2013 **9**:6.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

