

## R E V I E W

# Selenium homeostasis and clustering of cardiovascular risk factors: a systematic review

*Mojgan Gharipour<sup>1</sup>, Masoumeh Sadeghi<sup>2</sup>, Mehrdad Behmanesh<sup>3</sup>, Mansour Salehi<sup>4</sup>, Pouya Nezafati<sup>5</sup>, Amin Gharipour<sup>6</sup>*

<sup>1</sup> Cardiac Rehabilitation Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medicine Sciences, Isfahan, Iran; <sup>2</sup> Cardiac Rehabilitation Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran; <sup>3</sup> Department of Genetics, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran; <sup>4</sup> Dept. of Genetics and Molecular Biology Medical School, Isfahan University of Medical Sciences; <sup>5</sup> Department of Cardiac Surgery, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran; <sup>6</sup> School of Information and Communication Technology, Gold Coast Campus, Griffith University, QLD4222, Australia

**Summary.** Selenium is a trace element required for a range of cellular functions. It is widely used for the biosynthesis of the unique amino acid selenocysteine [Sec], which is a structural element of selenoproteins. This systematic review focused on the possible relation between selenium and metabolic risk factors. The literature was searched via PubMed, Scopus, ISI Web of Science, and Google Scholar. Searches were not restricted by time or language. Relevant studies were selected in three phases. After an initial quality assessment, two reviewers extracted all the relevant data, whereas the third reviewer checked their extracted data. All evidence came from experimental and laboratory studies. Selenoprotein P is the best indicator for selenium nutritional levels. In addition, high levels of selenium may increase the risk of metabolic syndrome while the lack of sufficient selenium may also promote metabolic syndrome. Selenium supplementation in subjects with sufficient serum selenium levels has a contrary effect on blood pressure, LDL, and total cholesterol. According to the bioavailability of different types of selenium supplementation such as selenomethionine, selenite and selenium-yeast, it seems that the best nutritional type of selenium is selenium-yeast. Regarding obtained results of longitudinal studies and randomized controlled trials, selenium supplementation should not be recommended for primary or secondary cardio-metabolic risk prevention in populations with adequate selenium status. ([www.actabiomedica.it](http://www.actabiomedica.it))

**Key words:** selenium, hypertension, diabetes, obesity, cardiovascular disease, risk factors, metabolic syndrome

## Introduction

As a medicine with toxicity, selenium (Se), was first doubted to be used immediately after being discovered in 1817. It especially involves those who work in industries (1). Selenium has been shown to be present in some biological systems since more than 60 years ago (2). Being found a few years later, selenocysteine was learned to be encoded by the codon UGA

during the 1980s (3). Today, selenium is mainly applied in redox reactions, as well as in the biosynthesis of a unique amino acid, selenocysteine (Sec), which is a structural element of selenoproteins (3). Nearly 35 selenoproteins have been already introduced (4).

Se was first reported by Seale et al. to be as the main part of a diet and thus its necessary biological amount was suggested for human health (5). A significant geographical variation is seen in its dietary intake

from plants, which is low in Europe due to differences in its bioavailability. In some parts of the world, such as the UK, China, and Russia, its considerably reduced intake can be mainly caused by the lowered imports of wheat grown in the northern selenium-rich high-protein soils for making bread. Its deficiency has been frequently reported in regions where most food is locally grown and consumed (6).

Se has been also reported to be involved in the brain (7), immune system, and thyroid functions (8), as well as infertility (9), prevention of cancer (10), diabetes (11), and cardiovascular diseases (12). Therefore, it seems to play a vital role in the metabolic syndrome incidence and progress, i.e., clustering of cardio-metabolic risk factors like glucose intolerance, dyslipidemia, abdominal obesity, and hypertension (13-14). The prevalence of the mentioned syndrome is rather high among Iranian people, especially women with increasing ages (15). The enhanced production of reactive oxygen and nitrogen species in human body would provide several components of metabolic syndrome. Therefore, oxidative stress relevant to metabolic syndrome seems to be the most important pathway (16).

The oxidative process can be reduced by numerous antioxidants including  $\beta$ -carotene, vitamin C, and micro-elements, such as zinc and copper via inactivation of free radicals, and thus the organism can be protected against metabolic syndrome complications (17). This study aimed to find the relationship between selenium homeostasis and cardiometabolic risk factors, specially metabolic syndrome.

## Materials and method

### *Data Sources and search strategy*

Literature research included recorded studies conducted from January 1966 to December 2015. To find relevant articles, searches were made in MEDLINE via PubMed, Scopus, ISI Web of Science, and Google Scholar. Searches were not restricted by language. Keywords included: "selenium" and "metabolic syndrome", "selenium and cardiovascular disease", "selenium" and "diabetes", selenium" and "hypertension", and "selenium" and "abdominal obesity".

### *Inclusion criteria*

To identify the relevant studies focused on selenium bioavailability and its levels associated with cardiovascular diseases and risk factors and metabolic syndrome, the following inclusion criteria were applied in the current review.

## Discussion and conclusion

### *Selenium involvement in biological systems*

In various pathophysiological conditions, selenium plays an essential role. Via the synthesis of selenoproteins containing at least one selenocysteine (Sec) as an amino acid, it is involved in biological pathways. Most selenoproteins have oxidoreductase functions. Sec is co-translated into a growing polypeptide in response to UGA codon, which ultimately terminates the translation. The Sec insertion machinery is developed by the biosystems to decode UGA as Sec. This machinery allows this amino acid to be assimilated at exclusive UGA codons with the help of a cis-acting element of Sec Insertion Sequence (SECIS) (18-22).

Transcription of selenoproteins in tissues differs based on a response to dietary intake of selenium or reactive oxygen species (23) as environmental signals. There are lower mRNA levels of several selenoproteins under selenium absence conditions (24). This process has been shown to occur through the decay of transcripts mediated by nonsense in-frame termination codons (25). However, selenoprotein translation is regulated by reduced mRNA levels, which are less considerable than the decrease in selenoenzyme activities (26-27).

Selenoprotein synthesis regulation can be controlled via a suitable UGA codon (Sec) recognition. Some studies have demonstrated that UGA translation is not responsive to selenium alone, while selenium supplementation is more efficient as UGA codon is read by reducing polysome loading of selenoprotein mRNAs (28). According to the results of reporter assays based on UGA or UGU at an appropriate position, Sec translation efficiency has been estimated from 1% to 15% (30).

tRNA[Ser]Sec may have a regulatory role in this process though SECIS elements individually support UGA recognition as Sec (29). Selenium appears as selenocysteine at the active site of many selenoproteins, such as glutathione peroxidase enzymes of classical GPx1, gastrointestinal GPx2, plasma GPx3, and phospholipid hydroperoxide GPx4 as a major class of selenoproteins with important functions.

Thioredoxin reduction, which is dependent on NADPH, is catalyzed by Thioredoxin reductase (TR) enzyme recently identified to contain selenocysteine, thus regulating its metabolic activity. Selenoprotein P with 10 Se atoms per molecule as selenocysteine contains almost 60% of Se in plasma. It may act as a protein transporting selenium. However, many other tissues have selenoprotein-P expression. This suggests that facilitation of selenium distribution in the whole body is not its sole function.

Being involved in the catalysis of 5'-mono-deiodination of prohormonethyroxine (T4) to 3,3',5'-triiodothyronine (T3) as an active thyroid hormone, iodothyronine deiodinase enzymes make the second major class of selenoproteins. The mid-piece portion of spermatozoa is a site for sperm capsule selenoprotein where the sperm flagella integrity is stabilized. As reported to be necessary for muscle metabolism, selenoprotein W is affected by selenium intake in terms of tissue concentrations. Having no systematic investigation conducted on reduced Se status in the UK over the past 2 decades would be a great concern for health implications.

#### *Selenium homeostasis in patients with cardiovascular diseases*

Selenium role in cardiovascular diseases has not been proven and thus remained highly controversial though many studies have been conducted on this issue. Overall, a significant relationship between selenium and cardiovascular diseases can be found in the baseline selenium concentration as a determining factor (30).

On this basis, carotid artery atherosclerosis progression in men with high serum LDLs has been reported by Salonen et al. to be probably related to serum selenium concentrations (31). Additionally, my-

ocardial infarction risk and death caused by ischemic heart diseases were discovered by them to be resulted from low selenium content of the Finnish soil in the eastern area of Finland as a determining factor (32). Lower serum selenium than 79 µg/L was identified by another cohort study in Denmark to enhance coronary heart disease risks among subjects with no relevant pre-existing histories (33). In contrast, no relationship between selenium concentration and ischemic heart diseases was found by Miettinen et al. (34). Moreover, in their study in the countryside areas of eastern and southwestern Finland, Virtamo et al. were not able to find a significant association between coronary heart disease risk and serum selenium (35). The association between cardiovascular diseases and serum selenium could not be explained by the studies carried out in other parts of Europe as well (36-37).

Higher mean serum selenium concentrations were found in all the above studies compared to those reported in the Finnish studies (31-33). Correspondingly, no relationship was seen between serum selenium and myocardial infarction risk in the US population. This can be due to high selenium intake level among them (38). In this research, very few subjects were found with less serum selenium than 80 µg/L, while the mean plasma levels were high among the rest and controls.

Several studies have dealt with the association between coronary disease extents and selenium status. As quantified by angiography, coronary atherosclerosis severity and serum selenium concentration were reported by these studies to be inversely correlated (39-40).

#### *Selenium and cardiometabolic risk factors*

##### *a. Selenium and Diabetes*

In different regions with high (the US) and low (China and the UK) selenium statuses, few data on the relationship between cardiometabolic risk factors and selenium status are available among the general population (41). An enhanced prevalence of diabetes was discovered by Laclaustra et al. to have occurred with increasing selenium levels in American adults (42-45).

The relationship between glycosylated hemoglobin levels, fasting plasma glucose, and diabetes and serum selenium concentrations was evaluated by La-

claustra et al. with the help of regression models and their prevalence was found to have augmented with increasing selenium concentrations up to 160 micro G/L (42). No relationship between diabetes or glucose levels and selenium was observed in countries with lower selenium levels, such as Singapore and France (42). Also, an association between blood selenium levels and metabolic syndrome and higher fasting blood sugar levels was discovered among Chinese adults (46).

#### *b. Selenium and dyslipidemia*

Higher cholesterol, triglyceride, and fasting glucose were found to be associated with higher selenium levels in a study on elderly Taiwanese (46). The relationship between blood lipids and plasma selenium concentration in British adults was observed with a mean plasma selenium concentration of  $1.10 \pm 0.19$  micromol/L by Stranges et al. Contrary to HDL levels, enhanced non-HDL and total cholesterol levels were discovered to be associated with higher plasma selenium ( $> \text{or} = 1.20$  micromol/L) in the UK adult population (47). In another research among US adults with sufficient selenium uptakes, elevated serum concentrations of HDL and LDL cholesterol, total cholesterol, triacylglycerol, apo A-I, and apo B were demonstrated by Bleys et al. to be associated with increased serum selenium levels (48). HDL cholesterol rise among US population was reported by the National Health and Nutrition Examination Survey [NHANES] to only occur at low selenium levels in 2003–2004 (42).

#### *c. Selenium and hypertension*

An underestimated risk factor of selenium deficiency was declared by Nawrot et al. to probably lead to the development of high blood pressure among European men. In this regard, 2.2 and 1.5 mmHg lower systolic and diastolic blood pressures [ $P = 0.017$ ] were shown to be associated with a higher blood selenium of 20  $\mu\text{g/L}$  by them by performing multivariate-adjusted cross-sectional analyses on men, respectively. Although no significant relationship was found between selenium and hypertension in women, a lower risk of developing high-normal blood pressure (37% [ $P = 0.001$ ]) was discovered to be associated with a higher baseline blood selenium of 20  $\mu\text{g/L}$  in their prospective analyses in men (49).

#### *d. Selenium and obesity*

No directly sufficient data on the relationship between abdominal obesity and selenium are available; yet, the role of different regimes containing high selenium resources of fruits and legumes as obesity indicators has been considered by few studies. Through insulin-resistance mechanisms, selenium seems to play a role in obesity pathophysiology. Improvement of the Body Mass Index (BMI) levels in obese children has been stated by Velázquez-López et al. to be resulted from a high uptake of mineral antioxidants like selenium through a Mediterranean eating regime (50). Based on a combined antioxidant supplementation with selenium, vitamin C, and vitamin E in obese children and adolescents participating in a modification program, various types of obesity were claimed by Mureret al. to be probably linked with inflammatory pathways and oxidative stresses (51).

Improved metabolic characteristics and reduced proinflammatory status in overweight and obese people was revealed in another study carried out by Hermsdorff et al. to be resulted from a hypocaloric diet based on legumes (52). A great advantage was shown by Savini et al. to be provided by foods containing antioxidants in obese subjects via redox state modulation (53).

#### *Selenium supplementation in different target groups and cardiometabolic risk factors*

Cardiometabolic risk factors in different target groups have been investigated by several studies. The efficiencies and biological activities of various types of selenium and improper forms for selenium supplementation in the mentioned groups have raised many questions. Plasma selenium was significantly modified after at least one week. After 2 to 11 weeks of supplementation, a stable status was achieved with the inorganic forms. Higher net enhancement was observed in the population groups of low selenium areas compared to those with a higher intake of basal selenium (54). On the contrary, increased levels of plasma selenium were seen to be caused by selenomethionine even after supplementation for 8.5 weeks (55).

The bioavailability of various selenium types, including organic-derivative forms of selenium, selenite, selenomethionine, food selenium, and yeast were in-

investigated by another study (56). Although yeast selenium produced inconsistent results, food (wheat and meat) selenium resulted in similar findings. In the case of yeast selenium, either a plateau in plasma selenium after 2, 11, and 16 weeks (57-59) was obtained or in a steady enhancement in selenium levels was caused until the end of the experiment at 11 to 12 weeks (60-61). Changes in plasma selenium remarkably differed from those of erythrocyte selenium. Only Chinese subjects with very low Se showed increased selenium levels of erythrocyte triggered by inorganic forms (62). Otherwise, even after 32 weeks, no modification was seen to have been caused by inorganic selenium supplementation.

Contrarily, all the studies revealed selenium enhancement in red blood cells caused by organic forms except for a study on yeast selenium (63). The activity of erythrocyte GSH-Px was shown as a less sensitive indicator of changing selenium levels due to changes in plasma, red blood cells, and glutathione peroxidase activities of platelets (64). A significant augmentation necessarily occurred at a slow rate of 4 to 6 weeks for a modification (65). GSH-Px activity of red blood cells in the Finish population underwent no changes induced by selenium intake of about 40 mg/day as reported by Levander et al. Also, a significant rise or change in GSH-Px activity of erythrocytes was reported by them to incumbently occur at least 1 to 3 weeks of supplementation (64).

Butler et al. (65) noted that selenate taken by women in New Zealand was associated with greater percentages of selenium and GSH-Px in plasma and red blood cells compared to those taking selenomethionine. They concluded higher blood selenium levels occur for selenomethionine compared to selenite when GSH-Px activity of platelets is saturated (65).

No protective effect of antioxidant supplementation in a well-nourished population was revealed by another clinical trial conducted on 5,220 participants. Nevertheless, subjects at risk for metabolic syndrome could be protected by baseline concentrations of serum antioxidants. This was highly dependent on lifestyle and particularly dietary patterns (66).

Glucose intolerance risk has not been shown to be significantly increased by high doses of supplemental selenium. However, the ineffectiveness of se-

lenium supplementation on the prevalence of type 2 diabetes was revealed in the subjects participating in a Selenium and Vitamin E Cancer Prevention Trial (SELECT) via a follow-up analysis for 10 years (67). Thus, our understanding of selenium regulation, especially its influence on energy metabolism is limited by these findings.

A nutritional dose of nearly 50 to 200  $\mu\text{g}/\text{day}$  of selenium was suggested by Joseph et al. based on the relationship between selenium and various disorders, such as myocardial ischemia/infarction, cardiomyopathy, and reperfusion injury (68).

## Conclusion

Several notable facts about selenium were revealed by this systematic study. First, selenium exact effect on cardiometabolic risk factors depends on serum selenium level. Lack of adequate selenium may develop metabolic syndrome, while high selenium levels may also increase the risk of metabolic syndrome. People with sufficient selenium levels should not take selenium supplements as cardiometabolic risk factors in a population with adequate selenium levels of serum are adversely affected by selenium treatment. For instance, selenium supplementation in the mentioned groups had a reverse influence on their blood pressures, LDLs, and total cholesterols.

Second, selenium nutritional levels seem to be indicated by selenoprotein P in the best way. Thus, selenium deficiency can be determined by selenoprotein P measurement among different populations.

Third, the best nutritional type of selenium seems to be selenium yeast as examined from different types of selenium supplementations, such as selenite, selenomethionine, and yeast selenium based on bioavailability.

Finally, primary or secondary cardiometabolic risks cannot be prevented by selenium supplementation in populations with sufficient selenium status considering the results of longitudinal studies and randomized controlled trials. However, plasma concentration of selenoprotein P can be optimized in a population with lower selenium concentrations by recommending a nutritional dose of 50 to 200  $\mu\text{g}/\text{day}$  of selenium.

## References

- da Silva AP. Selenium. Its physiologic and clinical importance. *Acta Med Port* 1993 Jul; 6(7): 347-60. Review. Portuguese. Erratum in: *Acta Med Port* 1993 Oct; 6(10): 493. PubMed PMID: 8379356.
- Burk RF. Biological activity of selenium. *Annu Rev Nutr* 1983; 3: 53-70. Review. PubMed PMID: 6357242.
- Stadtman TC. Selenium biochemistry. *Science* 1974 Mar 8; 183(4128): 915-22. Review. PubMed PMID: 4605100.
- Vadgama JV, Wu Y, Shen D, Hsia S, Block J. Effect of selenium in combination with Adriamycin or Taxol on several different cancer cells. *Anticancer Res* 2000 May-Jun; 20(3A): 1391-414. PubMed PMID: 10928049.
- Seale LA, Gilman CL, Hashimoto AC, Ogawa-Wong AN, Berry MJ. Diet-Induced Obesity in the Selenocysteine-Lyase Knockout Mouse. *Antioxid Redox Signal* 2015 Oct 1; 23(10): 761-74. doi: 10.1089/ars.2015.6277.
- Reiter R, Wendel A. Selenium and drug metabolism--I. Multiple modulations of mouse liver enzymes. *Biochem Pharmacol* 1983 Oct 15; 32(20): 3063-7. PubMed PMID: 6639674.
- Stadtman TC. Selenium biochemistry. *Science* 1974 Mar 8; 183(4128): 915-22. Review. PubMed PMID: 4605100.
- Kabata-Pendias A. Geochemistry of selenium. *J Environ Pathol Toxicol Oncol* 1998; 17(3-4): 173-7.
- Seale LA, Hashimoto AC, Kurokawa S, Gilman CL, Seydali A, Bellinger FP, Raman AV, Berry MJ. Disruption of the selenocysteine lyase-mediated selenium recycling pathway leads to metabolic syndrome in mice. *Mol Cell Biol* 2012 Oct; 32(20): 4141-54.
- Dodig S, Cepelak I. The facts and controversies about selenium. *Acta Pharm* 2004 Dec; 54(4): 261-76. Review. PubMed PMID: 15634611.
- Brigelius-Flohé R, Banning A, Schnurr K. Selenium-dependent enzymes in endothelial cell function. *Antioxid Redox Signal* 2003 Apr; 5(2): 205-15. Review. PubMed PMID: 12716480.
- Rayman MP. The importance of selenium to human health. *Lancet* 2000 Jul 15; 356(9225): 233-41. Review. PubMed PMID: 10963212.
- Reaven P. Metabolic syndrome. *J Insur Med* 2004; 36(2): 132-42. Review. PubMed PMID: 15301226.
- Gharipour M, Sadeghi M, Hosseini M, Andalib E, Boroujeni MB, Sarrafzadegan N. Effect of age on the phenotype of metabolic syndrome in developing country. *Adv Biomed Res* 2015 May 29; 4: 103.
- Sarrafzadegan N, Gharipour M, Sadeghi M, Khosravi AR, Tavassoli AA. Metabolic syndrome in Iranian elderly. *ARYA Atheroscler* 2012 Winter; 7(4): 157-61.
- Sarrafzadegan N, Gharipour M, Ramezani MA, Rabiei K, Zolfaghari B, Tavassoli AA, Boshtam M, Zarfeshani S, Khosravi A, Yousefi A. Metabolic syndrome and health-related quality of life in Iranian population. *J Res Med Sci* 2011 Mar; 16(3): 254-61.
- Valko M, Morris H, Cronin MT. Metals, toxicity and oxidative stress. *Curr Med Chem* 2005; 12(10): 1161-208.
- Labunskyy VM, Hatfield DL, Gladyshev VN. Selenoproteins: molecular pathways and physiological roles. *Physiol Rev* 2014 Jul; 94(3): 739-77. doi:10.1152/physrev.00039.2013. Review. PubMed PMID: 24987004
- Diamond AM, Choi IS, Crain PF, Hashizume T, Pomerantz SC, Cruz R, Steer CJ, Hill KE, Burk RF, McCloskey JA, Hatfield DL. Dietary selenium affects methylation of the wobble nucleoside in the anticodon of selenocysteine tRNA[[Ser]Sec]. *J Biol Chem* 1993 Jul 5; 268(19): 14215-23.
- Zhou J, Liu WJ, Peng SW, Sun XY, Frazer I. Papillomavirus capsid protein expression level depends on the match between codon usage and tRNA availability. *J Virol* 1999 Jun; 73(6): 4972-82. PubMed PMID: 10233959.
- Jameson RR, Diamond AM. A regulatory role for Sec tRNA[Ser]Sec in selenoprotein synthesis. *RNA* 2004 Jul; 10(7): 1142-52. PubMed PMID: 15208449.
- Jornot L, Junod AF. Hyperoxia, unlike phorbol ester, induces glutathione peroxidase through a protein kinase C-independent mechanism. *Biochem J* 1997 Aug 15; 326 (Pt 1): 117-23. PubMed PMID: 9337858.
- Bermano G, Nicol F, Dyer JA, Sunde RA, Beckett GJ, Arthur JR, Hesketh JE. Tissue-specific regulation of selenoenzyme gene expression during selenium deficiency in rats. *Biochem J* 1995 Oct 15; 311 (Pt 2): 425-30. PubMed PMID: 7487877
- Hadley KB, Sunde RA. Selenium regulation of thioredoxin reductase activity and mRNA levels in rat liver. *J Nutr Biochem*. 2001 Dec; 12(12): 693-702. PubMed PMID: 12031252.
- Maquat LE. Evidence that selenium deficiency results in the cytoplasmic decay of GPx1 mRNA dependent on pre-mRNA splicing proteins bound to the mRNA exon-exon junction. *Biofactors* 2001; 14(1-4): 37-42. Review. PubMed PMID: 11568438.
- Di Cosmo C, McLellan N, Liao XH, Khanna KK, Weiss RE, Papp L, Refetoff S. Clinical and molecular characterization of a novel selenocysteine insertion sequence-binding protein 2 [SBP2] gene mutation [R128X]. *J Clin Endocrinol Metab* 2009 Oct; 94(10): 4003-9. doi: 10.1210/jc.2009-0686. Epub 2009 Jul 14. PubMed PMID: 19602558; PubMed Central PMCID: PMC2758735.
- Weiss Sachdev S, Sunde RA. Selenium regulation of transcript abundance and translational efficiency of glutathione peroxidase-1 and -4 in rat liver. *Biochem J* 2001 Aug 1; 357(Pt 3): 851-8. PubMed PMID: 11463357; PubMed Central PMCID: PMC1222016.
- Fletcher JE, Copeland PR, Driscoll DM. Polysome distribution of phospholipid hydroperoxide glutathione peroxidase mRNA: evidence for a block in elongation at the UGA/selenocysteine codon. *RNA* 2000 Nov; 6(11): 1573-84. PubMed PMID: 11105757.
- Martin GW 3rd, Berry MJ. Selenocysteine codons decrease polysome association on endogenous selenoprotein mRNAs. *Genes Cells* 2001 Feb; 6(2): 121-9. PubMed PMID: 11260257.

30. Kollmus H, Hentze MW, Hauser H. Regulated ribosomal frameshifting by an RNA-protein interaction. *RNA* 1996 Apr; 2(4): 316-23. PubMed PMID: 8634912; PubMed Central PMCID: PMC1369374.
31. Salonen JT, Salonen R, Seppanen K, Kantola M, et al. Interactions of serum copper, selenium and LDL-C in atherogenesis. *Br Med J* 1991; 302: 756-60.
32. Salonen JT, Salonen R, Pentilla J, et al. Serum fatty acids, apolipoproteins, selenium and vitamin antioxidants and the risk of death from CAD. *Am J Cardiol* 1985; 56: 226-31
33. Suadicani P, Hein HO, Gyntelberg F. Serum selenium concentration and risk of ischaemic heart disease in a prospective cohort study of 3000 males. *Atherosclerosis* 1992; 96: 33-42.
34. Miettinen TA, Alfthan G, Huttunen JK, et al. Serum selenium concentration related to MI and fatty acid content of serum lipids. *Br Med J* 1983; 287: 517-9.
35. Virtamo J, Valkeila E, Alfthan G, et al. Serum selenium and the risk of CHD and stroke. *Am J Epidemiol* 1985; 122: 276-82.
36. Ringstad J, Thelle DS. Risk of MI in relation to serum concentration of selenium. *ActaPharmacolToxicol* 1986; 59(7): 336-9.
37. Kok FJ, deBruijnAM, Vermeeren R, et al. Serum selenium, vitamin antioxidants and cardiovascular mortality: a nine year follow up study in the Netherlands. *Am J Clin Nutr* 1987; 45: 462-8.
38. Salvini S, Hennekens CH, Morris JS, et al. Plasma levels of the antioxidant selenium and risk of MI among US physicians. *Am J Cardiol* 1995; 76: 1218-21.
39. Moore JA, Noiva R, Wells IC. Selenium concentration in plasma of patients with arteriographically defined coronary atherosclerosis. *Clin Chem* 1984; 30: 1171-3.
40. Kok FJ, Vanpoppel G, Melse J, et al. Do antioxidants and PUFA have a combined association with coronary atherosclerosis? *Atherosclerosis* 1991; 86: 85-90.
41. Haug A, Graham RD, Christophersen OA, Lyons GH. How to use the world's scarce selenium resources efficiently to increase the selenium concentration in food. *Microb Ecol Health Dis* 2007 Dec; 19(4): 209-228.
42. Laclaustra M, Navas-Acien A, Stranges S, Ordovas JM, Guallar E. Serum selenium concentrations and diabetes in U.S. adults: National Health and Nutrition Examination Survey [NHANES] 2003-2004. *Environ Health Perspect* 2009 Sep; 117(9): 1409-13.
43. Hurst R, Hooper L, Norat T, Lau R, Aune D, Greenwood DC, Vieira R, CollingsR,Harvey LJ, Sterne JA, Beynon R, Savović J, Fairweather-Tait SJ. Selenium and prostate cancer: systematic review and meta-analysis. *Am J Clin Nutr* 2012 Jul; 96(1): 111-22.
44. Hurst R, Armah CN, Dainty JR, Hart DJ, Teucher B, Goldson AJ, Broadley MR, Motley AK, Fairweather-Tait SJ. Establishing optimal selenium status: results of a randomized, double-blind, placebo-controlled trial. *Am J Clin Nutr* 2010 Apr; 91(4): 923-31.
45. Rayman MP. The importance of selenium to human health. *Lancet* 2000 July 15; 356(9225): 233-41. Review. PubMed PMID: 10963212.
46. Yang KC, Lee LT, Lee YS, Huang HY, Chen CY, Huang KC. Serum selenium concentration is associated with metabolic factors in the elderly: a cross-sectional study. *Nutr Metab [Lond]* 2010 May 6; 7: 38.
47. Stranges S, Tabák AG, Guallar E, Rayman MP, Akbaraly TN, Laclaustra M, Alfthan G, Mussalo-Rauhamaa H, Viikari JS, Raitakari OT, Kivimäki M. Selenium status and blood lipids: the cardiovascular risk in Young Finns study. *J Intern Med* 2011 Nov; 270(5): 469-77.
48. Bleys J, Navas-Acien A, Laclaustra M, Pastor-Barriuso R, Menke A, Ordovas J, Stranges S, Guallar E. Serum selenium and peripheral arterial disease: results from the national health and nutrition examination survey, 2003-2004. *Am J Epidemiol* 2009 Apr 15; 169(8): 996-1003. doi: 10.1093/aje/kwn414. Epub 2009 Feb 16. PubMed PMID: 19221120; PubMed Central PMCID: PMC2727225.
49. Nawrot TS, Staessen JA, Roels HA, Den Hond E, Thijs L, Fagard RH, Dominiczak AF, Struijker-Boudier HA. Blood pressure and blood selenium: a cross-sectional and longitudinal population study. *Eur Heart J* 2007 Mar; 28(5): 628-33.
50. Velázquez-López L, Santiago-Díaz G, Nava-Hernández J, Muñoz-Torres AV, Medina-Bravo P, Torres-Tamayo M. Mediterranean-style diet reduces metabolic syndrome components in obese children and adolescents with obesity. *BMC Pediatr* 2014 Jul 5; 14: 175.
51. Murer SB, Aeberli I, Braegger CP, Gittermann M, Hersberger M, Leonard SW, Taylor AW, Traber MG, Zimmermann MB. Antioxidant supplements reduced oxidative stress and stabilized liver function tests but did not reduce inflammation in a randomized controlled trial in obese children and adolescents. *J Nutr* 2014 Feb; 144(2): 193-201.
52. Hermsdorff HH, Puchau B, Volp AC, Barbosa KB, Bressan J, Zulet MÁ, Martínez JA. Dietary total antioxidant capacity is inversely related to central adiposity as well as to metabolic and oxidative stress markers in healthy young adults. *NutrMetab [Lond]* 2011 Aug 22; 8: 59. doi: 10.1186/1743-7075-8-59. PubMed PMID:21859453; PubMed Central PMCID: PMC3179702.
53. Savini I, Catani MV, Evangelista D, Gasperi V, Avigliano L. Obesity-associated oxidative stress: strategies finalized to improve redox state. *Int J Mol Sci* 2013 May 21; 14(5): 10497-538.
54. Clausen J, Nielsen SA. Comparison of whole blood selenium values and erythrocyte glutathione peroxidase activities of normal individuals on supplementation with selenate, selenite, L selenomethionine, and high selenium yeast. *Biol Trace Elem Res* 1988 Jan-Apr; 15: 125-38.
55. Xia Y, Hill KE, Byrne DW, Xu J, Burk RF. Effectiveness of selenium supplements in a low-selenium area of China. *Am J Clin Nutr* 2005 Apr; 81(4): 829-34.
56. Bleys J, Navas-Acien A, Guallar E. Serum selenium levels and all-cause, cancer, and cardiovascular mortality among US adults. *Arch Intern Med* 2008 Feb 25; 168(4): 404-

10. doi: 10.1001/archinternmed.2007.74. PubMed PMID: 18299496.
57. Bugel SH, Sandstrom B, Larsen EH. Absorption and retention of selenium from shrimps in man. *J Trace Elem Med Biol* 2001; 14: 198-204.
58. Bleys J, Navas-Acien A, Stranges S, Menke A, Miller ER 3rd, Guallar E. Serum selenium and serum lipids in US adults. *Am J Clin Nutr* 2008 Aug; 88(2): 416-23. PubMed PMID: 18689378; PubMed Central PMCID: PMC2553708.
59. Hurst R, Armah CN, Dainty JR, Hart DJ, Teucher B, Goldson AJ, Broadley MR, Motley AK, Fairweather-Tait SJ. Establishing optimal selenium status: results of a randomized, double-blind, placebo-controlled trial. *Am J Clin Nutr* 2010 Apr; 91(4): 923-31.
60. Daniels LA. Selenium metabolism and bioavailability. *Biol Trace Elem Res* 1996; 54: 185-199.
61. Bleys J, Navas-Acien A, Guallar E. Selenium and diabetes: more bad news for supplements. *Ann Intern Med* 2007 Aug 21; 147(4): 271-2. Epub 2007 Jul 9. PubMed PMID: 17620657.
62. van der Torre HW, Van Dokkum W, Schaafsma G, Wedel M, Ockhuizen T. Effect of various levels of selenium in wheat and meat on blood Se status indices and on Se balance in Dutch men. *Br J Nutr* 1991 Jan; 65(1): 69-80. PubMed PMID: 1997131.
63. Bleys J, Navas-Acien A, Guallar E. Serum selenium and diabetes in U.S. adults. *Diabetes Care* 2007 Apr; 30(4): 829-34. PubMed PMID: 17392543.
64. Levander OA. Considerations on the assessment of selenium status. *Fed Proc* 1985 Jun; 44(9): 2579-83. PubMed PMID: 3996613.
65. Butler JA, Thomson CD, Whanger PD, Robinson MF. Selenium distribution in blood fractions of New Zealand women taking organic or inorganic selenium. *Am J Clin Nutr* 1991 Mar; 53(3): 748-54.
66. Czernichow S, Vergnaud AC, Galan P, Arnaud J, Favier A, Faure H, Huxley R, Hercberg S, Ahluwalia N. Effects of long-term antioxidant supplementation and association of serum antioxidant concentrations with risk of metabolic syndrome in adults. *Am J Clin Nutr* 2009 Aug; 90(2): 329-35.
67. Lippman SM, Goodman PJ, Klein EA, Parnes HL, Thompson IM Jr, Kristal AR, et al. Designing the Selenium and Vitamin E Cancer Prevention Trial [SELECT]. *J Natl Cancer Inst* 2005; 97: 94-102.
68. Joseph J, Loscalzo J. Selenistasis: epistatic effects of selenium on cardiovascular phenotype. *Nutrients* 2013 Jan 31; 5(2): 340-58.

Received: 22 December 2016

Accepted: 1 February 2017

Correspondence:

Dr. Masoumeh Sadeghi

Associate Professor in Cardiology, Cardiac Rehabilitation

Research Center, Cardiovascular Research Institute,

Isfahan University of Medical Sciences, Isfahan, Iran

E-mail: sadeghimasoumeh@gmail.com