

RESEARCH ARTICLE

Open Access

Reversal of TGF- β_1 stimulation of α -smooth muscle actin and extracellular matrix components by cyclic AMP in Dupuytren's - derived fibroblasts

Latha Satish^{1*}, Phillip H Gallo¹, Mark E Baratz², Sandra Johnson¹ and Sandeep Kathju¹

Abstract

Background: Myofibroblasts, a derived subset of fibroblasts especially important in scar formation and wound contraction, have been found at elevated levels in affected Dupuytren's tissues. Transformation of fibroblasts to myofibroblasts is characterized by expression of alpha- smooth muscle actin (α -SMA) and increased production of extracellular matrix (ECM) components, both events of relevance to connective tissue remodeling. We propose that increasing the activation of the cyclic AMP (cAMP)/protein kinase A signaling pathway will inhibit transforming growth factor-beta1 (TGF- β_1)-induced ECM synthesis and myofibroblast formation and may provide a means to blunt fibrosis.

Methods: Fibroblasts derived from areas of Dupuytren's contracture cord (DC), from adjacent and phenotypically normal palmar fascia (PF), and from palmar fascia from patients undergoing carpal tunnel release (CTR; CT) were treated with TGF- β_1 (2 ng/ml) and/or forskolin (10 μ M) (a known stimulator of cAMP). Total RNA and protein extracted was subjected to real time RT-PCR and Western blot analysis.

Results: The basal mRNA expression levels of fibronectin- extra domain A (FN1-EDA), type I (COL1A2) and type III collagen (COL3A1), and connective tissue growth factor (CTGF) were all significantly increased in DC- and in PF-derived cells compared to CT-derived fibroblasts. The TGF- β_1 stimulation of α -SMA, CTGF, COL1A2 and COL3A1 was greatly inhibited by concomitant treatment with forskolin, especially in DC-derived cells. In contrast, TGF- β_1 stimulation of FN1-EDA showed similar levels of reduction with the addition of forskolin in all three cell types.

Conclusion: In sum, increasing cAMP levels show potential to inhibit the formation of myofibroblasts and accumulation of ECM components. Molecular agents that increase cAMP may therefore prove useful in mitigating DC progression or recurrence.

Background

Dupuytren's contracture (DC) is a fibroproliferative disease of the hand's palmar fascia, which can cause permanent and irreversible flexion contracture of the digits [1]. It is the most common inherited disease of connective tissues in humans [2]. Although DC is not rare, debate over its etiology has been ongoing since before its modern-day description over 120 years ago [3]. DC is known to result from changes occurring in the dermis and palmar fascia [4]. Fibroblasts are the major cell population associated with DC in all stages (both during

the formation of nodules and cords) and represent an important target for therapeutic intervention. Importantly, differentiation of fibroblasts into myofibroblasts, identified by their expression of alpha-smooth muscle actin (α -SMA) [5-9], is considered to be responsible for the development of typical clinical symptoms and offers an opportunity for molecular intervention.

Myofibroblast formation is controlled by a variety of growth factors, cytokines and even mechanical stimuli [8,10]. Transforming growth factor-beta1 (TGF- β_1) is the most important of these and has been demonstrated in Dupuytren's tissue using various techniques [11,12] along with its receptors [4]. Berndt et al. [13] showed a greater intensity of staining for TGF- β_1 protein in proliferative nodules and colocalization of TGF- β_1 synthesis

* Correspondence: lsatish@wpahs.org

¹Center for Genomic Sciences, Allegheny-Singer Research Institute, Pittsburgh, PA-15212, USA

Full list of author information is available at the end of the article

with the myofibroblast phenotype to these regions. Furthermore, addition of TGF- β_1 resulted in significant up-regulation of cells staining for α -SMA in primary cultures of fibroblasts derived from Dupuytren's nodule and cord tissue. It therefore seems likely that this growth factor plays a central function in the development and progression of the disease.

Surgical intervention remains the mainstay of treatment for DC, but there is a high recurrence rate after surgery [14-16]. TGF- β_1 release might also play a significant role in the recurrence of the disease after surgical treatment. The local trauma of surgical excision and the resultant natural wound healing response will typically lead to the release of growth factors which include TGF- β_1 . Any residual tissue with a disease or pre-disease phenotype will be susceptible to stimulation, myofibroblast transformation, collagen synthesis and the formation of recurrent disease. Some studies have correlated recurrence of DC with the presence of myofibroblasts [17].

In this context, it is reasonable to hypothesize that a means of counter-acting the signaling mechanisms of TGF- β -mediated up-regulation of α -SMA and ECM gene expression in Dupuytren's tissue may provide novel approaches to the therapy of DC disease. Accordingly, we have focused our attention on cyclic AMP (cAMP), a signal transduction mediator that may interfere with TGF- β -initiated functions. The second messenger cAMP regulates fibroblast physiology in many tissues. Intracellular cAMP levels are the result of a balance between synthesis, which is regulated by G-protein-coupled receptors that stimulate (via G_s) or inhibit (via G_i) adenylyl cyclase (AC), and degradation, which occurs via cyclic nucleotide phosphodiesterase (PDE). Increases in cAMP influence cell growth, cell death, and differentiated cell functions, primarily (although not exclusively) by promoting phosphorylation of proteins via the activation of cAMP-dependent protein kinase A (PKA) [18]. PKA-mediated phosphorylation of cAMP-response element-binding protein (CREB) and CREB-mediated regulation of transcription via interaction with cAMP-response elements is a major pathway that alters cellular gene expression [19].

One mechanism by which cAMP may regulate fibrogenicity is via interaction with the TGF- β signaling pathway. Recent work suggests that activation of the cAMP/PKA signaling pathway inhibits TGF β_1 -induced collagen synthesis and myofibroblast formation in cardiac and pulmonary fibroblasts [20,21]. These results suggest that overproduction of cAMP may provide a means to blunt fibrosis.

To our knowledge there have been no studies that investigate the relationship between cAMP signaling and TGF- β -mediated effects in DC disease. In this study we

sought to establish the baseline functioning of cAMP and the effects of its elevation in DC-derived fibroblasts. We specifically examined alpha-smooth muscle actin, connective tissue growth factor (CTGF), as well as important components of the extracellular matrix.

Methods

Cell Culture

Primary cultures of fibroblasts were obtained from the surgically resected Dupuytren's contracture samples (DC), from matching specimens of normal appearing palmar fascia in DC patients (PF), and from specimens of normal palmar fascia of patients undergoing carpal tunnel surgery (CT) as previously described [22,23]. All samples were collected with the informed consent of the patient and the study protocol conformed to the ethical guidelines of the 1975 Declaration of Helsinki. All specimens were collected with the approval of the Allegheny-Singer Research Institute's institution review board involving Human Subjects (IRB protocol RC-4040) and all the patients signed the written informed consent under institutional review board approval. The cultures were maintained in MEM- α medium (Invitrogen Corporation, Carlsbad, CA) supplemented with 10% fetal bovine serum (FBS, Gemini Bioproducts, West Sacramento, CA) and 1% antibiotic-antimycotic solution (Sigma, St Louis, MO). All cultures were used at passage levels between 3-6 with no changes evident in cell morphology.

CT-, PF-, and DC- derived fibroblasts were plated onto 6-well Falcon tissue culture plates and grown until 80% confluence. Cells were quiesced for 24 hours in

MEM- α medium supplemented with 0.1% dialyzed fetal bovine serum (Gemini Bioproducts) and 1% antibiotic-antimycotic solution. After 24 hours the cells were then treated or not with TGF- β_1 (2 ng/ml) (Peprotech, Inc. Rockyhill, NJ) and/or forskolin (Sigma) (10 μ M) and incubated for 37°C for 24 hours. Cells were then washed with phosphate buffered saline (PBS) and lysed using M-PER obtained from Thermo Fisher Scientific (Rockford, IL) for protein extraction and RLT-lysis buffer (Qiagen Inc. Valencia, CA) for RNA isolation according to the manufacturer's instructions. RNA quality was assessed by A260/280 ratio using an ND-1000 spectrophotometer (Nanodrop Technologies Inc, Wilmington, DE) and by capillary electrophoresis with the Agilent 2100 bioanalyzer (Agilent technologies, Inc. Palo Alto, CA). At least three independent primary cell cultures of CT-, PF- and DC- derived fibroblasts were used in experiments involving treatment with TGF- β_1 or forskolin. Six independent sets of CT-, PF-, and DC-derived fibroblasts were used in establishing the basal mRNA expression of specific extracellular matrix (ECM) proteins.

Quantitative Real time RT-PCR

Total RNA isolated (RNeasy Micro Kit, Qiagen Inc., Valencia, CA) from untreated DC-, PF- and CT-derived fibroblasts was subjected to real time RT-PCR to determine the relative mRNA expression levels at baseline for fibronectin (FN1-EDA), type I collagen (COL1A2), type III collagen (COL3A1) and connective tissue growth factor (CTGF). RNA isolated from cells treated with TGF- β_1 , forskolin, and with both agents was also subjected to real time RT-PCR to determine the changes in the mRNA levels of α -SMA (ACTA2), FN1-EDA, COL1A2, COL3A1 and CTGF.

Real-time RT-PCR was performed using kits obtained from Applied Biosystems (Foster City, CA) that utilize FAM™Taqman®MGB probes and a Taqman® Universal PCR Master Mix. Assays were performed on the above noted gene products using human GAPDH as an endogenous normalizing control. Reverse transcription was performed on 30 ng of total RNA with random primers (100 ng), gene specific primer for FN1-EDA (10 pmole) and with M-MLV-reverse transcriptase (Invitrogen Corporation, Carlsbad, CA). The primers (forward primer 5'-TAAAGGACTGGCATTCACTGATGT-3'; reverse primer - 3'-GTGCAAGGCAACCACACTGA-5') and probe (5'6 FAM-CCCTGAGGATGGAATCCATGAGC-TATTCC-TAMRA 3') used for human FN1-EDA [GenBank: X07718] were designed using Primer Express software (Applied Biosystems). Primers were obtained from Integrated DNA Technologies (Coralville, IA) and Taqman probes were purchased from Applied Biosystems. In all assays the primer sets were first tested to verify that amplifiers of the expected molecular weight resulted before their employment in real time RT-PCR.

Subsequent PCR amplification and detection of template was carried out using Applied Biosystems transcript-specific assays including: COL1A2 (ID-Hs01028971_m1), COL3A1 (ID-Hs00943793), ACTA2 (ID-Hs00426835_g1) and CTGF (ID-Hs00170014_m1) using 15 ng of cDNA and 20x final concentration of Gene Expression Mix, which contains both forward and reverse primers adjusted to final volume of 15.0 μ l. Identical reaction mixes were prepared with human FN1-EDA primers and probes. The reaction set up and the thermal cycling protocol were as previously described (23). Using the comparative critical cycle (C_t) method the expression levels of the target genes were normalized to the GAPDH endogenous control (ID-HS99999905_m1) and the relative abundance was calculated. Data were analyzed using the 7900 HT SDS software version 2.1 provided by Applied Biosystems.

Immunoblotting

Proteins extracted were subjected to Bradford assay to determine the protein concentration. Equal quantities of

proteins were separated on SDS-PAGE, transferred to a Whatman™ Protran pure nitrocellulose immobilization membrane (GE Health Care, Piscataway, NJ) and probed with antibodies specific to α -SMA (Abcam, Cambridge, MA) and fibronectin (Santa Cruz Biotechnology, Inc. Santa Cruz, CA) using GAPDH (Abcam, Cambridge, MA) as loading control. The membranes were conjugated with HRP-labeled secondary antibody, and the signals were detected using SuperSignal® West Femto Trial Kit Prod #34094 (Thermo Scientific, Rockford, IL). The intensity of the protein bands was quantitated using NIH Image J 1.44p, available in the public domain at <http://imagej.nih.gov/ij>.

Statistical Analysis

Statistical analyses were performed using two-way ANOVA utilizing GraphPad Prism 5 for Windows Version (5.04) from Graph Pad Software Inc. Utilizing the same program Bonferroni post-test to compare replicate means by row was also performed to determine the p values. P value less than 0.05 was considered significant.

Results

Basal mRNA expression levels of ECM proteins were significantly increased in Dupuytren-derived fibroblasts

We first examined the message levels of ECM proteins, namely COL1A2, COL3A1, FN1-EDA and CTGF, a matricellular protein, by qRT-PCR. Our results identified increased mRNA expression levels of all the above gene products in DC- derived fibroblasts relative to CT- derived fibroblasts (Figure 1a, b, c, d). Interestingly, PF- derived fibroblasts express these ECM components in a similar fashion to fibroblasts from active disease, suggesting that even apparently normal fascia in DC patients may harbor an incipient disease phenotype.

Forskolin inhibited the TGF- β_1 stimulation of α -SMA mRNA and protein

Our previous findings have demonstrated an elevation at baseline of α -SMA mRNA and protein levels in DC- in comparison to CT- and PF- derived fibroblasts (Satish et al., manuscript in preparation). The present study shows that addition of TGF- β_1 greatly augments the levels of α -SMA mRNA in CT-, PF- and DC- derived fibroblasts. To determine if increased levels of cAMP could reduce the TGF- β_1 induced levels of α -SMA, forskolin, a well-established adenylyl cyclase (AC) activator and an inducer of cAMP in fibroblasts [20,24,25] was utilized. We found that by increasing cAMP levels there was a substantial reduction in TGF- β_1 induced mRNA levels of α -SMA in DC- derived fibroblasts compared to TGF- β_1 treatment alone. Although apparent reductions in TGF- β_1 -induced α -SMA mRNA levels were also observed in CT-derived fibroblasts and PF-derived fibroblasts




Figure 1 Basal mRNA expression of ECM proteins was significantly elevated in DC-derived fibroblasts. Real time RT-PCR was performed on RNA extracted from 6 independent primary cultures derived from CT-, PF- and DC- tissues to determine the mRNA expression levels of COL1A2 (a), COL3A1 (b), FN1-EDA (c), and CTGF (d). Values are means \pm SEM of three independent experiments performed in duplicate. Statistical analyses were performed using two-way ANOVA.

compared with TGF- β_1 treatment alone, the extent of these cAMP effects was significantly less than in DC-derived cells (Figure 2a). Similar significant reductions in TGF- β_1 -induced α -SMA protein levels were seen in all three-cell types by Western blot (Figure 3a-d). Forskolin by itself did not have any significant effect on α -SMA mRNA or protein levels in any cell type. These results strongly suggest that myofibroblast formation (as evidenced by α -SMA accumulation) can be significantly inhibited in DC- derived cells by increasing cAMP levels.

Forskolin reduced the TGF- β_1 induction of fibronectin mRNA and protein

Extracellular matrix deposition likely plays a crucial role in the fibrosis noted in DC, and previous studies have observed increased deposition of an oncofetal isoform of fibronectin (IIIICS spliced variant) in DC lesional tissues and in DC-derived primary cell cultures [22]. In this study we examined FN1-extra domain A (EDA), as this isoform has shown differential expression between fibrotic versus scarless healing seen in mucosal and skin wound healing [26]. Forskolin treatment alone had no significant effect on FN1-EDA mRNA levels in any of our three cell types (Figure 2b), nor were fibronectin

protein levels affected in CT- and PF-derived cells, but we did observe a significant decrease in fibronectin protein in DC- derived fibroblasts on forskolin treatment by Western blot (Figure 3a-c, e), the mechanism for which may be post-transcriptional.

We found that forskolin inhibited TGF- β_1 -induction of fibronectin mRNA to a similar degree in CT-, PF- and DC- derived fibroblasts when measured against TGF- β_1 treatment alone (Figure 2b). This is in contrast to α -SMA, where DC-derived cells were uniquely and especially susceptible to this forskolin effect. Fibronectin protein levels in all three cell types also showed relative decrease when forskolin was added compared to TGF- β_1 alone (Figure 3a-c, e).

Forskolin inhibited the TGF- β_1 induction of CTGF mRNA in PF- and DC- derived cells but not CT-derived cells

We next determined the effect of increased cAMP levels on another TGF- β_1 target gene, CTGF. Since TGF- β may induce CTGF through several pathways, including SMAD, ras/raf/MEK/ERK, Ets-1, JNK, and protein kinase C, CTGF has long been thought to be an important mediator of its fibrotic effects [27-30]. The TGF- β_1 induction of CTGF mRNA increase was substantially reduced by combined incubation with forskolin in PF-




Figure 2 Forskolin effectively reduced TGF- β_1 stimulation of α -SMA, FN1-EDA, CTGF, COL1A2 and COL3A1. Fibroblast cultures from CT-, PF- and DC- tissues were left untreated or were stimulated with forskolin (10 μ M) in the presence or absence of TGF- β_1 (2 ng/ml). Twenty-four hours later, mRNA expression levels of α -SMA (a), FN1-EDA (b), CTGF (c), COL1A2 (d) and COL3A1 (e) were analyzed by real time RT-PCR. In each experiment at least three independent cultures obtained from all the three cell types were used. Values are means \pm SEM of six independent studies performed in duplicate. Statistical analyses were performed using two-way ANOVA.

and DC- derived fibroblasts compared to TGF- β_1 alone (Figure 2c). As with α -SMA, these results again suggest that the biology of fibroblasts from DC patients is exquisitely sensitive to the mitigating actions of cAMP.

Forskolin reduced the TGF- β_1 stimulation of Type I and Type III collagen

We next investigated the effect of increased cAMP (via forskolin treatment) on collagen expression as TGF- β is a known stimulator of collagen production [31]. We specifically examined if increased cAMP levels can abrogate TGF- β_1 induction of type I (α -2 chain; COL1A2) and type III collagen (α -1 chain; COL3A1) expression. Forskolin alone did not have any significant effect on the relative levels of COL1A2 and COL3A1 mRNAs in any of the three cell types. Forskolin did, however, suppress the TGF- β_1 induction of COL1A2 and COL3A1 mRNAs in CT-, PF- and DC -derived fibroblasts (Figure 2d, e). Of note, the degree of inhibition seen when TGF- β_1 was co-incubated with forskolin was significantly greater in DC -derived cells than in the CT- or PF-cells. Since increased collagen deposition is a hallmark of DC disease, these results again suggest that mechanisms to elevate cAMP may be useful adjunctive therapies to counteract the fibrotic phenotypes of DC cells.

Discussion

Dupuytren's contracture, fibrosis in the palmar fascia of the hand, is a fibroproliferative disorder that can impose severe functional damage eventually leading to disability of the hand in affected individuals [32]. Efforts have been made to control the fibrosis seen in DC using various non-surgical treatment strategies but with limited success [33]. Injectable collagenase clostridium histolyticum [34] to treat DC shows potential promise but its clinical application has thus far elicited a varied response among hand surgeons. Alternative treatment options including non-surgical molecular therapeutic agents to prevent progression and recurrence of DC disease are still wanting.

Because myofibroblast formation and activity have been linked to the etiology of both primary and recurrent DC, molecular interventions that interfere with myofibroblastic functions may offer a novel avenue of therapy. A number of such interventions have been proposed and essayed. Glucocorticoids have been shown to increase apoptosis of Dupuytren's-associated fibroblasts, and to reduce the abundance of TGF- β_1 and fibronectin CS1 in myofibroblast-populated stroma in DC nodules injected with depomedrone [35,36]. Repeated intraleisional injection of DC nodules (not cords) with triamcinolone did show some regression of the nodules [37] but some 50% of patients developed recurrence or




Figure 3 TGF- β_1 stimulated α -SMA and FN1-EDA protein expression were substantially reduced by forskolin in CT-, PF- and DC-derived fibroblasts. CT-, PF- and DC-derived fibroblasts (a, b, c) grown on 6-well culture dishes were stimulated with forskolin (10 μ M), TGF- β_1 (2 ng/ml), with both agents, or were left untreated for 24 hours in MEM- α medium containing 0.1% dialyzed FBS. Whole cell lysates were collected after 24 hours and the samples were processed for Western immunoblotting with specific antibodies for α -SMA and FN1-EDA (20 μ g/lane). Specificity of the modulation and identical protein loading was confirmed with a loading control GAPDH antibody. Densitometry analysis was done on protein bands obtained from three independent experiments performed in triplicate using two independent primary cultures of CT-, PF- and DC-derived fibroblasts (Figure 3d and 3e). A representative immunoblot is shown here.

progression of the disease within the window of the study. Whether such an approach would succeed in more advanced disease with actual cord formation is unclear.

Another agent that acts against myofibroblasts that has been used in DC is 5-fluorouracil (5-FU). Treatment of DC-derived fibroblasts with 5-FU inhibited their proliferation and their differentiation to myofibroblasts [38]. However, clinical use of 5-FU at the time of surgery resulted in no difference between treated and untreated digits as determined by joint angle measurements [39], leaving its clinical utility open to question.

It has been observed in rat cardiac fibroblasts and in a human pulmonary fibroblast-derived cell line that elevation of cAMP can inhibit cellular proliferation and differentiated functions (such as collagen synthesis). These observations suggested that a similar approach might favorably alter fibroblast/myofibroblast behavior in the setting of Dupuytren's contracture. We therefore sought to determine if increased cAMP levels could inhibit TGF- β_1 -induced myofibroblast formation (as indicated by α -SMA accumulation) and ECM production in DC-derived cells. TGF- β_1 was chosen as a test stimulatory cytokine as it has been implicated in the pathogenesis of DC [2,4,40].

Multiple interesting observations have arisen from these experiments. When assaying for basal levels of expression of α -SMA and ECM proteins in our three cell types, it is clear that PF-derived cells more closely resemble DC-derived cells than control CT-derived cells in all four gene products tested. This suggests that, although obtained from phenotypically normal fascia, PF-derived cells may already exhibit a disease phenotype at the cellular level. Such an observation is consistent with our total expressomic analyses of DC- and PF- versus CT-derived fibroblasts, wherein we find that global gene expression patterns of PF-cells closely resemble (but are not identical to) DC-derived cells and vary sharply from CT-derived cells (Satish et al., manuscript in preparation).

We also found that TGF- β_1 , as expected, increased expression levels of all gene products assayed significantly, whereas cAMP elevation (as induced by forskolin treatment) alone had minimal effect. cAMP was, however, in all instances able to dramatically blunt the effects of TGF- β_1 . DC-derived cells were particularly susceptible to cAMP action, generally exhibiting more inhibition of gene expression by cAMP action than PF- or CT-cells. These observations suggest that agents to elevate cAMP may well be able to suppress the differentiation of DC-fibroblasts to a myofibroblast phenotype, and to mitigate the abnormal ECM deposition that would then typically ensue. Although forskolin (or other

similar agents) may be impractical to deliver directly to DC-affected tissues over the long periods of time in which the disease develops or progresses, we postulate that molecular therapeutic approaches administering activated adenylyl cyclase, possibly by a gene therapy approach, may accomplish the same effects. Successful use of adenylyl cyclase to inhibit myofibroblast formation and function has been demonstrated in cardiac and pulmonary cells [20,21].

A particular point of interest in this study is the examination of the behavior of CTGF in our three cell types. CTGF has been described as a co-factor to TGF- β by enhancing ligand-receptor binding in activated cells [41]. Studies in various cell populations have also demonstrated roles for CTGF in the TGF- β -dependent induction of fibronectin, collagen and tissue inhibitor of metalloproteinase-1 (TIMP-1) [42-44]. A recent study by Sisco et al. [45] showed that antisense inhibition of CTGF could limit hypertrophic scarring *in vivo* without affecting the outcome of wound closure. To our knowledge this report for the first time demonstrates increased basal expression levels of CTGF in PF- and in DC-derived fibroblasts compared to CT-derived cells, and this relative increase is enhanced by addition of TGF- β_1 . Further, we also find that elevated cAMP levels most successfully reduce this increased CTGF mRNA expression in DC-derived fibroblasts. This report thus points to a potential role for CTGF in the etiopathology of DC, and suggests that measures to target its expression or function (including agents that elevate cAMP) may usefully limit fibrosis in Dupuytren's contracture.

The observations reported herein do not directly identify the precise mechanisms by which increased cAMP levels inhibit myofibroblast formation. Recent data indicate that cAMP acts in a PKA-dependent manner to inhibit TGF- β /Smad signaling and gene activation by disruption of transcriptional cofactor binding in human keratinocytes [46]; it is possible that similar mechanisms are at work in DC-fibroblasts, and are being investigated. Moreover, we are in the process of delineating the migratory and contractile behavior of DC-derived fibroblasts when cAMP levels are increased. Demonstration of a change in these mechanocellular properties would provide even more evidence of the utility of a cAMP-based approach as an anti-fibrotic measure in Dupuytren's contracture.

Conclusion

In summary, increasing cAMP levels show potential to inhibit the formation of myofibroblasts and accumulation of ECM components. Molecular agents that increase cAMP may therefore prove useful in mitigating DC progression or recurrence.

Acknowledgements

The authors thank Allegheny-Singer Research Institute, The Pittsburgh Foundation and Pennsylvania Department of Health for their financial support towards this study. We extend our thanks to Dr. J Christopher Post and Dr. Garth D Ehrlich for their support. We thank Ms. Mary O'Toole for her assistance with the preparation of this manuscript.

Author details

¹Center for Genomic Sciences, Allegheny-Singer Research Institute, Pittsburgh, PA-15212, USA. ²Division of Upper Extremity Surgery, Department of Orthopaedics, Allegheny General Hospital, Pittsburgh, PA-15212, USA.

Authors' contributions

LS conceived the study. LS and SK discussed and designed the study. LS, PG, SJ performed the experiments. LS and SK drafted the manuscript. LS, SK and MEB critically reviewed manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 27 December 2010 Accepted: 25 May 2011

Published: 25 May 2011

References

1. Bayat A, Cunliffe EJ, McGrouther DA: Assessment of clinical severity in Dupuytren's disease. *Br J Hosp Med* 2007, **68**:604-609.
2. Tomasek JJ, Vaughan MB, Haakma CJ: Cellular structure and biology of Dupuytren's disease. *Hand Clin* 1999, **15**:21-34.
3. Dupuytren G: Permanent retraction of the fingers, produced by an affection of the palmar fascia. Clinical lectures on surgery. *Lancet* 1884, **2**:222.
4. Kloen P, Jennings CL, Gebhardt MC, Springfield DS, Mankin HJ: TGF-beta: possible roles in Dupuytren's contracture. *J Hand Surg* 1995, **20A**:101-108.
5. Luck JV: Dupuytren's contracture: a new concept of the pathogenesis correlated with surgical management. *J Bone Joint Surg [Am]* 1959, **41-A**:635-664.
6. Chiu HF, McFarlane RM: Pathogenesis of Dupuytren's contracture: a correlative clinic-pathological study. *J Hand Surg* 1978, **3**:1-10.
7. Schürch W, Skalli O, Gabbiani G: Cellular biology. In *Dupuytren's disease: biology and treatment*. Edited by: McFarlane RM, McGrouther DA, Flint MH. London: Churchill Livingstone; 1990:31-47.
8. Serini G, Gabbiani G: Mechanisms of myofibroblast activity and phenotypic modulation. *Exp Cell Res* 1999, **250**:273-283.
9. Hindman HB, Marty-Roix R, Tang JB, Jupiter JB, Simmons BP, Spector M: Regulation of expression of alpha-smooth muscle actin in cells of Dupuytren's contracture. *J Bone Joint Surg Br* 2003, **85**:448-455.
10. Powell DW, Mifflin RC, Valentich JD, Crowe SE, Saada JL, West AB: Myofibroblasts. II. Intestinal subepithelial myofibroblasts. *Am J Physiol* 1999, **277**:C1-C9.
11. Badalamente MA, Sampson SP, Hurst LC, Dowd A, Miyasaka K: The role of TGF-beta in Dupuytren's disease. *J Hand Surg* 1996, **21A**:210-215.
12. Zamaro RL, Heights R, Kraemer BA, Ehrlich HP, Groner JP: Presence of growth factors in palmar and plantar fibromatoses. *J Hand Surg* 1994, **19A**:435-441.
13. Berndt A, Kosmehl H, Mandel U, Gabler U, Leo X, Celeda D, Zardi L, Katenkamp D: TGF-beta and bFGF synthesis and localization in Dupuytren's disease (nodular palmar fibromatosis) relative to cellular activity, myofibroblast phenotype and oncofetal variants of fibronectin. *Histochem J* 1995, **27**:1014-1020.
14. Rodrigo JJ, Niebauer JJ, Brown RL, Doyle JR: Treatment of Dupuytren's contracture-long-term results after fasciotomy and fascial excision. *J Bone Joint Surg* 1976, **58A**:380-387.
15. Dias JJ, Braybrooke J: Dupuytren's contracture: an audit of the outcomes of surgery. *J Hand Surg* 2006, **31B**:514-521.
16. Badalamente MA, Hurst LC: Efficacy and safety of injectable missed collagenase subtypes in the treatment of Dupuytren's contracture. *J Hand Surg* 2007, **32A**:767-774.
17. Gelberman RH, Amiel D, Rudolph RM, Vance RM: Dupuytren's contracture. An electron microscopic, biochemical, and clinical correlative study. *J Bone Joint Surg* 1980, **62A**:425-432.
18. Francis SH, Corbin JD: Structure and function of cyclic nucleotide-dependent protein kinases. *Ann rev Physiol* 1994, **56**:237-272.
19. Montminy M: Transcriptional regulation by cyclic AMP. *Annu Rev Biochem* 1997, **66**:807-822.
20. Liu X, Ostrom RS, Insel PA: cAMP-elevating agents and adenylyl cyclase overexpression promote an antifibrotic phenotype in pulmonary fibroblasts. *Am J Physiol* 2004, **286**:C1089-C1099.
21. Swaney JS, Roth DM, Olson ER, Naugle JE, Meszaros GJ, Insel PA: Inhibition of cardiac myofibroblast formation and collagen synthesis by activation and overexpression of adenylyl cyclase. *Proc Natl Acad Sci USA* 2005, **102**:437-442.
22. Howard JC, Varallo VM, Ross DC, Faber KJ, Roth JH, Seney S, Gan BS: Wound healing-associated proteins Hsp47 and fibronectin are elevated in Dupuytren's contracture. *J Surg Res* 2004, **117**:232-238.
23. Satish L, Laframboise WA, O'Gorman DB, Johnson S, Janto B, Gan BS, Baratz ME, Hu FZ, Post JC, Ehrlich GD, Kathju S: Identification of differentially expressed genes in fibroblasts derived from patients with Dupuytren's Contracture. *BMC Med Genomics* 2008, **1**:10.
24. Böhm M, Raghunath M, Sunderkötter C, Schiller M, Ständer S, Brzoska T, Cauvet T, Schiöth HB, Schwartz T, Luger TA: Collagen metabolism is a novel target of the neuropeptide alpha-melanocyte-stimulating hormone. *J Biol Chem* 2004, **279**:6959-6966.
25. Schiller M, Dennler S, Anderegg U, Kokot A, Simon JC, Luger TA, Mauviel A, Böhm M: Increased cAMP levels modulate TGF-β/Smad-induced expression of extracellular matrix components and other key fibroblast functions. *J Biol Chem* 2010, **285**:409-421.
26. Li-Korotky HS, Hebda PA, Lo CY, Dohar JE: Age-dependent differential expression of fibronectin variants in skin and airway mucosal wounds. *Arch Otolaryngol Head Neck Surg* 2007, **133**:919-924.
27. Pannu J, Nakerakanti S, Smith E, ten Dijke P, Trojanowska M: Transforming growth factor-beta receptor type I-dependent fibrogenic gene program is mediated via activation of Smad1 and ERK1/2 pathways. *J Biol Chem* 2007, **282**:10405-10413.
28. Leask A, Holmes A, Black CM, Abraham DJ: Connective tissue growth factor gene regulation. Requirements for its induction by transforming growth factor-beta 2 in fibroblasts. *J Biol Chem* 2003, **278**:13008-13015.
29. Verrecchia F, Chu ML, Mauviel A: Identification of novel TGF-beta/SMAD gene targets in dermal fibroblasts using a combined cDNA microarray/promoter transactivation approach. *J Biol Chem* 2001, **276**:17058-17062.
30. Holmes A, Abraham DJ, Sa S, Shiwen X, Black CM, Leask A: CTGF and SMADs, maintenance of scleroderma phenotype is independent of SMAD signaling. *J Biol Chem* 2001, **276**:10594-10601.
31. Alioto RJ, Rosier RN, Burton RL, Puzas JE: Comparative effects of growth factors on fibroblasts of Dupuytren's tissue and normal palmar fascia. *J Hand Surg Am* 1994, **19**:442-452.
32. Hindocha S, Stanley JK, Watson JS, Bayat A: Revised tubiana's staging system for assessment of disease severity in dupuytren's disease-Preliminary clinical findings. *Hand* 2008, **3**:80-86.
33. Hurst LC, Badalamente MA: Nonoperative treatment of Dupuytren's disease. *Hand Clin* 1999, **15**:97-107.
34. Hurst LC, Badalamente MA, Hertz VR, Hotchkiss RN, Kaplan FT, Meals RA, Smith TM, Rodzivila J, CORD I Study Group: Injectable collagenase clostridium histolyticum for Dupuytren's contracture. *N Engl J Med* 2009, **361**:968-979.
35. Meek RM, McLellan S, Crossan JF: Dupuytren's disease. A model for the mechanism of fibrosis and its modulation by steroids. *J Bone Joint Surg Br* 1999, **81**:732-738.
36. Meek RM, McLellan S, Reilly J, Crossan JF: The effect of steroids on Dupuytren's disease: role of programmed cell death. *J Hand Surg Br* 2002, **27**:270-273.
37. Ketchum LD, Donahue TK: The injection of nodules of Dupuytren's disease with triamcinolone acetonide. *J Hand Surg Am* 2000, **25**:1157-1162.
38. Jemec B, Linge C, Grobelaar AO, Smith PJ, Sanders R, McGrouther DA: The effect of 5-fluorouracil on Dupuytren fibroblast proliferation and differentiation. *Chir Main* 2000, **19**:15-22.
39. Bulstrode NW, Bisson M, Jemec B, Pratt AL, McGrouther DA, Grobelaar AO: A prospective randomised clinical trial of the intra-operative use of 5-fluorouracil on the outcome of dupuytren's disease. *J Hand Surg Br* 2004, **29**:18-21.
40. Baird KS, Crossan F, Ralston SH: Abnormal growth factor and cytokine expression in Dupuytren's contracture. *J Clin Pathol* 1993, **46**:425-428.

41. Abreu JG, Kotpura NI, Reversade B, De Robertis EM: Connective-tissue growth factor (CTGF) modulates cell signaling by BMP and TGF-beta. *Nature Cell Biol* 2002, 4:599-604.
42. Arnott JA, Nugloze E, Rico MC, Arango-Hisijara I, Odgren PR, Safadi FF, Popoff SN: Connective tissue growth factor (CTGF/CCN2) is a downstream mediator for TGF-beta1-induced extracellular matrix production in osteoblasts. *J Cell Physiol* 2007, 210:843-852.
43. Chujo S, Shirasaki F, Kawara S, Inagaki Y, Kinbara T, Inaoki M, Takigawa M, Takehara K: Connective tissue growth factor causes persistent proalpha2(I) collagen gene expression induced by transforming growth factor-beta in a mouse fibrosis model. *J Cell Physiol* 2005, 203:447-456.
44. Grotendorst GR: Connective tissue growth factor: a mediator of TGF-beta action on fibroblasts. *Cytokine Growth Factor Rev* 1997, 8:171-179.
45. Sisco M, Kyger ZB, O'Shaughnessy KD, Kim PS, Schultz GS, Ding XZ, Roy NK, Dean NM, Mustoe TA: Antisense inhibition of connective tissue growth factor (CTGF/CCN2) mRNA limits hypertrophic scarring without affecting wound healing in vivo. *Wound Repair Regen* 2008, 16:661-673.
46. Schiller M, Verrecchia F, Mauviel A: Cyclic adenosine 3',5'-monophosphate-elevating agents inhibit transforming growth factor-beta-induced SMAD3/4-dependent transcription via a protein kinase A-dependent mechanism. *Oncogene* 2003, 22:8881-8890.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1471-2474/12/113/prepub>

doi:10.1186/1471-2474-12-113

Cite this article as: Satish et al.: Reversal of TGF- β_1 stimulation of α -smooth muscle actin and extracellular matrix components by cyclic AMP in Dupuytren's - derived fibroblasts. *BMC Musculoskeletal Disorders* 2011 12:113.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

