


RESEARCH ARTICLE

Open Access



Investigation of cortical thickness and volume during spontaneous attacks of migraine without aura: a 3-Tesla MRI study

Faisal Mohammad Amin^{1*} , Roberto De Icco^{2,3}, Mohammad Al-Mahdi Al-Karagholi¹, Jayachandra Raghava^{4,5}, Frauke Wolfram⁶, Henrik B. W. Larsson⁴ and Messoud Ashina¹

Abstract

Background: Structural imaging has revealed changes in cortical thickness in migraine patients compared to healthy controls is reported, but presence of dynamic cortical and subcortical changes during migraine attack versus inter-ictal phase is unknown. The aim of the present study was to investigate possible changes in cortical thickness during spontaneous migraine attacks. We hypothesized that pain-related cortical area would be affected during the attack compared to an inter-ictal phase.

Methods: Twenty-five patients with migraine without aura underwent three-dimensional T1-weighted imaging on a 3-Tesla MRI scanner during spontaneous and untreated migraine attacks. Subsequently, 20 patients were scanned in the inter-ictal phase, while 5 patients did not show up for the inter-ictal scan. Four patients were excluded from the analysis because of bilateral migraine pain and another one patient was excluded due to technical error in the imaging. Longitudinal image processing was done using FreeSurfer. Repeated measures ANOVA was used for statistical analysis and to control for multiple comparison the level of significance was set at $p = 0.025$.

Results: In a total of 15 patients, we found reduced cortical thickness of the precentral ($p = 0.023$), pericalcarine ($p = 0.024$), and temporal pole ($p = 0.017$) cortices during the attack compared to the inter-ictal phase. Cortical volume was reduced in prefrontal ($p = 0.018$) and pericalcarine ($p = 0.017$) cortices. Hippocampus volume was increased during attack ($p = 0.007$). We found no correlations between the pain side or any other clinical parameters and the reduced cortical size.

Conclusion: Spontaneous migraine attacks are accompanied by transient reduced cortical thickness and volume in pain-related areas. The findings constitute a fingerprint of acute pain in migraine patients, which can be used as a possible biomarker to predict antimigraine treatment effect in future studies.

Trial registration: The study was registered at [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02202486) (NCT02202486).

Keywords: Migraine cortex, Pain cortex, Migraine brain, Cortical volume, Migraine attack

* Correspondence: faisal@dadlnet.dk

¹Danish Headache Center, Department of Neurology, Faculty of Health and Medical Sciences, Rigshospitalet Glostrup, University of Copenhagen, Valdemar Hansens Vej 5, 2600 Glostrup, Denmark

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Structural neuroimaging studies in migraine had focus on unveiling underlying pathophysiological mechanisms and radiological biomarkers of the disease. Several data supported the presence of cortical and sub-cortical morphological alterations in the migraine brain, although conflicting evidences were published [1]. When compared to healthy controls, migraine patients were characterized by an increased cortical thickness of somatosensory, frontal, occipital, and temporal areas [2–4]. In contrast, other studies reported a decreased thickness of frontal as well as somatosensory areas [4–6], or even no differences with healthy controls [7]. Another way to investigate the migraine brain is represented by the comparison of different phases of the migraine cycle. Resting-state functional magnetic resonance imaging was widely used in the past years to explore this topic. During spontaneous migraine attacks, an altered thalamocortical network was described, namely an increased connectivity with orbitofrontal and parietal cortices, and a decreased connectivity with primary somatosensory cortex [8]. Moreover, the hypothalamus showed dynamic connections during the migraine cycle namely a coupling with the spinal trigeminal nuclei in the inter-ictal phase, that shifts to a stronger connection with the dorsal rostral pons in the ictal phase [9]. Nonetheless, cortical thickness and volume could exhibit rapid changes when studied in different conditions, or after exposure to specific stimuli [10]. The pathophysiology of this acute modulation is not completely elucidated, and fast-adjusting processes, such as regulation of glial cells volumes, interstitial cortical space as well as synaptic elements, probably play a major role [10]. Despite these observations, evidences about morphologic changes between different phases of the migraine cycle are lacking.

The aim of this study is to investigate the morphological changes of cortical and sub-cortical brain structures during spontaneous migraine attacks of migraine without aura compared to outside of the attack. We hypothesized that migraine attacks would be accompanied by morphological changes in pain-related areas in the brain.

Methods

Study design and participants

Patients were recruited from 2013 to 2014 among those attending the outpatient clinic of the Danish Headache Center (Copenhagen, Denmark) and through web a page (<http://www.forsoegsperson.dk>). Inclusion criteria were: age 18 to 65 years; diagnosis of episodic migraine without aura according to the ICHD criteria [11]. Exclusion criteria were: history of any other primary or secondary headache (only infrequent episodic tension-type headache less than 5 days a month was allowed); pregnant or

breast-feeding women; contraindication for MRI; signs or verified diagnosis of any cardiovascular or cerebrovascular disease; active psychiatric disorder and/or drug abuse.

Procedures

The patients underwent two MRI scans during two separate sessions: session 1 (S1) was performed during a spontaneous migraine attack, while session 2 (S2) was performed in the inter-ictal period.

During S1, patients were asked to call an investigator of the study (FMA) as soon as possible after the onset of a migraine attack. All patients were previously instructed about the International Headache Society's migraine without aura criteria [12]. After the call, patients travelled to the hospital by taxi or public transport (fare reimbursed) between 4 and 24 h from attack onset. After hospital arrival, the patients underwent the first brain MRI scan during the spontaneous migraine attack. Patients were not allowed to treat their attacks before the MRI scan. During S2, the same patients were scanned during an inter-ictal migraine-free period. The inter-ictal period was defined as no headache in the 48 h, and no migraine attack in the 72 h before the MRI scan. Analgesic as well as migraine-specific drugs were not allowed in the 48 h before and 24 h after S1 and S2. All T1-weighted images were assessed by a neuroradiologist (FW) for pathological abnormalities, which were not found in any of the patients included in the present study.

MRI data analysis

We obtained the MRI recordings from a Philips Achieva 3 T MRI scanner (Philips Healthcare, Best, Netherlands) using a 32-channel SENSE head coil. To minimize motion artefacts due to head movement, we placed foam pads in both temple regions in the head coil. The three-dimensional high-resolution T1-weighted MP-RAGE (magnetization-prepared rapid gradient-echo) images of the brain were acquired with the following parameters: repetition time 6.9 ms, echo time 2.78 ms, field-of-view 263 mm × 281 mm × 150 mm, matrix size 256 mm × 256 mm, 137 sagittal slices, and voxel size 1.1 mm × 1.1 mm × 1.1 mm. FreeSurfer version 5.3.0 was used to process the acquired T1-weighted images using the longitudinal processing pipeline described in detail elsewhere [11, 13, 14]. First, all scans were processed cross-sectionally at each time point with the default automated processing pipeline that includes motion correction, intensity normalization, skull stripping, coregistration, segmentation, spherical registration and cortical/subcortical parcellations [15, 16]. Then, an unbiased within-subject template is generated for each participant by processing the scans from both the time points. Several following

processing steps, such as brain extraction, atlas registration, generation of cortical surface maps and parcellations are then initialized from the within-subject template to obtain longitudinally processed data. This significantly increased sensitivity, reliability, statistical power and ensures inverse consistency [14]. The Desikan–Killiany atlas [17] in FreeSurfer was used to extract regional measures of cortical thickness and volume [16, 18].

Statistical analysis

The statistical analysis was performed with SPSS version 21.0 for Windows. All clinical and demographic variables and cortical thickness values, including volume of cortical and subcortical structures showed a normal distribution using the Kolmogorov-Smirnov test. The analysis was performed with a two factors ANOVA for repeated measures: factor SESSION (2 levels: S1 scan vs. S2 scan) and factor SIDE (pain side vs. non-pain side), followed by a post-hoc correction for multiple comparison according to Bonferroni. Association between clinical and demographic variables and MRI data were performed by mean of specifically designed linear regression models. The experimental conditions before S1 and S2 scans were compared with a Student t-test for paired samples. Continuous variables are presented as “mean \pm standard deviation”, while categorical variables are presented as “absolute numbers with percentages”. Finally, considering that every ROI was tested twice (ANOVA and correlation analysis), to further control for multiple comparison we set the level of significance at $\alpha = 0.025$.

Results

Study population

Twenty-five patients were enrolled in the study and completed the S1 session (i.e. all 25 patients were scanned during on-going migraine headache). Of these, 10 patients were excluded for the following reasons: 5 patients were lost to follow-up and did not complete the S2 session; 4 patients had a bilateral migraine attack during the S1 scan; 1 patient was excluded because it was impossible to analyze the S1 MRI scan for the presence of movement artefacts.

A complete dataset was available for 15 patients (age 40.1 ± 10.3 years, 13 female, height 169.3 ± 5.9 cm, weight 86.3 ± 11.1 Kg, 12 right-handed). The study population had an average disease duration of 19.7 ± 10.1 years, and a migraine frequency of 3.3 ± 1.9 days per month. Four patients (26.7%) were on a stable preventive drug therapy during the overall study period (data are summarized in Table 1).

S1 and S2 scans were performed 30.2 ± 15.9 days apart. The number of hours of sleep before MRI scan ($p =$

Table 1 Clinical and demographic features of study population

	Study Population	
N	15	
Age (years)	40.1 ± 10.3	
Female sex	13 (86.7%)	
Height (cm)	169.3 ± 5.9	
Weight (kg)	86.3 ± 11.1	
Right handedness	12 (80.0%)	
Disease duration (years)	19.7 ± 10.1	
Number of migraine attack per month	3.3 ± 1.9	
Number of migraine days per month	6.6 ± 3.7	
Migraine prophylaxis	4 (26.7%)	
Usual side of migraine	Left	8 (53.3%)
	Right	3 (20.0%)
	Shifting	4 (26.7%)
Days between S1 and S2	30.2 ± 15.9	

Legend: S1: MRI scan performed during spontaneous migraine attack; S2: MRI scan performed during inter-ictal period

0.954) and the days elapsed since the last menses ($p = 0.145$) were comparable between S1 and S2 MRI sessions. In contrast, the days elapsed since last migraine attacks were higher before S1 scan ($p = 0.015$) (Table 2).

S1 scan was performed after 7.9 ± 3.7 h from migraine onset. The clinical phenotype of the migraine attack recorded during S1 are summarized in Table 2.

Structural changes during attack (S1 scan) compared to the inter-ictal period (S2 scan)

When compared to S2 (inter-ictal scan), during S1 (migraine attack scan) we found a significant reduction of thickness and volume of precentral cortex ($p = 0.023$ for thickness, and $p = 0.018$ for volume – Fig. 1) and pericalcarine cortex ($p = 0.024$ for thickness, and $p = 0.017$ for volume – Fig. 2). Also, the temporal pole cortex thickness was significantly lower at S1 when compared to S2 ($p = 0.017$). Regarding subcortical structures, we found an increase of hippocampus volume during S1 ($p = 0.007$).

Correlation of structural changes with pain

For all the results described above, we did not find a significant association with the side of migraine attack pain. Only for the temporal pole cortex thickness there was a significant interaction session and side ($p = 0.009$), but the post-hoc analysis was not statistically significant. Descriptively, it showed a trend toward a reduction of its thickness ipsilaterally to the migraine attack pain side ($p = 0.080$), and toward an increase of thickness contralaterally to the pain side ($p = 0.056$). Exploratively, a linear regression model analysis showed a positive correlation between the number of days elapsed since

Table 2 Comparison of experimental conditions between S1 and S2 scans, and clinical phenotype of migraine attack recorded at S1

	S1 Spontaneous migraine attack	S2 Inter-ictal period	p-value
N	15	15	–
Days elapsed since last migraine attack	20.1 ± 16.3	9.3 ± 6.8	0.015
Sleep hours before MRI scan	6.8 ± 2.0	6.9 ± 1.3	0.954
Days elapsed since last menses (for female patients, N = 13)	29.1 ± 27.3	42.3 ± 49.1	0.145
Side of migraine attack during S1	Left Right	8 (53.3%) 7 (46.7%)	
Hours from migraine onset to scan	7.9 ± 3.7		
NRS of migraine pain during S1	6.8 ± 2.1		
Throbbing pain	14 (93.3%)		
Nausea	15 (100.0%)		
Vomit	2 (13.3%)		
Aggravated by physical activity	13 (86.7%)		
Photophobia	11 (73.3%)		
Phonophobia	5 (33.3%)		

Legend: S1: MRI scan performed during spontaneous migraine attack; S2: MRI scan performed during inter-ictal period; NRS: nociceptive rating scale


the last migraine attack and the rostral anterior cingulate cortex thickness ipsilateral to the migraine pain side recorded during S1 ($R^2 = 0.727$). When corrected for age, sex, disease duration, and migraine frequency, the thickness of rostral anterior cingulate cortex showed an increase of 0.014 mm for every day elapsed since last migraine attack ($p = 0.014$) (Table 3).

Discussion

This is the first study specifically designed to evaluate whether cortical and sub-cortical brain structures show morphological changes during a spontaneous migraine attack when compared to the inter-ictal period. The main findings of our study are that thickness and volume of precentral and pericalcarine cortices, and thickness of the temporal pole cortex were reduced during a spontaneous migraine attack. At subcortical level, the hippocampus volume was increased during migraine attack. All these structures are involved in pain modulation and perception as part of the pain matrix [19, 20].

Cortical thickness and volume are measures of the size of brain cortex, which is progressively reduced with the age. However, cortical thickness can also be modified by various stimuli, including for instance transcranial magnetic stimulation (TMS) [21] and electroconvulsive therapy (ECT) [22]. A longitudinal MRI study of ECT in depressive patients reported increased cortical thickness immediately after ECT, but which was normalized to baseline after a 6 months follow-up period [22]. Thus, transient changes in cortical thickness can also occur. In the present study, cortical thickness was reduced during the attacks compared to inter-ictal phase in the patients. Due to the lack of a healthy control group it is not


known whether the cortical thickness in the inter-ictal phase was normal. However, previous studies comparing the inter-ictal phase in migraine patients with healthy controls reported reduced cortical volume of several areas, including the prefrontal cortex [23]. Moreover, there seems to be a relationship between the total load of attacks during life and decreased gray matter volume [24–29], suggesting that reduced cortical thickness may be a consequence of attacks. The thickness of the temporal pole [30] and pericalcarine cortex [31] is suggested to play an important role in migraine. Compared to healthy controls, Coppola and colleagues reported ictal increase and interictal reduction of gray matter density of the temporal pole in migraine patients [32]. However, this study was not longitudinal, but investigated two groups of patients. We cannot explain the opposite findings in this region. Despite the discrepancy, it is interesting that the change in the temporal pole was reported among several investigated areas. Coppola and colleagues defined the ictal phase as 12 h before or after beginning of an attack, while in our study none of the patients were scanned before onset of headache attack. The timing of the scans may be plausible explanation of the discrepancy. The mean time between the attack and inter-ictal phase scans was 30 days with the shortest interval of 12 days, but there was no correlation between cortical thickness or volume and the time between the scans. It is therefore highly likely that the changes in thickness was rapid onset and short-lasting, which can reflect short-term brain morphology but also changes in blood flow appearing as cortical thickness changes [33]. In either case, a reduced activity of these areas relative to other areas of the pain network would be expected.



The dorsolateral prefrontal cortex has an inhibitory role in pain pathways [34] and TMS of the prefrontal cortex is an effective therapy in chronic migraine [35]. In depressive patients TMS induced increase of cortical thickness, which was related to good effect of the therapy [21]. In addition to its efficacy in chronic migraine [36], TMS can also be used as acute treatment of the migraine attack. A randomized double-blind, parallel-group, sham-controlled trial of 164 patients reported significant higher responder rates after 2 h in the group treated with single-pulse TMS in the occipital region (visual cortex) compared with sham in the same region

[37]. Like the prefrontal cortex, the visual cortex may have an inhibitory role in pain processing, which is affected during migraine attacks. It is known that cortical thickness of the visual cortex is reduced inter-ictal [3, 4, 38] and here we report a further reduction during the attack.

The role of hippocampus and its morphological alterations were also evaluated in migraine patients [39]. It is noteworthy that hippocampal volume seemed to strictly relate to headache frequency, as it was greater in low-frequency migraine when compared to high-frequency migraine [40]. Moreover, hippocampal volume correlated



with the migraine frequency, showing a positive plastic adaption at low migraine frequency, and negative adaptation at higher headache frequency [41].

The biological significance of the structural changes observed in the present study is yet to be elucidated but could be a consequence of hypoperfusion rather than shrinking of the neurons and glial cells. In this view, it is worth noting that even if these brain areas showed a significant modulation during a spontaneous migraine attack in our cohort of patients, we were not able to find an association with the migraine pain side as well as significant correlations with other

clinical features of the disease. For these reasons, our results must be interpreted with caution, because they could represent a fast plastic adaptation to a non-specific stimulus, more than a specific reaction of the brain to the migraine attack. Reduced cortical thickness of the visual cortex and the prefrontal cortex may most likely reflect trigeminovascular and general pain respectively [42, 43]. In which order of the migraine cycle, cortical thickness change occurs is yet unknown, as we have no pre-ictal MRI scans in our patients. Repeated MRI scans in the individual patients may explain this in future.

Table 3 Multivariate linear regression model; dependent Variable: rostral anterior cingulate cortex thickness ipsilateral to the migraine pain side recorded during S1

	Non standardized coefficients		Standardized coefficients Beta	t	p-value	95% confidence interval fo B	
	B	Standard deviation error				Lower limit	Upper limit
Constant	2.885	0.440		6.562	0.000	1.890	3.879
Gender	-0.250	0.144	-0.328	-1.732	0.117	-0.577	0.077
Age	0.003	0.008	0.093	0.314	0.761	-0.015	0.020
Disease duration (years)	0.015	0.000	0.581	1.841	0.099	-0.004	0.034
Number of migraine attacks per month	-0.027	0.040	-0.187	-0.667	0.521	-0.117	0.064
Days elapsed since last migraine attack	0.014	0.005	0.820	3.034	0.014	0.004	0.024

R² of multivariate linear regression model: 0.727

Another interesting, but explorative finding was that the cortical thickness recorded during the spontaneous migraine attack of rostral anterior cingulate cortex ipsilateral to the migraine pain side, positively correlated with number of days elapsed since the previous migraine attack. This result was consistent after correction for age, sex, disease duration, and migraine frequency. The plastic adaptation of the cingulate cortex is not specific to migraine as it was reported in several pain and non-pain conditions [6]. In our cohort, the gradual increase in thickness of the cingulate cortex could represent a compensatory mechanism that starts and develops after a painful stimulus. On the other hand, the anterior cingulate cortex thickness was not significantly reduced during a spontaneous migraine attack, and for this reason this result must be interpreted with caution.

The low sample size of the present cohort limits further analyses in subgroups of patients (e.g. those with nausea versus without nausea etc.). Moreover, this study was designed to investigate the pain phase, why we waited at least 4 h from onset to scan. Thus, we cannot examine the early phase of the attack which is a limitation of this study.

Conclusions

We have presented the first study on morphological MRI changes during spontaneous migraine attacks compared to the inter-ictal phase. Acute attacks are accompanied by reduced cortical thickness and volume in pain and previously migraine-related areas. Although, we do not ascribe present findings as being migraine specific, these findings constitute a fingerprint of acute pain in migraine patients. In future studies, this fingerprint can be used as a biomarker in order to evaluate antimigraine treatment effect.

Abbreviations

ANOVA: Analysis of variance; ECT: Electroconvulsive therapy; ICHD: International Classification of Headache Disorders;

MP-RAGE: Magnetization-prepared rapid gradient-echo; MRI: Magnetic resonance imaging; TMS: Transcranial magnetic stimulation

Acknowledgements

Not applicable.

Authors' contributions

FMA: Design, patient recruitment, acquisition, data analysis and interpretation and drafting. RDI: Data analysis, interpretation, drafting and critical review of the manuscript. MAK: Data analysis, interpretation and critical review of the manuscript. JMR: Design, data analysis and critical review of the manuscript. FW: Design, data evaluation and critical review of the manuscript. HBWL: Design, acquisition, and critical review of the manuscript. MA: Design, interpretation, funding and critical review of the manuscript. All authors have approved the submitted version.

Funding

Authors received financial support for this study from the Lundbeck Foundation (R155-2014-171).

Availability of data and materials

The data that support the findings of this study are available, but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of data protection department of Rigshospitalet, Copenhagen, Denmark.

Declarations

Ethics approval and consent to participate

All participants gave their written consent after receiving detailed oral and written information and the study was conducted in accordance with the Helsinki II Declaration of 1964, with later revisions. The Ethical Committee of Copenhagen (H-3-2012-073) approved the study. All participants signed a written informed consent after a thorough explanation of study procedures. The study was registered at [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02202486) (NCT02202486). Other sub-studies under the same umbrella protocol have been published elsewhere [8, 44].

Consent for publication

Not applicable.

Competing interests

FMA has received lecturing fee and/or served as advisory board member for Teva, Lundbeck, Eli Lilly and Novartis. FMA is principal investigator for phase 4 trials for Novartis and Teva. RDI reported no conflicts of interests. MAK reports no conflicts of interests. JMR reports no conflicts of interests. FW reports no conflicts of interests.

HBWL reports no conflicts of interests.

MA reports receiving consulting fees and advisory boards fees from Alder, Allergan, Amgen, Eli Lilly, Lundbeck, Novartis, and Teva, fees for serving as a principal investigator, paid to his institution, from Alder, Allergan, Amgen, Electro-Core, Eli Lilly, Lundbeck, Novartis, and Teva, and grant support, paid to his institution, from Novo Nordisk Foundation, Novartis, and Lundbeck Foundation.

Author details

¹Danish Headache Center, Department of Neurology, Faculty of Health and Medical Sciences, Rigshospitalet Glostrup, University of Copenhagen, Valdemar Hansens Vej 5, 2600 Glostrup, Denmark. ²Headache Science & Neurorehabilitation Center, IRCCS Mondino Foundation, Pavia, Italy. ³Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy. ⁴Functional Imaging Unit, Department of Clinical Physiology, Nuclear Medicine and PET, Faculty of Health and Medical Sciences, Rigshospitalet, University of Copenhagen, Glostrup, Denmark. ⁵Centre for Neuropsychiatric Schizophrenia Research, CNSR and Centre for Clinical Intervention and Neuropsychiatric Schizophrenia Research, CINS, Mental Health Centre Glostrup, University of Copenhagen, 2600 Glostrup, Denmark. ⁶Department of Radiology, Herlev-Gentofte Hospital, University of Copenhagen, Herlev, Denmark.

Received: 29 April 2021 Accepted: 3 August 2021

Published online: 21 August 2021

References

- Hougaard A, Amin FM, Ashina M (2014) Migraine and structural abnormalities in the brain. *Curr Opin Neurol* 27(3):309–314. <https://doi.org/10.1097/WCO.0000000000000086>
- DaSilva AFM, Granziera G, Snyder J et al (2007) Thickening in the somatosensory cortex of patients with migraine. *Neurology*. 69(21):1990–1995. <https://doi.org/10.1212/01.wnl.0000291618.32247.2d>
- Granziera C, DaSilva AFM, Snyder J et al (2006) Anatomical alterations of the visual motion processing network in migraine with and without aura. *PLoS Med* 3(10):e402. <https://doi.org/10.1371/journal.pmed.0030402>
- Messina R, Rocca MA, Valsasina P (2013) Cortical abnormalities in patients with migraine: a surface-based analysis. *Radiology*. 268(1):170–180. <https://doi.org/10.1148/radiol.13122004>
- Hougaard A, Amin FM, Hoffmann MB, Larsson HBW, Magon S, Sprenger T, Ashina M (2015) Structural gray matter abnormalities in migraine relate to headache lateralization, but not aura. *Cephalalgia*. 35(1):3–9. <https://doi.org/10.1177/0333102414532378>
- Hougaard A, Amin FM, Arngriim N, Vlachou M, Larsen VA, Larsson HBW, Ashina M (2016) Sensory migraine aura is not associated with structural gray matter abnormalities. *NeuroImage Clin* 11:322–327. <https://doi.org/10.1016/j.nicl.2016.02.007>
- Datta R, Detre JA, Aguirre GK, Cucchiara B (2011) Absence of changes in cortical thickness in patients with migraine. *Cephalalgia*. 31(14):1452–1458. <https://doi.org/10.1177/0333102411421025>
- Amin FM, Hougaard A, Magon S, Sprenger T, Wolfram F, Rostrup E, Ashina M (2018) Altered thalamic connectivity during spontaneous attacks of migraine without aura: a resting-state fMRI study. *Cephalalgia*. 38(7):1237–1244. <https://doi.org/10.1177/0333102417729113>
- Schulte LH (2016) May a the migraine generator revisited: continuous scanning of the migraine cycle over 30 days and three spontaneous attacks. *Brain*. 139(7):1987–1993. <https://doi.org/10.1093/brain/aww097>
- Elvsåshagen T, Zak N, Norbom LM, Pedersen PØ, Quraishi SH, Bjørnerud A, Alnæs D, Doan NT, Malt UF, Groote IR, Westlye LT (2017) Evidence for cortical structural plasticity in humans after a day of waking and sleep deprivation. *NeuroImage*. 156:214–223. <https://doi.org/10.1016/j.neuroimage.2017.05.027>
- Reuter M, Rosas HD, Fischl B (2010) Highly accurate inverse consistent registration: a robust approach. *NeuroImage*. 53(4):1181–1196. <https://doi.org/10.1016/j.neuroimage.2010.07.020>
- Headache Classification Subcommittee of the International Headache Society (2004) The International Classification of Headache Disorders: 2nd edition. *Cephalalgia* 24(Suppl 1):9–160
- Reuter M, Fischl B (2011) Avoiding asymmetry-induced Bias in longitudinal image processing. *NeuroImage*. 57(1):19–21. <https://doi.org/10.1016/j.neuroimage.2011.02.076>
- Reuter M, Schmansky NJ, Rosas HD, Fischl B (2012) Within-subject template estimation for unbiased longitudinal image analysis. *NeuroImage*. 61(4):1402–1418. <https://doi.org/10.1016/j.neuroimage.2012.02.084>
- Collins DL, Neelin P, Peters TM, Evans AC (1994) Automatic 3D inter-subject registration of MR volumetric data in standardized talairach space. *J Comput Assist Tomogr* 18(2):192–205. <https://doi.org/10.1097/00004728-199403000-00005>
- Dale AM, Fischl B, Sereno MI (1999) Cortical surface-based analysis. I. Segmentation and surface reconstruction. *NeuroImage* 9(2):179–194. <https://doi.org/10.1006/nimg.1998.0395>
- Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM, Maguire RP, Hyman BT, Albert MS, Killiany RJ (2006) An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*. 31(3):968–980. <https://doi.org/10.1016/j.neuroimage.2006.01.021>
- Fischl B, Dale AM (2000) Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc Natl Acad Sci U S A* 97(20):11050–11055. <https://doi.org/10.1073/pnas.200033797>
- Davis KD, Moayed M (2013) Central mechanisms of pain revealed through functional and structural MRI. *J Neuroimmune Pharmacol* 8(3):518–534. <https://doi.org/10.1007/s11481-012-9386-8>
- Peyron R, Laurent B, García-Larrea L (2000) Functional imaging of brain responses to pain. A review and meta-analysis (2000). *Neurophysiol Clin* 30(5):263–288. [https://doi.org/10.1016/S0987-7053\(00\)00227-6](https://doi.org/10.1016/S0987-7053(00)00227-6)
- Boes AD, Uitermarkt BD, Albazron FM, Lan MJ, Liston C, Pascual-Leone A, Dubin MJ, Fox MD (2018) Rostral anterior cingulate cortex is a structural correlate of repetitive TMS treatment response in depression. *Brain Stimul* 11(3):575–581. <https://doi.org/10.1016/j.brs.2018.01.029>
- Gbyl K, Rostrup E, Raghava JM, Carlsen JF, Schmidt LS, Lindberg U, Ashraf A, Jørgensen MB, Larsson HBW, Rosenberg R, Videbech P (2019) Cortical thickness following electroconvulsive therapy in patients with depression: a longitudinal MRI study. *Acta Psychiatr Scand* 140(3):205–216. <https://doi.org/10.1111/acps.13068>
- Jia Z, Yu S (2017) Grey matter alterations in migraine: a systematic review and meta-analysis. *NeuroImage Clin* 14:130–140. <https://doi.org/10.1016/j.nicl.2017.01.019>
- Kim JH, Suh SI, Seol HY, Oh K, Seo WK, Yu SW, Park KW, Koh SB (2008) Regional grey matter changes in patients with migraine: a voxel-based morphometry study. *Cephalalgia*. 28(6):598–604. <https://doi.org/10.1111/j.1468-2982.2008.01550.x>
- Liu J, Lan L, Li G, Yan X, Nan J, Xiong S, Yin Q, von Deneen KM, Gong Q, Liang F, Qin W, Tian J (2013) Migraine-related gray matter and white matter changes at a 1-year follow-up evaluation. *J Pain* 14(12):1703–1708. <https://doi.org/10.1016/j.jpain.2013.08.013>
- Maleki N, Becerra L, Brawn J, Bigal M, Burstein R, Borsook D (2012) Concurrent functional and structural cortical alterations in migraine. *Cephalalgia*. 32(8):607–620. <https://doi.org/10.1177/0333102412445622>
- Rocca MA, Ceccarelli A, Falini A, Colombo B, Tortorella P, Bernardoni L, Comi G, Scotti G, Filippi M (2006) Brain gray matter changes in migraine patients with T2-visible lesions: a 3-T MRI study. *Stroke*. 37(7):1765–1770. <https://doi.org/10.1161/01.STR.0000226589.00599.4d>
- Schmidt-Wilcke T, Leinisch E, Straube A, Kampfe N, Draganski B, Diener HC et al (2005) Gray matter decrease in patients with chronic tension type headache. *Neurology*. 65(9):1483–1486. <https://doi.org/10.1212/01.wnl.0000183067.94400.80>
- Valfre W, Rainero I, Bergui M, Pinessi L (2008) Voxel-based morphometry reveals gray matter abnormalities in migraine. *Headache*. 48(1):109–117. <https://doi.org/10.1111/j.1526-4610.2007.00723.x>
- Schwedt TJ, Berisha V, Chong CD (2015) Temporal lobe cortical thickness correlations differentiate the migraine brain from the healthy brain. *PLoS One* 10(2):e0116687. <https://doi.org/10.1371/journal.pone.0116687>
- Schwedt TJ, Chong CD, Wu T, Gaw N, Fu Y, Li J (2015) Accurate classification of chronic migraine via brain magnetic resonance imaging. *Headache*. 55(6):762–777. <https://doi.org/10.1111/head.12584>
- Coppola G, Di Renzo A, Tinelli E, Iacovelli E, Lepre C, Di Lorenzo C et al (2015) Evidence for brain morphometric changes during the migraine cycle: a magnetic resonance-based morphometry study. *Cephalalgia*. 35(9):783–791. <https://doi.org/10.1177/0333102414559732>
- Franklin TR, Wang Z, Shin J, Jagannathan K, Suh JJ, Detre JA, O'Brien CP, Childress AR (2013) A VBM study demonstrating ‘apparent’ effects of a

- single dose of medication on T1-weighted MRIs. *Brain Struct Funct* 218(1): 97–104. <https://doi.org/10.1007/s00429-012-0385-6>
34. Lorenz J, Minoshima S, Casey KL (2003) Keeping pain out of mind: the role of the dorsolateral prefrontal cortex in pain modulation. *Brain*. 126(5):1079–1091. <https://doi.org/10.1093/brain/awg102>
 35. Brighina F, Piazza A, Vitello G, Aloisio A, Palermo A, Daniele O, Fierro B (2004) rTMS of the prefrontal cortex in the treatment of chronic migraine: a pilot study. *J Neurol Sci* 227(1):67–71. <https://doi.org/10.1016/j.jns.2004.08.008>
 36. Martelletti P, Jensen RH, Antal A, Arcioni R, Brighina F, De Tommaso M et al (2013) Neuromodulation of chronic headaches: position statement from the European headache federation. *J Headache Pain* 14(1):86. <https://doi.org/10.1186/1129-2377-14-86>
 37. Lipton RB, Dodick DW, Silberstein SD, Saper JR, Aurora SK, Pearlman SH, Fischell RE, Ruppel PL, Goadsby PJ (2010) Single-pulse transcranial magnetic stimulation for acute treatment of migraine with aura: a randomised, double-blind, parallel-group, sham-controlled trial. *Lancet Neurol* 9(4):373–380. [https://doi.org/10.1016/S1474-4422\(10\)70054-5](https://doi.org/10.1016/S1474-4422(10)70054-5)
 38. Magon S, May A, Stankewitz A, Goadsby PJ, Schankin C, Ashina M, Amin FM, Seifert CL, Mallar Chakravarty M, Müller J, Sprenger T (2019) Cortical abnormalities in episodic migraine: a multi-center 3T MRI study. *Cephalalgia*. 39(5):665–673. <https://doi.org/10.1177/0333102418795163>
 39. Liu HY, Chou KH, Chen WT (2018) Migraine and the Hippocampus. *Curr Pain Headache Rep* 22(2):13. <https://doi.org/10.1007/s11916-018-0668-6>
 40. Maleki N, Becerra L, Brawn J, McEwen B, Burstein R, Borsook D (2013) Common hippocampal structural and functional changes in migraine. *Brain Struct Funct* 218(4):903–912. <https://doi.org/10.1007/s00429-012-0437-y>
 41. Liu HY, Chou KH, Lee PL, Fuh JL, Niddam DM, Lai KL et al (2016) Hippocampus and amygdala volume in relation to migraine frequency and prognosis. *Cephalalgia*. 37:1329–1336
 42. Sava SL, de Pasqua V, Magis D, Schoenen J (2014) Effects of visual cortex activation on the nociceptive blink reflex in healthy subjects. *PLoS One* 9(6): e100198. <https://doi.org/10.1371/journal.pone.0100198>
 43. Sokal P, Harat M, Zieliński P, Furtak J, Paczkowski D, Rusinek M (2015) Motor cortex stimulation in patients with chronic central pain. *Adv Clin Exp Med* 24(2):289–296. <https://doi.org/10.17219/acem/40452>
 44. Amin FM, Hougaard A, Cramer SP, Christensen CE, Wolfram F, Larsson HBW, Ashina M (2017) Intact blood-brain barrier during spontaneous attacks of migraine without aura: a 3T DCE-MRI study. *Eur J Neurol* 24(9):1116–1124. <https://doi.org/10.1111/ene.13341>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

