

Gene expression profiling analysis of MENX-associated rat pituitary adenomas contributes to understand molecular mechanisms of human pituitary adenomas

HONGZHI ZHANG¹, CHUAN XU², NINGYANG SUN², YINTING ZHOU²,
XIAOFEI YU¹, XUE YAN² and QIUJUAN ZHANG²

¹Department of Neurology, Shuguang Hospital Affiliated to Shanghai University of Traditional Chinese Medicine, Shanghai 201203; ²Department of Neurology, Yueyang Hospital of Integrated Chinese and Western Medicine Affiliated to Shanghai University of Traditional Chinese Medicine, Shanghai 200437, P.R. China

Received October 17, 2014; Accepted July 10, 2015

DOI: 10.3892/ol.2015.3904

Abstract. The present study aimed to screen potential genes associated with pituitary adenomas to obtain further understanding with regard to the pathogenesis of pituitary adenomas. The microarray GSE23207 dataset, containing 16 pituitary adenoma samples from multiple endocrine neoplasia syndrome-associated rats and 5 normal pituitary tissue samples, was downloaded from Gene Expression Omnibus. The Linear Models for Microarray Data package was used to identify the differentially-expressed genes (DEGs) with the cut-off criteria of a \log_2 fold change (FC) >1 and adjusted P-values of <0.05 . The potential functions of the DEGs were predicted by functional and pathway enrichment analysis with the Database for Annotation, Visualization and Integrated Discovery. Furthermore, the interaction associations of the up- and downregulated DEGs obtained from the Search Tool for the Retrieval of Interacting Genes database were respectively revealed by the protein-protein interaction networks visualized with Cytoscape. A total of 391 upregulated and 238 downregulated DEGs in were screened in the pituitary adenoma samples. The upregulated DEGs with a higher degree in the protein-protein interaction network (e.g., *CCNA2*, *CCNB1* and *CDC20*) were significantly involved in cell cycle and cell division. Notably, *PTTG1* was enriched in every functional term. These DEGs interacted with each other. The downregulated DEGs (e.g., *GABRA1*, *GABRA4* and *GABRB1*) also interacted with each other, and were relevant to neuroactive ligand-receptor interaction; the DEG *POU1F1*,

interacting with *POMC*, was correlated with the development of the pituitary gland, adenohipophysia and endocrine system. Certain DEGs, including *CCNB1*, *CCNA2*, *CDC20*, *GABRA1*, *GABRA4*, *GABRB1*, *POU1F1* and *POMC*, and particularly *PTTG1*, were shown to be closely involved in the pathogenesis of pituitary adenomas.

Introduction

Pituitary adenomas, accounting for ~15% of all diagnosed intracranial tumors, are benign monoclonal adenomas that originate from cells of the anterior pituitary gland (1). Surgical resection, with or without adjuvant radiotherapy, is always the first line of treatment for the majority of pituitary adenomas, with the exception of prolactinomas (2). However, these treatments cannot usually control invasive pituitary adenomas due to the limited understanding of the underlying molecular mechanisms. Thereby, further research into the tumorigenesis will contribute to identifying novel therapeutic targets, which will be conducive to the development of novel therapeutic approaches for pituitary adenomas.

In past years, considerable progress has been made in identifying the key players in pituitary adenomas. A previous study has shown that the phosphoinositide 3-kinase/AKT signaling pathway is activated and enhanced in pituitary adenomas, which may be due to the mutation and amplification of an oncogene, *PIK3CA* (3). Mutation in another oncogene, *GNAS*, which encodes the guanine nucleotide-activating α subunit has also been suggested to be involved in pituitary hyperplasia (4). Meanwhile, a tumor suppressor aryl hydrocarbon receptor-interacting protein has been demonstrated to function in modulating cellular signaling and cAMP signaling pathways via regulation of the localization of the aryl hydrocarbon receptor (5). Also, the absence of expression of another two tumor suppressors, growth arrest and DNA-damage-inducible β (*GADD45 β*) and γ (*GADD45 γ*), has been observed in human pituitary adenomas (6,7). Aberrant methylation of a number of genes, such as *DAPK* (8) and *FGFR2* (9) has been confirmed to have a momentous role in pituitary tumorigenesis. Additionally, certain cell cycle regulators, such as p16,

Correspondence to: Dr Qiujuan Zhang, Department of Neurology, Yueyang Hospital of Integrated Chinese and Western Medicine Affiliated to Shanghai University of Traditional Chinese Medicine, 110 Ganhe Road, Shanghai 200437, P.R. China
E-mail: qiujuan_zhang@163.com

Key words: pituitary adenomas, differentially-expressed genes, pathway, protein-protein interaction

p21, p27, cyclin D1 and cyclin E, have also been demonstrated to function in pituitary tumorigenesis (8,10). Recently, certain microRNAs (miRNA/miR) have been found to be crucial in pituitary adenomas. For instance, the expression levels of miR-431 and miR-770-5p have been found to be slightly higher in non-functioning pituitary adenomas compared to their levels in the normal pituitary gland (11). Recently, another study has shown that miRNA-dependent impairment of the HMGA/E2F1 pathway functions as pro-oncogene signaling in pituitary adenomas. Several miRNAs targeting *HMGA2* (miR-326, miR-570 and miR-432) or *E2F1* (miR-326 and miR-603) could inhibit the growth of pituitary cell lines (HP75 and GH3) (12).

Lee *et al.* demonstrated that gonadotroph adenomas in MENX-affected rats closely resemble their human counterparts (13). The study further found that *CYP11A1* and *NUSAP1*, two commonly dysregulated differentially-expressed genes (DEGs) in the gonadotroph adenomas of rats and humans, are upregulated in 77 and 95% of human gonadotroph adenomas, respectively. Using the microarray data deposited by Lee *et al.*, the present study aimed to further identify genes that were differentially expressed between pituitary adenomas samples and normal controls. Following Gene Ontology (GO) functional and pathway enrichment analysis of the screened DEGs, Protein-Protein Interaction (PPI) networks were constructed for the up- and downregulated DEGs, respectively, in order to learn more about the interaction of proteins encoded by DEGs, which may aid in our understanding of the molecular mechanisms of pituitary adenomas. The results are expected to assist in elucidating the etiology of pituitary adenomas, and provide novel insights for the clinical diagnosis of this disease.

Materials and methods

Affymetrix microarray data. The gene expression profile data of GSE23207 (13) were acquired from Gene Expression Omnibus (GEO; <http://www.ncbi.nlm.nih.gov/geo/>), which was based on the platform of the GPL6247 [RaGene-1_0-st] Affymetrix Rat Gene 1.0 ST Array. This dataset contains 16 samples of pituitary homozygous mutants (p27Kip1/Cdkn1) from MENX-associated rats, aged 7-8 months, with large tumors 1-2 mm in size, and 5 samples of normal pituitary tissues purchased from BioChain Inc. (Hayward, CA, USA).

Data preprocessing and screening of DEGs. CEL files and probe annotation files were downloaded, and the gene expression data of all the samples were preprocessed via the Robust Multichip Averaging background correction (14), quantile normalization and probe summarization methods using the Oligo package (15). The Linear Models for Microarray Data package (16) of R was used for the identification of genes that were significantly differentially expressed in pituitary adenomas samples. The raw P-value was adjusted by the Benjamin and Hochberg method (17), and only the genes meeting the cut-off criteria of a \log_2 fold change (FC) of >1 and an adjusted P-value of <0.05 were selected as DEGs.

GO and pathway enrichment analysis. The Database for Annotation, Visualization and Integrated Discovery (DAVID) gene functional classification tool now provides a comprehensive

set of novel and powerful tools for researchers to understand the biological meaning behind abundant genes (18). Pathway enrichment analysis was conducted to identify the significant metabolic pathways for the DEGs (19). $P < 0.05$ and a count number of >2 were used as the cut-off criteria for GO and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analyses by DAVID.

PPI network construction. The Search Tool for the Retrieval of Interacting Genes database was used to analyze the PPIs for DEGs by calculating the combined scores (20), and a score >0.4 was chosen as the cut-off. Next, PPI networks for up- and downregulated DEGs were visualized using Cytoscape (<http://cytoscape.org/>) (21). The highly connected nodes (hub proteins) were detected by calculating the degree of each node protein based on the scale-free property of interaction networks (22).

Results

Identification of DEGs. Based on the cut-off criteria, a total of 629 DEGs were screened from the pituitary adenomas samples, including 391 upregulated and 238 downregulated DEGs.

Enrichment analysis of up- and downregulated DEGs. According to GO functional annotation, the upregulated DEGs were mainly enriched in GO terms associated with the cell cycle and cell division. For example, DEGs such as *CCNA2*, *NUSAP1*, *CCNB1*, *CENPF*, *CDC20* and *SPC25* were significantly involved in the cell cycle ($P=1.08 \times 10^{-12}$); *NUSAP1*, *CENPF*, *SPC25*, *CDC20* and *CCNB1* were involved in the M phase ($P=3.09 \times 10^{-11}$); and DEGs such as *CCNB1*, *SPC25*, *TOP2A*, *CDC20* and *CCNA2* were correlated with cell division ($P=1.25 \times 10^{-7}$). Notably, *PTTG1* was found to be enriched in every GO term (Table I). The downregulated DEGs, such as *KCND3*, *GABRA1*, *GABRA4* and *GABRB1*, were markedly associated with ion transport ($P=6.05 \times 10^{-7}$); DEGs such as *KCNJ5*, *KCND3*, *KCNJ6* and *KCNT2* were relevant to metal ion transport ($P=3.87 \times 10^{-6}$) and potassium ion transport ($P=7.01 \times 10^{-5}$); DEGs such as *DRD2*, *POU1F1* and *GHRHR* were distinctly associated with the positive regulation of multicellular organism growth ($P=2.47 \times 10^{-4}$), pituitary gland development ($P=5.70 \times 10^{-4}$), adeno-hypophysis development ($P=8.52 \times 10^{-4}$), diencephalon development ($P=1.24 \times 10^{-3}$) and endocrine system development ($P=6.09 \times 10^{-3}$); and DEGs such as *NOTCH2*, *ERBB4* and *POU1F1* were involved in cell fate commitment ($P=3.22 \times 10^{-2}$) and the regulation of cell proliferation ($P=4.55 \times 10^{-2}$) (Table II).

According to the KEGG pathway enrichment analysis, the upregulated DEGs were mainly enriched in 10 pathways. For example, *CDC20*, *CCNB1*, *CCNB2*, *BUB1*, *CDKN1A* and *MCM3* were enriched in the pathway of the cell cycle ($P=5.01 \times 10^{-7}$); *CDC20*, *CCNB1*, *CCNB2*, *BUB1* and *PLK1* were distinctly enriched in the pathway of oocyte meiosis ($P=1.058 \times 10^{-3}$); *CCNB1*, *CCNB2*, *CCNA2*, *BUB1* and *PLK1* were significantly enriched in the pathway of progesterone-mediated oocyte maturation ($P=1.150 \times 10^{-3}$); and *CDKN1A*, *CCNB1*, *CCNB2* and *CASP8* were markedly enriched in the p53 signaling pathway ($P=3.4841 \times 10^{-2}$) (Table III). Meanwhile, the downregulated

Table I. Top two enriched GO biological process term clusters with the highest enrichment score for the upregulated differentially-expressed genes.

Enrichment score	Term	Description	Count	P-value	Genes	
7.432	GO:0007049	Cell cycle	31	1.08x10 ⁻¹²	SPC25, CCNA2, CENPF, NUSAPI, CDC20, NDC80, CCNB1, PTTG1, CCNB2, BUB1B...	
	GO:0022403	Cell cycle phase	23	1.10x10 ⁻¹²	NUSAPI, CENPF, NDC80, CDC20, CCNB1, PTTG1, SPC25, CDKN1A, PLK1, BUB1B...	
	GO:0022402	Cell cycle process	27	3.26x10 ⁻¹²	PTTG1, SPC25, CDKN2C, CENPF, NUSAPI, CDC20, NDC80, CCNB1, CDKN1C, BUB1B...	
	GO:0000279	M phase	19	3.09x10 ⁻¹¹	NUSAPI, CENPF, NDC80, CDC20, PTTG1, CCNB1, SPC25, PLK1, BUB1B, SKA3...	
	GO:0051301	Cell division	14	1.25x10 ⁻⁷	NUSAPI, CDC20, PTTG1, CCNB1, SPC25, CCNB2, PLK1, BUB1B, TOP2A, CCNA2...	
	GO:0000087	M phase of mitotic cell cycle	12	2.16x10 ⁻⁷	CCNB1, SPC25, PLK1, NUF2, NUSAPI, BUB1B, SKA3, CENPF, CDC20, PTTG1...	
	GO:0000278	Mitotic cell cycle	16	5.40x10 ⁻⁷	NUSAPI, CENPF, NDC80, CDC20, PTTG1, CCNB1, SPC25, CDKN1A, CDKN2C, BUB1B...	
	GO:0000280	Nuclear division	10	1.09x10 ⁻⁵	CCNB1, SPC25, KIF11, PLK1, NUF2, NUSAPI, BUB1B, SKA3, CDC20, PTTG1	
	GO:0007067	Mitosis	10	1.09x10 ⁻⁵	CCNB1, SPC25, KIF11, PLK1, NUF2, NUSAPI, BUB1B, SKA3, CDC20, PTTG1	
	GO:0048285	Organelle fission	10	1.74x10 ⁻⁵	CCNB1, SPC25, KIF11, PLK1, NUF2, NUSAPI, BUB1B, SKA3, CDC20, PTTG1	
5.335	GO:0051439	Regulation of ubiquitin-protein ligase activity during mitotic cell cycle	4	4.86x10 ⁻²	CCNB1, PLK1, BUB1B, CDC20	
	GO:0000279	M phase	19	3.09x10 ⁻¹¹	NUSAPI, CENPF, NDC80, CDC20, PTTG1, CCNB1, SPC25, PLK1, BUB1B, SKA3...	
	GO:0051327	M phase of meiotic cell cycle	7	2.32x10 ⁻⁴	ADCY3, KIF2C, MKI67, PLK1, SGOL2, PTTG1, RAD51	
	GO:0007126	Meiosis	7	2.32x10 ⁻⁴	ADCY3, KIF2C, MKI67, PLK1, SGOL2, PTTG1, RAD51	
	GO:0051321	Meiotic cell cycle	7	2.76x10 ⁻⁴	ADCY3, KIF2C, MKI67, PLK1, SGOL2, PTTG1, RAD51	
	GO, gene ontology.					

Table II. Top two enriched GO biological process term clusters with the highest enrichment score for the downregulated differentially-expressed genes.

Enrichment score	Term	Description	Count	P-value	Genes	
4.727	GO:0006811	Ion transport	16	6.05x10 ⁻⁷	KCNJ3, GABRA1, GABRA4, GABRB2, GABRB1, CACNG6, ATP2B4, KCNT2, KCNH7, CACNA1A,.....	
	GO:0030001	Metal ion transport	12	3.87x10 ⁻⁶	KCNJ5, KCND3, KCNJ6, ATP2B4, CACNG6, KCNH7, KCNH8, SCNN1G, KCNJ3, CACNA1A,.....	
	GO:0006812	Cation transport	12	2.54x10 ⁻⁵	KCNJ5, KCND3, KCNJ6, ATP2B4, CACNG6, KCNH7, KCNH8, SCNN1G, KCNJ3, CACNA1A,.....	
	GO:0006813	Potassium ion transport	7	7.01x10 ⁻⁵	KCNJ5, KCND3, KCNJ6, KCNT2, KCNH7, KCNH8, KCNJ3	
	GO:0015672	Monovalent inorganic cation transport	8	5.52x10 ⁻⁴	KCNJ5, KCND3, KCNJ6, KCNT2, KCNH7, KCNH8, SCNN1G, KCNJ3	
	2.321	GO:0040018	Positive regulation of multicellular organism growth	4	2.47x10 ⁻⁴	GHI, DRD2, POU1F1, GHRHR
		GO:0021983	Pituitary gland development	4	5.70x10 ⁻⁴	DRD2, POU1F1, GHRHR, TBX19
		GO:0021984	Adenohypophysis development	3	8.52x10 ⁻⁴	DRD2, POU1F1, GHRHR
		GO:0021536	Diencephalon development	4	1.24x10 ⁻³	DRD2, POU1F1, GHRHR, TBX19
		GO:0043567	Regulation of insulin-like growth factor receptor signaling pathway	3	1.82x10 ⁻³	GHI, POU1F1, GHRHR
GO:0040014		Regulation of multicellular organism growth	4	4.82x10 ⁻³	GHI, DRD2, POU1F1, GHRHR	
GO:0035270		Endocrine system development	4	6.09x10 ⁻³	DRD2, POU1F1, GHRHR, TBX19	
GO:0030900		Forebrain development	5	9.58x10 ⁻³	ERBB4, DRD2, POU1F1, GHRHR, TBX19	
GO:0045927		Positive regulation of growth	4	9.75x10 ⁻³	GHI, DRD2, POU1F1, GHRHR	
GO:0048732		Gland development	5	1.75x10 ⁻²	ERBB4, DRD2, POU1F1, GHRHR, TBX19	
GO:0045165	Cell fate commitment	4	3.22x10 ⁻²	NOTCH2, ERBB4, POU1F1, TBX19		
GO:0051240	Positive regulation of multicellular organismal process	5	3.49x10 ⁻²	GHI, ERBB4, DRD2, POU1F1, GHRHR		
GO:0042127	Regulation of cell proliferation	8	4.55x10 ⁻²	NOTCH2, ERBB4, DRD2, NR3C2, CDK6, POU1F1, GHRHR, TBX19		

GO, gene ontology.

Table III. Results of pathway enrichment analysis of the up- and downregulated differentially-expressed genes.

Category	Term	Description	Count	P-value	Genes
Upregulated	rno04110	Cell cycle	14	5.01x10 ⁻⁷	CDC20, MCM3, CDKN1C, CCNBI, CDKN1A, CCNB2, CDKN2C, BUBI, BUBIB, CCNA2.....
	rno04114	Oocyte meiosis	9	1.06x10 ⁻³	CCNBI, ADCY3, AR, CCNB2, MAPK12, PLK1, BUBI, CDC20, PTTGI
	rno00601	Glycosphingolipid biosynthesis	5	1.10x10 ⁻³	B4GALTI, B3GALT2, B3GALT5, FUT4, B4GALT4
	rno04914	Progesterone-mediated oocyte maturation	8	1.15x10 ⁻³	CCNBI, ADCY3, CCNB2, KRAS, MAPK12, PLK1, BUBI, CCNA2
	rno04062	Chemokine signaling pathway	9	1.46x10 ⁻²	ADCY3, KRAS, LYN, ARRBI, PREX1, GRK5, PRKCD, CCL6, SHC4
	rno00330	Arginine and proline metabolism	5	1.70x10 ⁻²	ARG1, GOT1, NOS1, ASS1, GAMT
	rno05219	Bladder cancer	4	2.94x10 ⁻²	CDKN1A, KRAS, PGF, VEGFA
	rno04115	p53 signaling pathway	5	3.48x10 ⁻²	CCNBI, CDKN1A, CCNB2, CASP8, IGFBP3
	rno04610	Complement and coagulation cascades	5	4.19x10 ⁻²	CIQA, AZM, CIS, CIQC, F2R
	rno00510	N-glycan biosynthesis	4	4.89x10 ⁻²	B4GALTI, MAN2A1, ALG5, MAN1A1
Downregulated	rno04080	Neuroactive ligand-receptor interaction	9	4.35x10 ⁻⁴	GHI, GABRA1, GABRA4, GABRB2, DRD2, GABRB1, TSHB, LHB, GHRHR
	rno00410	β-alanine metabolism	3	1.05x10 ⁻²	ALDH2, ALDH1A7, DPYD
	rno00340	Histidine metabolism	3	1.25x10 ⁻²	MAOB, ALDH2, ALDH1A7
	rno00982	Drug metabolism	4	1.44x10 ⁻²	GSTA4, MAOB, MGST1, GSTM7
	rno00380	Tryptophan metabolism	3	3.75x10 ⁻²	MAOB, ALDH2, ALDH1A7
	rno04020	Calcium signaling pathway	5	3.99x10 ⁻²	ATP2B4, ERBB4, PLCG2, PLCB1, CACNA1A
	rno00480	Glutathione metabolism	3	4.93x10 ⁻²	GSTA4, MGST1, GSTM7

Table IV. Upregulated DEGs with connection degrees of >30 and the downregulated DEGs with connection degrees of at least 3 in the protein-protein interaction networks.

Category	Degree
Upregulated DEGs	
<i>CDK1</i>	51
<i>CCNB1</i>	47
<i>CCNA2</i>	46
<i>BUB1</i>	44
<i>ECT2</i>	43
<i>TPX2</i>	42
<i>NDC80</i>	42
<i>PRC1</i>	42
<i>NUSAP1</i>	41
<i>TOP2A</i>	41
<i>CCNB2</i>	41
<i>PBK</i>	41
<i>RACGAP1</i>	41
<i>TTK</i>	41
<i>PIK1</i>	40
<i>SPC25</i>	40
<i>BUB1B</i>	40
<i>CENPF</i>	40
<i>CDKN3</i>	39
<i>NUF2</i>	38
<i>CDC20</i>	38
<i>KIF11</i>	38
<i>DLGAP5</i>	38
<i>SGOL2</i>	37
<i>DTL</i>	36
<i>KIF20A</i>	36
<i>CDCA2</i>	36
<i>KIF20B</i>	35
<i>ESCO2</i>	35
<i>RAD51</i>	34
<i>ARHGAP11A</i>	34
<i>CKAP2</i>	33
<i>HMMR</i>	33
<i>KIF2C</i>	32
<i>DEPDC1</i>	32
Downregulated DEGs	
<i>POMC</i>	6
<i>GSTA4</i>	5
<i>GSTA1</i>	5
<i>POU1F1</i>	4
<i>ERBB4</i>	4
<i>GABRA1</i>	3
<i>MGST1</i>	3
<i>GABRA4</i>	3
<i>GSTM7</i>	3
<i>ALDH2</i>	3
<i>GHI</i>	3
<i>NR4A2</i>	3
<i>MAOB</i>	3

Table IV. Continued.

Category	Degree
<i>KCND3</i>	3
<i>NOTCH2</i>	3
<i>GABRB1</i>	3
DEGs, differentially-expressed genes.	

DEGs were mainly enriched in 7 pathways. *GHI*, *GABRA1*, *GABRA4* and *GABRB1* were enriched in the pathway of neuroactive ligand-receptor interaction ($P=4.35 \times 10^{-4}$); *MAOB*, *ALDH2* and *ALDH1A7* were mainly enriched in the pathways of histidine metabolism ($P=1.2476 \times 10^{-2}$) and tryptophan metabolism ($P=3.7487 \times 10^{-2}$); *ATP2B4*, *ERBB4* and *PLCG2* were enriched in the calcium signaling pathway ($P=3.9919 \times 10^{-2}$); and *GSTA4*, *MGST1*, *GSTM7* were significantly enriched in the pathways of drug metabolism ($P=1.4397 \times 10^{-2}$) and glutathione metabolism ($P=4.9294 \times 10^{-2}$) (Table III).

Analysis of PPI network. The PPI networks constructed with the up- and downregulated DEGs consisted of 1,044 and 69 PPI pairs, respectively. In the former, *PTTG1*, along with *CCNB1*, *CCNA2*, *SPC25*, *CENPF*, *NUSAP1*, *CDC20*, *TOP2A* and *BUB1*, were observed to interact with each other (Fig. 1). Within the PPI network built with downregulated DEGs, *GABRA1*, *GABRA4*, *GABRB1* and *GABRB1* were observed to interact with each other; *GSTA3*, *GSTA4*, *GSTM7* and *MGST1* were also found to interact with each other, and *POU1F1* was observed to interact with *POMC* (Fig. 2). The connection degrees of the top 15% highly-connected upregulated DEGs were each >30, and those of *CDK1*, *CCNB1*, *CCNA2* and *BUB1* were 51, 47, 46 and 44, respectively (Table IV). The top 20% highly-connected downregulated DEGs all had connection degrees of at least 3, and the degrees of *POMC*, *GSTA4*, *POU1F1*, *ERBB4*, *KCND3* and *NOTCH2* were 6, 5, 4, 4, 3 and 3, respectively (Table IV).

Discussion

In the present study, 391 DEGs were identified to be significantly upregulated and 238 were significantly downregulated in the pituitary adenomas samples. According to the constructed PPI network with the upregulated DEGs, *PTTG1* interacted with other DEGs with higher connection degrees, such as *CCNB1*, *CCNA2*, *SPC25*, *CENPF*, *NUSAP1*, *CDC20*, *TOP2A* and *BUB1*.

PTTG1, a tumorigenic gene *in vivo* (23), is abundantly expressed in pituitary tumors (24). As a securin protein, *PTTG1* is correlated with the mitotic checkpoint that prevents abnormal chromosome segregation (25), and peaks at the G₂/M phase (26). The overexpression of *PTTG1* results in cell transformation and induces aneuploidy (27), and this exists in >90% of pituitary tumors (28). *PTTG1*, together with *CCNB1*, *CCNA2*, *BUB1*, *SPC25*, *CENPF*, *NUSAP1*, *TOP2A* and *CDC20*, were all found to be enriched in GO terms associated with the cell cycle or cell division, which are indispensable for tumor growth. It




Figure 1. Protein-protein interaction network constructed with the upregulated differentially-expressed genes. Different shades of nodes colors represent the degree of up- or downregulation.

has been reported that *CDC20* is involved in the degradation of *PTTG1*-encoding products (29). Meanwhile, previous studies have also reported the abnormal expression of *CCNB1* (30), *CCNA2* (31), and *BUB1* (32) in pituitary adenomas. Furthermore, *CCNB1* was enriched in the p53 signaling pathway. *PTTG1*-encoding protein can cooperate with p53 to take part in cell apoptosis and DNA damage/repair (33,34). Altered p53 expression has been reported in pituitary carcinomas (35). Also, *PTTG1* can activate β -fibroblast growth factor, cyclin D3 and c-myc to promote cell proliferation (36,37). Therefore, *PTTG1* may play a crucial role in the occurrence of pituitary adenomas via interaction with *CCNB1*, *CCNA2*, *CENPF*, *NUSAP1*, *CDC20*, *TOP2A*, *BUB1* and p53.

Within the PPI network constructed with downregulated DEGs, *GABRA1*, *GABRA4* and *GABRB1* had higher degrees of connection to other genes. These genes were enriched in the pathway of neuroactive ligand-receptor interaction. *GABRA1*, *GABRA4* and *GABRB1* encode γ -aminobutyric acid (GABA) receptors. GABA is the major inhibitory neurotransmitter in the mammalian brain and may act as a paracrine or autocrine regulating factor in the human pituitary gland and human pituitary growth hormone adenomas (38). It has been reported that GABA has a specific effect on the electrical

activity of a tumoral line of rat pituitary cells, and that it inhibits prolactin secretion directly at the pituitary level (39). Additionally, *POU1F1* was also observed to have a higher connection degree in the PPI network. This gene encoding a member of the POU family of transcription factors (40), was correlated with the development of the pituitary gland, adenohypophysis and endocrine system. In humans, *POU1F1* mutation has been shown to be associated with combined pituitary hormone deficiency (41). *POU1F1* is also implicated in cell growth and prevents cell apoptosis (42). In the present study, it was observed to interact with *POMC*, which encodes a polypeptide hormone precursor. The encoded polypeptide hormone precursor is synthesized mainly in corticotrophin cells of the anterior pituitary (43). A previous study has shown that in silent pituitary adenomas, *POMC* mRNA has a diffuse low level or is absent (44). Thus, *GABRA1*, *GABRA4*, *GABRB1*, *POU1F1* and *POMC* may also have critical roles in pituitary adenoma occurrence via close interaction.

In conclusion, upregulated DEGs, such as those associated with the cell cycle or cell division (e.g., *CCNB1*, *CCNA2*, *BUB1*, *CENPF*, *NUSAP1*, *CDC20*, *TOP2A* and particularly *PTTG1*) and downregulated DEGs, such as those relevant to neuroactive ligand-receptor interaction (e.g., *GABRA1*, *GABRA4* and




Figure 2. Protein-protein interaction network constructed with the downregulated differentially-expressed genes. Different shades of nodes colors represent the degree of up- or downregulation.

GABRB1), as well as those correlated with the development of the pituitary gland, adenohypophysis and endocrine system (e.g., *POU1F1*) may have essential roles in the pathogenesis of pituitary adenomas. The present study provides novel information for the clinical diagnosis of this disease.

Acknowledgements

This study was supported by grants from the National Natural Science Foundation of China (no. 81273732/H2708) and the Shanghai Educational Commission Funding Project (no. 2012JW68).

References

- Chesnokova V and Melmed S: Pituitary tumour-transforming gene (PTTG) and pituitary senescence. *Horm Res* 71 (Suppl 2): 82-87, 2009.
- Pereira AM and Biermasz NR: Treatment of nonfunctioning pituitary adenomas: What were the contributions of the last 10 years? A critical view. *Ann Endocrinol (Paris)* 73: 111-116, 2012.
- Lin Y, Jiang X, Shen Y, Li M, Ma H, Xing M and Lu Y: Frequent mutations and amplifications of the *PIK3CA* gene in pituitary tumors. *Endocr Relat Cancer* 16: 301-310, 2009.
- Vortmeyer AO, Glasker S, Mehta GU, Abu-Asab MS, Smith JH, Zhuang Z, Collins MT and Oldfield EH: Somatic *GNAS* mutation causes widespread and diffuse pituitary disease in acromegalic patients with McCune-Albright syndrome. *J Clin Endocrinol Metab* 97: 2404-2413, 2012.
- Vandeva S, Tichomirowa MA, Zacharieva S, Daly AF and Beckers A: Genetic factors in the development of pituitary adenomas. *Endocr Dev* 17: 121-133, 2010.
- Mezzomo LC, Gonzales PH, Pesce FG, Kretzmann Filho N, Ferreira NP, Oliveira MC and Kohek MB: Expression of cell growth negative regulators *MEG3* and *GADD45γ* is lost in most sporadic human pituitary adenomas. *Pituitary* 15: 420-427, 2012.
- Michaelis KA, Knox AJ, Xu M, Kiseljak-Vassiliades K, Edwards MG, Geraci M, Kleinschmidt-DeMasters BK, Lillehei KO and Wierman ME: Identification of growth arrest and DNA-damage-inducible gene beta (*GADD45beta*) as a novel tumor suppressor in pituitary gonadotrope tumors. *Endocrinology* 152: 3603-3613, 2011.
- Farrell WE: Epigenetic mechanisms of tumorigenesis. *Horm Metab Res* 37: 361-368, 2005.
- Ezzat S: Epigenetic control in pituitary tumors. *Endocr J* 55: 951-957, 2008.
- Melmed S: Pathogenesis of pituitary tumors. *Nat Rev Endocrinol* 7: 257-266, 2011.
- Cheunschon P, Zhou Y, Zhang X, Lee H, Chen W, Nakayama Y, Rice KA, Tessa Hedley-Whyte E, Swearingen B and Klibanski A: Silencing of the imprinted *DLK1-MEG3* locus in human clinically nonfunctioning pituitary adenomas. *Am J Pathol* 179: 2120-2130, 2011.
- D'angelo D, Palmieri D, Mussnich P, Roche M, Wierinckx A, Raverot G, Fedele M, Croce CM, Trouillas J, Fusco A, *et al*: Altered microRNA expression profile in human pituitary GH adenomas: Down-regulation of miRNA targeting *HMGA1*, *HMGA2* and *E2F1*. *J Clin Endocrinol Metab* 97: 2011-3482, 2012.
- Lee M, Marinoni I, Irmeler M, Psaras T, Honegger JB, Beschoner R, Anastasov N, Beckers J, Theodoropoulou M, Roncaroli F, *et al*: Transcriptome analysis of *MENX*-associated rat pituitary adenomas identifies novel molecular mechanisms involved in the pathogenesis of human pituitary gonadotroph adenomas. *Acta Neuropathol* 126: 137-150, 2013.
- Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U and Speed TP: Exploration, normalization and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249-264, 2003.

15. Carvalho BS and Irizarry RA: A framework for oligonucleotide microarray preprocessing. *Bioinformatics* 26: 2363-2367, 2010.
16. Smyth GK: Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* 3, 2004.
17. Benjamini Y and Hochberg Y: Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J R Stat Soc Ser B* 57: 289-300, 1995.
18. Huang da W, Sherman BT and Lempicki RA: Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44-57, 2008.
19. Huang DW, Sherman BT and Lempicki RA: Bioinformatics enrichment tools: Paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 37: 1-13, 2009.
20. Von Mering C, Huynen M, Jaeggi D, Schmidt S, Bork P and Snel B: STRING: A database of predicted functional associations between proteins. *Nucleic Acids Res* 31: 258-261, 2003.
21. Kohl M, Wiese S and Warscheid B: Cytoscape: Software for visualization and analysis of biological networks. *Methods Mol Biol* 696: 291-303, 2013.
22. He X and Zhang J: Why do hubs tend to be essential in protein networks? *PLoS genetics* 2: e88, 2006.
23. Zhang X, Horwitz GA, Prezant TR, Valentini A, Nakashima M, Bronstein MD and Melmed S: Structure, expression and function of human pituitary tumor-transforming gene (PTTG). *Mol Endocrinol* 13: 156-166, 1999.
24. Chen L, Puri R, Lefkowitz EJ and Kakar SS: Identification of the human pituitary tumor transforming gene (hPTTG) family: Molecular structure, expression and chromosomal localization. *Gene* 248: 41-50, 2000.
25. Quereda V and Malumbres M: Cell cycle control of pituitary development and disease. *J Mol Endocrinol* 42: 75-86, 2009.
26. Yu R, Ren SG, Horwitz GA, Wang Z and Melmed S: Pituitary tumor transforming gene (PTTG) regulates placental JEG-3 cell division and survival: Evidence from live cell imaging. *Mol Endocrinol* 14: 1137-1146, 2000.
27. Yu R, Lu W, Chen J, McCabe CJ and Melmed S: Overexpressed pituitary tumor-transforming gene causes aneuploidy in live human cells. *Endocrinology* 144: 4991-4998, 2003.
28. Zhang X, Horwitz GA, Heaney AP, Nakashima M, Prezant TR, Bronstein MD and Melmed S: Pituitary tumor transforming gene (PTTG) expression in pituitary adenomas. *J Clin Endocr Metab* 84: 761-767, 1999.
29. Zur A and Brandeis M: Securin degradation is mediated by *fzy* and *fzr* and is required for complete chromatid separation but not for cytokinesis. *EMBO J* 20: 792-801, 2001.
30. Raverot G, Wierinckx A, Dantony E, Auger C, Chapas G, Villeneuve L, Brue T, Figarella-Branger D, Roy P, Jouanneau E, *et al*: Prognostic factors in prolactin pituitary tumors: Clinical, histological and molecular data from a series of 94 patients with a long postoperative follow-up. *J Clin Endocr Metab* 95: 1708-1716, 2010.
31. Leone V, Langella C, D'Angelo D, Mussnich P, Wierinckx A, Terracciano L, Raverot G, Lachuer J, Rotondi S, Jaffrain-Rea ML, *et al*: Mir-23b and miR-130b expression is downregulated in pituitary adenomas. *Mol Cell Endocrinol* 390: 1-7, 2014.
32. Yoshida D, Nomura R and Teramoto A: Signalling pathway mediated by CXCR7, an alternative chemokine receptor for stromal-cell derived factor-1 α , in AtT20 mouse adrenocorticotrophic hormone-secreting pituitary adenoma cells. *J Neuroendocrinol* 21: 481-488, 2009.
33. Yu R, Heaney AP, Lu W, Chen J and Melmed S: Pituitary tumor transforming gene causes aneuploidy and p53-dependent and p53-independent apoptosis. *J Biol Chem* 275: 36502-36505, 2000.
34. Hamid T and Kakar SS: PTTG/securin activates expression of p53 and modulates its function. *Mol Cancer* 3: 18, 2004.
35. Thapar K, Scheithauer BW, Kovacs K, Pernicone PJ, Laws ER Jr: p53 expression in pituitary adenomas and carcinomas: Correlation with invasiveness and tumor growth fractions. *Neurosurgery* 38: 765-770, 1996.
36. McCabe C, Khaira J, Boelaert K, Heaney AP, Tannahill LA, Hussain S, Mitchell R, Olliff J, Sheppard MC, Franklyn JA and Gittoes NJ: Expression of pituitary tumour transforming gene (PTTG) and fibroblast growth factor-2 (FGF-2) in human pituitary adenomas: Relationships to clinical tumour behaviour. *Clin Endocrinol (Oxf)* 58: 141-150, 2003.
37. Pei L: Identification of c-myc as a down-stream target for pituitary tumor-transforming gene. *J Biol Chem* 276: 8484-8491, 2001.
38. End K, Gamel-Didelon K, Jung H, Tolnay M, Lüdecke D, Gratzl M and Mayerhofer A: Receptors and sites of synthesis and storage of gamma-aminobutyric acid in human pituitary glands and in growth hormone adenomas. *Am J Clin Pathol* 124: 550-558, 2005.
39. Israel J, Dufy B, Gourdji D and Vincent J: Effects of GABA on electrical properties of cultured rat pituitary tumor cells: An intracellular recording study. *Life Sci* 29: 351-359, 1981.
40. Cohen LE, Wondisford FE and Radovick S: Role of Pit-1 in the gene expression of growth hormone, prolactin and thyrotropin. *Endocrinol Metab Clin North Am* 25: 523-540, 1996.
41. Quentien MH, Barlier A, Franc JL, Pellegrini I, Brue T and Enjalbert A: Pituitary transcription factors: From congenital deficiencies to gene therapy. *J Neuroendocrinol* 18: 633-642, 2006.
42. Pellegrini I, Roche C, Quentien MH, Ferrand M, Gunz G, Thirion S, Bagnis C, Enjalbert A and Franc JL: Involvement of the pituitary-specific transcription factor pit-1 in somatotrope cell growth and death: An approach using dominant-negative pit-1 mutants. *Mol Endocrinol* 20: 3212-3227, 2006.
43. Millington GW: Proopiomelanocortin (POMC): The cutaneous roles of its melanocortin products and receptors. *Clin Exp Dermatol* 31: 407-412, 2006.
44. Stefaneanu L, Kovacs K, Horvath E and Lloyd RV: In situ hybridization study of pro-opiomelanocortin (POMC) gene expression in human pituitary corticotrophs and their adenomas. *Virchows Arch A Pathol Anat Histopathol* 419: 107-113, 1991.