

RESEARCH ARTICLE

Open Access

Detailed kinetics and regulation of mammalian 2-oxoglutarate dehydrogenase

Feng Qi^{1,2}, Ranjan K Pradhan¹, Ranjan K Dash¹ and Daniel A Beard^{1*}

Abstract


Background: Mitochondrial 2-oxoglutarate (α -ketoglutarate) dehydrogenase complex (OGDHC), a key regulatory point of tricarboxylic acid (TCA) cycle, plays vital roles in multiple pathways of energy metabolism and biosynthesis. The catalytic mechanism and allosteric regulation of this large enzyme complex are not fully understood. Here computer simulation is used to test possible catalytic mechanisms and mechanisms of allosteric regulation of the enzyme by nucleotides (ATP, ADP), pH, and metal ion cofactors (Ca^{2+} and Mg^{2+}).

Results: A model was developed based on an ordered ter-ter enzyme kinetic mechanism combined with conformational changes that involve rotation of one lipoic acid between three catalytic sites inside the enzyme complex. The model was parameterized using a large number of kinetic data sets on the activity of OGDHC, and validated by comparison of model predictions to independent data.

Conclusions: The developed model suggests a hybrid rapid-equilibrium ping-pong random mechanism for the kinetics of OGDHC, consistent with previously reported mechanisms, and accurately describes the experimentally observed regulatory effects of cofactors on the OGDHC activity. This analysis provides a single consistent theoretical explanation for a number of apparently contradictory results on the roles of phosphorylation potential, NAD (H) oxidation-reduction state ratio, as well as the regulatory effects of metal ions on OGDHC function.

Background

The 2-oxoglutarate (α -ketoglutarate; α KG) dehydrogenase complex (OGDHC, EC 1.2.4.2, EC 2.3.1.61, and EC 1.6.4.3) is a multi-enzyme complex which catalyzes the chemical reaction:



OGDHC is primarily located within the mitochondrial matrix and is a key regulatory enzyme complex in the TCA cycle, responsible for oxidative decarboxylation of 2-oxoglutarate, transferring a succinyl group to coenzyme A (CoASH^{4-}) and producing reducing equivalents (NADH^{2-}) for the electron transport system. Regulation of OGDHC not only affects the distribution of 2-oxoglutarate between the TCA cycle and malate-aspartate shuttle system, but also has effects on the oxidative

deamination of glutamate. OGDHC is a crucial target of reactive oxygen species (ROS) and also able to generate ROS, which make it distinctly important for bioenergetics [1]. The molecular organization of OGDHC is similar to that of the pyruvate dehydrogenase complex (PDHC) as it belongs to the same heterogeneous family of 2-oxo acid dehydrogenase multi-enzyme complexes [2]. It consists of multiple copies of three enzyme components: oxoglutarate dehydrogenase (E1), dihydro-lipoamide succinyltransferase (E2), and dihydro-lipoamide dehydrogenase (E3). Consecutive actions of these enzymes catalyze the oxidation of 2-oxoglutarate and reduction of NAD^+ , which results in the production of NADH^{2-} and Succinyl-CoA^{4-} (Figure 1A). Allosteric interactions associated with the E1 component are known to be the predominant target for controlling of OGDHC activity [3].

OGDHC was first purified from the pig heart mitochondria by Sanadi *et al.* [4] and subsequently studied by many researchers to examine its catalytic and regulatory properties within permeabilized, un-coupled, and coupled mitochondria from a variety of mammalian

* Correspondence: beardda@gmail.com

¹Biotechnology and Bioengineering Center and Department of Physiology, Medical College of Wisconsin, Milwaukee, WI-53226, USA
Full list of author information is available at the end of the article




Figure 1 Schematic representation of the proposed mechanism of 2-oxoglutarate dehydrogenase complex (OGDHC). (A) It consists of three component enzymes: oxoglutarate dehydrogenase (E1), dihydro-lipoamide succinyltransferase (E2), and dihydro-lipoamide dehydrogenase (E3). The schematic representation here does not describe true stoichiometry of the multiple copies of three enzymes in the complex. The binding domain and lipoyl domain of E2 polypeptide are connected to the complex core with flexible links (dotted lines), are used here to describe the mechanism in which a single lipoyic acid rotates between the three catalytic sites. In the catalytic cycle, the disulfide at the tip of the lipoyl can be in oxidized, reduced or semi-reduced lipoate forms, the later one is connected with succinyl residue transferred from oxoglutarate. (B) This schematic illustrates the proposed kinetic schemes along with the mechanism of conformational changes. The forward reaction is read in the clockwise direction. The complex has three binding sites: one for each composite enzyme. E1 binds to 2-oxoglutarate or corresponding product CO₂ (top), E2 binds to CoA or corresponding product Succinyl-CoA (bottom-right), E3 binds to NAD or corresponding product NADH (bottom-left). It is assumed that, in the process of conformational changes, the rotation of one lipoyic acid between three catalytic sites leads to transfer succinyl from E1 to E2 and proton from E2 to E3.

tissues [5-11]. A catalytic mechanism for the overall reaction of the enzyme complex was also first proposed by Sanadi *et al.* [4] which suggested that the coenzyme, NAD⁻, and 2-oxoglutaric acid participate in the reaction with the help of the cofactors thiamine pyrophosphate (TPP), lipoic acid, and FAD²⁻ [12,13]. Their proposed mechanism is a Hexa-Uni-Ping-Pong mechanism in Cleland's terminology [14] where it is assumed that the first product (CO₂) is released before the second substrate (CoASH⁴⁻) binds, and the second product (Succinyl-CoA⁴⁻) is released before the third substrate (NAD⁻) binds to the enzyme. Subsequently, Koike *et al.* [15] postulated another mechanism in which, the lipoic acids transfer intermediates by rotating between the three catalytic sites. Furthermore, experimental results of fluorescence resonance energy transfer and dynamic anisotropy showed that the lipoic acids in the E2 component undergo motion where they rotate between different catalytic sites [16-18]. The results of steady-state kinetic studies done by Hamada *et al.* [19] and Smith *et al.* [10] contradict each other, and not all results are compatible with the Sanadi mechanism [4]. This issue was addressed by Mcminn and Ottaway [20] with kinetic studies based on the Fromm method [21]. Mcminn and Ottaway [20] explained the observed nonlinearity in the reciprocal plots of the results and proposed a phenomenological mechanism with semi-random characteristic. A recent study by Aevansson *et al.* [22] on the crystal structure and architecture of 2-oxo acid dehydrogenase multi-enzyme complexes, provides interesting insights into the plausible kinetic mechanism of 2-oxo acid dehydrogenase family which includes OGDHC.

It has been consistently shown that the activity of OGDHC is controlled by various factors, including the variations of the NAD oxidation-reduction state, the state of phosphorylation of the nucleotide systems, and the ratio of succinyl-CoA to CoA-SH. Regulation by reversible phosphorylation has not been demonstrated. Experiments in isolated mitochondria of heart, liver, and kidney have shown that the OGDHC is regulated by Ca²⁺ ions with a marked decrease in the apparent K_m for 2-oxoglutarate in the presence of adenine nucleotides and minimal effect of Ca²⁺ at saturating concentration of 2-oxoglutarate [9,23-25]. Moreover, the apparent K_m for 2-oxoglutarate is lowered by a decrease in the ATP/ADP ratio, which can significantly increase the sensitivity of the enzyme to Ca²⁺ ions [23,25]. It has been reported that the maximum activity of OGDHC is unaffected by changes in pH, while the apparent K_m of the enzyme for 2-oxoglutarate is greatly altered by changes in pH over the range of 6.5 - 7.5 [23]. A number of studies have also demonstrated the possible role of Mg²⁺ ions in the regulation of OGDHC either by directly affecting the activity of the enzyme or by modulating

the Ca²⁺ effect on the enzyme. Mg²⁺ ion has been shown to increase [26-28] or to have no effect on the activity of OGDHC [29,30]. McCormack and Denton [23] studied isolated OGDHC from pig heart mitochondria and found that there is no effect of EDTA and 1 mM Mg²⁺ on the activity of OGDHC when Ca²⁺ concentration was effectively less than 1 μM. Panov and Scarpa [9] concluded that the effects of Mg²⁺ and Ca²⁺ ions on the OGDHC activity are additive only at relative low concentration of free cations which suggested that at high concentrations, each ion may compete each other for binding sites. It is also evident that, in the presence of low Ca²⁺ concentration, Mg²⁺ ion can strongly modify the enzyme's affinities for 2-oxoglutarate and NAD⁻ [9]. However, the kinetic mechanisms by which these divalent metal ions regulate the properties of mitochondrial OGDHC are not understood.

Although a number of attempts have been made to understand the catalytic mechanisms of OGDHC, both experimentally and theoretically, there is no mechanistic model that consistently explains the available experimental data on the kinetics of this enzyme complex and adequately describes the regulatory roles of nucleotides and other metal ion cofactors (Ca²⁺, Mg²⁺, etc.). Therefore a mechanistic model of OGDHC is needed to understand the orchestrated controlling of OGDHC by cofactors inside mitochondria under different physiological conditions. In the present work, a kinetic model of OGDHC is introduced to quantitatively understand the catalytic properties and regulation of OGDHC, based on the observations from a large number of independent experimental studies in mammalian tissues. The model accurately describes the catalytic properties of this enzyme complex observed experimentally, and clarifies many contradictory results reported in earlier studies.

Methods

In this section, we first present a general kinetic model for conformational changes in OGDHC, based on a presumed ter-ter enzyme mechanism via substrate channeling. The model is then used to characterize the kinetics of the 2-oxoglutarate dehydrogenase reaction (Equation 1) and further extended to describe the regulatory roles of cofactors, i.e., nucleotides and various metal ions. The kinetic parameters of the model are estimated using a wide variety of experimental data, available in the literature.

Kinetic scheme for conformational changes in a ter-ter enzyme mechanism

The kinetic equation of the proposed model for OGDHC reaction is derived from a ter-ter enzyme mechanism combined with a model of conformational changes that represent the rotation of the single lipoic

acid between different catalytic sites [15,18]. The derivation is inspired by a previously developed model for trans-carboxylase [31]. The assumption of the model is that the enzyme complex is composed of three sub-enzyme (E1, E2, and E3), each with one binding site: site 1 binds to 2-oxoglutarate (αKG^{2-}) or corresponding product CO_2 , site 2 binds to CoASH^{4-} or corresponding product Succinyl-CoA $^{4-}$, site 3 binds to NAD^- or corresponding product NADH^{2-} (Figure 1A). Furthermore, the basic mechanism involves conformational changes, where the rotation of one lipoic acid between three catalytic sites leads to transfer of succinyl from E1 to E2 and proton from E2 to E3. In the catalytic cycle, the disulfide at the tip of the lipoyl can be in oxidized, reduced or semi-reduced lipoate forms, the semi-reduced form is bound with succinyl residue transferred from 2-oxoglutarate (Figure 1B).

Each of the six conformational states shown in Figure 1B can involve any possible binding states associated with the enzyme. For example, the first site is either empty or bound to 2-oxoglutarate or CO_2 ; the second site is either empty or bound to CoASH^{4-} or Succinyl-CoA $^{4-}$; and the third site is either empty or bound to NAD^- or NADH^{2-} . Therefore, there are a total of 27 binding states for each one of the six conformational states, which gives rise to $27 \times 6 = 162$ distinct states in the model. Here we denote these 162 states as E_{xyz}^i , where $i \in \{1, 2, 3, 4, 5, 6\}$ represents the index for conformational states, and $x \in \{\emptyset, A, P\}$, $y \in \{\emptyset, B, Q\}$ and $z \in \{\emptyset, C, R\}$ represent the binding states of site 1, site 2, and site 3. The lower-case e_{xyz}^i is used to represent the fraction of each state. Therefore, the total fractional states can be expressed as

$$e_{total}^i = \sum_{x \in \{\emptyset, A, P\}, y \in \{\emptyset, B, Q\}, z \in \{\emptyset, C, R\}} e_{xyz}^i \quad (2)$$

We assume rapid equilibrium binding for all 27 binding states, implying that the binding processes are much faster than the conformational change processes. With this assumption, Equation (2) can be written as:

$$e_{total}^i = e_{free}^i \times \left(1 + \frac{[A]}{K_A} + \frac{[P]}{K_P}\right) \times \left(1 + \frac{[B]}{K_B} + \frac{[Q]}{K_Q}\right) \times \left(1 + \frac{[C]}{K_C} + \frac{[R]}{K_R}\right), \quad (3)$$

where $e_{free}^i \equiv e_{\emptyset\emptyset\emptyset}^i$ denotes the fraction of free enzyme complex that binds to the reactants; K_A , K_B , K_C , K_P , K_Q and K_R are the dissociation constants associated with the binding of reactants (A: αKG^{2-} , B: CoASH^{4-} and C: NAD^-) and products (P: CO_2 , Q: Succinyl-CoA $^{4-}$ and R: NADH^{2-}) to the enzyme complex. Here, we assumed that these constants do not depend on conformational

states of the enzyme complex and the bound reactant at one site does not influence the binding reaction at another site: all binding interactions are independent of one other. These assumptions are necessary to make the model tractable, and are validated by comparing the model predictions to the available experimental data.

We define f_i as the fractions in conformation state i ($i \in \{1, 3, 5\}$) which can undergo forward conformational transformation to state $i+1$ (lipoate changes in three different redox forms). Similarly, f_{-i} is the fraction in conformation state $i+1$ which can undergo conformational transformation in the reverse direction (Figure 1B). Specifically, the binding of 2-oxoglutarate at site 1 is necessary for transition from conformation state 1 to 2. Therefore, we can write

$$\begin{aligned} f_1 &= \left(\frac{[A]/K_A}{1 + [A]/K_A + [P]/K_P}\right), \\ f_{-1} &= \left(\frac{[P]/K_P}{1 + [A]/K_A + [P]/K_P}\right), \\ f_3 &= \left(\frac{[B]/K_B}{1 + [B]/K_B + [Q]/K_Q}\right), \\ f_{-3} &= \left(\frac{[Q]/K_Q}{1 + [B]/K_B + [Q]/K_Q}\right), \\ f_5 &= \left(\frac{[C]/K_C}{1 + [C]/K_C + [R]/K_R}\right), \\ f_{-5} &= \left(\frac{[R]/K_R}{1 + [C]/K_C + [R]/K_R}\right). \end{aligned} \quad (4)$$

The net turn-over (reaction velocity) for this mechanism can be expressed

$$v = \frac{V}{[E_{total}]} = f_5 k_5 [e_{total}^5] - f_{-5} k_{-5} [e_{total}^6]. \quad (5)$$

Applying the King and Altman method to the scheme shown in Figure 1B gives the following expression for the net reaction velocity:

$$v = \frac{f_1 f_3 f_5 k_1 k_2 k_3 k_4 k_5 k_6 - f_{-1} f_{-3} f_{-5} k_{-1} k_{-2} k_{-3} k_{-4} k_{-5} k_{-6}}{\left(\begin{aligned} & f_{-1} f_{-3} k_{-1} k_{-2} k_{-3} k_{-4} (k_6 + k_{-6}) + f_{-1} f_5 k_{-1} k_{-2} k_4 k_5 (k_6 + k_{-6}) \\ & + f_3 f_5 k_2 k_3 k_4 k_5 (k_6 + k_{-6}) + f_{-3} f_{-5} (k_2 + k_{-2}) k_{-3} k_{-4} k_{-5} k_{-6} \\ & + f_{-3} f_1 (k_2 + k_{-2}) k_{-3} k_{-4} k_1 k_6 + f_1 f_5 (k_2 + k_{-2}) k_1 k_4 k_5 k_6 \\ & + f_1 f_3 (k_4 + k_{-4}) k_1 k_2 k_3 k_6 + f_{-1} f_{-5} (k_4 + k_{-4}) k_{-5} k_{-6} k_{-1} k_{-2} \\ & + f_3 f_{-5} (k_4 + k_{-4}) k_{-5} k_{-6} k_2 k_3 + f_{-1} f_{-3} f_5 k_{-1} k_{-3} k_5 k_{-2} (k_6 + k_{-6}) \\ & + f_{-1} f_3 f_5 k_{-1} k_3 k_4 k_5 (k_6 + k_{-6}) + f_{-3} f_{-5} f_1 k_{-3} k_{-5} k_1 k_{-4} (k_2 + k_{-2}) \\ & + f_{-3} f_1 f_3 k_{-3} k_1 k_5 k_6 (k_2 + k_{-2}) + f_{-1} f_{-5} f_3 k_{-1} k_{-5} k_3 k_{-6} (k_{-4} + k_4) \\ & + f_{-5} f_1 f_3 k_{-5} k_1 k_3 k_2 (k_4 + k_{-4}) + f_1 f_3 f_5 k_1 k_3 k_5 (k_4 k_2 + k_6 k_4 + k_2 k_6) \\ & + f_{-1} f_{-3} f_{-5} k_{-1} k_{-3} k_{-5} (k_{-2} k_{-4} + k_{-4} k_{-6} + k_{-6} k_{-2}) \end{aligned} \right)} \quad (6)$$

This complex expression can also be obtained using our KAPattern package [32], available freely for the derivation of enzyme rate equations. Substituting the fractional occupancy distributions as defined in Equation (4), we obtain an expression for the reaction velocity in terms of the individual rate constants and dissociation

constants. The kinetic constants can be also expressed in terms of various rate constants. Using the Haldane relationship, the velocity equation can be written as:

$$v = \frac{V_f V_r ([A][B][C] - [P][Q][R]/K_{eq})}{\left(\begin{aligned} &V_r [A][B][C] + K_{mC} V_r [A][B][R]/K_{ir} \\ &+ K_{mC} V_r [A][B] + K_{mB} V_r [A][C][Q]/K_{iq} \\ &+ K_{mB} V_r [A][C] + K_{mP} V_f [A][Q][R]/K_{ia}/K_{eq} \\ &+ K_{ic} K_{mB} V_r [A][Q]/K_{iq} + K_{mA} V_r [B][C][P]/K_{ip} \\ &+ K_{mA} V_r [B][C] + K_{ia} K_{mC} V_r [B][P][R]/K_{ip}/K_{ir} \\ &+ K_{ia} K_{mC} V_r [B][R]/K_{ir} + K_{ib} K_{mA} V_r [C][P][Q]/K_{iq}/K_{ip} \\ &+ K_{ib} K_{mA} V_r [C][P]/K_{ip} + V_f [P][Q][R]/K_{eq} \\ &+ K_{mR} V_f [P][Q]/K_{eq} + K_{mQ} V_f [P][R]/K_{eq} + K_{mP} V_f [Q][R]/K_{eq} \end{aligned} \right)} \quad (7)$$

where the kinetic constants are defined as:

$$\begin{aligned} V_f &= num_1/Coef_{ABC}, V_r = num_2/Coef_{PQR}, K_{mA} = Coef_{BC}/Coef_{ABC}, K_{mB} = Coef_{AC}/Coef_{ABC}, \\ K_{mC} &= Coef_{AB}/Coef_{ABC}, K_{mP} = Coef_{QR}/Coef_{PQR}, K_{mQ} = Coef_{PR}/Coef_{PQR}, K_{mR} = Coef_{PQ}/Coef_{PQR}, \\ K_{ip} &= Coef_{AB}/Coef_{BPR} = Coef_{BC}/Coef_{BCP}, K_{iq} = Coef_{CP}/Coef_{CPQ} = Coef_{AC}/Coef_{ACQ}, \\ K_{ir} &= Coef_{AC}/Coef_{ACR} = Coef_{AB}/Coef_{ABR}, K_{ia} = Coef_{QR}/Coef_{AQR} = Coef_{BR}/Coef_{ABR}, \\ K_{ib} &= Coef_{PR}/Coef_{BPR} = Coef_{CP}/Coef_{BCP}, K_{ie} = Coef_{PQ}/Coef_{CPQ} = Coef_{AC}/Coef_{ACQ} \text{ and} \\ K_{eq} &= num_1/num_2. \end{aligned}$$

Here we used the shorthand notation similar to that of Segel [33] where $num = k_1 k_2 k_3 k_4 k_5 k_6 K_P K_Q K_R$, $num = k_1 k_2 k_3 k_4 k_5 k_6 K_A K_B K_C$, and $Coef_{AB} = K_C K_P K_Q K_R (k_1 k_2 k_3 k_6 (k_4 + k_{-4}))$, and so on. V_f and V_r have units of mass per unit time per mass of protein. Other kinetic parameters associated with the binding of reactants and products have the units of concentration (mass per unit volume).

In the ter-ter biochemical reaction, the fourteen unknown kinetic parameters in Equation (7) are related to the equilibrium constant K_{eq} (known) via the following equilibrium relationship:


$$\begin{aligned} K_{eq} &= \left(\frac{[P][Q][R]}{[A][B][C]} \right)_{eq} = \frac{num_1}{num_2} \\ &= \frac{K_{ip} K_{iq} K_{ir}}{K_{ia} K_{ib} K_{ic}} = \frac{V_f K_{mP} K_{iq} K_{ir}}{V_r K_{ia} K_{mB} K_{ic}} \\ &= \frac{V_f K_{ip} K_{mQ} K_{ir}}{V_r K_{ia} K_{ib} K_{mC}} = \frac{V_f K_{ip} K_{iq} K_{mR}}{V_r K_{mA} K_{ib} K_{ic}}, \end{aligned} \quad (8)$$

reducing the total number of independent unknown kinetic parameter to thirteen.

Kinetic model of OGDHC using a ter-ter enzyme mechanism

We apply the above general form of the ter-ter enzyme mechanism for the analysis of available experimental data on the kinetic of OGDHC to estimate the unknown kinetic parameters and to elucidate whether the proposed mechanism is able to explain the available kinetic data.

In the TCA cycle, OGDHC is primarily involved in the fifth step for oxidation of acetyl-CoA. The reference chemical reaction is given by Equation (1). The corresponding biochemical reaction is given by



Here biochemical reactants, such as αKG , correspond to ensemble chemical species, such as αKG^{2-} , $H \alpha KG^-$, etc. The chemical reaction in Equation (1) is unambiguously balanced in terms of mass and charge, whereas this biochemical reaction is not. In this reaction, the reactant $CO_{2,tot}$ represents the sum of aqueous carbon dioxide and bicarbonate species (CO_3^{2-} , HCO_3^- and H_2CO_3).

The equilibrium constant for the reference reaction can be written as:

$$\begin{aligned} K_{eq,ogdhc}^0 &= \left(\frac{[\text{Succinyl} - CoA^{4-}][CO_3^{2-}][NADH^{2-}][H^+]}{[\alpha KG^{2-}][CoASH^{4-}][NAD^-]} \right)_{eq} \\ &= \exp \left(-\frac{\Delta_r G_{ogdhc}^0}{RT} \right), \end{aligned} \quad (10)$$

where $\Delta_r G_{ogdhc}^0$ is the standard Gibbs free energy of the reference reaction which is computed using the basic thermodynamic data (298.15 K, I = 0.15 M) listed in Li et al. [34].

For the development of the kinetic model of OGDHC, we assume that the ter-ter enzyme mechanism proposed in the previous section along with the conformational changes (Figure 1B) can explain the observed kinetics of OGDHC. Because the kinetic data we used here to estimate the unknown kinetic parameters were all from the initial velocity studies in which only the products $NADH^{2-}$ and Succinyl-CoA^- were present in the reaction mediums, the forward flux of OGDHC can be reduced from Equation (7) as:

$$J_{ogdhc}^+ = \frac{V_f [A][B][C]}{\left(\begin{aligned} &[A][B][C] + K_{mC} [A][B][R]/K_{ir} \\ &+ K_{mC} [A][B] + K_{mB} [A][C][Q]/K_{iq} \\ &+ K_{mB} [A][C] + K_{ic} K_{mB} [A][Q]/K_{iq} \\ &+ K_{mB} K_{ic} [A][Q][R]/K_{iq}/K_{ir} \\ &+ K_{mA} [B][C] + K_{ia} K_{mC} [B][R]/K_{ir} \\ &+ K_{mB} K_{ic} K_{ia} [Q][R]/K_{iq}/K_{ir} \end{aligned} \right)}, \quad (11)$$

where [A], [B], [C], [R] and [Q] represent the concentrations of αKG^{2-} , $CoASH^{4-}$, NAD^- , $NADH^{2-}$, and Succinyl-CoA^{4-} , respectively. This kinetic expression for OGDHC reaction contains 8 unknown kinetic parameters. Thus, this expression was used first to estimate the 8 unknown kinetic parameters. Using the relationship $J_{ogdhc}^+ / J_{ogdhc}^- = \exp(\Delta G/RT)$ to determine the reverse flux [35], we obtain the full flux expression for OGDHC as:

$$J_{ogdhc} = J_{ogdhc}^+ - J_{ogdhc}^-$$

$$= J_{ogdhc}^+ \left(1 - \frac{1}{K_{eq,ogdhc}^0} \frac{([\text{Succinyl-CoA}^{4-}][\text{CO}_3^{2-}])}{([\text{NADH}^{2-}][\text{H}^+])} \right) \left(\frac{[\alpha\text{KG}^{2-}][\text{CoASH}^{4-}][\text{NAD}^-]}{[\text{NADH}^{2-}][\text{H}^+]} \right) \quad (12)$$

The kinetic expressions for the regulatory effects of various cofactors on OGDHC (which depend on 10 additional kinetic parameters) are parameterized in the Results section, and are estimated separately (see below). Because CO_2, tot dependent terms are not included in the denominator in Equation (11), CO_2, tot dependent product inhibition is not accounted for in Equation (12).

Parameter estimation

The developed kinetic model of OGDHC has 8 adjustable parameters for catalytic mechanism and 10 adjustable parameters for cofactor regulation (Table 1). Parameter values were estimated in a systematic modular manner in multiple steps by least-squares fitting of the model simulated steady-state flux to the available experimental data as detailed in Results section below. The FMINCON algorithm in MATLAB (The MathWorks, Natick, MA) was used to solve this non-linear optimization problem. In addition, sensitivity analysis was performed to estimate the sensitivity of the least square error to small changes in the optimal parameter values. The sensitivity was computed using:

$$S_i = \frac{\max(|E_i^*(x_i \pm 0.1x_i) - E_i^*(x_i)|)}{0.1E_i^*(x_i)}, \quad (13)$$

where E^* is the least square difference between model simulations and experimental data, and x_i is the optimized value of the i^{th} parameter.

Since all kinetic parameters in the model are measured relative to species concentration, we performed a

composition analysis to estimate the concentration of all ionic species all experiments analyzed here [36].

Results

Parameterization of basic kinetic mechanism of OGDHC


In this section, we present the detailed parameterization and validation of the proposed kinetic model based on the available experimental data on the kinetics of OGDHC, measured in a wide variety of experimental conditions. To study the catalytic mechanism of OGDHC, McMinn and Ottaway [20] investigated the kinetic properties of the OGDHC system, which was prepared from fresh pig heart mitochondria. Following the method of Fromm [21] in which it was concluded from initial velocity studies that the catalytic mechanism of OGDHC is not consistent with the Hexa-Uni-Ping-Pong mechanism. While their observations suggest a random order kinetic mechanism with respect to the binding of NAD^- and CoASH^{4-} and release of Succinyl- CoA^{4-} , the binding of 2-oxoglutarate and release of CO_2 is described as a Ping-Pong mechanism. Initial velocity kinetics measured by Smith *et al.* [10] with purified pig heart mitochondria OGDHC showed that Succinyl- CoA^{4-} and NADH^{2-} were inhibitors, but no inhibitory effects were observed with GTP or ATP. Their results also show that Succinyl- CoA^{4-} inhibition was competitive with CoASH^{4-} and independent of the NAD^- oxidation-reduction state. These data are used here to identify the kinetic parameters of our OGDHC model.

The experimental data in Figure 2 were used to estimate the values of unknown kinetic parameters that govern the basic catalytic mechanism of OGDHC (Equation (12)) based on the best fits of the model to the data (See Figure caption for details). Measured enzyme activity is expressed in $\mu\text{moles NADH}^{2-}$ formed/mg protein/min. We follow a systematic optimization procedure to estimate each kinetic parameter of the model using appropriate experimental

Table 1 Kinetic parameter values for 2-oxoglutarate dehydrogenase complex

Basic kinetic parameters			Kinetic parameters for regulatory cofactors		
Parameter:	Value	Sensitivity	Parameter:	Value	Sensitivity
V_{\max} $\mu\text{mol mg}^{-1} \text{min}^{-1a}$	-	-	K_{GCa} (μM)	0.893	0.293
K_{mA} (mM)	0.273	3.601	K_{IATP} (mM)	0.106	0.943
K_{mB} (μM)	6.96	0.934	K_{IADP} (mM)	0.305	2.073
K_{mC} (μM)	98.6	2.839	K_{GMg} (μM)	19.49	0.228
K_{iCa} (mM)	75.9	1.981	α_{Ca}^0	0.262	9.648
K_{iF} (mM)	2.4	1.995	α_{ATP}	6.694	1.628
K_{iC} (mM)	0.112	3.970	α_{ADP}	0.173	4.412
K_{iQ} (μM)	0.218	5.224	α_{Mg}	1.00	6.594
K_{GdH} (pH)	6.11	3.623	β_{Mg}	4.222	4.603

^a We cannot estimate the value of parameter $V_{F, \max}$ in the model without knowing the enzyme activity in the experimental setup. The value of V_{\max} reported in the table is equal $X \times V_{F, \max}$ where X is the enzyme activity. The optimization estimates that $V_{\max} \approx 2.16 \mu\text{mol mg}^{-1} \text{min}^{-1}$ for Figure 2, $V_{\max} \approx 11.033 \text{ unit/mg}$ for Figure 3 and Figure 4, $V_{\max} \approx 167 \text{ nmol mg}^{-1} \text{min}^{-1}$ for Figure 5.




data. In the first step, Figure one of McMinn and Ottaway [20] is used to determine the parameters K_{mA} , K_{mB} and K_{mC} . Because these data describe the changes in the initial rate of OGDHC activities in response to variations in the concentration of different substrates in the absence of products, they facilitate identifying the kinetic parameter associated with the binding of 2-oxoglutarate, CoASH^{4-} , and NAD^- . Next, the kinetic data from Figure three of McMinn and Ottaway [20] are used to estimate K_{ia} and K_{ir} . Finally, the kinetic results from Figure five of Smith *et al.* [10] are used to determine the last two parameters related to Succinyl-CoA⁴⁻ inhibition: K_{ic} and K_{iq} .

Henceforth, these kinetic parameters are fixed at their estimated values (Table 1).

Parameterization of the cofactor-dependent regulatory mechanisms

Denton *et al.* [23] conducted a number of experiments to study the effects of Ca^{2+} , pH, and adenine nucleotides on the activity of OGDHC from pig heart mitochondria. Their data are used here to identify the kinetic parameters that characterize the activation/inhibition mechanism of Ca^{2+} , pH, and adenine nucleotides (See Figure caption for details). Enzyme activities are expressed as units of enzyme activity per




mg of protein. (One unit of activity is the amount of enzyme which transforms 1 μmol of substrate per minute at 30°C).

To fit the data that describe the regulatory effects of Ca²⁺ from Figures 3 and 4 of Denton *et al.* [23], it is necessary to account for mechanisms of allosteric activation/inhibition of Ca²⁺ in the model. It has been shown that Ca²⁺ can significantly affect the function of OGDHC by modulating the enzyme affinity for 2-oxoglutarate, showing a sigmoidal kinetics. Here we propose a general scheme of nonessential and mixed-type activation to characterize the effects of Ca²⁺. Based on this scheme, we assume the presence of two binding sites for Ca²⁺ on OGDHC and modify V_{max} and K_m of the enzyme complex for 2-oxoglutarate as follows:

$$V_{max,1} = V_{max,0} \left(\frac{1 + \frac{\beta_{Ca}[Ca^{2+}]}{\alpha_{Ca}K_{aCa}} + \frac{\beta_{Ca}[Ca^{2+}]^2}{(\alpha_{Ca}K_{aCa})^2}}{1 + \frac{[Ca^{2+}]}{\alpha_{Ca}K_{aCa}} + \frac{[Ca^{2+}]^2}{(\alpha_{Ca}K_{aCa})^2}} \right), \quad (14)$$

$$K_{m,1} = K_{m,0} \left(\frac{1 + \frac{[Ca^{2+}]}{K_{aCa}} + \frac{[Ca^{2+}]^2}{K_{aCa}^2}}{1 + \frac{[Ca^{2+}]}{\alpha_{Ca}K_{aCa}} + \frac{[Ca^{2+}]^2}{(\alpha_{Ca}K_{aCa})^2}} \right).$$

The modified flux expression for OGDHC is obtained by substituting Equation (14) into Equation (11). Three adjustable parameters (α_{Ca} , β_{Ca} , and K_{aCa}) are estimated based on the data from Denton *et al.* [23]; the model fits are shown in Figure 3A-B. For model simulations, the kinetic constants for substrates: 2-oxoglutarate, NAD⁻, and CoASH⁴⁻ are fixed at their previously



estimated values, obtained from data of McMinn and Ottaway [20]. Based on the fits to these data we find that β_{Ca} is close to one. For simplicity, we fixed $\beta_{Ca} = 1$, meaning that Ca^{2+} affects only the K_m of 2-oxoglutarate, not the V_{max} . This mechanism is also consistent with the conclusions of Denton *et al.* [23]. (Statistical analysis of different model formulations supports the validity of the null hypothesis that adding the extra adjustable parameters, β_{Ca} , does not lead to any significant improvement in fitting results.) The estimated values of the kinetic parameters are summarized in Table 1.

Experimentally it has been shown that the maximal activity of OGDHC is largely unaffected by changes in pH over the range 6.6-7.4, whereas the K_m of the enzyme is markedly altered by pH in this range [23]. In our model, the effect of pH on the OGDHC activity was described based on the observations studies of Denton and colleagues [23]. Here, protons are treated as the essential activators of OGDHC which increase the binding affinity of the enzyme to 2-oxoglutarate. Therefore, the K_m of 2-oxoglutarate is modified by multiplying the term $K_{aH}/[H^+]$ such that $K_{mA,1} = (K_{mA,1}K_{aH})/[H^+]$ (in Equation (15)). Figure 3C illustrates the model fits to the data obtained from Table 2 in McCormack and Denton [23] where the activity of OGDHC was studied under varying pH in both presence and absence of Ca^{2+} in the buffer.

Denton and colleagues [23] also studied the effect of adenine nucleotides (ATP and ADP) on the OGDHC activity where it was shown that both ATP and ADP significantly impact the K_m of the enzyme for 2-oxoglutarate and that the regulations of OGDHC by Ca^{2+} and adenine nucleotides seem to be independent. Here the regulatory effects of nucleotides on OGDHC activity is modeled as similar to that of Ca^{2+} . Specifically, we assume that there are different binding sites on the

OGDHC that bind to ATP or ADP. (The available kinetic data cannot exclude the other possibility that ATP and ADP can bind at the same site.) Therefore, the V_{max} and K_m for 2-oxoglutarate was modified as a function of nucleotide concentrations as follows:

$$V_{max,2} = V_{max,1} \frac{\left(1 + \frac{\beta_{ATP}[ATP]_T}{\alpha_{ATP}K_{ATP}}\right) \left(1 + \frac{\beta_{ADP}[ADP]_T}{\alpha_{ADP}K_{ADP}}\right)}{\left(1 + \frac{[ATP]_T}{\alpha_{ATP}K_{ATP}}\right) \left(1 + \frac{[ADP]_T}{\alpha_{ADP}K_{ADP}}\right)}, \quad (15)$$

$$K_{mA,2} = K_{mA,1} \frac{\left(1 + \frac{[ATP]_T}{K_{iATP}}\right) \left(1 + \frac{[ADP]_T}{K_{iADP}}\right)}{\left(1 + \frac{[ATP]_T}{\alpha_{ATP}K_{iATP}}\right) \left(1 + \frac{[ADP]_T}{\alpha_{ADP}K_{iADP}}\right)} \frac{K_{aH}}{[H^+]}$$

where $[ATP]_T$ and $[ADP]_T$ represent the total concentrations of the nucleotides. Experimental results show that neither free nucleotides nor magnesium binding nucleotides are solely responsible for activation observed. More data are needed to quantitatively and qualitatively specify the activation effects of each nucleotide's ionic forms. Figure 4(A-C) are model fits to the data obtained from Figures 5 and 6 of Denton and colleagues [23]. Optimization results based on these data indicate that both ATP and ADP change the K_m of OGDHC for 2-oxoglutarate without altering the maximum activity, which suggests that both β_{ATP} and β_{ADP} are equal to one.

Mg^{2+} is known to regulate the activity of OGDHC. In a recent study, Rodriguez-Zavala *et al.* [37] examined the effects of ligands, such as ATP, ADP, Ca^{2+} , and Mg^{2+} on the activity of OGDHC in both isolated pig heart enzyme complex and mitochondrial extracts. These data facilitate the characterization of the regulatory effect of Mg^{2+} on the OGDHC activity and are used here to estimate the Mg^{2+} associated kinetic parameters. Enzyme





Figure 5 OGDHC activity regulated by magnesium (A) OGDHC activity of isolated enzyme as a function of ADP in the presence of 1.0 mM NAD, 0.25 mM CoASH, and 0.5 mM 2-oxoglutarate with additions of 0.0 (circle), 25 μ M (up-triangle), 50 μ M (down-triangle) and 200 μ M (square) Mg^{2+} . Kinetic data were obtained from Figure Two of [37]. (B-C) OGDHC activity as a function of 2-oxoglutarate in the presence of 1.0 mM NAD, 0.25 mM CoASH with 600 μ M (B) or 0 μ M (C) Mg^{2+} addition. (up-triangle) no further additions (control); (circle) +600 μ M ATP; (down-triangle) +600 μ M ADP. Kinetic data were obtained from Figure Four of [37]. V_{max} has been changed less than 10% to better fit the symbols. Solid lines are model fitting results for the data points represented by sym-bols. The experimental data were obtained at pH 7.35 and 30°C in 120 mM KCl, 20 mM MOPS-K, 0.5 mM EGTA buffer.



activity is measured in nmol NADH²⁺ formed per minute per mg protein.

Experimental data from purified OGDHC from pig heart mitochondria from Rodriguez-Zavala *et al.* [37] with zero Mg²⁺ were used to estimate the V_{max} of the enzyme. Figure 5B shows the model simulations (lines) using the parameter estimates obtained above, with the exception that V_{max} was adjusted to match these experimental data. Data in Figure 5A with non-zero Mg²⁺ concentration and the data in Figure 5C are then used to estimate the kinetic parameters associated with the binding of Mg²⁺. Our model fits to these data assume that Mg²⁺ not only increases the activity by binding to the enzyme complex, but also potentiates the 2-oxoglutarate affinity to the enzyme and decrease the K_m of OGDHC for 2-oxoglutarate. Specifically, we assume two binding sites for Mg²⁺ and modify the V_{max} and the K_m as

$$V_{\max,3} = V_{\max,2} \left(\frac{1 + \frac{\beta_{Mg}[Mg^{2+}]}{\alpha_{Mg}K_{aMg}} + \frac{\beta_{Mg}[Mg^{2+}]^2}{(\alpha_{Mg}K_{aMg})^2}}{1 + \frac{[Mg^{2+}]}{\alpha_{Mg}K_{aMg}} + \frac{[Mg^{2+}]^2}{(\alpha_{Mg}K_{aMg})^2}} \right), \quad (16)$$

$$K_{mA,3} = K_{mA,2} \left(\frac{1 + \frac{[Mg^{2+}]}{K_{aMg}} + \frac{[Mg^{2+}]^2}{K_{aMg}^2}}{1 + \frac{[Mg^{2+}]}{\alpha_{Mg}K_{aMg}} + \frac{[Mg^{2+}]^2}{(\alpha_{Mg}K_{aMg})^2}} \right)$$

The data shown in Figure 5 are used to estimate the adjustable kinetic parameters related to Mg²⁺ ions in our kinetic model for OGDHC. Fits to the data are plotted in Figure 5 and the parameter values

summarized in Table 1. The developed model is able to satisfactorily explain the effect of Mg²⁺ ions on the enzyme activity. These results, combined with those shown in Figures 2 and 3, imply that the matrix free Ca²⁺ and Mg²⁺ ions concentrations exert significant and distinct effects on the OGDHC activity.

Complete flux expression for the 2-oxoglutarate dehydrogenase complex

Based on the proposed mechanisms of allosteric activation and inhibition of various cofactors, the flux expression (Equation (11)) of the OGDHC can be further modified. Applying the catalytic and regulatory mechanisms of Equations 14-16, the final forward flux expression is

$$J_{ogdhc}^+ = \frac{V_f[A][B][C] \cdot N}{\left([A][B][C] + \frac{1}{K_{ir}}K_{mC}[A][B][R] + K_{mC}[A][B] + \frac{1}{K_{iq}}K_{mB}[A][C][Q] + K_{mB}[A][C] + \frac{K_{mB}K_{ic}}{K_{iq}K_{ir}}[A][Q][R] + \frac{K_{ic}}{K_{iq}}K_{mB}[A][Q] + K_{mA}[B][C]\alpha_A + \frac{K_{ia}}{K_{ir}}K_{mC}[B][R] + \frac{K_{mB}K_{ic}K_{ia}}{K_{iq}K_{ir}}[Q][R] \right)} \quad (17)$$

$$\text{where } \alpha_A = \frac{\left(1 + \frac{[Ca^{2+}]}{K_{aCa}} + \frac{[Ca^{2+}]^2}{K_{aCa}^2} \right) \times \left(1 + \frac{[ATP]_T}{K_{ATP}} \right) \times \left(1 + \frac{[ADP]_T}{K_{ADP}} \right) \times \left(1 + \frac{[Mg^{2+}]}{K_{aMg}} + \frac{[Mg^{2+}]^2}{K_{aMg}^2} \right)}{\left(1 + \frac{[Ca^{2+}]}{\alpha_{Ca}K_{aCa}} + \frac{[Ca^{2+}]^2}{\alpha_{Ca}^2K_{aCa}^2} \right) \times \left(1 + \frac{[ATP]_T}{\alpha_{ATP}K_{ATP}} \right) \times \left(1 + \frac{[ADP]_T}{\alpha_{ADP}K_{ADP}} \right) \times \left(1 + \frac{[Mg^{2+}]}{\alpha_{Mg}K_{aMg}} + \frac{[Mg^{2+}]^2}{\alpha_{Mg}^2K_{aMg}^2} \right)} \times \left(\frac{K_{aH}}{[H^+]} \right)$$

$$\text{and } N = \frac{\left(1 + \frac{\beta_{Mg}[Mg^{2+}]}{\alpha_{Mg}K_{aMg}} + \frac{\beta_{Mg}[Mg^{2+}]^2}{(\alpha_{Mg}K_{aMg})^2}\right)}{\left(1 + \frac{[Mg^{2+}]}{\alpha_{Mg}K_{aMg}} + \frac{[Mg^{2+}]^2}{(\alpha_{Mg}K_{aMg})^2}\right)}$$

Estimated values of K_{aHb} , K_{iATP} , K_{aADP} , K_{aMg} , α_{ADP} , α_{ATP} , α_{Ca} , α_{Mg} , and β_{Mg} are listed in Table 1.

Independent validation of the developed kinetic model of OGDHC

Finally, the model is independently validated (corroborated) by comparing the model predictions to the initial rate data of Rutter and Denton [38] on the kinetics of OGDHC obtained from permeabilized mitochondria and mitochondrial extracts (see Figure 6(A-C)). They studied the regulations of NAD-linked isocitrate dehydrogenase and 2-oxoglutarate dehydrogenase by Ca^{2+} , nucleotide and nicotinamide nucleotides in permeabilized rat heart mitochondria and in mitochondria extracts. Data from their study were not used for estimation of model parameters and used here to further validate the proposed mechanisms and regulation of OGDHC. Therefore, the flux expression of Equation (17) was used for simulations with the values of the kinetic parameters the same as estimated before (see Table 1). The model accurately describes the kinetics and regulation of OGDHC, observed experimentally, without having to re-estimate the model kinetic parameters, signifying the accuracy of the model and the associated model parameters.

To determine the degree to which the model simulations are sensitive to the estimated parameter values, the relative sensitivities are computed and listed in Table 1. A high sensitivity value indicates that a small change in a given parameter can lead to significant changes in model outputs, used to identify the parameter values. All of our adjustable parameters of the model have sensitivities over 30%. Two parameter estimates (K_{aCa} and K_{aMg}) show relatively low sensitivity compared to the others, indicating that predictions of the developed model are less sensitive to these two values. This implies that these two parameters may not be identified accurately by the present analysis, given the sparseness of the data sets analyzed in this work regarding the regulation of OGDHC by Ca^{2+} and Mg^{2+} . Further experiments are required to adequately establish the appropriate regulatory mechanisms and the robustness of each model parameters.

Discussion

A number of kinetic models have been previously developed to explain the basic catalytic mechanisms and regulations by cofactors of OGDHC. Sanadi *et al.* [12] first proposed a Hexa-Uni-Ping-Pong mechanism for the

overall reaction by studying various roles and locations of the cofactors: thiamine pyrophosphate, lipoic acid, and FAD^{2-} within this complex. Hemada *et al.* [19] conducted kinetic studies and proposed a similar mechanism to that of Sandi *et al.* and suggested that $NADH^{2-}$ is a competitive inhibitor of NAD^+ . Whereas Smith *et al.* [10] suggested a noncompetitive inhibition of $NADH^{2-}$ with NAD^+ , the catalytic mechanism was not consistent with Sandi *et al.* Later, McMinn and Ottaway [20] tested a series of possible alternate mechanisms using computer optimization techniques and initial velocity studies and concluded that the binding of NAD^+ and $CoASH^{4-}$ and the release of Succinyl- CoA^{4-} is a random order, whereas the binding of the substrate 2-oxoglutarate and release of the product CO_2 still follows a Ping-Pong mechanism. Besides above experimental studies, a number of integrated models of mitochondrial bioenergetics have been developed which used different type of OGDHC models. Cortassa *et al.* [39] describe the activity of OGDHC as a function of Ca^{2+} , Mg^{2+} and substrate concentrations using phenomenological terms. Wu *et al.* [40] used a simple kinetic model of OGDHC from Kohn and Garfinkel [41] in their integrated model of TCA cycle that does not incorporate the regulatory effect of metal ion cofactors. In a recent integrated study of mitochondrial bioenergetics, Bazil *et al.* [42] developed a kinetic model of OGDHC based on a Hexa-Uni-Ping-Pong mechanism with a general description of the cofactor dependency of OGDHC activity. In summary, there have been a wide variety of kinetic models of OGDHC with contrasting kinetic mechanism and cofactor regulations.

In this paper, we developed a unified mechanistic model of OGDHC, in which Ca^{2+} , Mg^{2+} , ADP^{3-} , and pH are treated as activators and ATP^{4-} as inhibitor of the OGDHC activity. The present model offers more realistic and meaningful explanations on the catalytic properties and regulation mechanisms of OGDHC than previous attempts. The analysis also provides a unique set of kinetic parameters that consistently describe a wide variety of experimental data sets on OGDHC function, obtained from diverse sources. Based on the assumed ter-ter mechanism and associated conformational changes, we are able to consistently reproduce the observed kinetics of OGDHC with a minimal number of model parameters. Thus, the proposed mechanism is found to be more appropriate compared to other alternate kinetic models [43].

Effects of nucleotides on the OGDHC activity

Energy-linked regulators, ADP and ATP, as well as inorganic phosphate, have been investigated for over two decades for their profound effects on kinetic properties of OGDHC. Kinetic studies of mammalian OGDHC,

isolated from varied sources, have shown that ADP causes activation of OGDHC [44-47]. This enzyme complex is sensitive to ADP, where ADP significantly decreases the K_m for 2-oxoglutarate without affecting the maximum rate (V_{max}) of the reaction via allosteric interactions. For example, studies on the OGDHC of rat heart mitochondria show a seven fold decrease in the K_m value for 2-oxoglutarate by ADP and thereby strongly increases the affinity of OGDHC for the substrate [47]. Other studies of OGDHC from human heart make similar conclusions on the activating effect of ADP [48]. It has also been shown that, at subsaturating concentration of 2-oxoglutarate the relationship between initial reaction rates of OGDHC and concentration of ADP is sigmoidal, suggesting a positive cooperativity in binding of ADP to the enzyme complex [49]. In contrast, at a suboptimal 2-oxoglutarate concentration, ATP was shown to inhibit OGDHC activity in pig heart and bovine kidney mitochondria [23,45]. In addition direct inhibition effect, recent investigations have shown the possible indirect inhibition OGDHC by ATP because of chelation of divalent ions which activate OGDHC, such as Ca^{2+} , Mg^{2+} [37]. The activating action of ADP and inhibiting action of ATP are in competitive opposition (Figure 3E). It is still unclear if these two effectors bind on the same site on the complex or not. Model analysis based on the available kinetic data cannot exclude either possibility. In our mechanistic model, the regulatory effect of ADP and ATP is incorporated by assuming different binding sites for ADP and ATP in the enzyme complex and the model satisfactorily describes the activating effect of ADP and inhibitory effect ATP observed in many experiments [46].

While Zavala *et al.* interpret the data of Figure 5A to indicate that $MgADP^-$ is the effective activator of OGDHC activity, our model analysis of the available data sets on Mg^{2+} , ADP, and ATP dependent kinetics (Figures 4 and Figure 5B and 5C) reveals that, magnesium and ADP have independent parallel effects on the OGDHC activity, the most parsimonious explanation of the data. However, alternative, more complex, models cannot be ruled out.

Like ADP, Pi has also been shown to decrease the K_m value for 2-oxoglutarate, without affecting the V_{max} of OGDHC reaction [44,50]. In a recent report, the Pi activation showed biphasic behavior, with pH dependence [37]. In the physiological concentrations range Pi exerts monophasic activation of OGDHC [37], which can be described by Equation (15) with three extra parameters. Due to the lack of consistent kinetic data for Pi effects, we do not integrate a Pi dependent regulation mechanism in our current model. However, this energy linked effector may be physiologically important. The overall rate of oxidative phosphorylation is largely determined

by phosphorylation potential [51,52]. In cells when ATP utilization increases, the production of ADP and Pi increase. Therefore, activation of OGDHC by ADP and Pi may represent a compensating effect.

Effect of pH on the OGDHC activity

Mitochondrial matrix proton (H^+ ion) concentration is known to affect the OGDHC properties. Specifically, studies on pig heart OGDHC showed that the change in pH in the range 6.6 to 7.4 can significantly alter K_m of the enzyme for 2-oxoglutarate, without affecting its maximal activity. McCormack and Denton illustrated the effect of pH on OGDHC activity both in the absence and presence of Ca^{2+} in their assay mediums [46]. Other experimental observations have shown that hydrogen ions favor the higher affinity of OGDHC for 2-oxoglutarate [45,53]. In our model we hypothesize that hydrogen ions are essential activators of OGDHC activity to describe the observed pH dependency of the OGDHC kinetics.

Effects of Ca^{2+} , Mg^{2+} , and EGTA on the OGDHC activity

Studies by McCormack and Denton demonstrate the activating effects of Ca^{2+} ions on intra-mitochondrial dehydrogenases: pyruvate (PDH), NAD-isocitrate (NAD-ICDH), and 2-oxoglutarate (OGDHC) [24,54,55]. Specifically, the rise in cytosolic Ca^{2+} concentration in response to extrinsic stimuli, such as hormones can enhance mitochondrial oxidative metabolism via direct activation of these three Ca^{2+} sensitive dehydrogenases. Such mechanisms may serve as a complementary way to stimulate ATP-synthesis to meet the increased energy demand of the cell [24,54,55].

Mg^{2+} ion has also been shown to regulate the OGDHC activity either by directly activating the enzyme or by modulating the Ca^{2+} effects on the enzyme. In some studies, Mg^{2+} shows no effects on OGDHC activity [23]. However, in other studies, Mg^{2+} is shown to increase the maximal activity of the enzyme and the affinity of OGDHC for 2-oxoglutarate by enhancing the Ca^{2+} stimulatory effects on the enzyme complex [9,25,37]. These different observations could be accounted for the different levels of endogenous Ca^{2+} and Mg^{2+} present in the purified enzyme complex prepared by different methods. Another possible explanation is that the stimulatory effects of Mg^{2+} is TPP-dependent, which is not explicitly considered in our model. Panov and Scarpa [9] found that Mg^{2+} only exerts its stimulatory effects in the presence of TPP, though exclusion of TPP from the reaction medium has no effect on the initial enzyme activity in the absence of Mg^{2+} . Also, it has been clearly shown that Mg^{2+} may affect the rate of oxidative phosphorylation in isolated mitochondria primarily via modulating the OGDHC

activity [25]. The site of action of Mg^{2+} ion on OGDHC is unknown. In the present model, we hypothesized a general scheme of nonessential activation of Ca^{2+} , by considering two Ca^{2+} and Mg^{2+} binding sites on OGDHC. The Mg^{2+} effect is incorporated in our model by exclusively modifying the enzyme activity and 2-oxoglutarate binding step. So the parameters V_{max} and K_{mA} are accordingly expressed as functions of Mg^{2+} (Equation (16)). Currently, our model assumes that the turnover rate of E1 is modified to same value for binding either one ion or two ions. And to make it simple, our model does not include possible interaction between Mg^{2+} and Ca^{2+} at high concentrations either. The effects are not additive [9] at high concentration, suggesting that Mg^{2+} and Ca^{2+} may compete for the binding site. Additional kinetic data set are necessary to test different mechanisms and refine our model to more accurately describe the nature of cation dependent kinetic of OGDHC.

EGTA, which is used in many studies to control Ca^{2+} ion concentration in reaction media, has been shown in experiments and theoretical analysis to inhibit the NAD-linked isocitrate dehydrogenases (ICDH) through the binding complex, $MgEGTA$ [36]. To date, it is still not clear if there is similar inhibition effect of EGTA or EDTA on the activity of OGDHC. McCormack and Denton [23] concluded that the OGDHC sensitivity to EGTA is very similar to that observed with ICDH [56], because addition of calcium chelators EGTA or EDTA is associated with a marked decrease in the activity of OGDHC at 0.2 mM 2-oxoglutarate. Panov and Scarpa [9], in ascribing the inhibition effect of EGTA to the complex formation between Ca^{2+} and chelators, concluded that the effect of Ca^{2+} and chelators is associated with different endogenous cation levels in different preparations. But this explanation cannot account for McCormack and Denton's observation that EGTA or EDTA causes a 40% decrease of activity of OGDHC after using Chelex remove much of the endogenous Ca^{2+} in the buffer. It is also noted that the K_M for 2-oxoglutarate in the absence of Ca^{2+} is 4 ± 1.1 mM measured by Panov and Scarpa [9] for commercially available enzyme (Sigma, St. Louis, lot 44H80801), which is almost 15 times the estimated value used in our model (Table 1) and that found by previous workers [19,27]. Only by using the reported K_M of Panov and Scarpa [9], can we reproduce their data using the same mechanism (Equation (17)). In the absence of clear experimental evidence and sufficient data set, our model does not explicitly account for an inhibition effect of EGTA or EDTA.

Conclusion

Our mechanistic OGDHC model based on a detailed catalytic mechanism successfully provides a single consistent theoretical explanation for many previously unresolved experimental observations on the kinetics and regulations of OGDHC. In particular, it suggests the most plausible physiological regulations of OGDHC by NAD(H) oxidation-reduction state, the nucleotide phosphorylation potential, pH and various metal ions (Mg^{2+} and Ca^{2+}). As a rise in NADH can reduce the OGDHC flux and thereby provides feedback regulation through the electron transport chain, it is important to ask how NAD oxidation-reduction state and oxidative phosphorylation state exert a coherent regulation of OGDHC in physiological context. Furthermore, how does the OGDHC respond to stimuli via the mitochondrial Ca^{2+} transport system? Such questions may be addressed by applying the present model in an integrated framework [43] along with other dehydrogenases [36], the oxidative phosphorylation system [57], electron transfer system [58], and cation transport systems [59-61].

Acknowledgements

The authors thank Doug Disabato for participating in the initial part of the model development and gathering available experimental data for the analysis, Fan Wu, Kalyan Vinnakota, and Jason Bazil for helpful discussion. This work was supported by NIH grants R01-HL072011 and R01-HL095122.

Author details

¹Biotechnology and Bioengineering Center and Department of Physiology, Medical College of Wisconsin, Milwaukee, WI-53226, USA. ²Sanford-Burnham Medical Research Institute at Lake Nona, 6400 Sanger Road, Orlando, FL, 32827, USA.

Authors' contributions

FQ conceived the basic idea, collected experimental data set, developed the model, analyzed data and drafted the manuscript. RKP helped to collect and analyze the data and drafted the manuscript. RKD and DAB advised the study and revised the manuscript. All authors read and approved the final manuscript.

Received: 7 June 2011 Accepted: 26 September 2011

Published: 26 September 2011

References

1. Tretter L, Adam-Vizi V: **Alpha-ketoglutarate dehydrogenase: a target and generator of oxidative stress.** *Philos Trans R Soc Lond B Biol Sci* 2005, **360**(1464):2335-2345.
2. Perham RN: **Domains, motifs, and linkers in 2-oxo acid dehydrogenase multienzyme complexes: a paradigm in the design of a multifunctional protein.** *Biochemistry* 1991, **30**(35):8501-8512.
3. Strumilo S: **Often ignored facts about the control of the 2-oxoglutarate dehydrogenase complex.** *Biochemistry and Molecular Biology Education* 2005, **33**(4):284-287.
4. Sanadi DR, Littlefield JW, Bock RM: **Studies on alpha-ketoglutaric oxidase. II. Purification and properties.** *J Biol Chem* 1952, **197**(2):851-862.
5. Denton RM, McCormack JG, Edgell NJ: **Role of calcium ions in the regulation of intramitochondrial metabolism. Effects of Na+, Mg2+ and ruthenium red on the Ca2+-stimulated oxidation of oxoglutarate and on pyruvate dehydrogenase activity in intact rat heart mitochondria.** *Biochem J* 1980, **190**(1):107-117.

6. Hansford RG, Castro F: Role of Ca²⁺ in pyruvate dehydrogenase interconversion in brain mitochondria and synaptosomes. *Biochem J* 1985, **227**(1):129-136.
7. McCormack JG, Bromidge ES, Dawes NJ: Characterization of the effects of Ca²⁺ on the intramitochondrial Ca²⁺-sensitive dehydrogenases within intact rat-kidney mitochondria. *Biochim Biophys Acta* 1988, **934**(3):282-292.
8. McCormack JG, Denton RM: The role of Ca²⁺ ions in the regulation of intramitochondrial metabolism and energy production in rat heart. *Mol Cell Biochem* 1989, **89**(2):121-125.
9. Panov A, Scarpa A: Independent modulation of the activity of alpha-ketoglutarate dehydrogenase complex by Ca²⁺ and Mg²⁺. *Biochemistry* 1996, **35**(2):427-432.
10. Smith CM, Bryla J, Williamson JR: Regulation of mitochondrial alpha-ketoglutarate metabolism by product inhibition at alpha-ketoglutarate dehydrogenase. *J Biol Chem* 1974, **249**(5):1497-1505.
11. Starkov AA, Fiskum G, Chinopoulos C, Lorenzo BJ, Browne SE, Patel MS, Beal MF: Mitochondrial alpha-ketoglutarate dehydrogenase complex generates reactive oxygen species. *J Neurosci* 2004, **24**(36):7779-7788.
12. Sanadi DR: *Enzymes*, 2 1963, 7: 307-344.
13. Massey V: The composition of the ketoglutarate dehydrogenase complex. *Biochim Biophys Acta* 1960, **38**:447-460.
14. Cleland WW: The kinetics of enzyme-catalyzed reactions with two or more substrates or products. I. Nomenclature and rate equations. *Biochim Biophys Acta* 1963, **67**:104-137.
15. Koike M, Reed LJ, Carroll WR: alpha-Keto acid dehydrogenation complexes. IV. Resolution and reconstitution of the Escherichia coli pyruvate dehydrogenation complex. *J Biol Chem* 1963, **238**:30-39.
16. Angelides KJ, Hammes GG: Mechanism of action of the pyruvate dehydrogenase multienzyme complex from Escherichia coli. *Proc Natl Acad Sci USA* 1978, **75**(10):4877-4880.
17. Waskiewicz DE, Hammes GG: Fluorescence polarization study of the alpha-ketoglutarate dehydrogenase complex from Escherichia coli. *Biochemistry* 1982, **21**(25):6489-6496.
18. Angelides KJ, Hammes GG: Structural and mechanistic studies of the alpha-ketoglutarate dehydrogenase multienzyme complex from Escherichia coli. *Biochemistry* 1979, **18**(25):5531-5537.
19. Hamada M, Koike K, Nakaula Y, Hiraoka T, Koike M: A kinetic study of the alpha-keto acid dehydrogenase complexes from pig heart mitochondria. *J Biochem* 1975, **77**(5):1047-1056.
20. McMinn CL, Ottaway JH: Studies on the mechanism and kinetics of the 2-oxoglutarate dehydrogenase system from pig heart. *Biochem J* 1977, **161**(3):569-581.
21. Fromm HJ: The use of competitive inhibitors in studying the mechanism of action of some enzyme systems utilizing three substrates. *Biochim Biophys Acta* 1967, **139**(2):221-230.
22. Aevarsson A, Seger K, Turley S, Sokatch JR, Hol WG: Crystal structure of 2-oxoisovalerate and dehydrogenase and the architecture of 2-oxo acid dehydrogenase multienzyme complexes. *Nat Struct Biol* 1999, **6**(8):785-792.
23. McCormack JG, Denton RM: The effects of calcium ions and adenine nucleotides on the activity of pig heart 2-oxoglutarate dehydrogenase complex. *Biochem J* 1979, **180**(3):533-544.
24. McCormack JG, Denton RM: Role of Ca²⁺ ions in the regulation of intramitochondrial metabolism in rat heart. Evidence from studies with isolated mitochondria that adrenaline activates the pyruvate dehydrogenase and 2-oxoglutarate dehydrogenase complexes by increasing the intramitochondrial concentration of Ca²⁺. *Biochem J* 1984, **218**(1):235-247.
25. Rodriguez-Zavala JS, Moreno-Sanchez R: Modulation of oxidative phosphorylation by Mg²⁺ in rat heart mitochondria. *J Biol Chem* 1998, **273**(14):7850-7855.
26. Hayakawa T, Kanzaki T, Kitamura T, Fukuyoshi Y, Sakurai Y, Koike K, Suematsu T, Koike M: Mammalian alpha-keto acid dehydrogenase complexes. V. Resolution and reconstitution studies of the pig heart pyruvate dehydrogenase complex. *J Biol Chem* 1969, **244**(13):3660-3670.
27. Hirashima M, Hayakawa T, Koike M: Mammalian alpha-keto acid dehydrogenase complexes. II. An improved procedure for the preparation of 2-oxoglutarate dehydrogenase complex from pig heart muscle. *J Biol Chem* 1967, **242**(5):902-907.
28. Patel MS: Inhibition by the branched-chain 2-oxo acids of the 2-oxoglutarate dehydrogenase complex in developing rat and human brain. *Biochem J* 1974, **144**(1):91-97.
29. Lai JC, Cooper AJ: Brain alpha-ketoglutarate dehydrogenase complex: kinetic properties, regional distribution, and effects of inhibitors. *J Neurochem* 1986, **47**(5):1376-1386.
30. Shylaja N, Maehara M, Watanabe K: Measurement of alpha-ketoglutarate dehydrogenase activity in tissue extracts and human platelets using reversed-phase high-performance liquid chromatography. *Anal Biochem* 1990, **191**(2):223-227.
31. Northrop DB: Transcarboxylase. VI. Kinetic analysis of the reaction mechanism. *J Biol Chem* 1969, **244**(21):5808-5819.
32. Qi F, Dash RK, Han Y, Beard DA: Generating rate equations for complex enzyme systems by a computer-assisted systematic method. *BMC Bioinformatics* 2009, **10**:238.
33. Segel IH: *Enzyme kinetics: behavior and analysis of rapid equilibrium and steady-state enzyme systems*. New York: Wiley; 1975.
34. Li X, Dash RK, Pradhan RK, Qi F, Thompson M, Vinnakota KC, Wu F, Yang F, Beard DA: A database of thermodynamic quantities for the reactions of glycolysis and the tricarboxylic acid cycle. *J Phys Chem B* 2010, **114**(49):16068-16082.
35. Beard DA, Qian H: Relationship between thermodynamic driving force and one-way fluxes in reversible processes. *PLoS ONE* 2007, **2**(1):e1144.
36. Qi F, Chen X, Beard DA: Detailed kinetics and regulation of mammalian NAD-linked isocitrate dehydrogenase. *Biochim Biophys Acta* 2008, **1784**(11):1641-1651.
37. Rodriguez-Zavala JS, Pardo JP, Moreno-Sanchez R: Modulation of 2-oxoglutarate dehydrogenase complex by inorganic phosphate, Mg(2+), and other effectors. *Arch Biochem Biophys* 2000, **379**(1):78-84.
38. Rutter GA, Denton RM: Regulation of NAD⁺-linked isocitrate dehydrogenase and 2-oxoglutarate dehydrogenase by Ca²⁺ ions within toluene-permeabilized rat heart mitochondria. Interactions with regulation by adenine nucleotides and NADH/NAD⁺ ratios. *Biochem J* 1988, **252**(1):181-189.
39. Cortassa S, Aon MA, Marban E, Winslow RL, O'Rourke B: An integrated model of cardiac mitochondrial energy metabolism and calcium dynamics. *Biophys J* 2003, **84**(4):2734-2755.
40. Wu F, Yang F, Vinnakota KC, Beard DA: Computer modeling of mitochondrial tricarboxylic acid cycle, oxidative phosphorylation, metabolite transport, and electrophysiology. *J Biol Chem* 2007, **282**(34):24525-24537.
41. Kohn MC, Garfinkel D: Computer simulation of metabolism in palmitate-perfused rat heart. II. Behavior of complete model. *Ann Biomed Eng* 1983, **11**(6):511-531.
42. Bazil JN, Buzzard GT, Rundell AE: Modeling mitochondrial bioenergetics with integrated volume dynamics. *PLoS Comput Biol* 2010, **6**(1):e1000632.
43. Beard DA: Simulation of cellular biochemical system kinetics. *Wiley Interdiscip Rev Syst Biol Med* 2010, **3**(2):136-146.
44. Lawlis VB, Roche TE: Regulation of bovine kidney alpha-ketoglutarate dehydrogenase complex by calcium ion and adenine nucleotides. Effects on S0.5 for alpha-ketoglutarate. *Biochemistry* 1981, **20**(9):2512-2518.
45. Lawlis VB, Roche TE: Inhibition of bovine kidney alpha-ketoglutarate dehydrogenase complex by reduced nicotinamide adenine dinucleotide in the presence or absence of calcium ion and effect of adenosine 5'-diphosphate on reduced nicotinamide adenine dinucleotide inhibition. *Biochemistry* 1981, **20**(9):2519-2524.
46. McCormack JG, Denton RM: A comparative study of the regulation of Ca²⁺ of the activities of the 2-oxoglutarate dehydrogenase complex and NAD⁺-isocitrate dehydrogenase from a variety of sources. *Biochem J* 1981, **196**(2):619-624.
47. Nichols BJ, Rigoulet M, Denton RM: Comparison of the effects of Ca²⁺, adenine nucleotides and pH on the kinetic properties of mitochondrial NAD⁺-isocitrate dehydrogenase and oxoglutarate dehydrogenase from the yeast *Saccharomyces cerevisiae* and rat heart. *Biochem J* 1994, **303**(Pt 2):461-465.
48. Ostrovtsova SA, Strumilo SA: Participation of adenosine diphosphate in regulation of the 2-oxoglutarate dehydrogenase complex from human heart. *Biomed Biochim Acta* 1990, **49**(6):515-517.
49. Strumilo SA, Taranda NI, Vinogradov VV: Peculiarities of the regulation of adrenal oxoglutarate dehydrogenase complex by NADH and adenosine diphosphate. *Biokhimiia* 1982, **47**(5):724-732.

50. Strumilo SA, Taranda NI, Vinogradov VV: **Role of phosphate and divalent metal ions in regulation of the activity of the alpha-ketoglutarate dehydrogenase complex from adrenal cortex.** *Biokhimiia* 1981, **46**(1):156-161.
51. Beard DA, Kushmerick MJ: **Strong inference for systems biology.** *PLoS Comput Biol* 2009, **5**(8):e1000459.
52. Wu F, Zhang EY, Zhang J, Bache RJ, Beard DA: **Phosphate metabolite concentrations and ATP hydrolysis potential in normal and ischaemic hearts.** *J Physiol* 2008, **586**(17):4193-4208.
53. Pawelczyk T, Angielski S: **Cooperation of Ca²⁺ and pH in regulation of the activity of the 2-oxoglutarate dehydrogenase complex and its components from bovine kidney cortex.** *Acta Biochim Pol* 1984, **31**(3):289-305.
54. McCormack JG, Denton RM: **Influence of calcium ions on mammalian intramitochondrial dehydrogenases.** *Methods Enzymol* 1989, **174**:95-118.
55. Denton RM, McCormack JG: **Ca²⁺ transport by mammalian mitochondria and its role in hormone action.** *Am J Physiol* 1985, **249**(6 Pt 1):E543-554.
56. Denton RM, Richards DA, Chin JG: **Calcium ions and the regulation of NAD⁺-linked isocitrate dehydrogenase from the mitochondria of rat heart and other tissues.** *The Biochemical journal* 1978, **176**(3):899-906.
57. Beard DA: **A Biophysical Model of the Mitochondrial Respiratory System and Oxidative Phosphorylation.** *PLoS Comput Biol* 2005, **1**(4):e36.
58. Chen X, Qi F, Dash RK, Beard DA: **Kinetics and regulation of mammalian NADH-ubiquinone oxidoreductase (Complex I).** *Biophys J* 2010, **99**(5):1426-1436.
59. Dash RK, Qi F, Beard DA: **A biophysically based mathematical model for the kinetics of mitochondrial calcium uniporter.** *Biophys J* 2009, **96**(4):1318-1332.
60. Pradhan RK, Beard DA, Dash RK: **A biophysically based mathematical model for the kinetics of mitochondrial Na⁺-Ca²⁺ antiporter.** *Biophys J* 2010, **98**(2):218-230.
61. Pradhan RK, Qi F, Beard DA, Dash RK: **Characterization of membrane potential dependency of mitochondrial Ca²⁺ uptake by an improved biophysical model of mitochondrial Ca²⁺ uniporter.** *PLoS One* 2010, **5**(10):e13278.

doi:10.1186/1471-2091-12-53

Cite this article as: Qi et al.: Detailed kinetics and regulation of mammalian 2-oxoglutarate dehydrogenase. *BMC Biochemistry* 2011 **12**:53.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

