

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jds.com

Correspondence

Orthokeratinized odontogenic cyst with foreign body giant cell reaction to luminal keratin materials at the exposed fibrous connective tissue wall

KEYWORDS

Orthokeratinized
odontogenic cyst;
Maxilla;
Foreign body reaction

Orthokeratinized odontogenic cyst (OOC) is a specific type of odontogenic cyst with a lower recurrence rate.¹ In this case report, we presented an OOC with foreign body giant cell reaction to luminal keratin materials at the exposed fibrous connective tissue wall in the right maxillary third molar region.

This 24-year-old male patient was referred to the dental department of our hospital for treatment of a radiolucent lesion around the crown of the impacted right maxillary third molar. Oral examination showed the missing of the right maxillary third molar in the oral cavity and a slight expansion of the buccal and palatal cortical plates of the right maxillary third molar region. Panoramic radiography revealed a unilocular radiolucent lesion surrounding the crown of the impacted right maxillary third molar. Therefore, a clinical diagnosis of a central type of dentigerous cyst was made. After discussing with the patient, the treatment plan was extraction of the impacted right maxillary third molar followed by total enucleation of the cystic lesion. After obtaining the signed informed consent, extraction of the impacted right maxillary third molar was performed under local anesthesia and the cystic lesion was totally removed. The excised soft tissue specimen was sent for histopathological examination. Microscopically, the specimen showed a cystic lesion lined by the orthokeratinized stratified squamous epithelium of 5–7 layers of

epithelial cells. The superficial granular cell layer was prominent and composed of one to two layers of granular epithelial cells with keratohyaline granules in the cytoplasm. Many keratin shreds were found in the cystic lumen (Fig. 1A, B, C and D). A daughter OOC was discovered in the fibrous cystic wall (Fig. 1B and D). In the subepithelial area with a moderate to severe lymphoplasmic cell infiltrate, loss of superficial granular cell layer and surface keratin layer from the lining stratified squamous epithelium was found (Fig. 1B). In a specific area with complete loss of the lining epithelium and exposure of the underlying fibrous connective tissue to the keratin shreds in the cystic lumen, foreign body reaction with formation of multinucleated foreign body giant cells at the surface of the fibrous connective tissue was noted (Fig. 1E, F, G and H). The multinucleated foreign body giant cells had foamy cytoplasm, indicating the phagocytosis of the cystic keratin materials into the cytoplasm of the surface multinucleated foreign body giant cells (Fig. 1G and H). The characteristic microscopic findings confirmed the histopathological diagnosis of an OOC with foreign body giant cell reaction to luminal keratin materials at the exposed fibrous connective tissue wall.

The keratin layer on the surface of oral mucosa or skin has its function to protect the underlying epithelial and connective tissue cells. However, when the keratin materials are exposed to the stromal connective tissue cells,

<https://doi.org/10.1016/j.jds.2023.08.028>

1991-7902/© 2023 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Figure 1 Histopathological photomicrographs of our orthokeratinized odontogenic cyst (OOC). (A, B, C and D) Low- and high-power photomicrographs showing a cystic lesion lined by the orthokeratinized stratified squamous epithelium of 5–7 layers of epithelial cells. The superficial granular cell layer was prominent and composed of one to two layers of granular epithelial cells with keratohyaline granules in the cytoplasm. Many keratin shreds were found in the cystic lumen. A daughter OOC was discovered in the fibrous cystic wall (B and D). In the subepithelial area with a moderate to severe lymphoplasmic cell infiltrate, loss of superficial granular cell layer and surface keratin layer from the lining stratified squamous epithelium was found (B). (E, F, G and H) In a specific area with complete loss of the lining epithelium and exposure of the underlying fibrous connective tissue to the keratin shreds in the cystic lumen, foreign body reaction with formation of multinucleated foreign body giant cells at the surface of the fibrous connective tissue wall was noted. The multinucleated foreign body giant cells had foamy cytoplasm, indicating the phagocytosis of the luminal keratin materials into the cytoplasm of the surface multinucleated foreign body giant cells (G and H) (Hematoxylin and eosin stain; original magnification; A, B and E, 4 × ; F, 10 × ; C, D and G, 20 × ; H, 40 ×).

such as in the cases of well-differentiated squamous cell carcinoma with formation of tumor keratin, squamous cell carcinoma with chemotherapeutic drug-induced keratin formation after adjuvant chemotherapy, and the ruptured epidermal cyst or ruptured OOC with exposure of the fibrous connective tissue wall to luminal keratin materials, foreign body reaction with phagocytosis of the keratin materials by the multinucleated foreign body giant cells may occur.^{2–5}

In this case, there was no inflammation in the most parts of the subepithelial fibrous connective tissue wall, suggesting that this OOC is a developmental odontogenic cyst. However, in the minor parts of the fibrous cystic wall, when there was moderate or severe inflammation in the subepithelial connective tissue areas, loss of characteristic superficial granular cell layer and surface keratin layer from the lining stratified squamous epithelium could be discovered.

Declaration of competing interest

The authors have no conflicts of interest relevant to this article.

Acknowledgments

None.

References

1. Wang YJ, Zhang JY, Dong Q, Li TJ. Orthokeratinized odontogenic cysts: a clinicopathologic study of 159 cases and molecular evidence for the absence of PTCH1 mutations. *J Oral Pathol Med* 2022;51:659–65.
2. Sánchez-Romero C, Carlos R, Soares CD, Almeida Op de. Unusual multinucleated giant cell reaction in a tongue squamous cell carcinoma: histopathological and immunohistochemical features. *Head Neck Pathol* 2018;12:580–6.
3. Cheng FC, Wang LH, Wang YP, Chang JYF, Chiang CP. Comparisons of histological features among primary oral squamous cell carcinomas before and after adjuvant chemotherapy and their lymph node metastatic cancer lesions after adjuvant chemotherapy. *J Dent Sci* 2022;17:49–56.
4. Gupta R, Verma P, Bansal N, Semwal T. A case of ruptured perineal epidermal cyst. *Cureus* 2020;12:e11099.
5. Brito LNS, Lima FJ, Alves PM, Nonaka CFW, Godoy GP. Orthokeratinized odontogenic cyst associated with multinucleated giant cell reaction: report of unusual findings. *Rev Gaucha Odontol* 2017;65:376–9.

Yi-Pang Lee¹

Ming-Jay Hwang¹

Department of Dentistry, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Hualien, Taiwan

Julia Yu-Fong Chang**

Department of Dentistry, National Taiwan University Hospital, College of Medicine, National Taiwan University, Taipei, Taiwan

Graduate Institute of Oral Biology, School of Dentistry, National Taiwan University, Taipei, Taiwan

Graduate Institute of Clinical Dentistry, School of Dentistry, National Taiwan University, Taipei, Taiwan

Chun-Pin Chiang*

Department of Dentistry, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Hualien, Taiwan

Department of Dentistry, National Taiwan University Hospital, College of Medicine, National Taiwan University, Taipei, Taiwan

Graduate Institute of Oral Biology, School of Dentistry, National Taiwan University, Taipei, Taiwan

Graduate Institute of Clinical Dentistry, School of Dentistry, National Taiwan University, Taipei, Taiwan

**Corresponding author. Department of Dentistry, National Taiwan University Hospital, No. 1, Chang-Te Street, Taipei 10048, Taiwan.

E-mail address: jyfchang@ntu.edu.tw (J.Y.-F. Chang)

*Corresponding author. Department of Dentistry, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, No. 707, Section 3, Chung-Yang Road, Hualien 970, Taiwan.

E-mail address: cpchiang@ntu.edu.tw (C.-P. Chiang)

Received 28 August 2023

Available online 6 September 2023

¹ These two authors contributed equally to this work.