

Cyclic mechanical stretch down-regulates cathelicidin antimicrobial peptide expression and activates a pro-inflammatory response in human bronchial epithelial cells

Harpa Karadottir^{1,*}, Nikhil Nitin Kulkarni^{1,*}, Thorarinn Gudjonsson^{2,3}, Sigurbergur Karason⁴ and Gudmundur Hrafn Gudmundsson¹

¹ Biomedical Center and Department of Life and Environmental Sciences, University of Iceland, Reykjavik, Iceland

² Stem Cell Research Unit, Biomedical Center, Department of Anatomy, Faculty of Medicine, School of Health Sciences, University of Iceland, Reykjavik, Iceland

³ Department of Laboratory Hematology, Landspítali University Hospital, University of Iceland, Reykjavik, Iceland

⁴ Department of Anaesthesia and Intensive Care and Faculty of Medicine, Landspítali University Hospital and University of Iceland, Reykjavik, Iceland

* These authors contributed equally to this work.

ABSTRACT

Mechanical ventilation (MV) of patients can cause damage to bronchoalveolar epithelium, leading to a sterile inflammatory response, infection and in severe cases sepsis. Limited knowledge is available on the effects of MV on the innate immune defense system in the human lung. In this study, we demonstrate that cyclic stretch of the human bronchial epithelial cell lines VA10 and BCi NS 1.1 leads to down-regulation of cathelicidin antimicrobial peptide (*CAMP*) gene expression. We show that treatment of VA10 cells with vitamin D3 and/or 4-phenyl butyric acid counteracted cyclic stretch mediated down-regulation of *CAMP* mRNA and protein expression (LL-37). Further, we observed an increase in pro-inflammatory responses in the VA10 cell line subjected to cyclic stretch. The mRNA expression of the genes encoding pro-inflammatory cytokines IL-8 and IL-1 β was increased after cyclic stretching, where as a decrease in gene expression of chemokines IP-10 and RANTES was observed. Cyclic stretch enhanced oxidative stress in the VA10 cells. The mRNA expression of toll-like receptor (*TLR*) 3, *TLR5* and *TLR8* was reduced, while the gene expression of *TLR2* was increased in VA10 cells after cyclic stretch. In conclusion, our *in vitro* results indicate that cyclic stretch may differentially modulate innate immunity by down-regulation of antimicrobial peptide expression and increase in pro-inflammatory responses.

Submitted 21 September 2015

Accepted 18 November 2015

Published 7 December 2015

Corresponding author

Gudmundur Hrafn Gudmundsson,
ghrafn@hi.is

Academic editor

Pankaj Goyal

Additional Information and
Declarations can be found on
page 15

DOI 10.7717/peerj.1483

© Copyright

2015 Karadottir et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Molecular Biology, Immunology, Respiratory Medicine

Keywords LL-37, Vitamin D3, Pro-inflammatory response, Innate immunity, Ventilator induced lung injury

INTRODUCTION

Mechanical ventilation (MV) is a lifesaving treatment for patients suffering from severe respiratory failure by alleviating the work of breathing and facilitating alveolar gas exchange (Slutsky & Ranieri, 2013). MV has, however, been associated with side effects including ventilator induced lung injury (VILI) coupled with injury on lung tissue, stress on epithelial and endothelial barriers, apoptosis, pro-inflammatory responses, increased oxidative stress and secondary infections like nosocomial bacterial pneumonia. This can be followed by sepsis or systemic inflammatory response syndrome and increased mortality (Baudouin, 2001; Uhlig, 2002; Syrkin et al., 2008). Success of treatment with MV requires limitation of VILI and associated side effects (Fan, Villar & Slutsky, 2013). This can be accomplished by either decreasing mechanical stress produced by MV or by increasing the endurance of lung tissues to such strain. Hence, it has become imperative to study the molecular mechanisms behind VILI in details to improve outcomes in patients treated with MV. Although poorly defined, down-regulation of innate immune responses has been proposed to favor bacterial growth and development of ventilator associated pneumonia (VAP) in the lungs of patients during MV (Santos et al., 2005).

Antimicrobial polypeptides (AMPs) constitute an important arm of the innate immune defense in the lungs and are expressed ubiquitously in epithelial cells, neutrophils and monocytes or macrophages (Laube et al., 2006). These cationic polypeptides are categorized into: (1) smaller processed peptides such as cathelicidins and defensins and (2) larger polypeptides like lactoferrin, lysozyme and secretory leukocyte peptidase inhibitor (SLPI) (Laube et al., 2006). LL-37 is the main cathelicidin antimicrobial peptide (CAMP) in humans, encoded by the *CAMP* gene (Dürr, Sudheendra & Ramamoorthy, 2006). LL-37 is stored as a pro-form (pro-LL-37) in cells and is activated upon secretion to the mature form LL-37 by specific proteases (Sørensen et al., 2001). LL-37 has direct antimicrobial activity against multiple pathogens and has been demonstrated to exhibit pro- and anti-inflammatory responses, wound healing and angiogenic properties (Cederlund, Gudmundsson & Agerberth, 2011). Inducers of AMPs like vitamin D3 (1, 25-dihydroxy vitamin D3 or 1,25D3) and 4-phenyl butyric acid (PBA) have been shown to increase *CAMP* gene expression via the vitamin D receptor (VDR) (Gombart, Borregaard & Koeffler, 2005; Kulkarni et al., 2015a; Kulkarni et al., 2015b). A recent clinical trial demonstrated that lower vitamin D3 levels and cathelicidin expression was associated with higher mortality in critically ill patients usually receiving MV (Leaf et al., 2015). The effects of MV on respiratory cells can be modeled *in vitro* by applying defined cyclic mechanical stretch mimicking the frequency and stretch conditions during MV (Pugin et al., 2008; Wu et al., 2013).

In this study, we demonstrate that cyclic mechanical stretch of human bronchial epithelial cells VA10 and BCi down-regulates the expression of antimicrobial peptide cathelicidin. Treatment with AMP inducers vitamin D3 and/or PBA counteracted cyclic stretch mediated down-regulation of cathelicidin expression in VA10 cells. We further demonstrate that cyclic stretching of VA10 cells activated a pro-inflammatory response by enhancing expression of pro-inflammatory cytokines and increasing oxidative stress.

MATERIALS AND METHODS

Cell culture, reagents and cyclic stretch

An E6/E7 viral oncogene immortalized human bronchial epithelial cell line VA10 was cultured as described previously ([Halldorsson et al., 2007](#)). Briefly, the cells were maintained in Bronchial/Tracheal Epithelial cell growth medium (Cell Applications, San Diego, CA, USA) with Penicillin-Streptomycin ((20 U/ml, 20 µg/ml, respectively) (Life Technologies, Carlsbad, CA, USA)) at 37 °C and 5% CO₂. BCI. NS 1.1 (henceforth referred to as BCI) is a human bronchial epithelial cell line was a kind gift from Dr. Matthew S. Walters, Weill Cornell Medical College, New York NY, USA ([Walters et al., 2013](#)) and was established by immortalization with retrovirus expressing human telomerase (hTERT). The BCI cells were cultured as described above for VA10 cell line. Equal amount of cells were seeded on each well in a 6 well collagen I coated Bioflex plates (Flexcell International Corporation, Burlington, CA, USA), and grown to approximately 80% confluence. These plates were then transferred to a base plate of the cell stretching equipment Flexcell FX-5000TM Tension System (Flexcell International Corporation, Burlington, CA, USA) in a humidified incubator at 37 °C and 5% CO₂. The cells were subjected to cyclic mechanical stretch with the following parameters: a stretching rate of 20% with a square signal, 0.33 Hz frequency (20 cycles/min) and a 1:1 stretch:relaxation ratio, as described previously ([Pugin et al., 2008](#)). The cells were stretched for 6 h and 24 h as described in the results. Control Bioflex plates were kept in the same incubator under static conditions as non-stretch controls. Vitamin D3 (1,25D3) and Sodium 4-phenyl butyric acid (PBA) were purchased from Tocris bioscience, UK. Vitamin D3 was reconstituted in 100% ethanol as per manufacturer's instructions. The final concentration of the solvent was kept at 0.2% v/v and did not affect gene and protein expression of target genes. PBA was reconstituted in ultrapure H₂O.

RNA isolation and quantitative real time PCR

Total RNA was isolated with NucleoSpin RNA kit (Macherey-Nagel, Düren, Germany) and quantified on a spectrophotometer (Nanodrop, Thermo Scientific, Waltham, MA, USA). One µg of total RNA was reverse transcribed into first strand cDNA for each sample with a RevertAid First strand cDNA synthesis kit (Thermo Scientific, Waltham, MA, USA) and modified with 100 unit of reverse transcriptase per reaction. Power SYBR[®] green Universal PCR master mix (Life technologies, Carlsbad, CA, USA) was used to quantify the cDNA on a 7500 Real time PCR machine (Life technologies, Carlsbad, CA, USA). The reference gene used for all experiments was *UBC* (Ubiquitin C) and *PPIA* (Peptidylprolyl Isomerase A) and an arithmetic mean of reference gene Ct values was used. Primers for *TLR1* and *TLR6* were designed with Pearl primer and used at a final concentration of 300 nM ([Marshall, 2004](#)). All other primers were purchased from Integrated DNA technologies (PrimeTime Predesigned qPCR Assay) and used at a final concentration of 500 nM as per manufacturer's instructions. The qPCR cycling conditions were as follows; (1) Hold stage: 95 °C for 10 min, followed by 40 cycles of (2) De-natured stage: 95 °C for 15 s and (3) Annealed/extended stage: 60 °C for 1 min. The $2^{(-\Delta\Delta CT)}$ Livak method was utilized for calculating fold differences over untreated control ([Livak & Schmittgen, 2001](#)). A detailed list of the primers used in the q-RT-PCR assay is shown in [Table 1](#).

Table 1 Primers used in the q-RT PCR assay.

Primer	Gene symbol	Ref. seq. number	Forward primer	Reverse primer
CAMP	<i>CAMP</i>	NM_004345	5'-GCA GTC ACC AGA GGA TTG TGA C-3'	5'-CAC CGC TTC ACC AGC CC-3'
DEFB1	<i>DEFB1</i>	NM_005218	5'-CCA GTC GCC ATG AGA ACT CC-3'	5'-GTG AGA AAG TTA CCA CCT GAG GC-3'
IL-8	<i>CXCL8</i>	NM_000584	5'-CTG GCA TCT TCA CTG ATT CTT G-3'	5'-TGT CTG GAC CCC AAG GAA-3'
IP-10	<i>CXCL10</i>	NM_001565	5'-CAG TTC TAG AGA GAG GTA CTC CT-3'	5'-GAC ATA TTC TGA GCC TAC AGC A-3'
RANTES	<i>CCL5</i>	NM_002985	5'-TGC CAC TGG TGT AGA AAT ACT C-3'	5'-GCT GTC ATC CTC ATT GCT ACT-3'
IL-1 β	<i>IL1B</i>	NM_000576	5'-GAA CAA GTC ATC CTC ATT GCC-3'	5'-CAG CCA ATC TTC ATT GCT CAA G-3'
PPIA	<i>PPIA</i>	NM_021130	5'-TCT TTC ACT TTG CCA AAC ACC-3'	5'-CAT CCT AAA GCA TAC GGG TCC-3'
TLR1	<i>TLR1</i>	NM_003263	5'-CAA GAC TGT AGC AAA TCT-3'	5'-GTT TCG CCA GAA TAC TTA-3'
TLR2	<i>TLR2</i>	NM_003264	5'-ATG ACC CCC AAG ACC CA-3'	5'-CCA TTG CTC TTT CAC TGC TTT C-3'
TLR3	<i>TLR3</i>	NM_003265	5'-GCA CTG TCT TTG CAA GAT GA-3'	5'-AGA CCC ATA CCA ACA TCC CT -3'
TLR4	<i>TLR4</i>	NM_003266	5'-ACC CCA TTA ATT CCA GAC ACA-3'	5'-GAG TAT ACA TTG CTG TTT CCT GTT G-3'
TLR6	<i>TLR6</i>	NM_006068	5'-TAT CCT ATC CTA TTG-3'	5'-AGT TGC CAA ATT CCT TAC-3'
TLR8	<i>TLR8</i>	NM_138636	5'-GAT CCA GCA CCT TCA GAT GAG-3'	5'-ACT TGA CCC AAC TTC GAT ACC-3'
TLR9	<i>TLR9</i>	NM_017442	5'-GGA GCT CAC AGG GTA GGA A-3'	5'-AGA CCC TCT GGA GAA GCC-3'
UBC	<i>UBC</i>	NM_021009	5'-GAT TTG GGT CGC AGT TCT TG-3'	5'-CCT TAT CTT GGA TCT TTG CCT TG-3'
LZY	<i>LZY</i>	NM_000239	5'-CTC CAC AAC CTT GAA CAT ACT GA-3'	5'-AGA TAA CAT CGC TGA TGC TGT AG-3'
LTF	<i>LTF</i>	NM_001199149	5'-AAT AGT GAG TTC GTG GCT GTC-3'	5'-TGT ATC CAG GCC ATT GCG-3'

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and western blot analysis

Supernatants from treated cells were enriched for proteins on Oasis HLB mini columns (Waters, Milford, MA, USA) and lyophilized (Speed Vac; Thermo Scientific, Waltham, MA, USA). Ten μ g of total lyophilized protein was loaded on a 4–12% Bis–Tris gradient SDS gel. Protein loading was checked with total protein stain for polyvinylidene fluoride (PVDF) membrane (Pierce MemCode reversible protein stain; Thermo Scientific, Waltham, MA, USA). Total cell lysate was prepared by addition of RIPA lysis buffer (Sigma Aldrich, St. Louis, MO, USA) supplemented with 1 \times Halt protease and phosphatase inhibitor cocktail (Thermo Scientific, Waltham, MA, USA). Cells were then washed three times with cold 1 \times PBS, incubated for 30 min on ice with RIPA buffer and centrifuged at 10,000 rpm for 5 min at 4 °C. Supernatants obtained were used for Western blot analysis. The protein content of supernatants was analyzed by utilizing a bio rad protein assay dye reagent based on the Bradford dye binding method (Catalog No. 500-0006, Bio-Rad, USA). SDS-PAGE with subsequent Western blot analysis was performed using the NuPage blotting kit (Life Technologies, Carlsbad, CA, USA). Ten μ g of each sample (cell lysates) was loaded on a 4–12% Bis–Tris gradient gel and run at 200 V for 30 min. The proteins were then transferred to a PVDF membrane (Millipore, Billerica, MA, USA) and blocked with 5% non-fat skimmed milk in 1 \times phosphate buffered saline (PBS) with 0.05% tween (Sigma Aldrich, St. Louis, MO, USA). Antibodies against LL-37 were purchased from Innovagen, Sweden (polyclonal Rabbit, Cat No. PA-LL37-100) and Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*) from Santa Cruz Biotechnology, USA (polyclonal Rabbit, Cat No. sc-25778). The primary antibodies were diluted 1:1,000 in 1% non-fat

skimmed milk powder (Blotto; Santa Cruz Biotechnologies, Santa Cruz, California, USA) in 0.05% Tween 1 × PBS and incubated overnight at 4 °C. Next the membrane was incubated with 1:10,000 Horseradish Peroxidase (HRP)-linked secondary anti-rabbit IgG antibody (Cat No. A0545; Sigma Aldrich, St. Louis, MO, USA) in 0.05% Tween 1 × PBS. The protein bands were visualized by chemiluminescence with Pierce ECLPlus Western blotting substrate (Thermo Scientific, Waltham, MA, USA) on Image Quant LAS 4000 station (GE Healthcare, Wauwatosa, WI, USA).

Immunofluorescence

VA10 cells were fixed with 3.5% paraformaldehyde (Sigma Aldrich, St. Louis, MO, USA) prepared in 1 × PBS for 15 min. The cells were then washed twice with 1 × PBS for 10 min at room temperature and blocked in immunofluorescence (IF) buffer (10% Fetal bovine serum (FBS) and 0.3% triton X-100 in 1 × PBS) for 30 min. The cells were incubated overnight with LL-37 primary antibody (polyclonal Rabbit, Cat No. PA-LL37-100; Innovagen, Sweden) diluted 1:100 in the IF buffer, at 4 °C. Following overnight incubation, the cells were washed twice with 1 × PBS. Next, the cells were incubated with secondary antibody anti-rabbit IgG Alexa Fluor® 488 (Catalog No. A11070)/546 conjugate (Catalog No. A11010) from Life Technologies, Carlsbad, CA, USA, 1:1,000 diluted in the IF buffer, for 1 h at room temperature. The cells were counterstained with nuclear stain 4, 6-diamidino-2-phenylindole (DAPI) at 1:5,000 dilution or 3 μM in IF buffer (Catalog No. D9564; Sigma Aldrich, St. Louis, USA). Finally, the cells were washed twice with 1 × PBS and ultrapure H₂O, and mounted in Fluormount-G solution (Southern Biotech, Birmingham, AL, USA) for microscopic analysis. The images were captured on Olympus fluoview Fv1200 confocal microscope at a 20× magnification. Olympus fluoview (FV) 1,000 software was used for processing the acquired images.

Oxidative stress measurement

After subjecting VA10 cells to cyclic stretch, CellROX green reagent (Life Technologies, Carlsbad, CA, USA) was added to the medium at a final concentration of 5 μM for 30 min. Next, the cells were washed twice with cold 1 × PBS (1,000 rpm for 5 min) and detached with a 1 × Accutase solution (Millipore, Billerica, MA, USA). Cells were then harvested and suspended in 100 μl of MACS buffer. (Miltenyi Biotec, San Diego, CA, USA) as per manufacturer's instructions. The samples were analyzed in MACSQuant flow cytometer (Miltenyi Biotec, San Diego, CA, USA), placing the CellROX green reagent signal in FL1. Intact cells were gated in the Forward Scatter/Side Scatter plot to exclude debris. The resulting FL1 data was plotted on a histogram and is represented as % CellROX positive cells before and after cyclic stretch.

ELISA

Sandwich enzyme-linked immunosorbent assays (ELISAs) were performed utilizing an interleukin 8 (IL-8) and interferon gamma-induced protein 10 (IP-10) assay kit according to manufacturer's instructions (Peprotech, London, UK). The results are represented from three independent experiments.

Statistical analysis

The q-RT PCR and ELISA results are represented as means \pm standard errors of the means (S: E.) from three independent experiments. An unpaired Student's *t*-test was used to compare two samples. $P < 0.05$ was considered statistically significant. All the statistical analysis for q-RT PCR and ELISA experiments was performed with the Prism 6 software (Graph Pad, USA). The Western blot and immunofluorescence data are represented from at least three independent experiments showing similar results.

RESULTS

Cyclic stretch down-regulates the expression of the cathelicidin antimicrobial peptide

We screened for the effect of mechanical stretch on AMP expression. VA10 cells were subjected to stretch for 6 and 24 h to analyze early and late changes in AMP mRNA expression. The mRNA expression of AMPs cathelicidin (*CAMP*), human beta defensin-1 (*DEFB1*), Lactoferrin (*LTF*) and Lysozyme (*LYZ*) was analyzed with quantitative real time PCR (qRT-PCR). The basal mRNA expression of *CAMP* was decreased at both 6 h and 24 h after cell stretching (Fig. 1A). *DEFB1* mRNA expression was reduced at 24 h after cell stretching but was unaffected after 6 h (Fig. S1). The basal mRNA expression of *LTF* and *LZY* was very low ($Ct > 32$) in the VA10 cells and was excluded from this study. The decrease in cathelicidin gene expression was further confirmed at protein level with Western blot (Fig. 1B) and immunofluorescence analysis (Fig. 1C). Western blot analysis of stretched VA10 cells showed a decrease in secreted pro-LL-37 (encoded by the *CAMP* gene) levels after 24 h of cyclic stretch (Fig. 1B). Further, immunofluorescence staining of stretched VA10 cells also showed a decrease in LL-37 protein expression at both 6 h and 24 h after stretching (Fig. 1C).

Treatment with vitamin D3 and/or 4-phenyl butyric acid (PBA) counteracts stretch mediated down-regulation of cathelicidin expression

We and others have demonstrated that treatment with vitamin D3 (1,25D3) and PBA enhances cathelicidin expression (Gombart, Borregaard & Koeffler, 2005; Kulkarni et al., 2015a; Kulkarni et al., 2015b). VA10 cells were treated with 100 nM 1,25D3 (Fig. 2A), 2 mM PBA (Fig. 2B) or co-treated with 2 mM PBA and 100 nM 1,25D3 (Fig. 2C). These cells were then stretched for either 6 h or 24 h and gene expression of *CAMP* was analyzed with q-RT PCR. Treatment of VA10 cells with 1,25D3 and/or PBA before stretch counteracted stretch mediated down-regulation of *CAMP* mRNA expression (Figs. 2A–2C). This counteraction was further confirmed at protein level with immunofluorescence (Fig. 2E) and Western blot analysis (Fig. 2F). Immunofluorescence staining of LL-37 confirmed that the 1,25D3 treatment prevented stretch mediated decrease in LL-37 protein expression with both at 6 h and 24 h stretch (Fig. 2E). Further, VA10 cells were treated with PBA and 1,25D3 as described above and stretched for 24 h. Protein expression of pro-LL-37 was analyzed with Western blot. Co-treatment with PBA and 1,25D3 predominantly enhanced




Figure 1 Cyclic mechanical stretch down-regulates cathelicidin antimicrobial peptide expression. (A) VA10 cells were stretched for 6 and 24 h. The mRNA expression of cathelicidin antimicrobial peptide (*CAMP*) was analyzed with q-RT PCR after cell stretching ($n = 3$, mean \pm S.E.). Relative expression levels (y -axis) in static cells were defined with an arbitrary value of '1' and changes relative to this value in stretched samples are represented. (B) VA10 cells were subjected to cyclic stretch for 24 h. Cultured supernatants from stretched cells were used for analysis of secreted cathelicidin (pro-LL-37) protein expression by Western blot. Total protein loading is shown by staining with MemCode blue protein stain. The Western blot is a representative of three independent experiments showing similar results. (C) VA10 cells were stretched for 6 h and 24 h. The cells were then stained with antibody against LL-37 (green) and protein expression was visualized with immunofluorescence confocal microscopy. The cells were counterstained with nuclear stain DAPI (blue). Data is representative of three independent experiments showing similar results. Bar = 40 μ m (ns indicates non-significant; $p < 0.01$, **; $p < 0.001$, ***).

pro-LL-37 expression in VA10 cells (Fig. 2F). This enhanced expression was lower in the stretched cells (Fig. 2F). Finally, we verified stretch mediated decrease of *CAMP* mRNA expression in another human bronchial epithelial cell line BCI. Similar to the VA10 cells, the BCI cells were treated with 100 nM 1,25D3 and stretched for 24 h. The basal mRNA expression of *CAMP* was reduced with 24 stretch, however was not significantly changed




Figure 2 Treatment with vitamin D3 (1,25D3) and 4-phenyl butyric acid (PBA) counteracts stretch mediated down-regulation of cathelicidin expression. (A) VA10 cells were stretched for 6 and 24 h with (+) or without (–) 100 nM 1,25D3, (B) 2 mM PBA and (C) co-treated with vitamin D3 and PBA as shown in the figure. The mRNA expression of *CAMP* was assessed with (continued on next page...)

Figure 2 (...continued)

qRT-PCR ($n = 3$, mean \pm S.E.). Relative expression levels (y -axis) in static cells were defined with an arbitrary value of '1' and changes relative to this value in stretched/treated samples are represented. (D) Similarly, the BCi cells were stretched for 6 h and 24 h with (+)/without (-) 100 nM 1,25D3 and the mRNA expression of *CAMP* was analyzed with q-RT PCR ($n = 3$, mean \pm S.E.). Relative expression levels (y -axis) in static cells were defined with an arbitrary value of '1' and changes relative to this value in stretched/treated samples are represented. (E) VA10 cells were treated with 20 nM 1,25D3 and stretched for 6 h and 24 h. LL-37 protein expression (red) was analyzed with immunofluorescence confocal microscopy. The cells were counterstained with nuclear stain DAPI (blue). The data is a representative of three independent experiments showing similar results. Bar = 100 μ m. (F) Protein expression of cellular pro-LL-37 from stretched cells was also analyzed by Western blot analysis. VA10 cells were treated with 2 mM PBA, 20 nM 1,25D3 or co-treated with PBA and 1,25D3, followed by stretching for 24 h. GAPDH was used as a loading control. The Western blot is a representative of three independent experiments showing similar results (ns indicates non-significant; $p < 0.05$, *; $p < 0.01$, **; $p < 0.001$, ***; $p < 0.0001$ =****).

after 6 h. Similar to the results in VA10 cells, treatment with 1,25D3 counteracted stretch mediated down-regulation of *CAMP* mRNA expression in BCi cells (Fig. 2D). Thus, we demonstrate that treatment with 1,25D3 and PBA counteracted cyclic stretch mediated down-regulation of cathelicidin AMP expression.

Cyclic stretch activates a pro-inflammatory response and modulates toll-like receptor expression

Previous studies have demonstrated that VILI and *in vitro* cyclic stretching of alveolar lung epithelial cells activates a pro-inflammatory response by enhanced secretion of pro-inflammatory cytokines and chemokines (Vlahakis *et al.*, 1999; Halbertsma *et al.*, 2005). We studied the effect of cyclic stretch on inflammation in VA10 cells by screening for stretch mediated changes in pro-inflammatory cytokines and chemokines expression (Figs. 3A–3F). The VA10 cells were stretched for 6 h and 24 h. The mRNA expression of pro-inflammatory cytokines *CXCL8* (encoding interleukin 8 or IL-8), *IL1B* (encoding interleukin 1 beta or IL-1 β) and chemokines *CXCL10* (encoding interferon gamma induced protein 10 or IP-10), *CCL5* (encoding regulated on activation, normal T cell expressed and secreted or RANTES) was analyzed with q-RT PCR (Figs. 3A–3D) after stretching. The mRNA expression of genes encoding IL-8 and IL-1 β was enhanced after 6 h and 24 h cyclic stretch (Figs. 3A–3C). Interestingly, mRNA expression of the gene encoding IP-10 was reduced following 6 h and 24 h stretch (Figs. 3A–3C). The mRNA expression of gene encoding RANTES was reduced after 24 h stretching and was not affected after 6 h stretch (Fig. 3D). Analysis of protein expression with ELISA demonstrated enhanced secretion of IL-8 after 24 h of cell stretching (Fig. 3E), whereas the total secreted levels of IP-10 was reduced (Fig. 3F). Increased oxidative stress through enhanced reactive oxygen species (ROS) has been shown to be involved in activation of a pro-inflammatory response (Mittal *et al.*, 2014). We observed increased oxidative stress following cyclic stretch with the ROS detector dye Cell ROX. VA10 cells were stretched for 24 h and Cell ROX dye (5 μ M) was added 30 min before end of cyclic stretching. The cells were then harvested and analyzed in a flow cytometer. A significant increase in percentage of Cell ROX positive




Figure 3 Cyclic stretch activates a pro-inflammatory response and enhances oxidative stress. (A–D) VA10 cells were subjected to stretch for 6 and 24 h. The mRNA expression of genes encoding pro-inflammatory cytokines IL-8 (A), IL-1 β (B) and chemokines IP-10 (C), RANTES (D) was measured with q-RT PCR ($n = 3$, mean \pm S.E.). Relative expression levels (y -axis) in static cells were defined with an arbitrary value of '1' and changes relative to this value in stretched (continued on next page...)

Figure 3 (...continued)

samples are represented. (E–F) The protein expression of IL-8 (E) and IP-10 (F) from cultured supernatants was measured with ELISA. VA10 cells were stretched for 24 h and ELISAs were performed ($n = 3$, mean \pm S.E.). (G) Oxidative stress was measured with CellROX green reagent. VA10 cells were subjected to stretch for 24 h. CellROX dye (5 μ M) was added 30 min before the end of stretching. The cells were then harvested and analyzed by flow cytometry. The data is represented as percentage positive CellROX (ROS) cells before and after cyclic stretch ($n = 3$, mean \pm S.E.) (ns indicates non-significant; $p < 0.05$, *; $p < 0.01$, **; $p < 0.001$, ***; $p < 0.0001$, ****).

cells was observed after stretching, indicating increased oxidative stress (Fig. 3G). Thus, we demonstrate that cyclic stretching activates a pro-inflammatory response in VA10 cells.

Next, we screened for stretch mediated changes in toll-like receptor (TLR) expression in VA10 cells (Figs. 4A–4H). TLRs play an important role in activation of pro-inflammatory responses and have been shown to be modulated by mechanical stretching of cells (Takeda & Akira, 2005; Shyu et al., 2010). VA10 cells were stretched for 6 h and 24 h as described above. The mRNA expression of *TLR1*, *TLR2*, *TLR3*, *TLR4*, *TLR5*, *TLR6*, *TLR8* and *TLR9* was analyzed with q-RT PCR. The mRNA expression of *TLR3*, *TLR5* and *TLR8* was reduced 6 h after stretch (Figs. 4C, 4E and 4G). Further, the mRNA expression of *TLR2* (Fig. 4B) was increased and gene expression of *TLR3* (Fig. 4C) and *TLR9* (Fig. 4H) was decreased after 24 h stretch.

Treatment with vitamin D3 and PBA differentially affects stretch mediated changes in pro-inflammatory cytokine IL-8 expression

Vitamin D3 and PBA have been shown to differentially modulate inflammatory responses, mainly having anti-inflammatory effect (Hansdottir et al., 2010; Roy et al., 2012). VA10 cells were treated with 20 nM 1,25D3 (Fig. 5A) and 2 mM PBA (Fig. 5B), followed by stretching for 6 h and 24 h as shown in Fig. 5. The mRNA expression of genes encoding the pro-inflammatory cytokine IL-8 was analyzed with q-RT PCR. Treatment with 1,25D3 and PBA increased basal expression of IL-8 after 6 h treatment in static cells (Figs. 5A and 5B). Interestingly, treatment with PBA significantly enhanced stretch mediated increase in IL-8 gene expression after 6 h stretch (Fig. 5B). This enhancement was not observed at 24 h after stretch (Fig. 5B).

DISCUSSION

Mechanical ventilation (MV) is necessary for maintenance of gas exchange and to prevent cardiorespiratory collapse in patients suffering from serious respiratory failure and the more severe condition of acute respiratory distress syndrome (ARDS), but may also result in increased mortality of patients by ventilator induced lung injury (VILI) (Slutsky & Ranieri, 2013). Cyclic stretch generated during MV has been implicated in modulation of innate immune responses leading to side effects that include biotrauma and secondary infections like pneumonia (Santos et al., 2005). We hypothesized that cyclic stretch modulates the expression of antimicrobial peptides (AMPs) that constitute an important arm of the innate immune system. MV can be modelled *in vitro* by cyclic mechanical stretching of respiratory epithelial cells (Vlahakis et al., 1999; Pugin et al., 2008). In this




Figure 4 Cyclic stretch modulates toll-like receptor (TLR) gene expression. (A–H) VA10 cells were subjected to cyclic stretch for 6 h and 24 h. The mRNA expression of *TLR1* (A), *TLR2* (B), *TLR3* (C), *TLR4* (D), *TLR5* (E), *TLR6* (F), *TLR8* (G) and *TLR9* (H) was analysed with q-RT PCR ($n = 3$, mean \pm S.E.). Relative expression levels (y -axis) in static cells were defined with an arbitrary value of '1' and changes relative to this value in stretched samples are represented (ns indicates non-significant; $p < 0.05$, *; $p < 0.01$, **; $p < 0.0001$, ****).




Figure 5 Treatment with 1,25D3 and PBA differentially affects stretch mediated changes in pro-inflammatory cytokine IL-8 gene expression. (A, B) VA10 cells were treated with (+) or without (-) 20 nM 1,25D3 (A) or 2 mM PBA (B) and subjected to cyclic stretch for 6 h and 24 h. The mRNA expression of genes encoding pro-inflammatory cytokine IL-8 was analyzed with q-RT PCR ($n = 3$, mean \pm S.E.). Relative expression levels (y -axis) in static cells were defined with an arbitrary value of '1' and changes relative to this value in stretched/treated samples are represented. (ns indicates non-significant; $p < 0.05$, *; $p < 0.01$, **; $p < 0.0001$, ****).

study we show that cyclic mechanical stretch of human bronchial respiratory epithelial cell line VA10 down-regulates gene expression of AMP cathelicidin and human beta defensin 1 (Figs. 1 and S1). To our knowledge, this is first report demonstrating direct effects of mechanical stretch on AMP expression *in vitro*. AMPs prevent the invasion of pathogens through the lung epithelium and an impaired AMP production could render the tissues more susceptible to infections (Bals *et al.*, 1998). In immunocompromised mice, CRAMP (homologue of human cathelicidin) knockout resulted in increased susceptibility to dermal and respiratory infections (Nizet *et al.*, 2001; Kovach *et al.*, 2012). Our group has previously demonstrated that the diarrhoeal pathogen *Shigella* down-regulates cathelicidin antimicrobial peptide (CAMP) expression as a strategy to subvert innate immune responses (Islam *et al.*, 2001). This down-regulation can be counteracted by treatment with inducers of cathelicidin like butyric acid and PBA (Raqib *et al.*, 2006; Sarker *et al.*, 2011). In this study, we demonstrate that treatment with vitamin D3 (1,25D3) and PBA counteracts cyclic stretch mediated down-regulation of cathelicidin expression in the VA10 cells (Fig. 2). Interestingly, patients with low vitamin D3 levels admitted to intensive care units receiving MV had higher mortality rate and longer hospital stay than patients with sufficient vitamin D3 (Parekh, Thickett & Turner, 2013; Leaf *et al.*, 2015; Quraishi *et al.*, 2015). Further, in animal models, MV has been shown to promote bacterial dissemination and infections (Schortgen *et al.*, 2004). Thus induction of AMPs with inducers like 1,25D3 and PBA may be a useful strategy to reduce these complications arising from MV, preventing infections and hence can be sought as adjunct therapeutics.

We further studied the effects of cyclic stretch on inflammatory responses in the VA10 cells. Lung injury in MV patients is aggravated due to activation of MV mediated sterile inflammatory response and may lead to biotrauma (Santos *et al.*, 2005; Dhanireddy *et al.*, 2006). This damage due to MV activated inflammatory responses is further enhanced

during infections. A recent meeting abstract noted that rhinovirus infection enhanced cyclic stretch mediated up-regulation of pro-inflammatory cytokine expression in human bronchial epithelial cells (Nikitenko et al., 2014). We noticed a significant increase in mRNA expression of genes encoding the pro-inflammatory cytokines IL-8 and IL-1 β in the VA10 cells subjected to cyclic stretch (Fig. 3). Interestingly, the gene expression of chemokine IP-10 reduced at both 6 and 24 h of stretch. Further the mRNA expression of chemokine RANTES was down-regulated after 24 h of stretch (Fig. 3). Others have similarly shown cyclic stretch mediated up-regulation of IL-8 and down-regulation of RANTES expression in BEAS-2B human bronchial epithelial cell line (Oudin & Pugin, 2002; Thomas et al., 2006). Stretch mediated increase in IL-8 levels was shown to be dependent on activation of mitogen activated protein kinase (MAPK) and rho-kinase signaling (Oudin & Pugin, 2002; Thomas et al., 2006). Further, secretion of IL-1 β was shown to be enhanced in mouse alveolar macrophages following MV induced cyclic stretch via caspase-1 and TLR 4 dependent activation of NLRP3 inflammasomes (Wu et al., 2013).

Increased oxidative stress following cyclic stretch further confirmed activation of a pro-inflammatory response in the VA10 cells (Fig. 3). Recently, cyclic stretch was shown to activate mitochondrial reactive oxygen species (ROS) production via activation of nuclear transcription factor NF κ B and NADPH oxidase (Nox) 4 signaling in pulmonary arterial smooth muscle cells (Wedgwood et al., 2015). Mitochondrial ROS drives production of pro-inflammatory cytokines (Naik & Dixit, 2011). We stained VA10 cells with the CellROX (ROS detector) dye and noticed a significant increase in percentage of CellROX positive VA10 cells after cyclic stretch, indicating enhancement of oxidative stress (Fig. 3). We further looked at the effects of cyclic stretch on expression of toll-like receptors (TLRs). TLRs sense pathogen associated molecular patterns (PAMPs) upon infection, leading to activation of down-stream signaling pathways and induction of pro-inflammatory cytokines and chemokines (Takeda & Akira, 2005). Cyclic stretch of cultured cardiomyocytes was shown to enhance TLR4 gene expression via activation of the p38 MAPK and NF- κ B pathway (Shyu et al., 2010). Interestingly, cyclic stretch of human alveolar epithelial cell line A549 enhanced TLR2 expression (Charles et al., 2011). Further it was demonstrated that cyclic stretch enhanced IL-6 and IL-8 secretion in response to Pam₃CSK₄, a classical TLR2 ligand (Charles et al., 2011). In our study, the mRNA expression of TLR3, TLR5, TLR8 and TLR9 was down-regulated after cyclic stretch, whereas the gene expression of TLR2 was increased in the VA10 cells (Fig. 4). A direct causal link between the stretch mediated changes in TLR gene expression and pro-inflammatory cytokine expression needs to be established. Vitamin D3 and PBA have been shown to have anti-inflammatory effects (Hansdottir et al., 2010; Roy et al., 2012). In our study, treatment of the VA10 cells with both 1,25D3 and PBA enhanced mRNA expression of the gene encoding IL-8 (Fig. 5). This is in correlation with our previous study in the VA10 cells (Kulkarni et al., 2015a; Kulkarni et al., 2015b). Further, treatment with PBA significantly enhanced stretch mediated increase in IL-8 gene expression 6 h after stretching in the VA10 cells (Fig. 5).

Our study has certain limitations. (1) The study was performed exclusively in cell lines. However, these respiratory cell lines (VA10 and BCi) have been shown to have primary

cell like characteristics and have differentiation potential when cultured at an air–liquid interface (*Halldorsson et al., 2007; Walters et al., 2013*). They represent the upper airway lung epithelia. Primary human bronchial epithelial cells did not grow properly on collagen I coated bioflex silastic membranes and had to be excluded from this study. (2) The mechanism behind cyclic stretch mediated down-regulation of AMP expression needs to be elucidated and is a future area of interest. We hypothesize that stretch activated stress pathways (e.g., hypoxia related HIF-1 α (*Eckle et al., 2013; Fan et al., 2015*)) could be involved in the observed down-regulation of AMP expression. Interestingly, acidification of cellular milieu upon cyclic stretch has been shown to promote bacterial growth in lung epithelial cells (*Pugin et al., 2008*). The relationship between stretch altered pH and its effects on AMP gene expression is also an area of interest.

In conclusion, our *in vitro* data shows that cyclic stretch down-regulates the expression of AMP cathelicidin in VA10 and BCi respiratory epithelial cells and activates a pro-inflammatory response in VA10 cells. These results could have clinical implications in regards to ventilator treatment of patients by identifying ways to increase the endurance of lung tissues to mechanical strain and preventing respiratory infections, encouraging further *in vivo* studies in this field.

ACKNOWLEDGEMENTS

We would like to thank Jon Thor Bergthorsson and Katrin Birna Petursdottir for the help with analyzing flow cytometry data. We would like to specially thank Arí Jon Arason for introduction to the Flexcell tension system.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This project was supported by grants from LSH (Landspítali University Hospital), Össur hf and the Oddur Olafsson fund to Sigurbergur Karason. In addition this work was supported by funds from the University of Iceland and RANNIS. Nikhil Nitin Kulkarni was supported by University of Iceland grant for PhD students and RANNIS. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Landspítali University Hospital.

Össur hf and the Oddur Olafsson fund.

University of Iceland.

Icelandic Research Fund (RANNIS).

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Harpa Karadottir and Nikhil Nitin Kulkarni conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables.
- Thorarinn Gudjonsson and Sigurbergur Karason analyzed the data, contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- Gudmundur Hrafn Gudmundsson conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper.

Data Availability

The following information was supplied regarding data availability:

The research in this article did not generate any raw data.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.1483#supplemental-information>.

REFERENCES

- Bals R, Wang X, Zasloff M, Wilson JM. 1998.** The peptide antibiotic LL-37/hCAP-18 is expressed in epithelia of the human lung where it has broad antimicrobial activity at the airway surface. *Proceedings of the National Academy of Sciences of the United States of America* **95**:9541–9546 DOI [10.1073/pnas.95.16.9541](https://doi.org/10.1073/pnas.95.16.9541).
- Baudouin SV. 2001.** Ventilator induced lung injury and infection in the critically ill. *Thorax* **56**:ii50–ii57.
- Cederlund A, Gudmundsson GH, Agerberth B. 2011.** Antimicrobial peptides important in innate immunity. *The FEBS Journal* **278**:3942–3951 DOI [10.1111/j.1742-4658.2011.08302.x](https://doi.org/10.1111/j.1742-4658.2011.08302.x).
- Charles P-E, Tissières P, Barbar SD, Croisier D, Dufour J, Dunn-Siegrist I, Chavanet P, Pugin J. 2011.** Mild-stretch mechanical ventilation upregulates toll-like receptor 2 and sensitizes the lung to bacterial lipopeptide. *Critical Care* **15**:R181 DOI [10.1186/cc10330](https://doi.org/10.1186/cc10330).
- Dhanireddy S, Altemeier WA, Matute-Bello G, O'Mahony DS, Glenny RW, Martin TR, Liles WC. 2006.** Mechanical ventilation induces inflammation, lung injury, and extra-pulmonary organ dysfunction in experimental pneumonia. *Laboratory Investigation; a Journal of Technical Methods and Pathology* **86**:790–799 DOI [10.1038/labinvest.3700440](https://doi.org/10.1038/labinvest.3700440).
- Dürr UHN, Sudheendra US, Ramamoorthy A. 2006.** LL-37, the only human member of the cathelicidin family of antimicrobial peptides. *Biophysics* **1758**:1408–1425.
- Eckle T, Brodsky K, Bonney M, Packard T, Han J, Borchers CH, Mariani TJ, Kominsky DJ, Mittelbronn M, Eltzschig HK. 2013.** HIF1A reduces acute lung injury by optimizing carbohydrate metabolism in the alveolar epithelium. *PLoS Biology* **11**:e1001665 DOI [10.1371/journal.pbio.1001665](https://doi.org/10.1371/journal.pbio.1001665).
- Fan D, Coughlin LA, Neubauer MM, Kim J, Kim MS, Zhan X, Simms-Waldrup TR, Xie Y, Hooper LV, Koh AY. 2015.** Activation of HIF-1 α and LL-37 by commensal bacteria inhibits *Candida albicans* colonization. *Nature Medicine* **21**:808–814 DOI [10.1038/nm.3871](https://doi.org/10.1038/nm.3871).
- Fan E, Villar J, Slutsky AS. 2013.** Novel approaches to minimize ventilator-induced lung injury. *BMC Medicine* **11**:85 DOI [10.1186/1741-7015-11-85](https://doi.org/10.1186/1741-7015-11-85).

- Gombart AF, Borregaard N, Koeffler HP. 2005.** Human cathelicidin antimicrobial peptide (CAMP) gene is a direct target of the vitamin D receptor and is strongly up-regulated in myeloid cells by 1,25-dihydroxyvitamin D₃. *FASEB Journal: Official Publication of the Federation of American Societies for Experimental Biology* **19**:1067–1077 DOI [10.1096/fj.04-3284com](https://doi.org/10.1096/fj.04-3284com).
- Halbertsma FJJ, Vaneker M, Scheffer GJ, Van der Hoeven JG. 2005.** Cytokines and biotrauma in ventilator-induced lung injury: a critical review of the literature. *The Netherlands Journal of Medicine* **63**:382–392.
- Halldorsson S, Asgrimsson V, Axelsson I, Gudmundsson GH, Steinarsdottir M, Baldursson O, Gudjonsson T. 2007.** Differentiation potential of a basal epithelial cell line established from human bronchial explant. *In Vitro Cellular & Developmental Biology. Animal* **43**:283–289 DOI [10.1007/s11626-007-9050-4](https://doi.org/10.1007/s11626-007-9050-4).
- Hansdottir S, Monick MM, Lovan N, Powers L, Gerke A, Hunninghake GW. 2010.** Vitamin D decreases respiratory syncytial virus induction of NF-kappaB-linked chemokines and cytokines in airway epithelium while maintaining the antiviral state. *Journal of Immunology* **184**:965–974 DOI [10.4049/jimmunol.0902840](https://doi.org/10.4049/jimmunol.0902840).
- Islam D, Bandholtz L, Nilsson J, Wigzell H, Christensson B, Agerberth B, Gudmundsson G. 2001.** Downregulation of bactericidal peptides in enteric infections: a novel immune escape mechanism with bacterial DNA as a potential regulator. *Nature Medicine* **7**:180–185 DOI [10.1038/84627](https://doi.org/10.1038/84627).
- Kovach MA, Ballinger MN, Newstead MW, Zeng X, Bhan U, Yu F, Moore BB, Gallo RL, Standiford TJ. 2012.** Cathelicidin-related antimicrobial peptide is required for effective lung mucosal immunity in Gram-negative bacterial pneumonia. *Journal of Immunology* **189**:304–311 DOI [10.4049/jimmunol.1103196](https://doi.org/10.4049/jimmunol.1103196).
- Kulkarni NN, Gunnarsson HI, Yi Z, Gudmundsdottir S, Sigurjonsson OE, Agerberth B, Gudmundsson GH. 2015a.** Glucocorticoid dexamethasone down-regulates basal and vitamin D₃ induced cathelicidin expression in human monocytes and bronchial epithelial cell line. *Immunobiology* Epub ahead of print DOI [10.1016/j.imbio.2015.09.001](https://doi.org/10.1016/j.imbio.2015.09.001).
- Kulkarni NN, Yi Z, Huehnken C, Agerberth B, Gudmundsson GH. 2015b.** Phenylbutyrate induces cathelicidin expression via the vitamin D receptor: linkage to inflammatory and growth factor cytokines pathways. *Molecular Immunology* **63**:530–539 DOI [10.1016/j.molimm.2014.10.007](https://doi.org/10.1016/j.molimm.2014.10.007).
- Laube DM, Yim S, Ryan LK, Kisich KO, Diamond G. 2006.** Antimicrobial peptides in the airway. *Current Topics in Microbiology and Immunology* **306**:153–182.
- Leaf DE, Croy HE, Abrahams SJ, Raed A, Waikar SS. 2015.** Cathelicidin antimicrobial protein, vitamin D, and risk of death in critically ill patients. *Critical Care* **19**:80 DOI [10.1186/s13054-015-0812-1](https://doi.org/10.1186/s13054-015-0812-1).
- Livak KJ, Schmittgen TD. 2001.** Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔC_T} method. *Gene Expression* **408**:402–408.
- Marshall OJ. 2004.** PerlPrimer: cross-platform, graphical primer design for standard, bisulphite and real-time PCR. *Bioinformatics* **20**:2471–2472 DOI [10.1093/bioinformatics/bth254](https://doi.org/10.1093/bioinformatics/bth254).
- Mittal M, Siddiqui MR, Tran K, Reddy SP, Malik AB. 2014.** Reactive oxygen species in inflammation and tissue injury. *Antioxidants & Redox Signaling* **20**:1126–1167 DOI [10.1089/ars.2012.5149](https://doi.org/10.1089/ars.2012.5149).
- Naik E, Dixit VM. 2011.** Mitochondrial reactive oxygen species drive proinflammatory cytokine production. *The Journal of Experimental Medicine* **208**:417–420 DOI [10.1084/jem.20110367](https://doi.org/10.1084/jem.20110367).

- Nikitenko S, Shariff S, Arnason J, Shelfoon C, Kooi C, Proud D, Leigh R. 2014.** Cyclic stretch augments human rhinovirus induced inflammatory responses in airway epithelial cells. *Allergy, Asthma & Clinical Immunology* **10**(Suppl 1):A71 DOI [10.1186/1710-1492-10-S1-A71](https://doi.org/10.1186/1710-1492-10-S1-A71).
- Nizet V, Ohtake T, Lauth X, Trowbridge J, Rudisill J, Dorschner RA, Pestonjamas V, Piraino J, Huttner K, Gallo RL. 2001.** Innate antimicrobial peptide protects the skin from invasive bacterial infection. *Nature* **414**:454–457 DOI [10.1038/35106587](https://doi.org/10.1038/35106587).
- Oudin S, Pugin J. 2002.** Role of MAP kinase activation in interleukin-8 production by human BEAS-2B bronchial epithelial cells submitted to cyclic stretch. *American Journal of Respiratory Cell and Molecular Biology* **27**:107–114 DOI [10.1165/ajrcmb.27.1.4766](https://doi.org/10.1165/ajrcmb.27.1.4766).
- Parekh D, Thickett DR, Turner AM. 2013.** Vitamin D deficiency and acute lung injury. *Inflammation & Allergy Drug Targets* **12**:253–261 DOI [10.2174/18715281113129990049](https://doi.org/10.2174/18715281113129990049).
- Pugin J, Dunn-Siegrist I, Dufour J, Tissières P, Charles P-E, Comte R. 2008.** Cyclic stretch of human lung cells induces an acidification and promotes bacterial growth. *American Journal of Respiratory Cell and Molecular Biology* **38**:362–370 DOI [10.1165/rcmb.2007-0114OC](https://doi.org/10.1165/rcmb.2007-0114OC).
- Quraishi SA, McCarthy C, Blum L, Cobb JP, Camargo CA. 2015.** Plasma 25-hydroxyvitamin D levels at initiation of care and duration of mechanical ventilation in critically ill surgical patients. *Journal of Parenteral and Enteral Nutrition* Epub ahead of print. pii: 0148607114566276.
- Raqib R, Sarker P, Bergman P, Ara G, Lindh M, Sack DA, Nasirul Islam KM, Gudmundsson GH, Andersson J, Agerberth B. 2006.** Improved outcome in shigellosis associated with butyrate induction of an endogenous peptide antibiotic. *Proceedings of the National Academy of Sciences of the United States of America* **103**:9178–9183 DOI [10.1073/pnas.0602888103](https://doi.org/10.1073/pnas.0602888103).
- Roy A, Ghosh A, Jana A, Liu X, Brahmachari S, Gendelman HE, Pahan K. 2012.** Sodium phenylbutyrate controls neuroinflammatory and antioxidant activities and protects dopaminergic neurons in mouse models of Parkinson's disease. *PLoS ONE* **7**:e38113 DOI [10.1371/journal.pone.0038113](https://doi.org/10.1371/journal.pone.0038113).
- Santos CC dos, Zhang H, Liu M, Slutsky AS. 2005.** Bench-to-bedside review: biotrauma and modulation of the innate immune response. *Critical Care* **9**:280–286 DOI [10.1186/cc3022](https://doi.org/10.1186/cc3022).
- Sarker P, Ahmed S, Tiash S, Rekha RS, Stromberg R, Bergman P, Gudmundsson GH, Agerberth B, Raqib R. 2011.** Phenylbutyrate counteracts shigella mediated downregulation of cathelicidin in rabbit lung and intestinal epithelia: a potential therapeutic strategy. *PLoS ONE* **6**:e20637 DOI [10.1371/journal.pone.0020637](https://doi.org/10.1371/journal.pone.0020637).
- Schortgen F, Bouadma L, Joly-Guillou M-L, Ricard J-D, Dreyfuss D, Saumon G. 2004.** Infectious and inflammatory dissemination are affected by ventilation strategy in rats with unilateral pneumonia. *Intensive Care Medicine* **30**:693–701 DOI [10.1007/s00134-003-2147-7](https://doi.org/10.1007/s00134-003-2147-7).
- Shyu K-G, Wang B-W, Lin C-M, Chang H. 2010.** Cyclic stretch enhances the expression of toll-like receptor 4 gene in cultured cardiomyocytes via p38 MAP kinase and NF-kappaB pathway. *Journal of Biomedical Science* **17**:15 DOI [10.1186/1423-0127-17-15](https://doi.org/10.1186/1423-0127-17-15).
- Slutsky AS, Ranieri VM. 2013.** Ventilator-induced lung injury. *The New England Journal of Medicine* **369**:2126–2136 DOI [10.1056/NEJMra1208707](https://doi.org/10.1056/NEJMra1208707).
- Sørensen OE, Follin P, Johnsen AH, Calafat J, Tjabringa GS, Hiemstra PS, Borregaard N. 2001.** Human cathelicidin, hCAP-18, is processed to the antimicrobial peptide LL-37 by extracellular cleavage with proteinase 3. *Blood* **97**:3951–3959 DOI [10.1182/blood.V97.12.3951](https://doi.org/10.1182/blood.V97.12.3951).
- Syrkina O, Jafari B, Hales CA, Quinn DA. 2008.** Oxidant stress mediates inflammation and apoptosis in ventilator-induced lung injury. *Respirology* **13**:333–340 DOI [10.1111/j.1440-1843.2008.01279.x](https://doi.org/10.1111/j.1440-1843.2008.01279.x).

- Takeda K, Akira S. 2005.** Toll-like receptors in innate immunity. *International Immunology* **17**:1–14 DOI [10.1093/intimm/dxh186](https://doi.org/10.1093/intimm/dxh186).
- Thomas RA, Norman JC, Huynh TT, Williams B, Bolton SJ, Wardlaw AJ. 2006.** Mechanical stretch has contrasting effects on mediator release from bronchial epithelial cells, with a rho-kinase-dependent component to the mechanotransduction pathway. *Respiratory Medicine* **100**:1588–1597 DOI [10.1016/j.rmed.2005.12.008](https://doi.org/10.1016/j.rmed.2005.12.008).
- Uhlig S. 2002.** Ventilation-induced lung injury and mechanotransduction: stretching it too far? *American Journal of Physiology. Lung cellular and Molecular Physiology* **282**:L892–L896 DOI [10.1152/ajplung.00124.2001](https://doi.org/10.1152/ajplung.00124.2001).
- Vlahakis NE, Schroeder MA, Limper AH, Hubmayr RD. 1999.** Stretch induces cytokine release by alveolar epithelial cells *in vitro*. *The American Journal of Physiology* **277**:L167–L173.
- Walters MS, Gomi K, Ashbridge B, Moore MAS, Arbelaez V, Heldrich J, Ding B-S, Rafii S, Staudt MR, Crystal RG. 2013.** Generation of a human airway epithelium derived basal cell line with multipotent differentiation capacity. *Respiratory Research* **14**:135 DOI [10.1186/1465-9921-14-135](https://doi.org/10.1186/1465-9921-14-135).
- Wedgwood S, Lakshminrusimha S, Schumacker PT, Steinhorn RH. 2015.** Cyclic stretch stimulates mitochondrial reactive oxygen species and Nox4 signaling in pulmonary artery smooth muscle cells. *American Journal of Physiology. Lung Cellular and Molecular Physiology* **309**:L196–L203 DOI [10.1152/ajplung.00097.2014](https://doi.org/10.1152/ajplung.00097.2014).
- Wu J, Yan Z, Schwartz DE, Yu J, Malik AB, Hu G. 2013.** Activation of NLRP3 inflammasome in alveolar macrophages contributes to mechanical stretch-induced lung inflammation and injury. *Journal of Immunology* **190**:3590–3599 DOI [10.4049/jimmunol.1200860](https://doi.org/10.4049/jimmunol.1200860).