

Invited Review: Factors associated with atypical brain development in preterm infants: insights from magnetic resonance imaging

J. P. Boardman*  and S. J. Counsell† 

*MRC Centre for Reproductive Health, University of Edinburgh, Edinburgh and †Centre for the Developing Brain, School of Imaging Sciences and Biomedical Engineering, King's College London, London, UK

J. P. Boardman and S. J. Counsell (2020) *Neuropathology and Applied Neurobiology* 46, 413–421

Factors associated with atypical brain development in preterm infants: insights from magnetic resonance imaging

Preterm birth (PTB) is a leading cause of neurodevelopmental and neurocognitive impairment in childhood and is closely associated with psychiatric disease. The biological and environmental factors that confer risk and resilience for healthy brain development and long-term outcome after PTB are uncertain, which presents challenges for risk stratification and for the discovery and evaluation of neuroprotective strategies. Neonatal magnetic resonance imaging reveals a signature of PTB that includes dysconnectivity of neural networks and atypical development of cortical and deep grey matter structures. Here we provide a brief review of perinatal factors that are associated with the MRI signature of PTB. We con-

sider maternal and foetal factors including chorioamnionitis, foetal growth restriction, socioeconomic deprivation and prenatal alcohol, drug and stress exposures; and neonatal factors including co-morbidities of PTB, nutrition, pain and medication during neonatal intensive care and variation conferred by the genome/epigenome. Association studies offer the first insights into pathways to adversity and resilience after PTB. Future challenges are to analyse quantitative brain MRI data with collateral biological and environmental data in study designs that support causal inference, and ultimately to use the output of such analyses to stratify infants for clinical trials of therapies designed to improve outcome.

Keywords: brain, development, magnetic resonance imaging, neonate, preterm birth

Introduction

Preterm delivery, defined as birth at <37 weeks of gestation, is estimated to affect 10.6% of all live births around the world, which equates to 14.84 million births per annum [1]. In resource rich settings, advances in perinatal care and service delivery have led to improved survival over the past two decades: around 25% of infants born at 22 weeks who are offered stabilization at birth will survive and this

number increases to around 80% for births at 26 weeks [2]. However, early exposure to extrauterine life can impact brain development and is closely associated with long-term intellectual disability, cerebral palsy, autism spectrum disorder, attention deficit hyperactivity disorder, psychiatric disease and problems with language, behaviour and socioemotional functions [3].

Computational magnetic resonance imaging of brain development after preterm birth

The neuroimaging signature of preterm birth (PTB) includes alterations in white and grey matter

Correspondence: James P. Boardman, MRC Centre for Reproductive Health, Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh EH16 4TJ, UK. Tel: +44 131 242 2567; Fax: +44 (0)131 242 6441; E-mail: james.boardman@ed.ac.uk

© 2019 The Authors. *Neuropathology and Applied Neurobiology* published by John Wiley & Sons Ltd on behalf of British Neuropathological Society.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

microstructure, impaired cortical folding and disturbances to regional brain growth (Figure 1), for review see [4]. Advances in foetal imaging enable direct comparisons between healthy foetal and preterm brain development at equivalent gestations and, although brain growth is rapid between 25 and 40 weeks of gestation in preterm infants [5], growth trajectories are slower in preterm infants than in healthy foetuses [6]. At term equivalent age, regional brain volumes are reduced in preterm infants compared to healthy control infants and there is a reduction in cortical surface area, which may contribute to the neural basis of subsequent adverse neurodevelopmental outcome [5,7,8].

Diffusion-weighted magnetic resonance imaging (dMRI) studies have provided valuable insights into the effects of maturation and injury on microstructural brain development. Biological inference from dMRI is rooted in the premise that water molecules move with Brownian motion in an environment without restrictions and change direction following collisions with other particles. In highly structured tissue such as brain, water movement is restricted by the presence of

axons, neuronal cell bodies, glial cells and macromolecules, which supports inference about water content, axonal density, axonal calibre, myelination, dendritic arborization and synapse formation (for review see [9]).

In general, anisotropy increases and mean diffusivity (MD) decreases with increasing maturation in the developing white matter of the preterm brain [10,11] representing a combination of decreasing tissue water content and increasing complexity of white matter structures with age. Lower fractional anisotropy (FA) and increased MD are observed throughout the white matter in preterm infants compared with term-born infants [12,13] and increased prematurity is associated with lower FA and higher MD [14,15]. Diffusion tensor imaging metrics, such as FA, are nonspecific and reflect many underlying properties of brain tissue including neuronal density, fibre orientation dispersion, degree of myelination, free-water content and axonal diameter. New approaches to analyse dMRI data, including those based on biophysical models such as neurite orientation dispersion and density imaging (NODDI) [16], are




Figure 1. (A) (i) T1- and (ii) T2-weighted images of an infant at 26 weeks gestational age (GA) and (iii) T1- and (iv) T2-weighted images of an infant at 42 weeks GA at the level of the basal ganglia. (B) Diffusion magnetic resonance imaging maps at the level of the basal ganglia (i) fractional anisotropy, FA (ii) mean diffusivity, MD (iii) orientation dispersion index, ODI and (iv) neurite density index, NDI. (C) Brain segmentation in an infant born at 27⁺⁴ weeks gestational age and imaged at 41⁺² weeks postmenstrual age. Key: Green = cortical grey matter, blue = white matter, grey = deep grey matter, pink = extracerebral cerebrospinal fluid. (D) Correlation between gestational age at birth and FA measures in white matter assessed using tract-based spatial statistics. Mean FA skeleton (green) overlaid on mean FA map in the axial plane. Voxels showing a significant correlation ($P < 0.05$) between GA at birth FA are shown in red. (E) Diffusion MR tractography (i) arcuate fasciculus and (ii) optic radiations

adding to our understanding of the preterm neuroimaging phenotype. The NODDI model aims to disentangle these different factors by separating the influence of neurite density and orientation dispersion from each other, to provide indices of orientation dispersion index (ODI), which captures the degree of dispersion of axonal fibre orientations (e.g. through fanning, bending, crossing) or dendrite orientations, and neurite density index (NDI), represented by the intracellular volume fraction [16]. NDI increases with maturation in developing white matter and, at term equivalent age, NDI throughout the white matter is negatively associated with degree of prematurity at birth [17].

Unlike the changes observed in white matter, anisotropy and diffusivity in the developing cortical grey matter decrease with maturity and ODI increases reflecting dendritic growth from cell bodies, in-growth of thalamocortical afferents, synapse formation and proliferation of glial cells [18]. In comparison with term-born infants, preterm infants at term-equivalent age have increased cortical FA and cortical MD suggesting impaired cortical development [19], whereas lower gestational age (GA), lower birthweight and slower weight gain have been associated with higher FA in the preterm cortex [20].

Perinatal factors associated with altered brain development in preterm infants

MRI of the brain in early life has opened opportunities to investigate maternal and infant factors associated with risk and resilience for healthy brain development (Figure 2).

Maternal and foetal factors

Histologic chorioamnionitis Chorioamnionitis is infection of the amniotic fluid, membranes, placenta and/or decidua, and it affects around 40–80% of very preterm deliveries. It can initiate a foetal inflammatory response that is injurious to the developing brain [21], and epidemiological evidence suggests an association between chorioamnionitis, cystic periventricular leukomalacia and cerebral palsy in preterm infants [22]. We have shown histologically confirmed chorioamnionitis is associated with diffuse white matter disease at term equivalent age [23], although it does not appear to contribute to intraventricular haemorrhage or

punctate white matter lesions on conventional imaging [24]. This suggests that the pathway to atypical brain development begins *in utero* for some preterm infants.

Foetal growth restriction Foetal growth restriction (FGR) refers to the foetus who does not achieve expected *in utero* growth potential due to genetic or environmental factors. FGR is closely associated with childhood sensory and motor deficits, cognitive impairment and cerebral palsy [25]. MRI studies report atypical brain development in preterm infants affected by FGR, including reduced total and cortical grey matter volumes, reduced cortical complexity, reduced myelination, altered hippocampal and cerebellar development, changes in FA within the white matter skeleton and structural connectivity of specific brain networks [25]. These data suggest that FGR preterm infants have a pattern of atypical development that is distinct from that seen in appropriately grown preterm infants.

Socioeconomic deprivation Among the general population, brain tissue development and neurodevelopmental outcome are both patterned by socioeconomic gradients that operate in early life [26], and there is growing evidence that social disadvantage may exert additive risk to low GA for brain injury and impaired cognitive outcome in children born preterm [27,28]. Further work is required to understand the biological mechanisms that may link socioeconomic deprivation in the perinatal period with atypical brain development; plausible mechanisms include gestational immune dysregulation [29], alterations to the maternal hypothalamic-pituitary adrenal axis [30,31] and epigenomic variation associated with adversity in pregnancy [32,33].

Maternal alcohol and drugs Many studies report that prenatal alcohol exposure (PAE) is associated with atypical white matter in childhood, adolescence and adulthood [34], but studying the brain in later life introduces possible confounding by postnatal events and circumstances. MRI studies of neonates with PAE have reported altered dMRI parameters in white matter tracts, which suggests that atypical development is already established by the time of birth [35,36]. Maternal tobacco smoking is associated with lower global and regional foetal brain growth, after adjustment for somatic growth restriction [37].




Figure 2. Maternal, foetal and neonatal factors associated with brain development in preterm infants

Prenatal exposure to prescribed medications, specifically selective serotonin reuptake inhibitors, may influence neonatal brain structure and function [38,39], including among preterm infants [40]; and prenatal exposure to methadone for treatment of heroin addiction is associated with atypical white matter development [41]. These observations from patients with depression and opioid use disorder raise urgent questions about the safety of maternal prescribed and non-prescribed drugs on the developing foetal brain. Neonatal MRI biomarkers may be useful for studies designed to disambiguate disease from treatment effects, and for investigating maternal pharmacotherapies that are safest for mother and foetus.

Maternal stress An increasing body of evidence suggests that maternal prenatal stress exposure (PNSE) and anxiety/depression is associated with increased risk for a range of adverse behavioural outcomes in offspring including anxiety disorders [42], externalizing behaviour [43] and attention deficit hyperactivity disorder [44].

Recent studies provide evidence that the developing white matter is vulnerable to maternal prenatal adversity. Maternal anxiety is associated with reduced FA in key regions that are associated with anxiety, cognition and emotion regulation in later childhood including amygdala, cingulum, inferior temporal and frontal regions, angular gyrus, uncinate fasciculus, dorsolateral prefrontal cortex, cerebellum and inferior fronto-occipital fasciculus, in term-born infants [45]. Dean and colleagues reported higher diffusivity and lower NDI in frontal white matter of term-born infants of mothers experiencing prenatal symptoms of depression and anxiety [46], and we have observed higher diffusivity in the uncinate fasciculus in preterm infants at term equivalent age who experienced PNSE, even when controlling for GA at birth, socioeconomic status and the number of days on parenteral nutrition [47]. Defining neonatal brain image markers of maternal stress offers new opportunities for investigating the biological pathways that link maternal well-being with foetal brain development.

Neonatal factors

Co-morbidities of PTB Bronchopulmonary dysplasia (BPD), defined as the need for supplemental oxygen and/or respiratory support after 36 weeks GA, complicates the postnatal course of around 30% of infants born with very low birth weight, and it is an independent predictor of poor neurodevelopmental outcome [48]. Neonatal brain MRI studies of patients with severe respiratory morbidity, for example those with BPD or a requirement for prolonged mechanical ventilation, have reduced global and local brain volume [8], and reduced FA in white matter tracts [14] compared with age matched preterm infants without this complication.

Necrotizing enterocolitis (NEC; ischaemic necrosis of the intestinal mucosa) and blood stream infection in preterm infants often lead to a protracted systemic inflammatory response, and both are associated with neurodevelopmental impairment in early childhood. MRI studies suggest that severe NEC is associated with white matter injury, which might mediate the relationship between NEC and adverse neurodevelopmental outcome [49–51].

Retinopathy of prematurity is associated with reduced brain volume and altered white matter microstructure [52,53], and the preterm infant, like the term infant, is susceptible to brain injury from bilirubin toxicity, hypocapnia and severe hypoglycaemia, so clinical policies designed to prevent these complications during neonatal intensive care are important.

Postnatal nutrition Nutritional factors play an important role in preterm brain development and neuroimaging is a useful tool for investigating tissue effects of nutritional exposures. Optimal protein and energy intake in the first weeks after PTB are associated with increased brain growth, improved white matter microstructure and neurodevelopmental performance [54–56], and breast milk, as opposed to formula feed, during the weeks to discharge from Neonatal intensive care unit leads to improved structural connectivity of developing networks and greater FA in major white matter fasciculi [57].

Pain and medication Very preterm infants are exposed to repeated painful stimuli as part of intensive care. The burden of painful exposures is associated with

volume reduction in thalamic nuclei, altered thalamic metabolic function (decreased N-acetylaspartate/Choline), reduced FA in thalamocortical networks and reduced functional connectivity, which implies that pain during this critical period of human development influences development of the somatosensory system [58,59]. Neonates who require intensive care sometimes require analgesic and/or sedative medications. Midazolam appears to have a dose-dependent association with reduced hippocampal volume and microstructure, independent of procedural pain exposure burden [60]. These studies raise important hypotheses about the possible roles of pain and medication in modifying preterm brain development, and they signal the MRI techniques that are likely to be most useful in future studies designed to evaluate the safety of medicines during neonatal intensive care.

Genomics and epigenomics Imaging-genomics methods are beginning to be used to investigate the contribution of genomic variation and epigenetic modifications to preterm brain development. For example single-nucleotide polymorphisms at fatty acid desaturase 2, the 22q.11 locus, discs large MAGUK scaffold protein 4, and in the peroxisome proliferator-activated receptor pathway are associated with altered FA in white matter, and polygenic risk for psychiatric disease is associated with abnormal deep grey matter development in preterm infants [61–64]. These early observations suggest that genetic variants may contribute to neuroanatomic variation after PTB, and that PTB might expose susceptibility to psychiatric disease.

DNA methylation (DNAm) provides a molecular link between early life stress and neuropsychiatric disease in adulthood. PTB is a profound physiological stressor that is associated with alterations in the methylome at sites that influence neural development and function, and exploratory analyses suggest that differential DNAm is associated with white matter development in preterm infants [32].

Integrated analysis of genomic data, differential DNAm and quantitative MRI offers new opportunity for understanding genetic and epigenetic bases of preterm brain injury, and the biological pathways that contribute to susceptibility and repair after PTB.

Conclusions and future directions

MRI can be used to characterize brain development in terms of macro-, and microstructure, function and metabolism. Combining features from neuroimaging with biological and/or clinical information has identified several maternal and neonatal factors that are associated with susceptibility to atypical brain development. Furthermore, analysis of data across different scales provides a framework for investigating whether and how determinants of brain development that operate in the general population such as maternal well-being, drug exposures and socioeconomic gradients may interact with PTB to modify risk.

The observation that multiple types of exposure and genomic/epigenomic variants contribute to atypical brain development after PTB presents challenges for understanding causal pathways to injury and repair, and therefore for designing neuroprotective strategies targeted to the right infants at the right time. These challenges could be addressed by replication studies to assess generalizability, and by pooling image data from different centres to enhance study population sizes because scale-up is needed to address issues of power and sensitivity, and to enable study designs that support causal inference.

Acknowledgements

Figure 2 was created with BioRender.com. Image panel of postnatal brain growth reprinted from NeuroImage, (59)3, Serag A et al, Construction of a consistent high-definition spatio-temporal atlas of the developing brain using adaptive kernel regression, 2255-2265, Copyright 2012, with permission from Elsevier [65]. Figure 1E ii is reprinted from Brain, (131)2, Bassi L et al, Probabilistic diffusion tractography of the optic radiations and visual function in preterm infants at term equivalent age, 573-582, Copyright 2008, with permission from Oxford University Press [66].

References

- 1 Chawanpaiboon S, Vogel JP, Moller AB, Lumbiganon P, Petzold M, Hogan D et al. Global, regional, and national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis. *Lancet Glob Health* 2019; 7: e37–46

- 2 Myrhaug HT, Brurberg KG, Hov L, Markestad T. Survival and Impairment of extremely premature infants: a meta-analysis. *Pediatrics* 2019; 143: e20180933
- 3 Johnson S, Marlow N. Early and long-term outcome of infants born extremely preterm. *Arch Dis Child* 2017; 102: 97–102
- 4 Batalle D, Edwards AD, O'Muircheartaigh J. Annual research review: not just a small adult brain: understanding later neurodevelopment through imaging the neonatal brain. *J Child Psychol Psychiatry* 2018; 59: 350–71
- 5 Makropoulos A, Aljabar P, Wright R, Huning B, Merchant N, Arichi T et al. Regional growth and atlas of the developing human brain. *NeuroImage* 2016; 125: 456–78
- 6 Bouyssi-Kobar M, du Plessis AJ, McCarter R, Brossard-Racine M, Murnick J, Tinkleman L et al. Third trimester brain growth in preterm infants compared with in utero healthy fetuses. *Pediatrics* 2016; 138: e20161640
- 7 Kapellou O, Counsell SJ, Kennea N, Dyet L, Saeed N, Stark J et al. Abnormal cortical development after premature birth shown by altered allometric scaling of brain growth. *PLoS Med* 2006; 3: e265
- 8 Inder TE, Warfield SK, Wang H, Huppi PS, Volpe JJ. Abnormal cerebral structure is present at term in premature infants. *Pediatrics* 2005; 115: 286–94
- 9 Pecheva D, Kelly C, Kimpton J, Bonthron A, Batalle D, Zhang H et al. Recent advances in diffusion neuroimaging: applications in the developing preterm brain. *F1000Research* 2018; 7: 1326
- 10 Huppi PS, Maier SE, Peled S, Zientara GP, Barnes PD, Jolesz FA et al. Microstructural development of human newborn cerebral white matter assessed in vivo by diffusion tensor magnetic resonance imaging. *Pediatr Res* 1998; 44: 584–90
- 11 Miller SP, Vigneron DB, Henry RG, Bohland MA, Ceppi-Cozzio C, Hoffman C et al. Serial quantitative diffusion tensor MRI of the premature brain: development in newborns with and without injury. *J Magn Reson Imaging* 2002; 16: 621–32
- 12 Anjari M, Srinivasan L, Allsop JM, Hajnal JV, Rutherford MA, Edwards AD et al. Diffusion tensor imaging with tract-based spatial statistics reveals local white matter abnormalities in preterm infants. *NeuroImage* 2007; 35: 1021–7
- 13 Thompson DK, Inder TE, Faggian N, Johnston L, Warfield SK, Anderson PJ et al. Characterization of the corpus callosum in very preterm and full-term infants utilizing MRI. *NeuroImage* 2011; 55: 479–90
- 14 Ball G, Counsell SJ, Anjari M, Merchant N, Arichi T, Doria V et al. An optimised tract-based spatial statistics protocol for neonates: applications to prematurity and chronic lung disease. *NeuroImage* 2010; 53: 94–102

- 15 Partridge SC, Mukherjee P, Henry RG, Miller SP, Berman JI, Jin H et al. Diffusion tensor imaging: serial quantitation of white matter tract maturity in premature newborns. *NeuroImage* 2004; **22**: 1302–14
- 16 Zhang H, Schneider T, Wheeler-Kingshott CA, Alexander DC. NODDI: practical in vivo neurite orientation dispersion and density imaging of the human brain. *NeuroImage* 2012; **61**: 1000–16
- 17 Bataille D, Hughes EJ, Zhang H, Tournier JD, Tuszó N, Aljabar P, et al. Early development of structural networks and the impact of prematurity on brain connectivity. *NeuroImage* 2017; **149**: 379–92
- 18 Eaton-Rosen Z, Melbourne A, Orasanu E, Cardoso MJ, Modat M, Bainbridge A et al. Longitudinal measurement of the developing grey matter in preterm subjects using multi-modal MRI. *NeuroImage* 2015; **111**: 580–9
- 19 Ball G, Srinivasan L, Aljabar P, Counsell SJ, Durighel G, Hajnal JV et al. Development of cortical microstructure in the preterm human brain. *Proc Natl Acad Sci U S A* 2013; **110**: 9541–6
- 20 Vinall J, Grunau RE, Brant R, Chau V, Poskitt KJ, Synnes AR et al. Slower postnatal growth is associated with delayed cerebral cortical maturation in preterm newborns. *Sci Transl Med* 2013; **5**: 168ra8
- 21 Leviton A, Paneth N, Reuss ML, Susser M, Allred EN, Dammann O et al. Maternal infection, fetal inflammatory response, and brain damage in very low birth weight infants. Developmental Epidemiology Network Investigators. *Pediatr Res* 1999; **46**: 566–75
- 22 Shatrov JG, Birch SC, Lam LT, Quinlivan JA, McIntyre S, Mendz GL. Chorioamnionitis and cerebral palsy: a meta-analysis. *Obstet Gynecol* 2010; **116**: 387–92
- 23 Anblagan D, Pataky R, Evans MJ, Telford EJ, Serag A, Sparrow S et al. Association between preterm brain injury and exposure to chorioamnionitis during fetal life. *Sci Rep* 2016; **6**: 37932
- 24 Bierstone D, Wagenaar N, Gano DL, Guo T, Georgio G, Groenendaal F et al. Association of histologic chorioamnionitis with perinatal brain injury and early childhood neurodevelopmental outcomes among preterm neonates. *JAMA Pediatr* 2018; **172**: 534–41
- 25 Miller SL, Huppi PS, Mallard C. The consequences of fetal growth restriction on brain structure and neurodevelopmental outcome. *J Physiol* 2016; **594**: 807–23
- 26 Johnson SB, Riis JL, Noble KG. State of the art review: poverty and the developing brain. *Pediatrics* 2016; **137**: e20153075
- 27 Benavente-Fernandez I, Synnes A, Grunau RE, Chau V, Ramraj C, Glass T et al. Association of Socioeconomic Status and Brain Injury With Neurodevelopmental Outcomes of Very Preterm Children. *JAMA Netw Open* 2019; **2**: e192914
- 28 Ene D, Der G, Fletcher-Watson S, O'Carroll S, MacKenzie G, Higgins M et al. Associations of socioeconomic deprivation and preterm birth with speech, language, and communication concerns among children aged 27 to 30 months. *JAMA Netw Open* 2019; **2**: e1911027
- 29 Gilman SE, Hornig M, Ghassabian A, Hahn J, Cherkertzian S, Albert PS et al. Socioeconomic disadvantage, gestational immune activity, and neurodevelopment in early childhood. *Proc Natl Acad Sci U S A* 2017; **114**: 6728–33
- 30 O'Donnell KJ, Meaney MJ. Fetal origins of mental health: the developmental origins of health and disease hypothesis. *Am J Psychiatry* 2017; **174**: 319–28
- 31 Robinson R, Lahti-Pulkkinen M, Heinonen K, Reynolds RM, Raikkonen K. Fetal programming of neuropsychiatric disorders by maternal pregnancy depression: a systematic mini review. *Pediatr Res* 2019; **85**: 134–45
- 32 Sparrow S, Manning JR, Cartier J, Anblagan D, Bastin ME, Piyasena C et al. Epigenomic profiling of preterm infants reveals DNA methylation differences at sites associated with neural function. *Transl Psychiatr* 2016; **6**: e716
- 33 Suarez A, Lahti J, Czamara D, Lahti-Pulkkinen M, Knight AK, Girchenko P et al. The epigenetic clock at birth: associations with maternal antenatal depression and child psychiatric problems. *J Am Acad Child Adolesc Psychiatry* 2018; **57**: 321–8.e2
- 34 Ghazi Sherbaf F, Aarabi MH, Hosein Yazdi M, Haghshomar M. White matter microstructure in fetal alcohol spectrum disorders: A systematic review of diffusion tensor imaging studies. *Hum Brain Mapp* 2019; **40**: 1017–36
- 35 Donald KA, Roos A, Fouche JP, Koen N, Howells FM, Woods RP et al. A study of the effects of prenatal alcohol exposure on white matter microstructural integrity at birth. *Acta Neuropsychiatr* 2015; **27**: 197–205
- 36 Taylor PA, Jacobson SW, van der Kouwe A, Molteno CD, Chen G, Wintermark P et al. A DTI-based tractography study of effects on brain structure associated with prenatal alcohol exposure in newborns. *Hum Brain Mapp* 2015; **36**: 170–86
- 37 Ekblad M, Korkeila J, Lehtonen L. Smoking during pregnancy affects foetal brain development. *Acta Paediatr* 2015; **104**: 12–18
- 38 Jha SC, Meltzer-Brody S, Steiner RJ, Cornea E, Woolson S, Ahn M et al. Antenatal depression, treatment with selective serotonin reuptake inhibitors, and neonatal brain structure: a propensity-matched cohort study. *Psychiatry Res Neuroimaging* 2016; **253**: 43–53
- 39 Rotem-Kohavi N, Williams LJ, Virji-Babul N, Bjornson BH, Brain U, Werker JF et al. Alterations in resting-state networks following in utero selective serotonin reuptake inhibitor exposure in the neonatal brain. *Biol Psychiatry Cogn Neurosci Neuroimaging* 2019; **4**: 39–49
- 40 Podrebarac SK, Duerden EG, Chau V, Grunau RE, Synnes A, Oberlander TF et al. Antenatal exposure to antidepressants is associated with altered brain

- development in very preterm-born neonates. *Neuroscience* 2017; **342**: 252–62
- 41 Monnelly VJ, Anblagan D, Quigley A, Cabez MB, Cooper ES, Mactier H et al. Prenatal methadone exposure is associated with altered neonatal brain development. *Neuroimage Clin* 2018; **18**: 9–14
 - 42 Davis EP, Sandman CA. Prenatal psychobiological predictors of anxiety risk in preadolescent children. *Psychoneuroendocrinology* 2012; **37**: 1224–33
 - 43 Robinson M, Mattes E, Oddy WH, Pennell CE, van Eekelen A, McLean NJ et al. Prenatal stress and risk of behavioral morbidity from age 2 to 14 years: the influence of the number, type, and timing of stressful life events. *Dev Psychopathol* 2011; **23**: 507–20
 - 44 Grizenko N, Fortier ME, Zadorozny C, Thakur G, Schmitz N, Duval R et al. Maternal stress during pregnancy, ADHD symptomatology in children and genotype: gene-environment interaction. *J Can Acad Child Adolesc Psychiatry* 2012; **21**: 9–15
 - 45 Rifkin-Graboi A, Meaney MJ, Chen H, Bai J, Hameed WB, Tint MT et al. Antenatal maternal anxiety predicts variations in neural structures implicated in anxiety disorders in newborns. *J Am Acad Child Adolesc Psychiatry* 2015; **54**: 313–21.e2
 - 46 Dean DC 3rd, Planalp EM, Wooten W, Kecskemeti SR, Adluru N, Schmidt CK et al. Association of prenatal maternal depression and anxiety symptoms with infant white matter microstructure. *JAMA Pediatr* 2018; **172**: 973–81
 - 47 Lautarescu A, Pecheva D, Nosarti C, Nihouarn J, Zhang H, Victor S et al. Maternal prenatal stress is associated with altered uncinate fasciculus microstructure in premature neonates. *Biol Psychiatry* 2019. <https://doi.org/10.1016/j.biopsych.2019.08.010>
 - 48 Schmidt B, Roberts RS, Davis PG, Doyle LW, Asztalos EV, Opie G et al. Prediction of late death or disability at age 5 years using a count of 3 neonatal morbidities in very low birth weight infants. *J Pediatr* 2015; **167**: 982–6.e2
 - 49 Lee I, Neil JJ, Huettner PC, Smyser CD, Rogers CE, Shimony JS et al. The impact of prenatal and neonatal infection on neurodevelopmental outcomes in very preterm infants. *J Perinatol* 2014; **34**: 741–7
 - 50 Shah DK, Doyle LW, Anderson PJ, Bear M, Daley AJ, Hunt RW, Inder TE. Adverse neurodevelopment in preterm infants with postnatal sepsis or necrotizing enterocolitis is mediated by white matter abnormalities on magnetic resonance imaging at term. *J Pediatr* 2008; **153**: 170–5.e1
 - 51 Shin SH, Kim EK, Yoo H, Choi YH, Kim S, Lee BK et al. Surgical necrotizing enterocolitis versus spontaneous intestinal perforation in white matter injury on brain magnetic resonance imaging. *Neonatology* 2016; **110**: 148–54
 - 52 Sveinsdottir K, Ley D, Hovel H, Fellman V, Huppi PS, Smith LEH et al. Relation of retinopathy of prematurity to brain volumes at term equivalent age and developmental outcome at 2 years of corrected age in very preterm infants. *Neonatology* 2018; **114**: 46–52
 - 53 Glass TJA, Chau V, Gardiner J, Foong J, Vinall J, Zwicker JG, et al. Severe retinopathy of prematurity predicts delayed white matter maturation and poorer neurodevelopment. *Arch Dis Child Fetal Neonatal Ed* 2017; **102**: F532–7
 - 54 Schneider J, Fischer Fumeaux CJ, Duerden EG, Guo T, Foong J, Graz MB et al. Nutrient intake in the first two weeks of life and brain growth in preterm neonates. *Pediatrics* 2018; **141**: e20172169
 - 55 Coviello C, Keunen K, Kersbergen KJ, Groenendaal F, Leemans A, Peels B et al. Effects of early nutrition and growth on brain volumes, white matter microstructure, and neurodevelopmental outcome in preterm newborns. *Pediatr Res* 2018; **83**: 102–10
 - 56 Beauport L, Schneider J, Faouzi M, Hagmann P, Huppi PS, Tolsa JF et al. Impact of early nutritional intake on preterm brain: a magnetic resonance imaging study. *J Pediatr* 2017; **181**: 29–36.e1
 - 57 Blesa M, Sullivan G, Anblagan D, Telford EJ, Quigley AJ, Sparrow SA et al. Early breast milk exposure modifies brain connectivity in preterm infants. *NeuroImage* 2019; **184**: 431–9
 - 58 Duerden EG, Grunau RE, Guo T, Foong J, Pearson A, Au-Young S et al. Early procedural pain is associated with regionally-specific alterations in thalamic development in preterm neonates. *J Neurosci* 2018; **38**: 878–86
 - 59 Schneider J, Duerden EG, Guo T, Ng K, Hagmann P, Bickle Graz M et al. Procedural pain and oral glucose in preterm neonates: brain development and sex-specific effects. *Pain* 2018; **159**: 515–25
 - 60 Duerden EG, Guo T, Dodbiba L, Chakravarty MM, Chau V, Poskitt KJ et al. Midazolam dose correlates with abnormal hippocampal growth and neurodevelopmental outcome in preterm infants. *Ann Neurol* 2016; **79**: 548–59
 - 61 Boardman JP, Walley A, Ball G, Takousis P, Krishnan ML, Hughes-Carre L et al. Common genetic variants and risk of brain injury after preterm birth. *Pediatrics* 2014; **133**: e1655–63
 - 62 Krishnan ML, Wang Z, Silver M, Boardman JP, Ball G, Counsell SJ et al. Possible relationship between common genetic variation and white matter development in a pilot study of preterm infants. *Brain Behav* 2016; **6**: e00434
 - 63 Krishnan ML, Van Steenwinckel J, Schang AL, Yan J, Arnadottir J, Le Charpentier T et al. Integrative genomics of microglia implicates DLG4 (PSD95) in the white matter development of preterm infants. *Nat Commun* 2017; **8**: 428
 - 64 Cullen H, Krishnan ML, Selzam S, Ball G, Visconti A, Saxena A et al. Polygenic risk for neuropsychiatric

- disease and vulnerability to abnormal deep grey matter development. *Sci Rep* 2019; **9**: 1976
- 65 Serag A, Aljabar P, Ball G, Counsell SJ, Boardman JP, Rutherford MA. Construction of a consistent high-definition spatio-temporal atlas of the developing brain using adaptive kernel regression. *NeuroImage* 2012; **59**: 2255–65
- 66 Bassi L, Ricci D, Volzone A, Allsop JM, Srinivasan L, Pai A, et al. Probabilistic diffusion tractography of the

optic radiations and visual function in preterm infants at term equivalent age. *Brain* 2008; **131**: 573–582

Received 14 October 2019

Accepted after revision 13 November 2019

Published online Article Accepted on 20 November 2019