






BMJ Open Parenting, child development and primary care – ‘Crescer em Grande!’ intervention (CeG!) based on the Touchpoints approach: a cluster-randomised controlled trial protocol

Filipa Fareleira ^{1,2}, Maria Raul Xavier ³, Julia Velte ⁴,
Andreia Teixeira ^{1,2,5}, Carlos Martins ^{1,2}

To cite: Fareleira F, Xavier MR, Velte J, *et al.* Parenting, child development and primary care—‘Crescer em Grande!’ intervention (CeG!) based on the Touchpoints approach: a cluster-randomised controlled trial protocol. *BMJ Open* 2021;**11**:e042043. doi:10.1136/bmjopen-2020-042043

► Prepublication history is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-042043>).

Received 28 June 2020
Revised 08 February 2021
Accepted 01 March 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Filipa Fareleira;
up201607919@g.uporto.pt

ABSTRACT

Introduction Despite support for parenting being already recognised as a priority, there remains a paucity of evidence on how to facilitate its adoption in regular visits of maternal and child health primary care (PC). We describe the protocol for a study to assess the effect of an innovative universal Touchpoints-based intervention—‘Crescer em Grande!’ (CeG!) - when supporting the process of transition to parenthood and early infancy, at multiple PC units.

Methods and analysis A cluster-randomised trial will be conducted in 12 PC units (clusters) from the Lisbon metropolitan area, Portugal. Participants will be a minimum of three family physicians and one nurse/unit, as well as 216 expecting parents and future babies until 18 months who are using the PC services. Sites will be randomised to either the CeG! or usual care. The CeG! will consist of: (1) the integration of the Touchpoints approach in PC maternal and well-child visits, with the support of 28 leaflets for parents to file in a folder; plus (2) training for PC providers on how to perform the CeG! into existing practice. Parents will be required to fill in questionnaires at point throughout their child’s 18-month, mostly online. The primary outcome will be the self-perception of parental competence (Parenting Sense of Competence Scale). Other outcomes include: family functioning, couple dynamics, mental health, well-being/quality of life, psychological experience of pregnancy, attachment, child development. Acceptability, satisfaction and feasibility of CeG! will also be obtained from providers’ and parents’ perspectives. Costs associated with delivering the CeG! will be calculated. Study analyses will be under the principle of intention-to-treat.

Ethics and dissemination Approval was obtained from the Ethics Committee of the Regional Health Administration. The results will be shared with participants and disseminated via peer-reviewed published papers, presentations at scientific and professional conferences.

Trial registration number ISRCTN90692907

INTRODUCTION

The family, despite structural changes it has been suffering, maintains a prominent place

Strengths and limitations of this study

- We designed a challenging primary care (PC) cluster-randomised trial to assess the effect of a preventative, innovative Touchpoints-based intervention - ‘Crescer em Grande!’ (CeG!) - at multiple PC units from the Portuguese National Health Service, its impact on perceived competence of parents of children under 18 months of age and relation with other dimensions like parents’ mental health, child development and family well-being, that remain to be studied.
- The CeG! intervention was developed to be implemented in a real-world setting, in a non-stigmatising context (with no identifying entrance criteria), as part of routine care (no additional schedule), along with well-known and reliable family health professionals, in order to support the process of transition to parenthood and early infancy that has well-known implications in future developmental pathways.
- The CeG! includes Touchpoints-based written materials (leaflets) developed by a group of PC providers, including doctors and one nurse; leaflets’ content, organisation and design were also reviewed with academic experts and later by PC providers and parents using a focus group approach to reach the final version.
- The trial will take place in 12 PC units from the Lisbon metropolitan area, Portugal; therefore, our conclusions may potentially not be generalisable to other realities.
- We will mainly use self-report measures instead of structured observations and independent assessments of the intervention, which can influence responses as a result of social convenience.

in society as an elementary unit of organisation and emotional support.¹ In the contemporary era, when new multiple challenges arise for families, it is imperative to ensure the welfare of the different elements and to promote the empowerment of families. This

is particularly true in the transition between the well-identified family life cycle stages. An example of the latter is the transition to parenthood²³ and the parental task of the early years, a period identified as crucial for shaping children's future.⁴⁻⁷

Support for parenting is already recognised as a necessity and a right of families and should be presented as a worldwide priority by raising awareness of positive parenting among professionals working with children and parents.¹⁸ The literature shows that valued parents, supported, knowledgeable of child development and sensitised to positive strategies, use fewer punitive strategies, can better understand the behaviour of the child, feel less stressed. This way, they can foster a better parent-child relationship and family welfare, feeling more competent to face any approaching parental challenge.⁹⁻¹²

Indeed, an essential part of being a 'good parent' is how parents perceive themselves as capable of performing the varied tasks associated with parenting: parental self-efficacy beliefs¹³ and their satisfaction derived from the parenting role.^{13,14} This feeling of parental competence is a central construct to understand the dynamic processes of the family system. It has been widely studied as having implications on parenting behaviour, mental health and well-being; parent-child interactions; and children's development and behaviour.¹⁵⁻²¹

The Touchpoints (TP) model began to emerge after Brazelton's seminal work in the 1960s to support the families' needs and worries, valuing its strengths, promoting parental competence and giving empowerment back to families. Nowadays, the TP Network comprises communities and systems of care (health, education and social services) over 100 sites in 33 states in the USA and at several other places internationally, including Portugal.²² This well-established model of development and intervention assumes its importance given that it is dynamic and presupposes predictable periods of disorganisation (touchpoint definition)—from the prenatal period throughout the child's development in which the anticipatory care by the primary care (PC) professionals has recognised importance.^{23,24} It is also a relational, inclusive, non-prescriptive approach based on the construction of an alliance between professionals and parents, parents and child, professionals and children, valuable for teams.²⁵ We believe this kind of approach with parents needs to be integrated into preventive and universal care plans to support families and thereby to improve their quality of life. In 2015, an American Parent National Survey²⁶ with 2200 parents of children birth to 5 years from a wide range of backgrounds showed that fully half of the parents are not getting the support they need not only when they feel overwhelmed or stressed but also in making everyday decisions. Those parents share a universal desire to improve their parenting skills and do want guidance from child development professionals, mainly from the ones who know their child and situation. They believe that good parenting can be learnt, and most say that if they knew more positive parenting strategies,

they would use them—'needs and demands' also found in other studies.²⁷⁻²⁹

Presently, there is well-established evidence that parenting programmes are effective in improving both parents and child outcomes,^{9-12,20,30-36} but few are universal and start during pregnancy.^{10,20,32,34} Parenting programmes are generally built to work jointly with families who have children at risk or with behavioural, emotional or social difficulties. Therefore, it brings advantages to a relatively small group of parents. Relevant difficulties in recruitment and sustained parental adherence have also been described, sometimes compromising effectiveness.³⁷⁻³⁹ Universal approaches to parenting support are an additional offer that assumes itself as a public health intervention.³⁴ The general aim is to enhance the quality of the early family environment as it could benefit those at risk of adverse parenting but also other families in the population at large.

One promising strategy for reaching families widely is the PC setting—a non-stigmatising, favourable setting for parenting interventions not only throughout the child's development but also for the perinatal period to support parents' transition into parenthood. The main reasons are due to PC provider's rapport and credibility with families and a well-established mechanism for consistent contact through regular visits during pregnancy and the first 3 years of a child's life.^{35,40,41} Thus, the PC offers an ideal setting to promote positive parenting and reduce income-related developmental disparities.³⁶ Despite these advantages, PC providers often lack the time needed, so research is required within the PC setting to study different solutions, optimising this setting.

To this end, a parental intervention based on the TP model—'Crescer em Grande!' (CeG!) ('Grow Big!', English translation merely indicative)—was designed to support parents in prenatal period and early childhood. To the best of our knowledge, no preventive, universal family-based parenting interventions using the TP approach were previously developed worldwide for PC maternal and well-child visits.

AIM

We designed a PC cluster-randomised trial (CRT) with the aim of assessing the effect of CeG! on parenting sense of competence of parents of children under 18 months of age, compared with usual care and its relation with other dimensions: parental mental health and well-being (stress, depression, anxiety, psychological experience of pregnancy, attachment, quality of life), child development (physical/sensorimotor, psychosocial) and family well-being (family and couple functioning).

METHODS AND ANALYSIS

Trial design

Sites will be randomised to either the CeG! intervention or usual care arms in a parallel, multicentre CRT. The

clusters will be PC units. Participants will be comprised of PC providers (of each PC unit) and a subset of parents and their newborn babies attending these units. The unit of analysis will be the parents.

We decided on a cluster design as it is the most rigorous to prevent eventual contamination within any site if providers were chosen as the unit of randomisation (since it would not be possible to do parents' randomisation because patients are attached to their family physician).

Our protocol is under the Standard Protocol Items: Recommendations for Interventional Trials.⁴²

Study setting

The trial will take place in 12 PC units from the northern region of the Lisbon metropolitan area, Portugal, permitting the recommended minimum of four clusters per trial arm to minimise potential confounding from cluster effect.⁴³ Clusters will be from the two types of PC units of the National Health Service (NHS): custom healthcare units and family health units. These public local PC units cover the entire Portuguese territory, and are organised in small group-based practices, with family physicians, similar number of nurses and fewer secretaries. In some units, health providers are paid a fixed salary (custom healthcare units), while in family health units, they can have performance related pay.⁴⁴

Intervention

The CeG! intervention is based on the TP model. The TP model is a new paradigm of human development and clinical intervention inspired by a relational model. It assumes that favouring the bond of the baby to his family has significant repercussions from the perinatal period and throughout the children's development. The model focuses on a non-linear process of development and how it is experienced by all involved. It presupposes well-identified and predictable periods of disorganisation—the TP—when the child's behaviour seems to fall apart, typically preceding a spurt in development. Touchpoints are often accompanied by parental frustration and self-doubt. Therefore, they offer unique opportunities to make a difference in the lives of children and families because each TP outlines universal themes that might be coming up for the family, opportunities for practice to support the child/family system and suggestions for anticipatory guidance to help to predict the next TP. This model is anchored in the premise that the enhancement of a child's ability to reach their developmental potential cannot happen without supporting and enhancing the family's emotional and relational functioning, so children can have parents who are convinced of their skills and competencies. Thus, the TP relational framework presupposes a transformation in working with families by choosing strategies to support families' positive attributes, values and desires for raising their child on a collaborative attitude and empathic involvement instead of the traditional prescriptive and objective approach. This 'paradigm shift' (table 1) is supported by a set of TP guiding principles and parent and provider assumptions—tools to use and be intentional when working

Table 1 The Touchpoints paradigm shift

From:	To:
Deficit	Positive
Linear development	Multidimensional development
Prescriptive	Collaborative
Objective involvement	Empathetic involvement
Strict discipline boundaries	Flexible discipline boundaries

Source: adapted from Touchpoints Reference Guide and Participant Training Materials, 2016.⁴⁵

with families across articulated work between professionals from health, education and social services to build a common language. The principles of TP must be seen as general guidelines 'for using the developmental knowledge and relational skill at each of the touchpoints in making judgements that support good parenting'.⁴⁵ The TP parent and provider assumptions represent a set of beliefs providers should hold and strive to act on in working with families.^{45–47}

Thus, the CeG! presents itself as a universal, preventative intervention to contribute to positive parenting, consisting of two components: (1) the integration of the TP approach in PC maternal and well-child visits, with the support of 28 leaflets for parents to file in a folder, plus (2) training for PC providers on how to deliver, perform and integrate the CeG! intervention into existing practice.

The leaflets were developed with the collaboration of 13 PC providers (doctors and one nurse) to be used by the PC team and families as part of routine care. Each will be delivered unitarily in scheduled consultations from 26 weeks of gestation (first TP in the third trimester) until the age of 18 months, anticipating the TP and parents' needs, worries, fears, maybe not always addressed in healthcare visits (table 2). Following the Portuguese NHS recommendations, during the study period (table 3), each family will receive approximately six prenatal surveillance visits, and nine well-child visits.^{40 41}

Providers in the CeG! arm will complete a 5-hour training designed to enhance their motivation, skills and self-efficacy to use different elements of the strength-based TP model as guidance to promote parental competence. For the sessions, we developed a set of presentation slides for clinic staff that elucidates on the intervention concept and philosophy, implementation strategies and health gains. The focus will be on reflecting on the developmental process and challenges faced by the different elements of the system: child, parents and health providers, to start getting into a practical perspective through the TP relational framework. The aim is to help PC providers rethink their practice with families of young children. Specific strategies (TP principles and assumptions) and examples of implementation will be provided as a guide to adopting a TP approach. Different presentation techniques will be used: interactive activities (individual and group), videos, brainstorming, reflective practice activities. The written material (leaflets) reinforces the TP principles and elaborates on the developmental sequence of main

Table 2 The CeG! topics by Touchpoints and scheduled routine care in primary care

Healthcare visits	Touchpoints*		Topics
<i>Pregnancy</i>			
26–29 weeks	Prenatal	The Ideal Baby	Touchpoints definition Mummy and Daddy—I just arrived, what now? Baby and feeding
30–32 weeks			
33–35 weeks			
36–40 weeks	Newborn	The Real Baby	Role of caregivers in development
<i>Child</i>			
Home visit	–	–	Child Safety Vaccination
1st visit	3 weeks	The Energy Sink	Baby's language
1st month	6–8 weeks	The Rewarding Baby	Sleeping
2nd month	4 months	Looking Outward	Cognitive development (1)
4th month	7 months	Up at Night	Feeding
6th month	9 months	The Pointer	Frequent symptoms in childhood Breath-holding Stranger and separation anxiety
9th month	12 months	The Walker	Conquering autonomy/Walking
12th month	15 months	The Clinger	Child and discipline (1)
15th month	18 months	Rebel With a Cause	Toilet training
18th month	24 months	Getting to 'No!'	Aggression/Most common discipline challenges (2)†
2 years	3 years	'Why?'	Pacifier/Thumb sucking/Transitional object
3 years	4 years	'What I Do Matters'	Fears and nightmares Imaginary friend
4 years	5 years	'Who I Am Matters'	Most common discipline challenges (3)‡
5 years	6 years	Entering the Real World	Cognitive development (2)
Other	When necessary		Bedwetting Sibling rivalry

*Source: adapted from Touchpoints reference guide for healthcare providers, 2012.⁴⁶

†Most common discipline challenges (2)—tantrums, hitting, biting.

‡Most common discipline challenges (3)—whimpering, lying, retorting, complaining.

expected changes in a child and family's life until the age of six. One or more unit representatives will be charged with encouraging and enabling other professionals in their unit to review the leaflets folder throughout the study.

The control will consist of routine, maternal and well-child 'standard-of-care' in PC (usual care arm). Physicians and nurses in the control arm provide the usual management plan based on their assessment of the family members and can deliver/suggest any written information to help parents if they intended, as they usually did in the past, before entering the study. At the end of the trial data collection, we will offer clinical staff in the control arm the same CeG! training previously given to the intervention arm.

Patient and public involvement

CeG! design and different stakeholders' involvement

Both CeG! components were previously reviewed and discussed with academic experts from the Fundação Brazelton/Gomes-Pedro, Lisbon, Portugal. To further strengthen the process, we also engaged parents and PC providers in the CeG! development through eight focus group discussions (11 parents and 9 doctors and nurses) where they

looked at the general concept and written material and were invited to give feedback. Focus group suggestions were analysed by the research team and considered before the study's beginning. No relevant changes were deemed necessary, and all the participants revealed good acceptance.

No patients were involved in the CRT design, nor will be involved in recruitment, conduction and burden of the intervention.

Study description and recruitment

Units

Participating PC units will be selected as a convenience sample pertaining to the same geographical area (northern region of the Lisbon metropolitan area), which includes nine groups of health centres (management structures of PC units). The principal investigator (PI) met with acquainted executive and research directors from three groups of health centres; together they identified the eligible PC units accordingly to inclusion/exclusion criteria; a letter of introduction was sent to each PC unit, via an invitation email; first responders will be recruited consecutively until the required sample number is reached (n=12).

PC units are eligible if they have: 20–30 min of sequential nurse and physician consultations each; at least 80% of PC providers working as a stable team during the last year before the start of the trial; at least three family physicians and one nurse willing to participate for the unit to be included in the study.

To minimise contamination, we will only recruit one PC unit per building, and no unit may employ a study investigator.

Participants

Primary care providers: At least three family physicians and one family nurse from each site will be recruited if they maintain a regular clinical practice in maternal and well-child visits in each PC unit. After a letter of introduction from the PI, the investigators will discuss the project with each PC unit team, its expected benefits and time commitments (study briefing) and ask each PC provider to sign an informed consent approved by their Regional Health Administration's Ethics Committee. If there is a team composed by a family physician and a family nurse for each list of users, both will need to enter the study. In case of an occasional PC providers' absence, each PC unit guarantees that another PC provider enrolled in the study will hold maternal and well-child visits. Within each clinic, prenatal and paediatric healthcare delivered during the study may not be provided by any person without relevant training (eg, a first-year family physician resident, trainee nurse). Primary care providers will be ineligible if they are planning to retire during the study period or abandon the unit for other reasons.

Parents: Participants will be recruited through the physicians' list or a list of patients waiting for physician assignment (custom healthcare units), at each of the 12 sites. Parents (at least one parent per expecting baby, ie, fathers and/or mothers) are eligible if they: (1) have a confirmed pregnancy in the early second trimester; (2) do prenatal surveillance with the family doctor and wish to maintain paediatric surveillance of their baby in the PC unit; (3) are at least 18 years old; (4) are fluent in Portuguese as judged by the consenting provider; and (5) can understand all aspects of the study and provide informed consent. Mothers or fathers are included if interested regardless of their partners' decline. Parents will be ineligible if they: (1) wish to do concomitant regular paediatric surveillance at another health unit (public or private) not motivated by the need for disease surveillance in secondary healthcare (eg, hospital follow-up for cardiac disease treated by a paediatric cardiologist); or 2) intend to relocate during the study period.

Baby/child: Children of the participating parents that are born during the study.

There will be no financial incentive for study participation.

Study description

Overall CRT: Once the PC providers complete training for the CeG! intervention or get the study briefing on the usual care arm, a group of parents (of 216 babies) will be recruited and followed longitudinally for the CRT. All eligible parents

will be invited to participate in the trial during pregnancy, in person, during the first clinic visit of the second trimester (usually performed between 14 and 20 weeks of pregnancy⁴¹). At this initial contact, a PC provider will discuss the purpose of the study and study procedures with the parent and assess his interest. If interested, the PC provider will obtain the parent's written informed consent to participate. Parents will be recruited consecutively until the required sample number is reached.

Parents will be asked to complete questionnaires in multiple scheduled assessment points until the child's 18-month well-child visit (table 3).

We will also develop two other complementary studies along with the CRT (figure 1), as explained below:

- ▶ Before the CRT beginning, we will do a cross-sectional study to assess the primary outcome (parenting sense of competence) in a first group of parents of a month-old child who attend participating units. The objective of this data collection is that we can assess equivalence between units in the usual parenting sense of competence before initiating the intervention, and statistically adjust for it in relevant data analyses, if needed. For 4–6 weeks, all that parents will be approached in units by a PC provider from each site, accordingly with the same inclusion criteria (see above, Parents). If parents agree to participate, informed consent will be obtained and they will be asked to fill in the Parenting Sense of Competence Scale, along with the sociodemographic questionnaire and Parental Stress Inventory (as expected in the trial, for the first-month assessment point). Although in a much simpler way, this first moment of data collection will replicate one trial data collection and, thus, allow clinic staff to become proficient in recruitment and data collection processes.
- ▶ To establish if there is an improvement following the PC providers' training period, a pre-test and post-test self-developed Practice and Knowledge Questionnaire (PKQ) will be used to collect the data (single-group pre-test–post-test study) along with a sociodemographic and professional questionnaire. The post-test PKQ will be delivered in two separate moments: following the training period and at the end of the CRT. PKQ will focus on child development, TP model (concept and practice), self-efficacy, needs and willingness to learn more about how to better support children and families. We have chosen to do it also to obtain a pretraining understanding of the providers' strengths and weaknesses when taking care of babies and their families (prenatal and postnatal). For this reason, the PKQ will also be applied to PC providers in the control arm (only one time at the study briefing).

Outcomes

Primary outcome

For parents

- ▶ Difference in mean change (from the first month to the 18-month well-child visit) in parenting sense of

Table 3 Assessment points and related evaluation tools

Primary healthcare visits														
Maternal healthcare						Child healthcare								
2nd trimester (1st visit)*	26–29 weeks	30–32 weeks	33–35 weeks	36–40 weeks	Newborn	1st month	2nd month	4th month	6th month	9th month	12th month	15th month	18th month	Post study
Assessment tools (parents)	SDQ RDAS QOL	DASS-21 PMAST AAS	FES	RDAS QOL	BF†	PSOC PSI	DASS-21	–	FES PAS	PSOC PSI	–	DASS-21 RDAS QOL	PSOC PSI CBCL CF‡	SQ

*First visit of second trimester with parents in primary care unit, after the study start; after cross-sectional study in the control arm; after the training period in Crescer em Grande! arm (parents' recruitment).

†Only performed by mother.

‡Forms filled by primary care provider.

AAS, Maternal/Paternal Antenatal Attachment Scale; BF, Baby form; CBCL, Child Behavior Checklist 1.5–5; CF, Child form; DASS-21, Depression, Anxiety and Stress Scale-21; FES, Family Environment Scale; PAS, Maternal/Paternal Postnatal Attachment Scale; PMAST, Pregnancy and Motherhood Attitudes Scale; PSI, Parental Stress Inventory; PSOC, Parenting Sense of Competence scale; QOL, EUROHIS-QOL-8; RDAS, Revised Dyadic Adjustment Scale; SDQ, sociodemographic questionnaire; SQ, satisfaction questionnaire.

competence between the two trial arms, using the Parenting Sense of Competence (PSOC) scale.

Secondary outcomes

For parents

- ▶ Difference in mean change (from the first to the last research assessment for each instrument—table 3) between the two trial arms for family functioning (Family Environment Scale (FES)), couple dynamics (Revised Dyadic Adjustment Scale (RDAS)), mental health including depression/anxiety and stress (Depression, Anxiety and Stress Scale-21 (DASS-21)); Parental Stress Inventory (PSI)) and well-being/quality of life (EUROHIS-QOL-8 (QOL)), at prenatal and postnatal periods.
- ▶ Differences in the psychological experience of pregnancy (Pregnancy and Motherhood Attitudes Scale (PMAS)) between the two trial arms, only at the prenatal period.
- ▶ Differences in maternal/paternal attachment (Antenatal Attachment Scale; Postnatal Attachment Scale) between the two trial arms, at prenatal and postnatal periods.

For the child

- ▶ Differences in child development (physical/sensori-motor, psychosocial) between the two trial arms (Baby form; Child form including The Modified Mary Sheridan Scale; Child Behaviour Checklist for Ages 1.5–5 (CBCL 1.5–5); parent version).

For PC providers

- ▶ Effect of the training period, namely on providers' knowledge on the TP model, self-efficacy and intention to use it to better support children and families (PKQ).

About the CeG! intervention

- ▶ Comparison of acceptability, satisfaction and feasibility of delivering the CeG! with respect to usual care (Satisfaction Questionnaire (SQ); PC providers/parents' version).
- ▶ Evaluation of costs associated with delivering the CeG!

Sample size calculation

The primary outcome will be analysed as the mean change of the PSOC score between the first month and the 18-month well-child visit in each parent. There will be an average change in each arm, and the effect of the intervention will be the difference between these averages.

A sample size of parents of 18 babies (fathers and/or mothers) per cluster, in 12 clusters, was calculated, considering a power of 80%, a type I error of 5%, an intra-cluster correlation coefficient=0.03⁴⁸ and a medium effect size of 0.5.⁴⁹

All calculations are based on the formula for comparison of two means in CRT and assume a 20% loss to follow-up. We used the University of California San Francisco (UCSF) Clinical and Translational Science




Figure 1 Data collection schematic.

CeG!, Crescer em Grandel; CRT, cluster-randomised trial; PCP, primary care provider; Pre-PKQ, Pre-test Practice and Knowledge Questionnaire; Post-PKQ, Post-test Practice and Knowledge Questionnaire. Note: The study was interrupted on March 2020 due to the COVID-19 pandemic. CRT (recruitment of expecting parents) will start only when the PC visits are reinitiated in Portugal on a face-to-face basis.

Institute—Sample Size Calculators, available at <http://www.sample-size.net>.

Since the average number of births per unit, per year, is approximately 150⁵⁰ and assuming conservatively that 50% of parents choose to participate in the research project, we estimate that recruitment can be completed in 3 months, enrolling a minimum of one to two participants a week, per PC unit.

Assignment/randomisation

Randomisation units are clusters—PC units—that will be allocated (Microsoft Excel random number generator) to intervention or control groups using simple randomisation in a 1:1 ratio, stratified by type of institution (custom healthcare units and family health units). The randomisation list will be generated by an independent statistician.

Allocation concealment will be ensured by the following procedures: the PI will assign a meaningless random alphabet letter to each unit as participation forms are received; the final anonymised list of the participating units will be sent to the independent statistician so he can blindly allocate units to each trial arm and then return allocation information to the PI.

As a CRT, only limited blinding could be enforced. Although it is not possible to blind participants and the research team to the trial arm of each assigned unit, consent will not identify other study arms, avoiding participants to know they are in a trial.⁴³ The study statistician will remain blinded during the study period until the database is locked, all analyses are completed and the study unblinded.

Data collection, management and analysis

Assessment

Although the leaflets are designed to accompany pregnancy and the development of children from 0 to 6 years of age, in this study it is chosen to include only assessment points until the child reaches 18 months of age due to: the time available for the project, namely for data collection; a higher concentration of the number of TP in the first 18 months of the child's life, as well as the number of well-child visits; also the first 18 months as a phase

of significant family adaptation with the birth of a new element.

Parents' enrolment and baseline assessment will be done at the first visit of their second trimester of pregnancy (usually performed between 14 and 20 weeks of pregnancy⁴¹) at the PC unit after the study begins (after cross-sectional study in the control arm; after the training period in CeG! arm). Once individual informed consent is obtained, a PC provider in each arm will deliver parents a self-completion sociodemographic questionnaire to collect baseline data, namely: age, gender, marital status, level of education, family's elements and income perception, occupation, support network, among others. Two additional tools assessing couple dynamics (RDAS) and health-related quality of life (QOL) will be delivered.

Relevant sociodemographic information on eligible individuals who decide not to enrol in the study will be anonymously collected. In addition to these data, the provider involved in the recruitment will ask decliners about the reason for the decline.

Multiple assessment points will continue until 18 months well-child visit (table 3), using the instruments below—all validated for the Portuguese population (except forms developed for the study).

- ▶ PSOC—this is a self-completion questionnaire addressed to parents (Cronbach alpha of 0.76⁵¹) designed to assess self-perception of parental competence regarding two dimensions: satisfaction and efficacy. The PSOC consists of 16 items (9 for satisfaction and 7 for efficacy) answered on a 5-point scale ranging from 'strongly disagree' to 'strongly agree'. Results are obtained by adding the quotations of individual items, after the inversion of some; a higher result indicates higher levels of confidence in parenting capacities. The total score ranges from 16 to 80 (no cutoffs).
- ▶ Sociodemographic and Professional Questionnaire—demographic and professional characteristics (eg, number of years of practice after vocational training, currently training residents) will be examined for PC providers at each unit. The instrument

- was developed for the study and will be applied after the study briefing, along with the Pre-test PKQ.
- ▶ Sociodemographic Questionnaire for parents—demographic, lifestyle and pregnancy questionnaire. The instrument was developed for the study.
 - ▶ Baby form; Child form—data on birth and baby/child clinical characteristics (weight, length, Apgar index, feeding and health status, including The Modified Mary Sheridan Scale) collected by PC provider. The instruments were developed for the study.
 - ▶ RDAS⁵²—evaluates couple relationship adjustment; the scale is organised in three subscales: Dyadic Consensus, Dyadic Satisfaction and Dyadic Cohesion.
 - ▶ QOL⁵³—evaluates the perception of the quality of life of adults (physical, psychological, social and environmental domains).
 - ▶ DASS-21⁵⁴—negative emotional states evaluation scale, short form of the DASS-42,⁵⁵ with 21 items distributed by three factors: depression, anxiety and stress.
 - ▶ PMAS⁵⁶—evaluates the psychological experience of pregnancy organised in seven dimensions: the imagined son, good mother, pregnancy as a factor of change/personal growth, difficult aspects of pregnancy/maternity, relationship with one's own mother, husband/companion support and body image and need of dependency. It will be performed only by the mother of the child.
 - ▶ Attachment Scale (antenatal and postnatal)^{57,58}—evaluates two dimensions: bonding quality and the time spent in the bonding mode or intensity of concern (maternal/paternal version).
 - ▶ FES^{59,60}—examines each family member's perceptions of the family in three ways—as it is (real), as it would be in a perfect situation (ideal) and as it will probably be in new situations (expected). We will use the relation dimension of the FES.⁶¹ This dimension includes three subscales: cohesion, expressiveness and conflict.
 - ▶ PSI⁶²—measures the level of parental stress specifically in the postnatal period.
 - ▶ CBCL 1.5–5⁶³—is part of the Achenbach System of Empirically Based Assessment⁶⁴ that offers a comprehensive approach to assessing adaptive and maladaptive functioning. The CBCL evaluates the overall functioning of children and young people and has been described as one of the most comprehensive and psychometrically sound measures for parent report of infant and toddler social-emotional development.⁶⁵
 - ▶ SQ—a satisfaction questionnaire related to satisfaction with care, parents–providers relationship, research involvement, acceptability and perspectives on intervention effectiveness will be organised (closed+open questions). It will be applied to clinical staff and parents (PC providers/parents' version).

Data collection and management

We will collect study data via longitudinal parents' assessments, through self-completion questionnaires,

administered mainly after each healthcare visit, as shown in [table 3](#).

When a parent agrees to participate, a PC provider will register the name in an electronic platform developed for the purpose, and a code including the letter randomly allocated to the PC unit will be automatically assigned. Identification will be kept until the end of the data analysis. Only the clinical staff from each unit will know the parents who participate in the study.

Except for the first assessment of the study (on paper support), questionnaires will be completed online (via email correspondence and phone message). Data will be collected through the electronic platform, which will allow the automatic submission of questionnaires to the parents in the timings defined, facilitating the entire data collection and storage process. Regular email reminders will be sent to parents in case of non-response; 48 hours before a clinical visit, an email will also be sent to each participants' PC providers.

The electronic platform was developed in full compliance with the new General Data Protection Regulation. The servers will be housed in the European Union, certified with compliance with all safety standards, and with regular backups and maintenance of duplicate databases in geographically distinct locations, protecting the information from any eventual loss. An administrator profile will be created only for the PI. The registration of the email/phone number of the users will be encrypted, with the decryption key only in possession of the PI. The individuals' visible identifier will be the random code assigned to them at the beginning of the study. Only the PI's profile will have access to the control panel of the study's evolution, manually register questionnaires on paper or reinforce the sending of emails to request the completion of missing questionnaires. After the end of the study, all data will be permanently erased from the servers. The data will be stored for 5 years on an external encrypted disk, which requires the use of a password.

If parents cannot answer electronically, there will also be a paper version of the study forms. In this case, the questionnaires must be completed in the unit. There will be a closed return box in each PC unit so that parents can place their completed coded questionnaires, ensuring their privacy and anonymity. At each consultation, PC providers should remind these participants of the placement of the questionnaires in the box available for this purpose. A team investigator will collect the questionnaires frequently from each unit. If necessary, the PI will contact the clinic representative, signalling a missing questionnaire for a specific code. Those site visits will also allow the study investigators to monitor the enrolment process, intervention delivery and protocol adherence. Email greetings on particular dates (eg, child's birthday, Christmas) will be sent to reinforce parents' adherence.

At each visit, information on attendance to health visit, eventual dropouts and reasons for dropout will be collected by a PC provider directly on the electronic platform. In each PC unit, there is also a form for the health

staff to record any other relevant occurrence during the trial.

To test the excellent functioning of the electronic platform and get feedback from PC providers and parents on the trial process, we will previously conduct a small pilot (without intervention) with three to four expecting parents within a PC unit that does not enter the CRT.

Statistical methods

The intention-to-treat principle will be adopted. Participants' data will not be included in any analyses if they withdraw consent. The study statistician will not be involved in intervention delivery or data collection and will remain blinded until all analyses are completed.

Initially, a descriptive statistical analysis of the variables studied for participants at each PC unit will be performed by study group (CeG! and control) and strata (type of unit: custom healthcare units and family health units). Categorical variables will be described by absolute and relative frequencies; continuous variables by means (and standard deviation) if they are normally distributed or by medians (and interquartile range) if not; ordinal variables will be presented by medians (and interquartile range).

Recognising that cluster randomisation might not achieve balance on descriptive variables between individuals at baseline, adjusting for these variables will be considered in the model. Therefore, both groups' outcomes will be compared using linear-mixed effect models adjusting for baseline measurements with a random intercept for the PC unit level to account for clustering effect. Possible missing patterns will be handled by multiple imputation and the linear-mixed models for the longitudinal analysis. Statistical significance will be set at $p < 0.05$.

Additionally, to determine if specific subpopulations respond differently to CeG! intervention, we will perform subgroup analyses based on baseline characteristics. This analysis will be conducted among gender-difference of parents, different types of family structures, families expecting their first child versus families with one or more children, mothers with above high school education versus mothers with high school or below and families experiencing financial strain versus families who are not. We will test for significance of subgroups using statistical tests for interaction. If required, multiple comparisons adjustments will be performed through alpha adjustment.

Reporting of results will follow the principles of Consolidated Standards of Reporting Trials statements and the extension for CRT.^{43 66}

Rates and reasons for missing data will be assessed and reported. Several aspects will be considered to ensure data consistency: verifying variables type, list all the possible values for each variable, check spelling errors and improper entries and investigate missing data. The out-of-range, unreliable or invalid values must be evaluated carefully. Furthermore, assessing the best method to deal with missing data must be probed. One or more imputation algorithms, such as the mean or median

values, or deep learning methods, must be implemented and assessed using different metrics such as distributional accuracy.

Process evaluation

Regular site visits will allow the study researchers to monitor the enrolment process, intervention delivery and protocol adherence. To enable early correction of any errors during data collection, there will be close supervision of data on the platform with subsequent contact with unit representative wherever needed; platform chat is available for any doubts during data insertion. Parents will be instructed to contact their healthcare team whenever any questions arise; a study's telephone contact and an email will be available.

We anticipate no harm arising from the intervention's implementation; therefore, no data monitoring committee was established. If we become aware of any harm or other adverse events, either through study data or other avenues, we will review and address these according to standard institutional processes, in consultation with the relevant ethics committee. We will also file regular reports on trial progress with the ethics committee. We have not devised any stopping guidelines, although we intend to carry out mid-term analyses of trial outcome data, according to the different measurement points. If any relevant health information is identified during the analysis of the participants' data, it shall be communicated to the attending family physician who will act in accordance.

Program costs

To help inform potential financing and adoption of the CeG! intervention, compared with usual care, we intend to estimate the incremental costs to deliver the intervention (eg, costs in printing leaflets, training costs) and overhead costs minus costs attributable only to research activities (eg, PC unit study monitoring visits).

ETHICS AND DISSEMINATION

This trial is conceived in accordance with the fundamental ethical principles of autonomy, beneficence, justice and non-maleficence and will be conducted under the principles expressed in the Declaration of Helsinki.

Research ethics approval

Approved by the Ethics Committee of the Regional Health Administration of Lisbon and Tagus Valley (Portugal); registration number: 013/CES/INV/2019 (date: 18 November 2019).

Consent

When adopting a CRT, it is not usually feasible to obtain participants' consent to randomisation. Instead, consent to randomisation is typically provided by a surrogate decision-maker at the cluster level. We obtained agreement to randomisation from a representative from each unit (the executive director of each group of health

centres involved). We will seek PC care providers' and eligible patients' written informed consent to participate in the study's data collection.

Privacy and confidentiality

Data confidentiality and anonymity will be ensured by the PI in full compliance with the new General Data Protection Regulation, both during the implementation phase of the study and in any resulting presentations or publications. The results will only be used in the framework of this study. All personal information will be safeguarded by assigning a meaningless random code, which will link the answers, without identification, across all the time points. Finally, restricted access to participant-level data and files will be ensured.

Dissemination policy

Members of the scientific community and the public will be able to access the full study protocol at <http://www.isrctn.com>. Substantive modifications to this protocol will be communicated to the relevant staff at participating units during regular communications; to others via written summaries, published modifications to the trial profile at <http://www.isrctn.com> or via statements in scientific papers arising from the study.

Findings from this study will be submitted for publication in peer-reviewed journals. The study results will also be shared with health professionals and other participants from participating units and through scientific conferences targeting primary and secondary care providers, and research community more widely. The full protocol will also be available on the PI's doctoral thesis. Once the study is completed and primary manuscripts published, data will be available on request to the PI.

Author affiliations

¹Center for Health Technology and Services Research (CINTESIS), Faculty of Medicine of the University of Porto, Porto, Portugal

²Department of Community Medicine, Information and Health Decision Sciences (MEDCIDS), Faculty of Medicine of the University of Porto, Porto, Portugal

³Universidade Católica Portuguesa, Faculty of Education and Psychology, Research Centre for Human Development, Porto, Portugal

⁴Unidade de Saúde Familiar AlphaMouro, Sintra, Portugal

⁵Instituto Politécnico de Viana do Castelo, Viana do Castelo, Portugal

Twitter Carlos Martins @mgfamiliarnet

Acknowledgements The authors would like to thank several persons: to Miguel Julião for his constant and unwavering support; to João Figueiredo and XERPA for developing the electronic platform; to Catarina Samorinha, and finally, to Fundação Brazelton/Gomes-Pedro in the person of João Gomes-Pedro, Ana Teresa Brito, and Joana Tinoco de Faria, for always supporting our project and believing in its future implications in children's and parents' lives. To the following colleagues for the collaboration in the leaflets' initial development: António Mendes dos Santos, Bertina Oliveira, Cristina André, Filipa Nóbrega, Jerusa Oliveira, Marcos Cabral, Mário Cruz, Maria Ana Sobral, Raquel Pedro, Susana Medeiros, Tânia Lima, Vânia Duarte, Vera Teles.

Contributors FF conceived the overall study, drafted the protocol and registered the trial. MRX contributed to the development of the study protocol and helped monitor the study process. CM provided overall guidance in cluster-randomised trial design. AT performed sample size calculations and statistical guidance in the development of the protocol. FF and JV participated in the trial methods adaptation

to the local context and have responsibilities for day-to-day running of the trial. All authors read and approved the final version manuscript.

Funding FF has a grant funded by national funds of the Ministry of Science, Technology and Higher Education (Portugal) - FCT (Fundação para a Ciência e a Tecnologia) grant: reference PD/BD/132860/2017 (since 2016). The Nestlé Company (Portugal) sponsored the printing of the leaflets that will be used to carry out this study, as an unrestricted grant. This article was supported by National Funds through FCT within CINTESIS, R&D Unit (reference UIDB/4255/2020). Funding sources had no role in the design or content of this study and will have no role during its execution, data analyses or decision to submit results.

Competing interests MRX is member of the Administrative Board (non-executive) of the Fundação Brazelton/Gomes-Pedro para as Ciências do Bebê e da Família, Portugal.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Filipa Fareira <http://orcid.org/0000-0002-7312-2178>

Maria Raul Xavier <http://orcid.org/0000-0002-2223-2000>

Julia Velte <http://orcid.org/0000-0003-1185-2500>

Andreia Teixeira <http://orcid.org/0000-0003-1199-2220>

Carlos Martins <http://orcid.org/0000-0001-8561-5167>

REFERENCES

- 1 OECD. Doing better for families 2011.
- 2 Petch J, Halford WK. Psycho-education to enhance couples' transition to parenthood. *Clin Psychol Rev* 2008;28:1125–37.
- 3 Cowan PA, Cowan CP. Normative family transitions, couple relationship quality, and healthy child development. In: Walsh F, ed. *Normal family processes: growing diversity and complexity*. New York: Guilford Press, 2012: 428–51.
- 4 Bowlby J. *A secure base parent child attachment and healthy human development*. New York: Basic Books, 1988.
- 5 Prevention FP. The appropriate target of infant psychotherapy. *Infant Mental Health Journal* 1998;19:124–50.
- 6 Shonkoff JP, Phillips DA. *From neurons to neighborhoods: the science of early childhood development*. Washington, DC, United Kingdom: National Academy Press, 2000.
- 7 University CotDcaH. From best practices to breakthrough impacts: a science-based approach to building a more promising future for young children and families 2016.
- 8 Recomendação do Conselho da Europa. *Política de apoio Parentalidade Positiva*. Lisboa, 2006.
- 9 Barlow J, Bergman H, Kornør H, et al. Group-Based parent training programmes for improving emotional and behavioural adjustment in young children. *Cochrane Database Syst Rev* 2016;8:CD003680.
- 10 Barlow J, Smailagic N, Huband N, et al. Group-Based parent training programmes for improving parental psychosocial health. *Cochrane Database Syst Rev* 2014;5:CD002020.
- 11 Barlow J, Smailagic N, Bennett C, et al. Individual and group based parenting programmes for improving psychosocial outcomes for teenage parents and their children. *Cochrane Database Syst Rev* 2011;3:CD002964.
- 12 Furlong M, McGilloway S, Bywater T, et al. Cochrane review: behavioural and cognitive-behavioural group-based parenting programmes for early-onset conduct problems in children aged 3 to 12 years (review). *Evid Based Child Health* 2013;8:318–692.
- 13 Coleman PK, Karraker KH. Self-efficacy & parenting quality: Findings & future applications. *Developmental Review* 1998;18:47–85.
- 14 Johnston C, Mash EJ. A measure of parenting satisfaction and efficacy. *J Clin Child Psychol* 1989;18:167–75.
- 15 Jones TL, Prinz RJ. Potential roles of parental self-efficacy in parent and child adjustment: a review. *Clin Psychol Rev* 2005;25:341–63.
- 16 Okagaki L, Bingham GE. Parents' social cognitions and their parenting behaviors. In: Luster T, Okagaki L, eds. *Parenting: an*

- ecologic perspective Monographs in parenting series. 2nd ed.. New Jersey: Lawrence Erlbaum Associates, Inc, 2005: 3–33.
- 17 Roskam I, Meunier J. The determinants of parental childrearing behavior trajectories: the effects of parental and child time-varying and time-invariant predictors. *International Journal of Behavioral Development* 2012;36:186–96.
 - 18 Salonen AH, Kaunonen M, Astedt-Kurki P, et al. Parenting self-efficacy after childbirth. *J Adv Nurs* 2009;65:2324–36.
 - 19 Ponomartchouk D, Bouchard G. New mothers' sense of competence: predictors and outcomes. *J Child Fam Stud* 2015;24:1977–86.
 - 20 Mihelic M, Morawska A, Filus A. Effects of early parenting interventions on parents and infants: a meta-analytic review. *Journal of Child and Family Studies* 2017;26:1507–26.
 - 21 Davidson Arad B, McLeigh JD, Katz C. Perceived collective efficacy and parenting competence: the roles of quality of life and hope. *Fam Process* 2020;59:273–87.
 - 22 Brazelton Touchpoints center, accessed 2020. Available: <https://www.brazeltontouchpoints.org/about/>
 - 23 Brazelton TB, Sparrow J. *Nurturing Children and Families - Building on the Legacy of T. Berry Brazelton*. Blackwell Publishing Ltd: Sussex, 2010.
 - 24 Sparrow J. Newborn behavior, parent-infant interaction, and developmental change processes: research roots of developmental, relational, and systems-theory-based practice. *J Child Adolesc Psychiatr Nurs* 2013;26:180–5.
 - 25 Gomes-Pedro J. Touchpoints. In: Gomes-Pedro J, ed. *Para Um sentido de coerência Na criança Mem-Martins: Publicações Europa-América, Lda*, 2005: 249–61.
 - 26 Zero to Three. Tuning in: parents of young children tell us what they think, know and need 2016.
 - 27 Young KT, Davis K, Schoen C, et al. Listening to parents. A national survey of parents with young children. *Arch Pediatr Adolesc Med* 1998;152:255–62.
 - 28 Bethell C, Peck C, Abrams M. Partnering with parents to promote the healthy development of young children enrolled in Medicaid results from a survey assessing the quality of preventive and developmental services for young children enrolled in Medicaid in three states. *New York* 2002:1–53.
 - 29 Patterson J, Mockford C, Barlow J, et al. Need and demand for parenting programmes in general practice. *Arch Dis Child* 2002;87:468–71.
 - 30 Kane GA, Wood VA, Barlow J. Parenting programmes: a systematic review and synthesis of qualitative research. *Child Care Health Dev* 2007;33:784–93.
 - 31 Dretzke J, Davenport C, Frew E, et al. The clinical effectiveness of different parenting programmes for children with conduct problems: a systematic review of randomised controlled trials. *Child Adolesc Psychiatry Ment Health* 2009;3:7.
 - 32 Kendrick D, Mulvaney CA, Ye L, et al. Parenting interventions for the prevention of unintentional injuries in childhood. *Cochrane Database Syst Rev* 2013;3:CD006020.
 - 33 Thomas RE, Baker PRA, Thomas BC, et al. Family-Based programmes for preventing smoking by children and adolescents. *Cochrane Database Syst Rev* 2015;2:CD004493.
 - 34 Liyana Amin NA, Tam WWS, Shorey S. Enhancing first-time parents' self-efficacy: a systematic review and meta-analysis of universal parent education interventions' efficacy. *Int J Nurs Stud* 2018;82:149–62.
 - 35 Shah R, Isaia A, Schwartz A, et al. Encouraging parenting behaviors that promote early childhood development among caregivers from low-income urban communities: a randomized static group comparison trial of a primary care-based parenting program. *Matern Child Health J* 2019;23:39–46.
 - 36 Shah R, Kennedy S, Clark MD, et al. Primary care-based interventions to promote positive parenting behaviors: a meta-analysis. *Pediatrics* 2016;137. doi:10.1542/peds.2015-3393. [Epub ahead of print: 19 Apr 2016].
 - 37 Spoth R, Redmond C. Research on family engagement in preventive interventions: towards improved use of scientific findings in primary prevention practice. *J Prim Prev* 2000;21:267–84. doi:10.1023/A:1007039421026
 - 38 Heinrichs N, Bertram H, Kuschel A, et al. Parent recruitment and retention in a universal prevention program for child behavior and emotional problems: barriers to research and program participation. *Prev Sci* 2005;6:275–86.
 - 39 Love JM, Kisker EE, Ross C, et al. The effectiveness of early head start for 3-year-old children and their parents: lessons for policy and programs. *Dev Psychol* 2005;41:885–901.
 - 40 Saúde D-Gda. Programa Nacional de Saúde Infantil e Juvenil. In: Saúde D-Gda, ed. Lisboa: Direção-Geral da Saúde, 2013: 1–116.
 - 41 Direção-Geral da Saúde. *Programa Nacional para a Vigilância da Gravidez de Baixo Risco*. Lisboa: Direção-Geral da Saúde, 2015: 5–103.
 - 42 Chan A-W, Tetzlaff JM, Altman DG, et al. Spirit 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013;158:200–7.
 - 43 Campbell MK, Piaggio G, Elbourne DR, et al. Consort 2010 statement: extension to cluster randomised trials. *BMJ* 2012;345:e5661.
 - 44 Pisco L. [Primary Healthcare Reform in Portugal on two fronts: autonomous family healthcare units and management of groupings of Health Centers]. *Cien Saude Colet* 2011;16:2841–52.
 - 45 Brazelton Touchpoints Center. *Touchpoints reference guide and participant training materials*. Boston: Brazelton Touchpoints Center, 2016.
 - 46 Brazelton Touchpoints Center. *Touchpoints reference guide for health care providers*. Boston: Brazelton Touchpoints Center, 2012.
 - 47 Fundação Brazelton/Gomes-Pedro, 2020. Available: <https://www.fundacaobgp.com/pt>
 - 48 Hemming K, Eldridge S, Forbes G, et al. How to design efficient cluster randomised trials. *BMJ* 2017;358:j3064.
 - 49 Cohen J. *Statistical power analysis for the behavioral sciences*. In: Hillsdale NJ, ed. 2nd edn. Lawrence Erlbaum Associates, 1988.
 - 50 Portuguese National Health Service. BI-CSP platform. Available: <https://www.sns.gov.pt/noticias/2018/10/01/plataforma-bi-csp-2/2019>
 - 51 Seabra-Santos MJ, Major S, Pimentel M. Escala de Sentido de Competência parental (PSOC): estudos psicométricos. *Avaliação Psicológica* 2015;14:97–106.
 - 52 Pereira MG. *Revised dyadic adjustment scale – Versão de investigação*. Braga, Portugal: Universidade do Minho, 2003.
 - 53 Pereira M, Melo C, Gameiro S. Estudos psicométricos da versão em Português Europeu do índice de qualidade de vida EUROHIS-QOL-8. *Laboratório de Psicologia* 2011;9:109–23.
 - 54 Pais Ribeiro J, Honrado A, Leal I. Contribuição para O estudo da adaptação portuguesa das escalas de ansiedade, depressão E stress (EADS) de 21 itens de Lovibond E Lovibond. *Psicologia, Saúde e Doenças* 2004;5:229–39.
 - 55 Lovibond PF, Lovibond SH. The structure of negative emotional states: comparison of the depression anxiety stress scales (DASS) with the Beck depression and anxiety inventories. *Behav Res Ther* 1995;33:335–43.
 - 56 Xavier MR, Paúl MC. *Construção E validação de uma escala de atitudes sobre a gravidez E a maternidade*. Avaliação Psicológica: Formas e Contextos, 1996: 419–24.
 - 57 Camarinho AP, Justo J. Padrões de vinculação pré-natal. Contributos para a adaptação da maternal and paternal antenatal attachment scale em casais durante O segundo trimestre de gestação Na região Centro de Portugal. *Revista Portuguesa de Pedopsiquiatria* 2010;28:7–22.
 - 58 Camarinho AP. *Vinculação pré-natal E organização psicológica do homem E da mulher durante a gravidez: relação CoM O tipo de parto E CoM a patologia obstétrica DOS II E III trimestres de gestação*. Universidade de Lisboa, 2011.
 - 59 Santos M, Fontaine A. Avaliação do ambiente familiar por crianças e pré-adolescentes: alguns aspectos da adaptação da FES de Moos & Moos. In: Almeida L, Simões M, Gonçalves M, eds. *Avaliação psicológica: formas E contextos*, 1995: 421–30.
 - 60 Matos PM, Fontaine M. *Family environment scale – FeS*. Adaptação portuguesa.: Faculdade de Psicologia e de Ciências da Educação da Universidade do Porto, 1992.
 - 61 Moos RH. Conceptual and empirical approaches to developing family-based assessment procedures: resolving the case of the family environment scale. *Fam Process* 1990;29:199–208.
 - 62 Gomez R, Leal I. Stress parental no período pós-parto: Adaptação do parental stress inventory para a população portuguesa. *Psicologia* 2009;50:361–72.
 - 63 Achenbach T, Rescorla L, Dias P. Manual do Sistema de Avaliação Empiricamente Validado (ASEBA) para O Período Pré-Escolar E Escolar 2014.
 - 64 Achenbach TM, Rescorla L. *Manual for the ASEBA school-age forms & profiles : an integrated system of multi-informant assessment*. Burlington, Vt.: ASEBA, 2001.
 - 65 Pontoppidan M, Niss NK, Pejtersen JH, et al. Parent report measures of infant and toddler social-emotional development: a systematic review. *Fam Pract* 2017;34:127–37.
 - 66 Schulz KF, Altman DG, Moher D, et al. Consort 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMJ* 2010;340:c332.