

# Transcriptome Analysis of the Chicken Follicular Theca Cells with miR-135a-5p Suppressed

Yan Zhou,<sup>\*,†,1</sup> Jie Liu,<sup>\*,†,1</sup> Qiuxia Lei,<sup>\*,†</sup> Haixia Han,<sup>\*,†</sup> Wei Liu,<sup>\*,†</sup> Tang Cunwei,<sup>‡</sup> Fuwei Li,<sup>\*,†,2</sup> and Dingguo Cao<sup>\*,†,2</sup>

<sup>\*</sup>Institute of Poultry Science, Academy of Agricultural Sciences of Shandong Province, Jinan, 250023, Shandong, China,

<sup>†</sup>Poultry Breeding Engineering Technology Center of Shandong Province, Jinan, 250023, Shandong, China, and <sup>‡</sup>Pingyi Yike poultry Co., Ltd, Linyi, 276000, Shandong, China

**ABSTRACT** As a class of transcription regulators, numerous miRNAs have been verified to participate in regulating ovary follicular development in chickens (*Gallus gallus*). Previously we showed that gga-miR-135a-5p has significant differential expression between high and low-yield chicken ovaries, and the abundance of gga-miR-135a-5p is significantly higher in follicular theca cells than in granulosa cells. However, the exact role of gga-miR-135a-5p in chicken follicular theca cells is unclear. In this study, primary chicken follicular theca cells were isolated and then transfected with gga-miR-135a-5p inhibitor. Transcriptome sequencing was performed in chicken follicular theca cells with or without transfection. Differentially expressed genes (DEGs) were analyzed using bioinformatics. A dual-luciferase reporter assay was used to verify the target relationship between gga-miR-135a-5p and predicted targets within the DEGs. Compared with the normal chicken follicle theca cells, 953 up-regulated and 1060 down-regulated genes were detected in cells with gga-miR-135a-5p inhibited. The up-regulated genes were significantly enriched in Gene Ontology terms and pathways involved in cell proliferation and differentiation. In chicken follicular theca cells, Krüppel-like factor 4 (KLF4), ATPase phospholipid transporting 8A1 (ATP8A1), and Complexin-1 (CPLX1) were significantly up-regulated when the expression of gga-miR-135a-5p was inhibited. In addition, KLF4, ATP8A1, and CPLX1 confirmed as targets of gga-miR-135a-5p by using a dual-luciferase assay *in vitro*. The results suggest that gga-miR-135a-5p may involve in proliferation and differentiation in chicken ovarian follicular theca cells by targeting KLF4, ATP8A1, and CPLX1.

## KEYWORDS

*Gallus gallus*  
ovarian theca cells  
gga-miR-135a-5p  
transcriptome sequencing

MicroRNAs (miRNAs) are a class of small noncoding RNAs of about 18-24 nucleotides in length (Lee *et al.* 1993; Bartel 2004) that function as regulators in post-transcriptional gene expression by targeting sequence-specific sites in the 3'-UTR of mRNA (Grosshans and Slack 2002; Ambros and Chen 2007; Krol *et al.* 2010). Studies indicate that miRNAs play key roles in ovarian

follicular development and function, including the formation of primordial follicles, follicular recruitment and selection, follicular atresia, oocyte-cumulus cell interaction, granulosa or theca cell function, and luteinization (Hawkins and Matzuk 2010; Hasuwa *et al.* 2013; Kang *et al.* 2013; Maalouf *et al.* 2015; Zhang *et al.* 2019). Mouse miR-145 and miR-181a (Yan *et al.* 2012; Zhang *et al.* 2013), bovinelet-7 families *et al.* (Salilew-Wondim *et al.* 2014), buffalo miR-210 (Shukla *et al.* 2018), porcine miR-26b *et al.* (Lin *et al.* 2012) and chicken miR-107 (Miao *et al.* 2016) were all validated to be involved in granulosa cell proliferation, apoptosis and other cell function. In ovary theca cells, studies have showed that miRNAs also play an important role in cell function. In bovine, miR-640 and miR-526b\* (Sohel *et al.* 2013), and bta-miR-335 (Gebremedhn *et al.* 2015) were proved to express more abundant in theca cells. Several predicted miRNA target interactions miR-155/miR-222-ETS1miR-199a-5p-JAG1, miR-155-MSH2 and miR-199a-5p/miR-150/miR-378-VEGFA in theca cells were putatively involved in follicular atresia (Donadeu *et al.* 2017). Another study showed that abundance of MIR-221 was

Copyright © 2020 Zhou *et al.*

doi: <https://doi.org/10.1534/g3.120.401701>

Manuscript received August 5, 2020; accepted for publication September 3, 2020; published Early Online September 8, 2020.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Supplemental material available at figshare: <https://doi.org/10.25387/g3.12899993>.

<sup>1</sup>These authors contributed equally to this work.

<sup>2</sup>Corresponding authors: Poultry Institute of Shandong Academy of Agricultural Science, No.202 North of Industry Road, Jinan, Shandong, P. R. China.250100. E-mail:jqsyzs@163.com; lifuwei1224@163.com

■ **Table 1 Information of the primers used for quantitative real-time PCR validation**

Genes	T <sub>m</sub> (°C)	Sequences (5'-3')	Products (bp)
IFIT5	60	F: GAAGAACCCAACAACCCAGA R: GTTTCAGTGCACGCTTCAGA	127
IFI6	60	F:AAGCCGGTTTCACTTCCTCT R: CTTTGGCACCCATTTCTTGT	135
SERPINB2	60	F:GCAGATCAGATGGCAAAGGT R: AGGCACGGTGATTTGATAGG	131
NSG1	60	F: TTTAGCGAGAAGAGCACAA R: TGACCACAACCTTGTCAGGA	119
GRK5	60	F: TGCTCCTACCAAAGATGTG R: CCAGCATTCTGCTTCTAGC	120
CXCL12	60	F: AGGGCCAACATTAAGCACCTC R: TTTAGCTTGGGATCAATGCACAC	110
FABP7	60	F: GGGAACGTGACTAAGCCCA R: TGTCTCCATCCAGGGTCACAAC	172
KLF4	60	F: CTGCGCAAGACCTACACCA R: AGTTCATCAGATCGGGCAAACCTTC	123
ATP8A1	60	F:TCTAACTGTGTGTTGAAGGCTGGA R: ATCGGGTGCCATAGGGATGA	145
STK35	60	F: GGATCGACCTGATGCCTTTGA R: CGTTGCACGTAATGGAGATGATG	137
FRMD4A	60	F: AAGGACAACGCCACCATTGAG R: TGCAGCCAGTTCAAACACCAC	102
OASL	60	F:AGATGTTGAAGCCGAAGTACCC R: CTGAAGTCCTCCCTGCCTGT	106
β-actin	60	F: TAAGCGTGTTATCATCTC R: GGGACTTGTATATTCT	86

66.sixfold greater ( $P < 0.001$ ) in TCs than in GCs in bovine large follicles, and thecal MIR-221 expression is increased by FGF9 (Robinson *et al.* 2018). In sheep, northern analyses showed that the expression of miR-199a-3p, miR-125b, miR-145, miR-31, miR-503, miR-21 and miR-142-3p in theca cells were higher than those in granulosa cells (McBride *et al.* 2012). In woman, two miRNAs-miR-92a and miR-92b were validated to be significantly downregulated in theca cells and might be involved in the pathogenesis of PCOS (Lin *et al.* 2015). In addition, miR-26a-5p was verified to facilitate theca cell proliferation in chicken ovarian follicles by targeting TNRC6A (Kang *et al.* 2017; Wu *et al.* 2019).

MiR-135a were proved to be overexpressed in GCs from PCOS patients, a study showed that miR-135a repressed ESR2 expression in GCs, which further inhibited CDKN1A expression, promoted GC proliferation and repressed GC apoptosis (Song *et al.* 2019). Furthermore, another finding indicates that miR-135a promotes


apoptosis and the DNA damage response in GCs in PCOS, likely via VEGFC signaling (Wei *et al.* 2020). To our knowledge, the function of gga-miR-135a-5p in chicken follicular theca cells has not been reported. Our previous study showed that gga-miR-135a-5p was differentially expressed, with a fold-change of 8.93, in high compared low-yield ovaries of a Chinese indigenous chicken breed, and it was expressed significantly higher in follicular theca cells than in granulosa cells (unpublished). Therefore, the overall results indicated that miR-135a-5p may play an important role in chicken follicular theca cells.

In this study, we first transfect the gga-miR-135a-5p inhibitor into chicken follicular theca cells. RNA sequencing was performed for transcriptome analysis using the Illumina HiSeq sequencing platform. A dual-luciferase report assay was used to verify the regulatory relationship between miR-135a-5p and the predicted differentially expressed gene (DEG) targets. We present evidence

■ **Table 2 Primers for the construction of dual-luciferase reporter plasmid**

Genes	Primers (5'-3')	Products (bp)
ATP8A1	ATP8A1_WT_F: GCGGCTCGAGAGCACTTGTAGTTCTGATG ATP8A1_WT_R: AATGCGGCCGCACATAGCGACCACCTTTCTG ATP8A1_mut_F: TAGAACTGTTCCGGTATAATGCACTAATATTGTTT ATP8A1_mut_R: AGTGCATTATACCGAACAGTTCTACAATAAAGAA	687
CPLX1	CPLX1_WT_F: GCGGCTCGAGTGAACAAGACCAAGATAA CPLX1_WT_R_R: AATGCGGCCGCACAGAGAGAGAAAACAAGC CPLX1_mut_F: TTTAGAGATTCCGGTATTGAAATCAGACTGCAATA CPLX1_mut_R: TGATTTCAATACCGAATCTCTAAAGATAGGTATG	781
KLF4	KLF4_WT_F: GCGGCTCGAGATCACCTCGCCTTACACAT KLF4_WT_R: AATGCGGCCGCTCAACCACTGACCAACATT KLF4_mut_F: CCTTCTAATTCCGGTATACAATGTTAAAAGAGGA KLF4_mut_R: ACATTGTATACCGAATTAGAAGGAAGAAAAAAA	1033

Note: WT, wild type; MUT, mutant type.



**Figure 1** The relative expression level of gga-miR-135a-5p in NG, TG and NC. \*\*  $P < 0.01$ . NG: follicular theca cells; TG: inhibitor transfected cells; NC: negative control.

that gga-miR-135a-5p is involved in the biological function of ovarian follicular development. This study provides a scientific basis for a mechanism of gga-miR-135a-5p regulation in the follicular development of poultry.

## MATERIALS AND METHODS

### Ethics statement

All experimental procedures were approved by the Animal Care Committee of the Academy of Agricultural Sciences, Shandong Province, Ji'nan, China. The care and use of experimental animals were carried out in accordance with the Directory Proposals on the Ethical Treatment of Experimental Animals, established by the Ministry of Science and Technology (Beijing, China).

### Birds and tissue harvest

Three single-comb white Leghorn hens were selected randomly from Shandong Poultry Breeding and Engineering Technology Research Center to be used in this study. All birds were reared in an environmentally controlled house. Fresh water and feed were provided according to the Feeding Standard established by the Ministry of Agriculture (Beijing, China). At 40 weeks of age, the F1-F5 follicles were removed carefully and placed in pre-cooled phosphate-buffered saline (PBS) for the next step of theca cell culture.

### Theca cell culture and transfection


The F1-F5 follicle theca layers were separated according to Kang *et al.* (Kang *et al.* 2017). The isolated theca layers were minced to 1mm<sup>3</sup> pieces and digested with collagenaseII (w/v, 0.2%, Gibco, Grand Island, New York, USA) at 37° for 30 min. Remove the supernatant, the cell precipitation digested with collagenaseII (w/v, 0.2%, Gibco) at 37° for 30 min again. The dispersed theca cells were filtrated with a sterilized 200-mesh filter and then centrifuged at 1800× rpm for 10 min. The cell precipitations were washed two times with cell culture medium containing M199

(HyClone, Logan, Utah, USA) supplemented with 10% (v/v) fetal bovine serum (Gibco) and 1% (v/v) penicillin-streptomycin solution (Solarbio, Beijing, China). The cells were then seeded in 24-well plates at a density of 2×10<sup>5</sup> per well and cultured at 37° in an atmosphere of 95% air and 5% CO<sub>2</sub>. The number of viable cells (>90%) was estimated using Trypan blue.


When the cells reached approximately 60–70% confluence, they were transfected with gga-miR-135a-5p inhibitor and a miRNA inhibitor negative control (NC) with Lipofectamine2000 (Invitrogen, Carlsbad, CA, USA) and Opti-MEM (Gibco) according to the manufacturer's instructions. Each transfection was performed at least in triplicate. The transfection efficiency was confirmed by real-time quantitative reverse transcription PCR (qRT-PCR). The primers used for gga-miR-135a-5p amplification were as follows: loop primer 5'-GTCGTATCCAGTGCAGGGTCCGAGGTATTTCGCACTGGAT-ACGACTCACATA -3'; forward: 5'-GCGAGAGGTATGGCTTT-TTATTCC-3', reverse: 5'-GTGCAGGGTCCGAGGT -3'. U6 was used as an internal control: loop primer: 5'-GTCGTATCCAGTGCAGGGTCCGAGGTATTTCGCACTGGATACGACTCACATA -3'; forward: 5'-CACTGGGCCATGCTAATCTTCTC -3'; reverse: 5'-GTGCAGGGTCCGAGGT -3'.

### RNA extraction, cDNA library construction and sequencing

A total of five samples, including two groups of normal chicken follicular theca cells groups T07 and T08 (NG) and 3 group transfected with gga-miR-135a-5p inhibitor T10, T11, and T12 (TG) were used for sequencing. The total RNA of each sample was extracted with Trizol (Aidlab, Beijing, China) according to the manufacturer's instructions. RNA integrity and concentration were checked using an Agilent 2100 Bioanalyzer (Agilent Technologies, Inc., Santa Clara, CA, USA). The mRNA was isolated by using NEBNext Poly (A) mRNA Magnetic Isolation Module (E7490, NEB, Ipswich, MA, USA). The cDNA library was constructed following the instructions of the NEBNext Ultra RNA Library Prep Kit for Illumina (NEB, E7530) and



**Figure 2** The expression level of the top 22 up-regulated genes with the most significant differential expression greater than fourfold between cells with normal and inhibited expression of gga-miR-135a-5p.



**Figure 3** Volcano plot of differentially expressed genes in cells with normal and inhibited expression of gga-miR-135a-5p. The X-axis represents  $\log_2$  (FC) and Y-axis represents  $-\log_{10}$  (FDR). The green dots indicate the down-regulated genes, the black dots indicate the genes with no significant differences, and the red dots indicate up-regulated genes.

NEBNext Multiplex Oligos for Illumina (NEB, E7500). In brief, the enriched mRNA was fragmented into approximately 200nt RNA inserts, which were used to synthesize the first-strand cDNA and the second cDNA. End-repair/dA-tail and adaptor ligation were performed on the double-stranded cDNA. Suitable fragments were isolated by AgencourtAMPure XP beads (Beckman Coulter, Inc.), and enriched by PCR amplification. Finally, the constructed cDNA libraries were sequenced on a flow cell using an Illumina HiSeq sequencing platform.

### Transcriptome analysis using reference genome-based reads mapping

Low quality reads, such as adaptor only, unknown nucleotides  $> 5\%$ , or  $Q_{20} < 20\%$  (percentage of sequences with sequencing error rates  $< 1\%$ ), were removed using *aperl* script. The clean reads filtered from the raw reads were mapped to the chicken genome (*Gallus gallus*, Galgal 4.75) using Tophat2 (Kim *et al.* 2013) software. The aligned records from the aligners in BAM/SAM format were further examined to remove potential duplicate molecules. Gene expression levels were estimated using fragments per kilobase of exon per million fragments mapped (FPKM) values by Cufflinks software (Trapnell *et al.* 2010).

### Identification of DEGs

DESeq and Q-value were employed and used to evaluate differential gene expression between cells expressing gga-miR-135a-5p (NG) and those with gga-miR-135a-5p inhibited (TG). After that, gene abundance differences between those samples were calculated based on the ratio of the FPKM values. The false discovery rate (FDR) control method was used to identify the threshold of the *P*-value in multiple tests in order to compute the significance of the differences. Here, only genes with an absolute value of  $\log_2$  fold change  $\geq 1$  and FDR significance score  $< 0.01$  were used for subsequent analysis.

### Functional annotation

The Database for Annotation, Visualization, and Integrated Discovery (DAVID v6.7) was used to annotate the Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes pathways of the DEGs. The GO includes biological process, molecular function, and cellular component categories. An online software analysis tool ([http://www.lc-bio.cn/overview/12?tools=GO\\_BarPlot](http://www.lc-bio.cn/overview/12?tools=GO_BarPlot)) was used to plot GO functional classification of the unigenes with a GO term hit to view the distribution of gene functions. Finally an online software analysis tool ([http://www.lc-bio.cn/overview/14?tools=KEGG\\_BarPlot](http://www.lc-bio.cn/overview/14?tools=KEGG_BarPlot)) was used to map the enriched pathways associated with the DEGs.

### Quantitative real-time PCR validation

To confirm the differential expression results, we conducted quantitative RT-PCR in a LightCycler 96 Real-Time PCR system (Roche, Switzerland) using a PrimeScript RT reagent Kit with a gDNA Eraser (Takara, Japan) and TB Green Premix Ex Taq II (TliRNaseH Plus, Takara, Japan) following the manufacturer's directions. A total of 12 genes were used in qPCR to determine the abundance of mRNAs.  *$\beta$ -actin* (Sangon Biotech, China) was used for normalization of the expression data. The relative mRNA expression level was calculated using the  $2^{-\Delta\Delta CT}$  method. All the primers for qRT-PCR are exhibited in Table 1. Three independent replications for each sample were used and data are presented as means  $\pm$  SD.

### Dual-luciferase report assay

The 3'-UTR sequence of KLF4, ATP8A1, and CPLX1 harboring the gga-miR-135a-5p binding sites were amplified with the primers ATP8A1\_WT, CPLX1\_WT, and KLF4\_WT (Table 2). The PCR products were cloned into the pmir-RB-REPORT (Ribibio, China) vector to construct the wild-type plasmid, designated ATP8A1\_WT, CPLX1\_WT and KLF4\_WT. The gga-miR-135a-5p binding sites were mutated in the WT vectors to construct the mutant luciferase

reporter vectors designated ATP8A1\_mut, CPLX1\_mut, and KLF4\_mut. Then, 293T cells were seeded into 24-well plates, and cotransfected with mimics or non-target control at a concentration of 50 nmol/L and 250 ng wild type or mutant luciferase reporter plasmids. After transfection for 48 hr, luciferase activities were measured using the Dual-GloLuciferase Assay System (Promega, USA).

### Data availability

The raw sequence data reported in this paper have been deposited in the Genome Sequence Archive (Genomics, Proteomics & Bioinformatics 2017) in Beijing Institute of Genomics (BIG) Data Center (Nucleic Acids Res 2019), Chinese Academy of Sciences, under accession number CRA001745 that are publicly accessible at <http://bigd.big.ac.cn/gsa>. Supplemental material available at figshare: <https://doi.org/10.25387/g3.12899993>.

## RESULTS

### Chicken follicle theca cells transfected successfully with gga-miR-135a-5P inhibitor

The expression level of gga-miR-135a-5p in chicken follicular theca cells, cells transfected with an inhibitor, and NC were detected by qRT-PCR, as shown in Figure 1, compared with the control and normal chicken follicular theca cells, the relative expression level of gga-miR-135a-5p decreased significantly in the cell inhibitor group ( $P < 0.01$ ).

### RNA-seq and data analysis


After stringent filtration quality control by removing adaptors and low quality reads, 34,217,686, 43,452,332, 31,645,116, 32,409,966, and 35,082,796 clean reads were obtained in the T07, T08, T10, T11, and T12 samples, respectively, of which 75.62%, 76.59%, 75.48%, 75.46%, and 76.30% were mapped to the chicken (*Gallus gallus*) reference sequence, respectively.

### RNA expression and differential analysis

Based on the filtering criteria of the gene abundance differences with an absolute value of  $\log_2$  fold change  $\geq 1$  and FDR significance score  $< 0.01$ , 2013 genes were found to express with a significant difference between cells with gga-miR-135a-5p expression (T07 and T08) and cells with gga-miR-135a-5p inhibited (T10, T11, and T12), while 953 genes were significantly up-regulated, and 1060 genes were significantly down-regulated in cells with gga-miR-135a-5p inhibited (Additional file: Table S1). Only 22 known up-regulated genes showed differences greater than  $\log_2$  fold change  $\geq 4$  between the groups (Figure 2). The volcano plot of DEGs in different groups is shown in Figure 3. Hierarchical clustering analysis of DEGs was performed, and the result of the heatmap is shown in Figure 4.

### Functional annotation of DEGs

The GO enrichment analysis for up-regulated DEGs showed that a total of 125 terms were enriched in biological processes, including cell proliferation, cell differentiation, cell division, regulation of transcription from RNA polymerase II promoters, etc. Among these, 39 terms were preferentially enriched in cell components, such as nucleus, cytoplasm, nucleoplasm, and plasma membrane and 34 terms were enriched in molecular functions including: ATP binding, DNA binding, and transcriptional activator activity (Figure 5).




**Figure 4** The heatmap of differentially expressed genes across all samples via Illumina sequencing. The depth of the color represents the level of gene expression in samples. Normal group: T07 and T08; Transfected group: T10, T11, and T12.

The KEGG analysis showed that 18 terms were enriched (Figure 6). The up-regulated DEGs were preferentially enriched in pathways associated with cellular functions such as the cell cycle (cell division), cytokine-cytokine receptor interaction (cell growth, differentiation, and cell death), the TGF-beta signaling pathway (cell proliferation, apoptosis, differentiation, and migration), Wnt signaling pathway (cell-fate specification, progenitor-cell proliferation, and the control of asymmetric cell division), and p53 signaling pathway (cell differentiation).


### Quantitative real-time PCR validation

To verify the RNA-seq data, 12 DEGs were selected randomly for relative expression analysis by quantitative RT-PCR (Figure 7). The results showed that the relative expression of Serpin family B member 2 (SERPINB2), C-X-C motif chemokine ligand 12 (CXCL12), fatty acid-binding protein 7 (FABP7), G protein-coupled receptor kinase 5 (GRK5), Krüppel-like factor 4 (KLF4), interferon-alpha inducible protein 6 (IFI6), serine/threonine kinase 35 (STK35), neuron specific gene family member 1 (NSG1), 2'-5'-oligoadenylate synthetase-like (OASL), ATPase phospholipid transporting 8A1 (ATP8A1) and interferon-induced protein with tetratricopeptide repeats 5 (IFIT5) in transfected cells were significantly higher than those in normal cells ( $P < 0.01$ ), whereas the relative expression of FERM domain-containing 4A (FRMD4A) between the two groups was different with





**Figure 7** Relative expression of 12 DEGs between cells with normal (NG) and inhibited (TG) expression of gga-miR-135a-5p.



regulate gene expression by binding to sites on the 3'-UTR of the three target genes, KLF4, ATP8A1, and CPLX1.

## DISCUSSION


The function of miR-135a in human disease has been widely studied (Tang *et al.* 2014; Zhang *et al.* 2017). It has been found to play an important role in several diseases including epithelial ovarian cancer (Tang *et al.* 2014), diabetes (Agarwal *et al.* 2013), malignant glioma (Zhang *et al.* 2016), sepsis (Zheng *et al.* 2017), and endometriosis lesions (Petracco *et al.* 2019). In addition, miR-135a-5p was found to be critical for exercise-induced adult neurogenesis (Pons-Espinal *et al.* 2019), by controlling NCX1 expression, miR-135a modulates cardiomyocyte automaticity, Ca<sup>2+</sup> extrusion, and arrhythmogenic Ca<sup>2+</sup> loading/spontaneous Ca<sup>2+</sup> release events to contribute to proarrhythmic remodeling after Complete atrioventricular block (Duong *et al.* 2017). As an important regulatory factor, miR-135a is involved in the regulation of 3T3-L1 preadipocyte differentiation and adipogenesis through the activation of canonical Wnt/ $\beta$ -catenin signaling by directly targeting Apc (Chen *et al.* 2014). In chicken, lncRNA-gga-miR-135a-mRNA interactions may promote the adipogenic differentiation of chicken preadipocytes (Chen *et al.* 2019). Our study provides evidence of gga-miR-135a-5p involvement in chicken follicular theca cells cytopoiesis.

We performed transfection with gga-miR-135a-5p inhibitor to suppress its expression in chicken follicular theca cells, and then performed transcriptome sequencing. Compared with the normal follicular theca cell group, 2013 differential expression genes composed of 953 up-regulated genes and 1060 down-regulated genes were identified. Bioinformatic analyses showed that the up-regulated

DEGs were enriched in the TGF- $\beta$  signaling pathway (gga04350), p53 signaling pathway (gga04115), and Wnt signaling pathway (gga04310), which are known to be involved in follicular development (Xu *et al.* 2018; Du *et al.* 2018). The up-regulated DEGs BMP2, BMP4, TGF $\beta$ 2, TGF $\beta$ 3, and BAMBI are enriched in the TGF- $\beta$  signaling pathway and have a promotive function in granulosa cell proliferation, follicle survival, and prevention of premature luteinization and/or atresia (Schmid *et al.* 1994; Souza *et al.* 2002; Knight and Glistler 2006). THBS1, CCNB2, and CDK1 are enriched in the p53 signaling pathway and have been found to play a role in granulosa cell proliferation in beef cattle (Dias *et al.* 2013), humans (Tremblay and Sirard 2017) and ovine species (Talebi *et al.* 2018). The Wnt family has been implicated in follicular development, and its components the up-regulated DEGs WNT2B, AXIN2, SFRP4, WNT6, NFATC2, WNT5B, and BAMBI have also been found to play important roles in ovarian follicle development (Li *et al.* 2014; Hatzirodos *et al.* 2014; Gupta *et al.* 2014; Drake *et al.* 2003; Chen *et al.* 2012). Thus, gga-miR-135a-5p may regulate the follicular theca cell development through the important signaling pathways mentioned above.

Bioinformatics analysis showed that the KLF4, CPLX1, and ATP8A1 DEGs were the target genes of gga-miR-135a-5p. The expression of KLF4, CPLX1, and ATP8A1 was significantly up-regulated in the gga-miR-135a-5p inhibitor group, and the direct binding relationships between them were further validated by a dual-luciferase reporter assay.

Krüppel-like factor 4 (KLF4) belongs to the KLF family of transcription factors, and exerts important biological effects on



**Figure 8** gga-miR-135a suppresses the expression of KLF4, ATP8A1, and CPLX1 in 293T cells. \*\*P < 0.01.

cellular proliferation, differentiation, and apoptosis (Ghaleb *et al.* 2005; Natesampillai *et al.* 2008; Black *et al.* 2001) in various cells types. Combined LH and IGF-I stimulation increased KLF4 mRNA expressed in porcine ovarian granulosa cells (Natesampillai *et al.* 2008). An H<sub>2</sub>O<sub>2</sub>-induced *in vitro* model and a 3-nitropropionic acid (NP)-induced *in vivo* model of mouse ovarian oxidative stress showed that miR-145 protects granulosa cells against oxidative stress-induced apoptosis by targeting KLF4 (Xu *et al.* 2017). The regulatory function of KLF4 on rat preovulatory granulosa cells also has been confirmed by Hyeonhae and Jaesook (Choi and Roh 2019). They found that KLF4 increases the susceptibility of preovulatory granulosa cells to apoptosis by down-regulating Bcl-2, and promotes an LH-induced cell cycle exit. Interestingly, a directly targeted gene of gga-miR-135a-5p, KLF4, was significantly enriched in the negative regulation of cell proliferation (GO:0008285) and regulation of cell differentiation (GO:0045595), which indicated that gga-miR-135a-5p may

up-regulate the expression of KLF4 to regulate the proliferation and differentiation of chicken follicular theca cells.

CPLX1 belongs to a highly conserved complexin protein family and encodes a neuronal protein (Fernandez and Dittman 2009). Studies have shown that *Cplx1*(*-/-*) mice have profound ataxia that limits their ability to perform co-ordinated motor tasks, and have pronounced deficits in social behaviors (Drew *et al.* 2007). The CPLX1 variants were involved in patients with ID, developmental delay, and myoclonic epilepsy (Brose 2008; Kielar *et al.* 2012; Redler *et al.* 2017). In chicken, there is a lack of data for CPLX1, it was only found to be correlated to earlobe color in Rhode Island Red chickens (Nie *et al.* 2016). Our results suggest that CPLX1 was not only strongly linked with the neurodevelopmental function of the neuronal cell body (GO:0043025), regulation of neurotransmitter secretion (GO:0046928), and synaptic growth at the neuromuscular junction (GO:0051124) as previously reported, but also involved in the



biological process of transmembrane transport (GO:0055085). We speculated that the gga-miR-135a-5p may participate in the regulation of the transmembrane transport function of chicken follicular theca cells by targeting CPLX1.

ATP8A1 is a member of the P4-ATPases subfamily. In mammalian cells, ATP8A1 has been implicated in the translocation of phospholipids (Daleke and Lyles 2000). In Chinese hamster ovary cells, the phospholipid flippase complex of ATP8A1 and CDC50A was found to play a major role in cell migration (Kato *et al.* 2013). Also, ATP8A1 was verified to play a role in regulating the growth and mobility of non-small-cell lung cancer cells (Dong *et al.* 2016). In this study, as confirmed by previous studies, ATP8A1 was enriched in the biological processes involved in phospholipid transport, such as: phospholipid-translocating ATPase activity (GO:0004012), phospholipid transport (GO:0015914) and amino phospholipid transport (GO:0015917). It was also enriched in the biological processes associated with cell development, such as positive regulation of multicellular organism growth (GO:0040018) and negative regulation of cell proliferation (GO:0008285). Interestingly, ATP8A1 was the direct target of gga-miR-135a-5p, this implies a regulation mechanism of gga-miR-135a-5p in chicken follicular theca cells by down-regulating ATP8A1.

In conclusion, gga-miR-135a-5p can directly target the 3'-UTR of KLF4, CPLX1, and ATP8A1 genes to inhibit their expression in chicken follicular theca cells. The data show that gga-miR-135a-5p may play an important role in regulating chicken ovarian follicular theca cells development. Our findings eliminate a gap in the knowledge of gga-miR-135a-5p regulation in chicken follicular theca cells. The exact regulation of apoptosis or proliferation by gga-miR-135a-5p in chicken follicular theca cells and the pathways involved, will be the focus of our future work.

## ACKNOWLEDGMENTS

The authors gratefully acknowledge D. L. Cao and H. L. Han for their help in managing the birds and collecting the data. We thank International Science Editing (<http://www.internationalscienceediting.com>) for editing this manuscript. This work was financially supported by the Agricultural Stock Breeding Project of Shandong Province (2019LZGC019), Natural Science Foundation of Shandong Province (ZR2019BC077), the earmarked fund for modern agro-industry technology research system (CARS-41-Z11), Jinan Layer Experiment Station of China Agriculture Research System (CARS-40-S12), and Construction of Subjects and Teams of Institute of Poultry Science (CXGC2018E11).

## LITERATURE CITED

- Agarwal, P., R. Srivastava, A.K. Srivastava, S. Ali, and M. Datta, 2013 miR-135a targets IRS2 and regulates insulin signaling and glucose uptake in the diabetic gastrocnemius skeletal muscle. *Biochimica et Biophysica Acta (BBA)* 1832: 1294–1303.
- Ambros, V., and X. Chen, 2007 The regulation of genes and genomes by small RNAs. *Development* 134: 1635–1641. <https://doi.org/10.1242/dev.002006>
- Bartel, D.P., 2004 MicroRNAs: genomics, biogenesis, mechanism, and function. *cell* 116: 281–297.
- Black, A. R., J. D. Black, and J. Azizkhan-Clifford, 2001 Sp1 and krüppel-like factor family of transcription factors in cell growth regulation and cancer. *J. Cell. Physiol.* 188: 143–160. <https://doi.org/10.1002/jcp.1111>
- Brose, N., 2008 Altered complexin expression in psychiatric and neurological disorders: cause or consequence? *Mol Cells.* 25: 7–19.
- Chen, C., Y. Peng, Y. Peng, J. Peng, and S. Jiang, 2014 miR-135a-5p inhibits 3T3-L1 adipogenesis through activation of canonical Wnt/ $\beta$ -catenin signaling. *J. Mol. Endocrinol.* 52: 311–320. <https://doi.org/10.1530/JME-14-0013>
- Chen, L., T. Zhang, S. Zhang, J. Huang, G. Zhang *et al.*, 2019 Identification of Long Non-Coding RNA-Associated Competing Endogenous RNA Network in the Differentiation of Chicken Preadipocytes. *Genes (Basel)* 10: 795. <https://doi.org/10.3390/genes10100795>
- Chen, X., H. Bai, L. Li, W. Zhang, R. Jiang *et al.*, 2012 Follicle characteristics and follicle developmental related Wnt6 polymorphism in Chinese indigenous Wanxi-white goose. *Mol. Biol. Rep.* 39: 9843–9848. <https://doi.org/10.1007/s11033-012-1850-2>
- Choi, H., and J. Roh, 2019 Role of Klf4 in the regulation of apoptosis and cell cycle in rat granulosa cells during the periovulatory period. *Int. J. Mol. Sci.* 20: 87. <https://doi.org/10.3390/ijms20010087>
- Daleke, D. L., and J. V. Lyles, 2000 Identification and purification of aminophospholipid flippases. *Biochimica et Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids* 1486: 108–127. [https://doi.org/10.1016/S1388-1981\(00\)00052-4](https://doi.org/10.1016/S1388-1981(00)00052-4)
- Dias, F. C. F., M. I. R. Khan, M. A. Sirard, G. P. Adams, J. Singh, 2013 Differential gene expression of granulosa cells after ovarian superstimulation in beef cattle. *Reproduction* 146: 181–191. <https://doi.org/10.1530/REP-13-0114>
- Dong, W., C. Yao, X. Teng, J. Chai, X. Yang *et al.*, 2016 MiR-140-3p suppressed cell growth and invasion by downregulating the expression of ATP8A1 in non-small cell lung cancer. *Tumour Biol.* 37: 2973–2985. <https://doi.org/10.1007/s13277-015-3452-9>
- Drake, J., R. Friis, and A. Dharmarajan, 2003 The role of sFRP4, a secreted frizzled-related protein, in ovulation. *Apoptosis* 8: 389–397. <https://doi.org/10.1023/A:1024181203729>
- Drew, C. J., R. J. Kyd, and A. J. Morton, 2007 Complexin 1 knockout mice exhibit marked deficits in social behaviours but appear to be cognitively normal. *Hum. Mol. Genet.* 16: 2288–2305. <https://doi.org/10.1093/hmg/ddm181>
- Du, L., T. Gu, Y. Zhang, Z. Huang, N. Wu *et al.*, 2018 Transcriptome profiling to identify key mediators of granulosa cell proliferation upon FSH stimulation in the goose (*Anser cygnoides*). *Br. Poult. Sci.* 59: 416–421. <https://doi.org/10.1080/00071668.2018.1459474>
- Duong, E., J. Xiao, X. Y. Qi, and S. Nattel, 2017 MicroRNA-135a regulates sodium-calcium exchanger gene expression and cardiac electrical activity. *Heart Rhythm* 14: 739–748. <https://doi.org/10.1016/j.hrthm.2017.01.045>
- Donadeu F. X., B. T. Mohammed, J. Ioannidis, 2017 A miRNA target network putatively involved in follicular atresia. *Domest. Anim. Endocrinol.* 58: 76–83. <https://doi.org/10.1016/j.domaniend.2016.08.002>
- Fernandez, J., and J. Dittman, 2009 The Role of Complexin in Neurotransmitter Release. *Ethn. Dis.* 19: 13.
- Gebremedhn, S., D. Salilew-Wondim, I. Ahmad, S. Sahadevan, M. M. Hossain *et al.*, 2015 MicroRNA Expression Profile in Bovine Granulosa Cells of Preovulatory Dominant and Subordinate Follicles during the Late Follicular Phase of the Estrous Cycle. *PLoS One* 10: e0125912. <https://doi.org/10.1371/journal.pone.0125912>
- Ghaleb, A. M., M. O. Nandan, S. Chanchevalap, W. B. Dalton, I. M. Hisamuddin *et al.*, 2005 Krüppel-like factors 4 and 5: the yin and yang regulators of cellular proliferation. *Cell Res.* 15: 92–96. <https://doi.org/10.1038/sj.cr.7290271>
- Grosshans, H., and F. J. Slack, 2002 Micro-RNAs: small is plentiful. *J. Cell Biol.* 156: 17–22. <https://doi.org/10.1083/jcb.200111033>
- Gupta, P., J. K. Folger, S. K. Rajput, L. Lv, J. Yao *et al.*, 2014 Regulation and regulatory role of WNT signaling in potentiating FSH action during bovine dominant follicle selection. *PLoS One* 9: e100201. <https://doi.org/10.1371/journal.pone.0100201>
- Hasuwa, H., J. Ueda, M. Ikawa, and M. Okabe, 2013 MiR-200b and miR-429 Function in Mouse Ovulation and Are Essential for Female Fertility. *Science* 341: 71–73. <https://doi.org/10.1126/science.1237999>
- Hatzirodos, N., K. Hummitzsch, H. F. Irving-Rodgers, and R. J. Rodgers, 2014 Transcriptome profiling of the theca interna in transition from small to large antral ovarian follicles. *PLoS One* 9: e97489. <https://doi.org/10.1371/journal.pone.0097489>

- Hawkins, S. M., and M. M. Matzuk, 2010 Oocyte-somatic cell communication and microRNA function in the ovary. *Annals Dendocrinologie* 71: 144–148. <https://doi.org/10.1016/j.ando.2010.02.020>
- Kang, L., X. Cui, Y. Zhang, C. Yang, and Y. Jiang, 2013 Identification of miRNAs associated with sexual maturity in chicken ovary by Illumina small RNA deep sequencing. *BMC Genomics* 14: 352. <https://doi.org/10.1186/1471-2164-14-352>
- Kang, L., C. Yang, H. Wu, Q. Chen, L. Huang *et al.*, 2017 miR-26a-5p Regulates TNRC6A Expression and Facilitates Theca Cell Proliferation in Chicken Ovarian Follicles. *DNA Cell Biol.* 36: 922–929.
- Kato, U., H. Inadome, M. Yamamoto, K. Emoto, T. Kobayashi *et al.*, 2013 Role for phospholipid flippase complex of ATP8A1 and CDC50A proteins in cell migration. *J. Biol. Chem.* 288: 4922–4934. <https://doi.org/10.1074/jbc.M112.402701>
- Kielar, C., S. J. Sawiak, P. N. Negredo, H. Desmond, and A. J. Morton, 2012 Tensor-based morphometry and stereology reveal brain pathology in the complexin1 knockout mouse. *PLoS One* 7: e32636. <https://doi.org/10.1371/journal.pone.0032636>
- Kim, D., G. Pertea, C. Trapnell, H. Pimentel, R. Kelley *et al.*, 2013 TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol.* 14: R36. <https://doi.org/10.1186/gb-2013-14-4-r36>
- Knight, P. G., and C. Glistler, 2006 TGF- $\beta$  superfamily members and ovarian follicle development. *Reproduction* 132: 191–206. <https://doi.org/10.1530/rep.1.01074>
- Krol, J., I. Loedige, and W. Filipowicz, 2010 The widespread regulation of microRNA biogenesis, function and decay. *Nat. Rev. Genet.* 11: 597–610. <https://doi.org/10.1038/nrg2843>
- Lee, R. C., R. L. Feinbaum, and V. Ambros, 1993 The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 75: 843–854. [https://doi.org/10.1016/0092-8674\(93\)90529-Y](https://doi.org/10.1016/0092-8674(93)90529-Y)
- Li, L., S.-Y. Ji, J.-L. Yang, X.-X. Li, J. Zhang *et al.*, 2014 Wnt/ $\beta$ -catenin signaling regulates follicular development by modulating the expression of Foxo3a signaling components. *Mol. Cell. Endocrinol.* 382: 915–925. <https://doi.org/10.1016/j.mce.2013.11.007>
- Lin, F., L. Ran, Z. X. Pan, B. Zhou, and D. B. Yu *et al.*, 2012 miR-26b Promotes Granulosa Cell Apoptosis by Targeting ATM during Follicular Atresia in Porcine Ovary. *PLoS One* 7: e38640. <https://doi.org/10.1371/journal.pone.0038640>
- Lin, L., T. Du, J. Huang, L.-L. Huang, and D.-Z. Yang, 2015 Identification of Differentially Expressed MicroRNAs in the Ovary of Polycystic Ovary Syndrome with Hyperandrogenism and Insulin Resistance. *Chin. Med. J. (Engl.)* 128: 169–174. <https://doi.org/10.4103/0366-6999.149189>
- Maalouf, S. W., W. S. Liu, and J. L. Pate, 2015 MicroRNA in ovarian function. *Cell Tissue Res.* 363: 7–18. <https://doi.org/10.1007/s00441-015-2307-4>
- McBride, D., W. Carre, S. D. Sontakke, C. O. Hogg, A. Law *et al.*, 2012 Identification of miRNAs associated with the follicular-luteal transition in the ruminant ovary. *Reproduction* 144: 221–233. <https://doi.org/10.1530/REP-12-0025>
- Miao, N., X. Wang, Y. Hou, Y. Feng, and Y. Gong, 2016 Identification of male-biased microRNA-107 as a direct regulator for nuclear receptor subfamily 5 group A member 1 based on sexually dimorphic microRNA expression profiling from chicken embryonic gonads. *Mol. Cell. Endocrinol.* 429: 29–40. <https://doi.org/10.1016/j.mce.2016.03.033>
- Natesampillai, S., J. Kerkvliet, P. C. Leung, and J. D. Veldhuis, 2008 Regulation of Kruppel-like factor 4, 9, and 13 genes and the steroidogenic genes LDLR, StAR, and CYP11A in ovarian granulosa cells. *Am. J. Physiol. Endocrinol. Metab.* 294: E385–E391. <https://doi.org/10.1152/ajpendo.00480.2007>
- Nie, C., Z. Zhang, J. Zheng, H. Sun, Z. Ning *et al.*, 2016 Genome-wide association study revealed genomic regions related to white/red earlobe color trait in the Rhode Island Red chickens. *BMC Genet.* 17: 115. <https://doi.org/10.1186/s12863-016-0422-1>
- Petracco, R., A. C. O. Dias, H. S. Taylor, Á. Petracco, M. Badalotti *et al.*, 2019 Evaluation of miR-135a/b expression in endometriosis lesions. *Biomed. Rep.* 11: 181–187.
- Pons-Espinal, M., C. Gasperini, M. J. Marzi, C. Braccia, A. Armirotti *et al.*, 2019 MiR-135a-5p Is Critical for Exercise-Induced Adult Neurogenesis. *Stem Cell Reports* 12: 1298–1312. <https://doi.org/10.1016/j.stemcr.2019.04.020>
- Redler, S., T. M. Strom, T. Wieland, K. Cremer, H. Engels *et al.*, 2017 Variants in CPLX1 in two families with autosomal-recessive severe infantile myoclonic epilepsy and ID. *Eur. J. Hum. Genet.* 25: 889–893. <https://doi.org/10.1038/ejhg.2017.52>
- Robinson, C. L., L. Zhang, L. F. Schutz, M. L. Totty, and L. J. Spicer, 2018 MicroRNA 221 expression in theca and granulosa cells: hormonal regulation and function. *J. Anim. Sci.* 96: 641–652. <https://doi.org/10.1093/jas/skx069>
- Salilew-Wondim, D., I. Ahmad, S. Gebremedhn, S. Sahadevan, M. M. Hossain *et al.*, 2014 The Expression Pattern of microRNAs in Granulosa Cells of Subordinate and Dominant Follicles during the Early Luteal Phase of the Bovine Estrous Cycle. *PLoS One* 9: e106795. <https://doi.org/10.1371/journal.pone.0106795>
- Schmid, P., D. Cox, H. Vanderputten, G. K. McMaster, and G. Bilbe, 1994 Expression of TGF- $\beta$ s and TGF- $\beta$  type II receptor mRNAs in mouse folliculogenesis: stored maternal TGF- $\beta$ 2 message in oocytes. *Biochem. Biophys. Res. Commun.* 201: 649–656. <https://doi.org/10.1006/bbrc.1994.1750>
- Shukla, A., S. Dahiya, S. K. Onteru, and D. Singh, 2018 Differentially expressed miRNA-210 during follicular-luteal transition regulates pre-ovulatory granulosa cell function targeting Hra and EFNA3. *J. Cell. Biochem.* 119: 7934–7943. <https://doi.org/10.1002/jcb.26508>
- Sohel, M. M. H., M. Hoelker, S. S. Noferesti, D. Salilew-Wondim, E. Tholen *et al.*, 2013 Exosomal and Non-Exosomal Transport of Extra-Cellular microRNAs in Follicular Fluid: Implications for Bovine Oocyte Developmental Competence. *PLoS One* 8: e78505. <https://doi.org/10.1371/journal.pone.0078505>
- Song, Y., G. Yu, Y. Xiang, Y. Li, L. Wan *et al.*, 2019 Altered miR-186 and miR-135a contribute to granulosa cell dysfunction by targeting ESR2: A possible role in polycystic ovary syndrome. *Mol. Cell. Endocrinol.* 494: 110478. <https://doi.org/10.1016/j.mce.2019.110478>
- Souza, C., B. Campbell, A. McNeilly, and D. Baird, 2002 Effect of bone morphogenetic protein 2 (BMP2) on oestradiol and inhibin A production by sheep granulosa cells, and localization of BMP receptors in the ovary by immunohistochemistry. *Reproduction* 123: 363–369. <https://doi.org/10.1530/rep.0.1230363>
- Talebi, R., A. Ahmadi, and F. Afraz, 2018 Analysis of protein-protein interaction network based on transcriptome profiling of ovine granulosa cells identifies candidate genes in cyclic recruitment of ovarian follicles. *J. Anim. Sci. Technol.* 60: 11. <https://doi.org/10.1186/s40781-018-0171-y>
- Tang, W., Y. Jiang, X. Mu, L. Xu, W. Cheng *et al.*, 2014 MiR-135a functions as a tumor suppressor in epithelial ovarian cancer and regulates HOXA10 expression. *Cell. Signal.* 26: 1420–1426. <https://doi.org/10.1016/j.cellsig.2014.03.002>
- Trapnell, C., B. A. Williams, G. Pertea, A. Mortazavi, G. Kwan *et al.*, 2010 Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* 28: 511–515. <https://doi.org/10.1038/nbt.1621>
- Tremblay, P. G., and M.-A. Sirard, 2017 Transcriptomic analysis of gene cascades involved in protein kinase A and C signaling in the KGN line of human ovarian granulosa tumor cells. *Biol. Reprod.* 96: 855–865. <https://doi.org/10.1093/biolre/iox024>
- Wei, Y., S. Lu, Y. Hu, L. Guo, X. Wu *et al.*, 2020 MicroRNA-135a Regulates VEGFC Expression and Promotes Luteinized Granulosa Cell Apoptosis in Polycystic Ovary Syndrome. *Reprod. Sci.* 27: 1436–1442.
- Wu, H., F. Fan, C. Liang, Y. Zhou, X. Qiao *et al.*, 2019 Variants of pri-miR-26a-5p polymorphisms are associated with values for chicken egg production variables and affects abundance of mature miRNA. *Anim. Reprod. Sci.* 201: 93–101. <https://doi.org/10.1016/j.anireprosci.2019.01.002>
- Xu, L., H. Sun, M. Zhang, Y. Jiang, C. Zhang *et al.*, 2017 MicroRNA-145 protects follicular granulosa cells against oxidative stress-induced apoptosis by targeting Kruppel-like factor 4. *Mol. Cell. Endocrinol.* 452: 138–147. <https://doi.org/10.1016/j.mce.2017.05.030>

- Xu, Y., J. Niu, G. Xi, X. Niu, Y. Wang *et al.*, 2018 TGF- $\beta$ 1 resulting in differential microRNA expression in bovine granulosa cells. *Gene* 663: 88–100. <https://doi.org/10.1016/j.gene.2018.04.036>
- Yan, G. J., L. X. Zhang, T. Fang, Q. Zhang, S. G. Wu *et al.*, 2012 MicroRNA-145 suppresses mouse granulosa cell proliferation by targeting activin receptor IB. *FEBS Lett.* 586: 3263–3270. <https://doi.org/10.1016/j.febslet.2012.06.048>
- Zhang, J., Y. Xu, H. Liu, and Z. Pan, 2019 MicroRNAs in ovarian follicular atresia and granulosa cell apoptosis. *Reprod. Biol. Endocrinol.* 17: 9. <https://doi.org/10.1186/s12958-018-0450-y>
- Zhang, Q., S. Haixiang, J. Yue, D. Lijun, W. Shaogen *et al.*, 2013 MicroRNA-181a Suppresses Mouse Granulosa Cell Proliferation by Targeting Activin Receptor IIA. *PLoS One* 8: e59667. <https://doi.org/10.1371/journal.pone.0059667>
- Zhang, T., Y. Shao, T.Y. Chu, H.S. Huang, Y.L. Liou *et al.*, 2016 MiR-135a and MRP1 play pivotal roles in the selective lethality of phenethyl isothiocyanate to malignant glioma cells. *Am J Cancer Res* 6: 957–972.
- Zhang, X., G. Feng, Z. Lei, W. Huaitao, S. Gang *et al.*, 2017 UCA1 Regulates the Growth and Metastasis of Pancreatic Cancer by Sponging miR-135a. *Oncology Research Featuring Preclinical & Clinical Cancer Therapeutics* 25: 1529–1541. <https://doi.org/10.3727/096504017X14888987683152>
- Zheng, G., M. Pan, W. Jin, G. Jin, and Y. Huang, 2017 MicroRNA-135a is up-regulated and aggravates myocardial depression in sepsis via regulating p38 MAPK/NF- $\kappa$ B pathway. *Int. Immunopharmacol.* 45: 6–12. <https://doi.org/10.1016/j.intimp.2017.01.029>

*Communicating editor: J. Birchler*