

# Crystal structure of methyl 2-(2*H*-1,3-benzodioxol-5-yl)-7,9-dibromo-8-oxo-1-oxaspiro[4.5]deca-2,6,9-triene-3-carboxylate

Lucimara Julio Martins,<sup>a</sup> Deborah de Alencar Simoni,<sup>b</sup> Ricardo Aparicio<sup>c\*</sup> and Fernando Coelho<sup>a</sup>

<sup>a</sup>Laboratory of Synthesis of Natural Products and Drugs, Institute of Chemistry, University of Campinas, PO Box 6154 - 13083-970, Campinas, SP, Brazil,

<sup>b</sup>Laboratory of Single Crystal X-Ray Diffraction, Institute of Chemistry, University of Campinas, PO Box 6154 - 13083-970, Campinas, SP, Brazil, and <sup>c</sup>Laboratory of Structural Biology and Crystallography, Institute of Chemistry, University of Campinas, PO Box 6154 - 13083-970, Campinas, SP, Brazil. \*Correspondence e-mail: [aparicio@iqm.unicamp.br](mailto:aparicio@iqm.unicamp.br)

Received 14 October 2014; accepted 11 November 2014

Edited by P. C. Healy, Griffith University, Australia

The title compound, C<sub>18</sub>H<sub>12</sub>Br<sub>2</sub>O<sub>6</sub>, was synthesized from Morita–Baylis–Hillman adducts. It incorporates the brominated spiro-hexadienone moiety typically exhibited by compounds of this class that exhibit biological activity. Both the brominated cyclohexadienone and the central five-membered rings are nearly planar (r.m.s. deviations of 0.044 and 0.016 Å, respectively), being almost perpendicularly oriented [interplanar angle = 89.47 (5)°]. With respect to the central five-membered ring, the brominated cyclohexadienone ring, the benzodioxol ring and the carboxylate fragment make C–O–C–C, O–C–C–C and C–C–C–O dihedral angles of –122.11 (8), –27.20 (11) and –8.40 (12)°, respectively. An intramolecular C–H···O hydrogen bond occurs. In the crystal, molecules are linked by non-classical C–H···O and C–H···Br hydrogen bonds resulting in a molecular packing in which the brominated rings are in a head-to-head orientation, forming well marked planes parallel to the *b* axis.


**Keywords:** Single-crystal X-ray study; spiro-hexadienone structure; Morita–Baylis–Hillman adducts.

**CCDC reference:** 1033626

## 1. Related literature

For compounds that contain a spiro-hexadienone moiety in their structures, related biological activities and examples of brominated spiro-hexadienones, see: König & Wright (1993); Lou (2012); Sorek *et al.* (2009). For strategies for the synthesis

of spiro-hexadienones from Morita–Baylis–Hillman adducts, see: Martins *et al.* (2014); Barontini *et al.* (2013).



## 2. Experimental

### 2.1. Crystal data

C <sub>18</sub> H <sub>12</sub> Br <sub>2</sub> O <sub>6</sub>	$\gamma = 78.077 (4)^\circ$
$M_r = 484.10$	$V = 852.7 (2) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.1929 (13) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.4811 (14) \text{ \AA}$	$\mu = 4.79 \text{ mm}^{-1}$
$c = 12.761 (2) \text{ \AA}$	$T = 100 \text{ K}$
$\alpha = 84.485 (4)^\circ$	$0.32 \times 0.17 \times 0.16 \text{ mm}$
$\beta = 80.007 (5)^\circ$	

### 2.2. Data collection

Bruker APEX CCD detector diffractometer	27788 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2010)	7098 independent reflections
$T_{\min} = 0.309$ , $T_{\max} = 0.515$	6288 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.017$

### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.018$	236 parameters
$wR(F^2) = 0.047$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\max} = 0.57 \text{ e \AA}^{-3}$
7098 reflections	$\Delta\rho_{\min} = -0.39 \text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
C3–H3···O5 <sup>i</sup>	0.95	2.61	3.4272 (13)	144
C7–H7···O5	0.95	2.34	2.9441 (13)	121
C10–H10A···O1 <sup>ii</sup>	0.99	2.55	3.3995 (13)	143
C12–H12A···Br1 <sup>iii</sup>	0.99	2.96	3.9411 (12)	173
C12–H12A···O1 <sup>iii</sup>	0.99	2.53	3.0852 (13)	116
C13–H13···O2 <sup>iv</sup>	0.95	2.63	3.3909 (13)	137
C16–H16C···O5 <sup>v</sup>	0.98	2.59	3.2434 (14)	124
C17–H17···Br1 <sup>vi</sup>	0.95	3.03	3.9069 (11)	153

Symmetry codes: (i)  $-x+2, -y+2, -z+2$ ; (ii)  $-x+2, -y+2, -z+1$ ; (iii)  $x, y-1, z+1$ ; (iv)  $-x+1, -y+2, -z+2$ ; (v)  $-x+3, -y+1, -z+2$ ; (vi)  $x, y-1, z$ .

Data collection: APEX2 (Bruker, 2010); cell refinement: SAINT (Bruker, 2010); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL (Hübschle *et al.*, 2011) and SHELXL2014 (Sheldrick, 2008); molecular graphics: Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: OLEX2 (Dolo-

manov *et al.*, 2003), *PLATON* (Spek, 2009) and *pubCIF* (Westrip, 2010).

### Acknowledgements

The authors acknowledge Dr Jorge Henrique Monteiro for preliminary structure refinement. This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP 2009/18390–4 and 2009/51602–5), the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). RA is the recipient of a research grant from CNPq.

---

Supporting information for this paper is available from the IUCr electronic archives (Reference: HG5413).

---

### References

- Barontini, M., Proietti Silvestri, I., Nardi, V., Crisante, F., Pepe, G., Pari, L., Gallucci, F., Bovicelli, P. & Righi, G. (2013). *Med. Chem. Res.* **22**, 674–680.
- Bruker (2010). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dolomanov, O. V., Blake, A. J., Champness, N. R. & Schröder, M. (2003). *J. Appl. Cryst.* **36**, 1283–1284.
- Hübschle, C. B., Sheldrick, G. M. & Ditttrich, B. (2011). *J. Appl. Cryst.* **44**, 1281–1284.
- König, G. M. & Wright, A. D. (1993). *Heterocycles*, **36**, 1351–1358.
- Lou, Y. (2012). *Acta Cryst.* **E68**, o1152.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Martins, L. J., Ferreira, B. V., Almeida, W. P., Lancellotti, M. & Coelho, F. (2014). *Tetrahedron Lett.* **55**, 5264–5267.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sorek, H., Rudi, A., Goldberg, I., Akin, M. & Kashman, Y. (2009). *J. Nat. Prod.* **72**, 784–786.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2014). E70, o1275–o1276 [doi:10.1107/S1600536814024763]

## Crystal structure of methyl 2-(2*H*-1,3-benzodioxol-5-yl)-7,9-dibromo-8-oxo-1-oxaspiro[4.5]deca-2,6,9-triene-3-carboxylate

Lucimara Julio Martins, Deborah de Alencar Simoni, Ricardo Aparicio and Fernando Coelho

### S1. Introduction

Compounds containing a spiro-hexadienone moiety typically exhibit biological activity, also sharing a structural architecture observed in some natural products (König & Wright, 1993; Sorek *et al.*, 2009). In view of the importance of this class of compounds, recent efforts resulted in a new synthetic strategy, which starts from Morita-Baylis-Hillman adducts as building blocks for organic synthesis (Martins *et al.*, 2014). Based on this methodology, it was possible to obtain a brominated spiro-hexadienone which resulted, to our knowledge, in the first report of a halogenated spiro-hexadienone crystal structure.

### S2. Experimental

#### S2.1. Synthesis and crystallization

Methyl 2-(2*H*-1,3-benzodioxol-5-yl)-7,9-dibromo-8-oxo-1-oxaspiro[4.5]deca-2,6,9-triene-3-carboxylate was prepared from a subset of  $\beta$ -ketoesters following the experimental protocol recently described by Barontini *et al.* (2013). A separable mixture of mono- and dibrominated (in majority) derivatives in good overall yields was obtained.

After chromatographic separation, the dibrominated compounds were easily transformed into halogenated spiro-hexadienones, in three steps procedure, starting with the Morita-Baylis-Hillman adducts.

Methyl 2-(2*H*-1,3-benzodioxol-5-yl)-7,9-dibromo-8-oxo-1-oxaspiro[4.5]deca-2,6,9-triene-3-carboxylate (48 mg, 0.1 mmol) was dissolved in absolute chloroform-*D*<sub>1</sub> (1 mL), followed by stirring until total dissolution was achieved. The solution was kept in the freezer. After two weeks, the resulting material was filtered under vacuum, washed with small portions of cold chloroform and dried in a desiccator to furnish pale yellow single crystals suitable for X-ray diffraction data collection.

#### S2.2. Refinement


The C-bound H atoms were positioned with idealized geometry and treated as riding atoms: phenyl, methyl and methylene C—H bond lengths were 0.95, 0.98 and 0.99 Å, respectively. The isotropic displacement parameters values ( $U_{\text{iso}}(\text{H})$ ) were fixed at  $1.5U_{\text{eq}}(\text{C})$  for methyl H atoms and  $1.2U_{\text{eq}}(\text{C})$  for all other attached H atoms.

### S3. Results and discussion

The title compound (Fig. 1) crystallized in the space group  $P\bar{1}$  assuming a conformational structure determined by non-classical intramolecular C—H—O and intermolecular C—H—O and C—H—Br bonding (Table 1, Fig. 2). The molecule contains one six-membered (brominated cyclohexadienone) and one central five-membered rings connected by a spiro-carbon C4. The mean plane of these planar rings, C1—C2—C4—C17—C18 (r.m.s. = 0.044 Å) and C4—C10—C11—C5—O2 (r.m.s. = 0.016 Å), are almost perpendicularly oriented, making a plane-plane angle of 90.53°. With respect to the


central five-membered ring, the brominated spiro-hexadienone and the benzodioxol rings make dihedral angles  $C5—O2—C4—C3 = -122.11(8)^\circ$  and  $O2—C5—C6—C14 = -27.20(11)^\circ$ , respectively, while the dihedral angle with the carboxylate fragment  $C10—C11—C15—O6$  is  $-8.40(12)^\circ$ .

In the hexadienone ring of the title compound, the  $C2—C3$ ,  $C17—C18$  and  $C1—O1$  bond lengths are  $1.3351(13)$ ,  $1.3354(13)$  and  $1.2139(11)$  Å, respectively, and the bond length between the spiro-carbon ( $C4$ ) and the oxygen atom  $C4—O2$  is  $1.4684(11)$  Å, similar to those reported for a related oxaspiro structure (Lou, 2012). In the latter, the plane-plane angle between the mean planes of the six-membered and the central five-membered rings in the spiro-carbon is  $96.06^\circ$ , slightly different from that observed in the title compound ( $90.53^\circ$ ). Further comparison also reveals different orientations of the carboxylate moiety, which makes a dihedral angle of  $170.6(1)^\circ$  ( $C10—C11—C15—O5$ ) in the title compound, with a corresponding angle equal to  $11.3(3)^\circ$  in the related structure (Lou, 2012). This large difference is consistent with an observed  $C7—H7—O5$  intramolecular hydrogen bond (Table 1) in the title compound, which results in a favourable conformation of the carboxylate group.



**Figure 1**

The molecular structure of the title compound with atom labels and 50% probability displacement ellipsoids.

**Figure 2**

Crystal packing of the title compound, showing hydrogen bonding interactions.

**2-(2*H*-1,3-Benzodioxol-5-yl)-7,9-dibromo-8-oxo-1-oxaspiro[4.5]deca-2,6,9-triene-3-carboxylate**

*Crystal data*

$C_{18}H_{12}Br_2O_6$

$M_r = 484.10$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 8.1929\ (13)\ \text{\AA}$

$b = 8.4811\ (14)\ \text{\AA}$

$c = 12.761\ (2)\ \text{\AA}$

$\alpha = 84.485\ (4)^\circ$

$\beta = 80.007\ (5)^\circ$

$\gamma = 78.077\ (4)^\circ$

$V = 852.7\ (2)\ \text{\AA}^3$

$Z = 2$

$F(000) = 476$

$D_x = 1.885\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 9887 reflections

$\theta = 3.2\text{--}35.1^\circ$

$\mu = 4.79\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Block, pale yellow

$0.32 \times 0.17 \times 0.16\ \text{mm}$

*Data collection*

Bruker APEX CCD detector

diffractometer

Radiation source: fine-focus sealed tube

Detector resolution:  $8.3333\ \text{pixels mm}^{-1}$

$\phi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2010)

$T_{\min} = 0.309$ ,  $T_{\max} = 0.515$

27788 measured reflections

7098 independent reflections

6288 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 34.3^\circ$ ,  $\theta_{\min} = 3.0^\circ$

$h = -12 \rightarrow 12$

$k = -13 \rightarrow 13$

$l = -20 \rightarrow 20$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.018$   
 $wR(F^2) = 0.047$   
 $S = 1.02$   
 7098 reflections  
 236 parameters  
 0 restraints

Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0245P)^2 + 0.2583P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.003$   
 $\Delta\rho_{\max} = 0.57 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.39 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.81611 (2)	1.44904 (2)	0.62579 (2)	0.01782 (3)
Br2	0.72560 (2)	0.85491 (2)	0.49155 (2)	0.01638 (3)
O1	0.75814 (11)	1.20677 (9)	0.48397 (6)	0.01915 (14)
O2	0.81234 (9)	0.93071 (9)	0.86474 (5)	0.01450 (12)
O3	0.57197 (10)	0.68379 (9)	1.33065 (6)	0.01608 (13)
O4	0.81596 (9)	0.50402 (9)	1.26861 (6)	0.01609 (13)
O5	1.24024 (10)	0.67487 (10)	1.01728 (6)	0.01916 (14)
O6	1.37041 (9)	0.72220 (9)	0.84968 (6)	0.01529 (12)
C1	0.79228 (12)	1.13898 (10)	0.56785 (7)	0.01214 (14)
C2	0.83286 (12)	1.22429 (10)	0.65381 (7)	0.01222 (14)
C3	0.88062 (12)	1.14902 (11)	0.74351 (7)	0.01357 (15)
H3	0.8963	1.2114	0.7977	0.016*
C4	0.91021 (12)	0.96878 (11)	0.76100 (7)	0.01264 (15)
C5	0.91926 (12)	0.83972 (10)	0.92957 (7)	0.01135 (14)
C6	0.82769 (11)	0.79788 (10)	1.03513 (7)	0.01106 (14)
C7	0.88494 (12)	0.65586 (11)	1.09690 (7)	0.01202 (14)
H7	0.9864	0.5828	1.0731	0.014*
C8	0.78592 (12)	0.62914 (11)	1.19325 (7)	0.01182 (14)
C9	0.63896 (12)	0.73620 (11)	1.23042 (7)	0.01254 (14)
C10	1.09810 (12)	0.89627 (12)	0.77135 (7)	0.01504 (16)
H10A	1.1501	0.8183	0.7164	0.018*
H10B	1.1655	0.9820	0.7656	0.018*
C11	1.08297 (12)	0.81321 (11)	0.88176 (7)	0.01190 (14)
C12	0.66357 (14)	0.52054 (12)	1.34563 (8)	0.01708 (17)
H12A	0.6912	0.5003	1.4188	0.020*
H12B	0.5948	0.4422	1.3344	0.020*
C13	0.58111 (12)	0.87530 (11)	1.17181 (8)	0.01435 (15)
H13	0.4809	0.9486	1.1975	0.017*
C14	0.67762 (12)	0.90331 (11)	1.07223 (7)	0.01288 (15)
H14	0.6401	0.9965	1.0287	0.015*

C15	1.23289 (12)	0.72953 (11)	0.92621 (7)	0.01239 (14)
C16	1.52864 (13)	0.64225 (14)	0.88280 (9)	0.02125 (19)
H16A	1.5418	0.6880	0.9478	0.032*
H16B	1.6222	0.6580	0.8261	0.032*
H16C	1.5291	0.5265	0.8969	0.032*
C17	0.85126 (12)	0.88535 (11)	0.68028 (7)	0.01319 (15)
H17	0.8524	0.7728	0.6918	0.016*
C18	0.79718 (12)	0.96291 (10)	0.59287 (7)	0.01170 (14)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.02471 (5)	0.00997 (4)	0.01954 (5)	-0.00466 (3)	-0.00492 (4)	0.00072 (3)
Br2	0.02188 (5)	0.01408 (4)	0.01473 (4)	-0.00300 (3)	-0.00654 (3)	-0.00306 (3)
O1	0.0307 (4)	0.0153 (3)	0.0107 (3)	-0.0017 (3)	-0.0063 (3)	0.0023 (2)
O2	0.0119 (3)	0.0197 (3)	0.0100 (3)	-0.0004 (2)	-0.0030 (2)	0.0048 (2)
O3	0.0186 (3)	0.0160 (3)	0.0112 (3)	-0.0020 (2)	0.0011 (2)	0.0023 (2)
O4	0.0173 (3)	0.0147 (3)	0.0138 (3)	-0.0009 (2)	-0.0019 (2)	0.0056 (2)
O5	0.0146 (3)	0.0278 (4)	0.0125 (3)	0.0000 (3)	-0.0034 (2)	0.0045 (3)
O6	0.0106 (3)	0.0196 (3)	0.0133 (3)	0.0005 (2)	-0.0009 (2)	0.0011 (2)
C1	0.0137 (4)	0.0115 (3)	0.0101 (3)	-0.0011 (3)	-0.0008 (3)	-0.0001 (3)
C2	0.0144 (4)	0.0101 (3)	0.0119 (4)	-0.0024 (3)	-0.0017 (3)	0.0000 (3)
C3	0.0153 (4)	0.0137 (3)	0.0123 (4)	-0.0032 (3)	-0.0034 (3)	-0.0009 (3)
C4	0.0137 (4)	0.0143 (3)	0.0091 (3)	-0.0018 (3)	-0.0025 (3)	0.0023 (3)
C5	0.0129 (4)	0.0113 (3)	0.0096 (3)	-0.0011 (3)	-0.0037 (3)	0.0010 (3)
C6	0.0123 (4)	0.0117 (3)	0.0093 (3)	-0.0021 (3)	-0.0028 (3)	0.0004 (3)
C7	0.0128 (4)	0.0113 (3)	0.0114 (3)	-0.0009 (3)	-0.0024 (3)	0.0003 (3)
C8	0.0135 (4)	0.0111 (3)	0.0109 (3)	-0.0020 (3)	-0.0036 (3)	0.0012 (3)
C9	0.0139 (4)	0.0133 (3)	0.0101 (3)	-0.0028 (3)	-0.0012 (3)	-0.0001 (3)
C10	0.0131 (4)	0.0196 (4)	0.0110 (4)	-0.0014 (3)	-0.0027 (3)	0.0036 (3)
C11	0.0122 (4)	0.0133 (3)	0.0095 (3)	-0.0010 (3)	-0.0024 (3)	0.0009 (3)
C12	0.0219 (5)	0.0149 (4)	0.0127 (4)	-0.0037 (3)	0.0000 (3)	0.0027 (3)
C13	0.0139 (4)	0.0140 (4)	0.0130 (4)	0.0005 (3)	-0.0006 (3)	0.0004 (3)
C14	0.0131 (4)	0.0127 (3)	0.0116 (4)	-0.0004 (3)	-0.0026 (3)	0.0015 (3)
C15	0.0115 (4)	0.0132 (3)	0.0117 (4)	-0.0005 (3)	-0.0023 (3)	-0.0003 (3)
C16	0.0119 (4)	0.0287 (5)	0.0201 (5)	0.0015 (4)	-0.0019 (3)	0.0012 (4)
C17	0.0151 (4)	0.0114 (3)	0.0126 (4)	-0.0018 (3)	-0.0031 (3)	0.0012 (3)
C18	0.0134 (4)	0.0110 (3)	0.0108 (3)	-0.0018 (3)	-0.0025 (3)	-0.0011 (3)

*Geometric parameters (Å, °)*

Br1—C2	1.8863 (9)	C6—C14	1.4001 (13)
Br2—C18	1.8887 (9)	C6—C7	1.4158 (12)
O1—C1	1.2139 (11)	C7—C8	1.3775 (13)
O2—C5	1.3772 (11)	C7—H7	0.9500
O2—C4	1.4684 (11)	C8—C9	1.3900 (13)
O3—C9	1.3721 (11)	C9—C13	1.3777 (13)
O3—C12	1.4444 (12)	C10—C11	1.5104 (13)

O4—C8	1.3746 (11)	C10—H10A	0.9900
O4—C12	1.4397 (13)	C10—H10B	0.9900
O5—C15	1.2158 (11)	C11—C15	1.4623 (13)
O6—C15	1.3520 (11)	C12—H12A	0.9900
O6—C16	1.4459 (13)	C12—H12B	0.9900
C1—C2	1.4887 (13)	C13—C14	1.4026 (13)
C1—C18	1.4904 (12)	C13—H13	0.9500
C2—C3	1.3351 (13)	C14—H14	0.9500
C3—C4	1.4985 (13)	C16—H16A	0.9800
C3—H3	0.9500	C16—H16B	0.9800
C4—C17	1.4996 (13)	C16—H16C	0.9800
C4—C10	1.5585 (14)	C17—C18	1.3354 (13)
C5—C11	1.3563 (13)	C17—H17	0.9500
C5—C6	1.4733 (12)		
C5—O2—C4	109.46 (7)	C11—C10—H10A	111.3
C9—O3—C12	104.80 (7)	C4—C10—H10A	111.3
C8—O4—C12	105.13 (7)	C11—C10—H10B	111.3
C15—O6—C16	115.44 (8)	C4—C10—H10B	111.3
O1—C1—C2	122.93 (8)	H10A—C10—H10B	109.2
O1—C1—C18	122.50 (8)	C5—C11—C15	128.64 (8)
C2—C1—C18	114.57 (8)	C5—C11—C10	110.13 (8)
C3—C2—C1	123.31 (8)	C15—C11—C10	121.07 (8)
C3—C2—Br1	121.81 (7)	O4—C12—O3	106.94 (7)
C1—C2—Br1	114.87 (6)	O4—C12—H12A	110.3
C2—C3—C4	121.73 (8)	O3—C12—H12A	110.3
C2—C3—H3	119.1	O4—C12—H12B	110.3
C4—C3—H3	119.1	O3—C12—H12B	110.3
O2—C4—C3	107.32 (7)	H12A—C12—H12B	108.6
O2—C4—C17	106.47 (7)	C9—C13—C14	116.62 (9)
C3—C4—C17	114.33 (8)	C9—C13—H13	121.7
O2—C4—C10	105.43 (7)	C14—C13—H13	121.7
C3—C4—C10	111.56 (8)	C6—C14—C13	121.91 (8)
C17—C4—C10	111.13 (8)	C6—C14—H14	119.0
C11—C5—O2	112.55 (8)	C13—C14—H14	119.0
C11—C5—C6	135.58 (8)	O5—C15—O6	122.92 (9)
O2—C5—C6	111.85 (8)	O5—C15—C11	127.60 (9)
C14—C6—C7	120.45 (8)	O6—C15—C11	109.47 (8)
C14—C6—C5	117.27 (8)	O6—C16—H16A	109.5
C7—C6—C5	122.27 (8)	O6—C16—H16B	109.5
C8—C7—C6	116.53 (8)	H16A—C16—H16B	109.5
C8—C7—H7	121.7	O6—C16—H16C	109.5
C6—C7—H7	121.7	H16A—C16—H16C	109.5
O4—C8—C7	127.79 (8)	H16B—C16—H16C	109.5
O4—C8—C9	109.53 (8)	C18—C17—C4	122.64 (8)
C7—C8—C9	122.61 (8)	C18—C17—H17	118.7
O3—C9—C13	128.16 (9)	C4—C17—H17	118.7
O3—C9—C8	109.94 (8)	C17—C18—C1	122.38 (8)



C13—C9—C8	121.85 (8)	C17—C18—Br2	121.91 (7)
C11—C10—C4	102.30 (7)	C1—C18—Br2	115.70 (6)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C3—H3...O5 <sup>i</sup>	0.95	2.61	3.4272 (13)	144
C7—H7...O5	0.95	2.34	2.9441 (13)	121
C10—H10 <i>A</i> ...O1 <sup>ii</sup>	0.99	2.55	3.3995 (13)	143
C12—H12 <i>A</i> ...Br1 <sup>iii</sup>	0.99	2.96	3.9411 (12)	173
C12—H12 <i>A</i> ...O1 <sup>iii</sup>	0.99	2.53	3.0852 (13)	116
C13—H13...O2 <sup>iv</sup>	0.95	2.63	3.3909 (13)	137
C16—H16 <i>C</i> ...O5 <sup>v</sup>	0.98	2.59	3.2434 (14)	124
C17—H17...Br1 <sup>vi</sup>	0.95	3.03	3.9069 (11)	153
C14—H14...O2	0.95	2.36	2.6937 (12)	100

Symmetry codes: (i)  $-x+2, -y+2, -z+2$ ; (ii)  $-x+2, -y+2, -z+1$ ; (iii)  $x, y-1, z+1$ ; (iv)  $-x+1, -y+2, -z+2$ ; (v)  $-x+3, -y+1, -z+2$ ; (vi)  $x, y-1, z$ .