



Microbial Ecology of Thailand Tsunami and Non-Tsunami Affected Terrestrials

Naraporn Somboonna^{1*}, Alisa Wilantho², Kruawun Jankaew³, Anunchai Assawamakin⁴, Duangjai Sangsrakru², Sithichoke Tangphatsornruang², Sissades Tongsim²

1 Department of Microbiology, Faculty of Science, Chulalongkorn University, Bangkok, Thailand, **2** Genome Institute, National Center for Genetic Engineering and Biotechnology, Pathumthani, Thailand, **3** Department of Geology, Faculty of Science, Chulalongkorn University, Bangkok, Thailand, **4** Department of Pharmacology, Faculty of Pharmacy, Mahidol University, Bangkok, Thailand

Abstract

The effects of tsunamis on microbial ecologies have been ill-defined, especially in Phang Nga province, Thailand. This ecosystem was catastrophically impacted by the 2004 Indian Ocean tsunami as well as the 600 year-old tsunami in Phra Thong island, Phang Nga province. No study has been conducted to elucidate their effects on microbial ecology. This study represents the first to elucidate their effects on microbial ecology. We utilized metagenomics with 16S and 18S rDNA-barcode pyrosequencing to obtain prokaryotic and eukaryotic profiles for this terrestrial site, tsunami affected (S_1), as well as a parallel unaffected terrestrial site, non-tsunami affected (S_2). S_1 demonstrated unique microbial community patterns than S_2 . The dendrogram constructed using the prokaryotic profiles supported the unique S_1 microbial communities. S_1 contained more proportions of archaea and bacteria domains, specifically species belonging to Bacteroidetes became more frequent, in replacing of the other typical florae like Proteobacteria, Acidobacteria and Basidiomycota. Pathogenic microbes, including *Acinetobacter haemolyticus*, *Flavobacterium* spp. and *Photobacterium* spp., were also found frequently in S_1 . Furthermore, different metabolic potentials highlighted this microbial community change could impact the functional ecology of the site. Moreover, the habitat prediction based on percent of species indicators for marine, brackish, freshwater and terrestrial niches pointed the S_1 to largely comprise marine habitat indicating-species.

Citation: Somboonna N, Wilantho A, Jankaew K, Assawamakin A, Sangsrakru D, et al. (2014) Microbial Ecology of Thailand Tsunami and Non-Tsunami Affected Terrestrials. PLoS ONE 9(4): e94236. doi:10.1371/journal.pone.0094236

Editor: Vasu D. Appanna, Laurentian University, Canada

Received: December 16, 2013; **Accepted:** March 13, 2014; **Published:** April 7, 2014

Copyright: © 2014 Somboonna et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The research was supported by Research Funds from the Faculty of Science, Chulalongkorn University, Under the A1B1-NS (RES-A1B1-NS-01), and Thai Aviation Refuelling Co., Ltd. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors received funding from Thai Aviation Refuelling Co., Ltd. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials.

* E-mail: Naraporn.S@chula.ac.th

Introduction

Phra Thong island, Phang Nga province of southern Thailand (Figure 1), represents a location for comparative studies of tsunami (S_1) and non-tsunami (S_2) affected terrestrial ecosystems. The S_1 and S_2 shared nearby geographies separated by a hill, whereby S_1 terrain was inundated by the Indian Ocean tsunami on 26 December 2004 and S_2 unaffected; otherwise both were comparable based on geological characteristics [1,2]. The tsunami left an Andaman Sea-facing, S_1 , distinguished terrestrial layer that was classified by geologist as a sand layer of 5–20 cm thick (layer A in Figure 1; [1]). Interestingly, geological evidence indicated three historic tsunamis also occurred prior to the 2004 tsunami at S_1 , and none to S_2 . The youngest recorded historic tsunami predating the 2004 tsunami was approximately 600 years ago (600yo) (layer B in Figure 1; [1]).

Each tsunami occurrence could affect the S_1 terrestrial characteristics due to the massive impact of seawater with marine organisms and garbage [1,3,4]. Studies comparing the 2004 tsunami affected versus non-affected (or pre-affected) terrestrials and terrestrial water reported the greater salinity, acidity, conductivity, turbidity and organic contents following the tsunami occurrence [5–7]. Studies also reported widespread disease-carrying vectors, such as mosquitoes, trematodes and snails, after

the 2004 tsunami [3,4]. Several bacterial and fungal infections involved skin and respiratory disorders were documented among repatriated tourists [8] and people working in the tsunami affected area [9]. In addition, the 2004 tsunami sediments consisted of higher concentrations of Mercury and Thallium [10,11]. Together, this chance of terrestrial characteristics could affect the microbial biodiversity and functional ecology.

Nonetheless, the impact of tsunamis on microbial diversity and ecology function remains ill-defined. The present study thereby analyzed the microbial biodiversity and their potential functional composites in the tsunami impacted S_1 terrain, in comparison to the non-affected S_2 site, using 16S and 18S rRNA genes pyrosequencing derived metagenomic DNA approach. For each site, the data included the prokaryotic and eukaryotic diversity profiles categorized into different depth levels corresponding to the terrestrial ages: 2004 tsunami, 1–300yo (pre-dating the 2004), 300–600yo, 600yo tsunami, and >600yo, respectively (starting from the top layer to a deeper layer), and also the amalgamated profiles for each site. Geologists determined the terrestrial age period from its depths below the land surface [1]. The overall results represent for the first time the use of metagenomics in analysing the prokaryotic and eukaryotic microbial biodiversity of the 2004 tsunami and non-tsunami affected terrestrials. Unlike





Figure 1. Index map of Phra Thong island relative to Phuket and terrestrial sites where samples were collected. The lower left photograph shows the pit wall of tsunami affected site (S_1). Light color sheets **A** and **B** represent 2004 tsunami and 600yo tsunami affected terrestrial layers, respectively [1]. The lower right photograph shows the pit wall of non-tsunami affected site (S_2) of the parallel geology, and samples of equivalent depths to those of S_1 were collected. Time period of the terrestrial is determined via sample depth [1]. doi:10.1371/journal.pone.0094236.g001

traditional biodiversity study that is conducted via cultivation technique and could reveal merely less than 1% of the true microbiota, metagenomics is a culture-independent technique that has been proved worldwide a robust, reliable and comprehensive tool for obtainment of entire microbiota from diverse environmental and clinical samples [12–17].

Materials and Methods

Sample collection

The owners of the lands gave permission to conduct the study on these sites. We confirm that the study did not involve endangered or protected species.

Phra Thong island provides a location for comparative tsunami (S_1 : N9.13194 E98.26250) and non-tsunami (S_2 : N9.07250 E98.27222) affected terrestrial studies based on geological evidences (Jankaew, personal communication) [1,2]. S_1 and S_2 are 6.73 km apart. S_1 is 0.40 km from the sea, and S_2 is 2.26 km from the sea (Figure 1). Approximately 1 kg samples were collected, each in sterile containers, between 11:00–15:00 hours during 23–24 March 2011. S_1 samples comprised: 2004 tsunami (14.5 cm), 1–300yo (22 cm), 300–600yo (29 cm), 600yo tsunami (38 cm), and >600yo (46 cm); S_2 samples comprised: S21 (14.5 cm), S22 (22 cm), S23 (29 cm), S24 (38 cm), and S25 (46 cm). The number in parenthesis represents the depth level where the sample was collected, and that entailed the approximate age of the sample relative to the year 2004. On-site records for color, texture and pH were done. All samples were transported in ice chest, stored in 4°C and processed for the next steps within 14 days.

Metagenomic DNA extraction and DNA quality examination

Each sample was mixed with a sterile spatula, and 15 g each was used for metagenomic DNA extraction [18]. Two independent metagenomic DNA extractions were performed per sample. The samples were dissolved in an extraction buffer (Epicentre, Wisconsin, USA) with Tween 20, low-speed centrifuged to remove large debris, and poured through four-layered sterile cheesecloth to remove particles and organisms of >30 μ m in size. Microorganisms between 0.22 and 30 μ m were collected by filtering over a sterile 0.22 μ m filter membrane (Merck Millipore, Massachusetts, USA) [15]. Total nucleic acid from each sample was extracted using Meta-G-Nome DNA Isolation Kit (Epicentre) following the manufacturer's protocols. Metagenomic DNA quality was assessed using agarose gel electrophoresis. The DNA concentration and purity was further analysed by A_{260} and the ratio of A_{260}/A_{280} spectrophotometry, respectively.

PCR generation of pyrotagged 16S and 18S rDNA libraries

Table 1 lists forward and reverse pyrotagged 16S and 18S rRNA gene primers. For broad-range 16S and 18S rRNA genes amplification, universal prokaryotic 338F (forward) and 803R (reverse) primers [19–21], and universal eukaryotic 1A (forward) and 516R (reverse) primers [15,22,23] were used. Italics denote the eight nucleotides pyrotag sequences, functioning to specify sample names [24]. A 50- μ l PCR reaction comprised 1 \times

EmeraldAmp GT PCR Master Mix (TaKaRa, Shiga, Japan), 0.3 μ M of each primer, and 100 ng of the metagenome. PCR conditions were 95°C for 4 min, and 30–35 cycles of 94°C for 45 s, 50°C for 55 s and 72°C for 1 min 30 s, followed by 72°C for 10 min. To generate the pyrotagged 16S or 18S rDNA libraries with minimized stochastic PCR biases, two to three independent PCRs were performed per extracted metagenomes, and two extracted metagenomes per sample, resulting in a minimum of four PCR products to be pooled for pyrosequencing per sample.

Gel purification and pyrosequencing

PCR products (~473 bp for 16S rDNAs; ~577 bp 18S rDNAs) were excised from agarose gels, and purified using PureLink Quick Gel Extraction Kit (Invitrogen, New York, USA). The 454-sequencing adaptors were ligated to all 16S and 18S rDNA fragments, the reactions were purified by MinElute PCR Purification Kit (Qiagen), and the samples were pooled for pyrosequencing on an eight-lane Roche picotiter plate, 454 GS FLX system (Roche, Branford, CT) at the in-house facility of the National Center for Genetic Engineering and Biotechnology, according to the recommendations of the supplier.

Sequence annotation and bioinformatic analyses

After removal of unreliable sequences, including sequences that failed the pyrosequencing quality cut-off and sequences shorter than 50 nucleotides, the sequences were categorized based on the appended pyrotag sequences. Sequences corresponding to the same sample category were inspected for domain and taxon compositions using mg-RAST [25,26] with default parameters. Species were identified by BLASTN [27] with E-value $\leq 10^{-5}$ against 16S rDNA databases including NCBI non-redundant [28], RDP [29] and Greengenes [30], and for 18S rDNAs the databases included NCBI non-redundant [28], EMBL [31,32] and SILVA [33]. Evolutionary distances and phylogenetic tree were computed with default thresholds (E-value $\leq 10^{-8}$, similarity score $\geq 80\%$). Species (or phylum) prevalence was determined by dividing the frequency of reads in the species (or phylum) by the total number of the identifiable reads. The differences in community structures were compared using Yue & Clayton theta similarity coefficients (Thetayc) and Morisita-Horn dissimilarity index, in mothur [34–36]. Low Thetayc and Morisita-Horn inferred high community similarity. An unweighted pair group method with arithmetic mean (UPGMA) clustering was constructed using Thetayc, in mothur [36]. Furthermore, functional subsystems and functional groups of the prokaryotic profiles were determined using SEED-based assignments in mg-RAST server [25,26,37]. For habitat classification, the data were compared against a World Register of Marine Species (WoRMS) database [38].

Results

Metagenome abundances and compositions at domain and kingdom levels

On-site records for physical characteristics of S_1 and S_2 were as shown in Figure 1. Alternating layers of black soil-like and grey sand-like comprised most of S_1 , whereby more homogeneous layers of grey sand-like predominated in S_2 . Differences in pH

Table 1. Pyrotagged16S and 18S rRNA genes universal primers.

Sample names	Forward primers (5'-3')	Reverse primers (5'-3')
16S rRNA gene universal primers		
2004 tsunami	TAGTAGCGACTCCTACGGGAGGCAGCAG	TAGTAGCGCTACCAGGGTATCTAATC
1-300yo	AGACGACGACTCCTACGGGAGGCAGCAG	AGACGACGCTACCAGGGTATCTAATC
300-600yo	ACTCGTAGACTCCTACGGGAGGCAGCAG	ACTCGTAGCTACCAGGGTATCTAATC
600yo tsunami	ACATCGAGACTCCTACGGGAGGCAGCAG	ACATCGAGCTACCAGGGTATCTAATC
>600yo	ACGCTATCACTCCTACGGGAGGCAGCAG	ACGCTATCTACCAGGGTATCTAATC
S21	TACTACGCACTCCTACGGGAGGCAGCAG	TACTACGCCTACCAGGGTATCTAATC
S22	AGCAGAGCACTCCTACGGGAGGCAGCAG	AGCAGAGCCTACCAGGGTATCTAATC
S23	TCAGCTACACTCCTACGGGAGGCAGCAG	TCAGCTACTACCAGGGTATCTAATC
S24	AGAGCGACTCCTACGGGAGGCAGCAG	AGAGCGACTACCAGGGTATCTAATC
S25	ATGCTCACACTCCTACGGGAGGCAGCAG	ATGCTCACCTACCAGGGTATCTAATC
18S rRNA gene universal primers		
2004 tsunami	AGATAGCGCTGGTTGATCCTGCCAGT	AGATAGCGACCAGACTTGCCCTCC
1-300yo	TGTAGACGCTGGTTGATCCTGCCAGT	TGTAGACGACCAGACTTGCCCTCC
300-600yo	TGCAGTAGCTGGTTGATCCTGCCAGT	TGCAGTAGACCAGACTTGCCCTCC
600yo tsunami	TCTGCGAGCTGGTTGATCCTGCCAGT	TCTGCGAGACCAGACTTGCCCTCC
>600yo	ATCAGCAGCTGGTTGATCCTGCCAGT	ATCAGCAGACCAGACTTGCCCTCC
S21	ATACAGTCCTGGTTGATCCTGCCAGT	ATACAGTACCAGACTTGCCCTCC
S22	ATCATATCCTGGTTGATCCTGCCAGT	ATCATATCACCAGACTTGCCCTCC
S23	TGCGATGCCTGGTTGATCCTGCCAGT	TGCGATGCACCAGACTTGCCCTCC
S24	ATCGCAGCCTGGTTGATCCTGCCAGT	ATCGCAGCACCAGACTTGCCCTCC
S25	TATACTACCTGGTTGATCCTGCCAGT	TATACTACACCAGACTTGCCCTCC

Italic sequence denotes the 8 nt-pyrotagged sequence.
doi:10.1371/journal.pone.0094236.t001

were also evident between the two ecosystems where S₁ ranged from 6–7 (more acidity), while S₂ ranged from 7–7.5.

Following total nucleic acids extraction of 0.22–30 μm sizes, S₁ and S₂ had average metagenomic concentration of 23.16 ng and 27.02 ng per gram of soil, respectively. Libraries of pyrotagged 16S and 18S rRNA gene fragments were constructed, and pyrosequenced to obtain the culture-independent prokaryotic and eukaryotic profiles of the sites. After removal of unreliable sequences, 21,592 reads for S₁ and 33,308 reads for S₂ remained for BLASTN species identification. Significant E-values ($\leq 10^{-5}$) were identified for 20,555 reads for S₁ (95.20%) and 31,946 reads for S₂ (95.91%). Reads with non-significant E-values ($> 10^{-5}$) were omitted from the analyses.

The domain compositions of S₁ indicated a lower proportion of eukaryotes and higher proportion of prokaryotes than S₂ (Figure 2). The proportion of eukaryotic species was reduced by almost half in S₁. Among prokaryotes, the greatest divergence between S₁ and S₂ was evident among archaea compared to bacteria where found increased in S₁ (Figure 3: 1.15-fold for bacteria, 4.07-fold archaea). These greater representations somewhat caused the reduced diversity of the other 4 kingdoms of lives.

Diversity of prokaryotic phyla and species

Major prokaryotic phyla for S₁ included Proteobacteria, Bacteroidetes, and Actinobacteria; and S₂ included Proteobac-

teria, Acidobacteria, and Actinobacteria, in diminishing order (Figure 4A). Consistent with Figure 3, S₁ demonstrated greater prokaryotic biodiversity than S₂. S₁ contained many new species belonging to uncultured species, i.e. OP3, GNO4 and SC3, whereas S₂ still contained high proportion of common environmental phyla, including Proteobacteria and Acidobacteria (Figure 4A). Different sample periods showed slight variation of prokaryotic phyla profiles in S₁, whereas in S₂ more variation among the phyla distributions was evident (Figure 4B). For instances, Proteobacteria comprised 62.23% in the S₁ layer of 2004 tsunami, 58.56% 1–300yo, 59.18% 300–600yo, 60.69% 600yo tsunami, and 62.54% >600yo; while S₂ comprised 46.07% in the S21 layer, 65.72% S22, 77.64% S23, 75.05% S24, and 67.41% S25. Actinobacteria comprised 14.48% in S₁ layer of 2004 tsunami, 13.22% 1–300yo, 13.53% 300–600yo, 13.82% 600yo tsunami, and 13.57% >600yo; while S₂ layers demonstrated 14.32% S21, 7.59% S22, 6.52% S23, 8.92% S24, and 10.47% S25 (Figure 4B). Further distinguished differences were diagnosed upon analysis based on species distributions: no parallel-age pairs of S₁ and S₂ showed similar species distribution pattern with the >600 yo and S25 pair showing the least differences. Examples of predominated species in S₁ were *Acinetobacter haemolyticus*, *Polynucleobacter* sp., *Polynucleobacter necessarius* and *Flavobacterium denitrificans*; and S₂ were *Burkholderia* sp. and *Silvimonas terrae* (Figure 5). Subsequently, the computed Thetayc and Morisita-Horn dissimilarity indices revealed dissimilar community structures between S₁




Figure 2. Percentages of prokaryotic and eukaryotic domains in S₁ and S₂.

doi:10.1371/journal.pone.0094236.g002

and S₂. As shown in UPGMA dendrogram in Figure 6A: terrestrial layers corresponding to S₁ site were clustered together separated from the S₂ layers. Although S21 was grouped with the S₁ layers, it was an outer related branch to the S₁ group.

Potential metabolic system analysis of prokaryotic communities

From a total of 28 possible metabolic subsystems by mg-RAST [25,26,37], the prokaryotic communities of S₁ contained 19 subsystems, and S₂ contained 11 subsystems. The predominant subsystems in the S₁ included: regulation and cell signalling (14.35%), cell wall and capsule (10.05%), protein metabolism (8.13%), and sulfur metabolism (3.83%) (Figure 7). Overall, S₁ inhabited prokaryotic communities with high metabolic potentials for cell wall and capsule (i.e. gram-negative cell wall components by TIdE/PmbA protein), protein metabolism (i.e. protein degradation by prolyl endopeptidase), regulation and cell signaling (i.e. regulation of virulence by two-component system response regulator), and carbohydrates (i.e. sugar alcohols and central carbohydrate metabolism by HPr kinase/phosphorylase). S₂ inhabited prokaryotic communities with high metabolic potentials for respiration (i.e. electron donating and accepting reactions by Type cbb3 cytochrome oxidase biogenesis protein), clustering based subsystems (i.e. copper-translocating P-type ATPase), and virulence, disease and defence (resistance to antibiotics and toxic compounds by acriflavin resistance protein).

Diversity of eukaryotic phyla and species

Dominant eukaryotic phyla for S₁ were in kingdom Animalia: Brachiopoda (47.82% in 2004 tsunami, 32.53% 1–300yo, 37.32% 300–600yo, 49.15% 600yo tsunami, 25.00% >600yo), and Mollusca (28.88% 2004 tsunami, 29.39% 1–300yo, 31.34% 300–600yo, 30.51% 600yo tsunami, 7.14% >600yo); and kingdom Protozoa: Dinophyta for particularly the >600yo layer (51.99%) (Figures 8A and 8B). For S₂, although Brachiopoda and Mollusca were dominant, fungal phylum Basidiomycota (0.08% S21, 66.51% S22, 7.47% S23, 12.82% S24, 12.86% S25) and animal phylum Arthropoda (3.05% S21, 3.42% S22, 28.33% S23, 7.70% S24, 2.86% S25) were the most prevalent. When analyzing the data into individual sample periods, similar finding to Figure 4B were found. Different sample periods of S₁ demonstrated less phyla pattern variation than those of S₂ (Figure 8B). Distinguished phyla pattern of S₂ from S₁ were displayed apparently in S22, S23 and S24 layers (Figure 8B), resulting in their divergence from S₁ and the other S₂ communities by the UPGMA dendrogram constructed using Thetayc dissimilarity indices (Figure 6B). Similar to prokaryotes (Figure 6A), the eukaryotic communities corresponding to the S₁ site were relatively clustered together (Figure 6B). Analysis at the species level identified a more diverse fungal and animal species among S₂ layers (Figure 9).

Habitat classification

The S₁ and S₂ prokaryotic and eukaryotic profiles were matched against WoRMS database [38] to further characterize their microbial ecology: how each is related to marine, brackish water, freshwater, and terrestrial species communities. Figure 10 exhibited a substantially higher abundance of marine species habitat with S₁ (S₁ = 24.11%, S₂ = 13.33%), and terrestrial species with S₂ (S₁ = 0.11%, S₂ = 1.42%). Examples of abundant marine prokaryotes in S₁ were: *Lutaonella thermophilus*, *Shevanella aquimarina*, *Erythrobacter ishigakiensis* and *Thalassobacter stenotrophicus*. Abundant marine eukaryotes in S₁ included: *Dinophysis acuminata*, *Ctenodrilidae sp.*, *Remanella sp.*, *Nemertinoidea elongatus*, *Skeletonema grethae*, *Crassostrea gigas*, *Hymenocotta mulli*, *Diplodasys ankei*, *Pinna muricata*, *Arenicola marina*, *Limopsis marionensis* and *Haliplanella lucia*.

Discussion

On-site records indicated the greater turbidity (Figure 1) and acidity of S₁ were in agreement with previous reports [1,2,5–7]. Together with many other tsunami studies, these different soil types suggested terrestrial component changes following tsunami inundation, which could affect the microbial ecology of the site. The terrestrial microbiome representing the 2004 tsunami-affected site has never been studied. Our findings represent the first to utilize metagenomics in gaining databases of these entire terrestrial microbiomes, including prokaryotes and eukaryotes, in tsunami-affected (S₁) and non-tsunami affected (S₂) sites of Phra Thong island, as of March 2011. The data helped characterize the microbial biodiversity and its impact by tsunami occurrence. This knowledge is essential for scientists and engineers involved with land management and environmental bio-improvement.

Diminished total nucleic acids from S₁ suggested a less populated microbial community. Although some fossil DNA and fragments of DNA from live animals (known as extracellular "dirt" DNA) could be included in the extracted metagenomes, and might partly complicate the analysis. Andersen et al. [39] found extracellular "dirt" DNA from the terrestrial surface could reflect an overall taxonomic richness and relative abundance of species of a site at the time of investigation. Hence, some extracellular "dirt"




Figure 3. Percentages of 6 kingdoms of lives in S₁ and S₂.
doi:10.1371/journal.pone.0094236.g003

DNA in our extracted metagenome should also reflect an overall biodiversity.

Libraries of pyrotagged 16S and 18S rRNA gene fragments were successfully constructed and pyrosequenced: 21,592 reads for S₁ and 33,308 reads for S₂ were retrieved after removal of unreliable sequences. For BLASTN species identification, greater than 95% of the S₁ and S₂ reads were identified with $\leq 10^{-5}$ E-values. The amount of reads should be sufficient to recapture the relationships among the samples, as Caporaso et al. [40] reported 2,000 reads could recapture the same relationships among samples as did with the full dataset. Additionally, many studies discovered variable regions 3 and 4 of 16S rRNA gene analyses were more effective than random sequence reads analyses in estimating the biodiversity and relationships among the samples [41–43].

In S₁, domain of prokaryotes became highly present (Figure 2). In particular, S₁ had a richer archaeal population (4.07-fold increase), meanwhile fungi, animals, plants, and protists were decreased (Figure 3). This finding was consistent with the fact that

tsunami inundation might leave a terrestrial site inhospitable, causing archaea and bacteria to be more common due to their flexible life activities and requirements [44–46]. The greater biodiversity of kingdoms in S₂ supported the more hospitable terrestrial habitat than S₁.

Phyla and species distribution patterns between S₁ and S₂ were different. Changing the prokaryotic pattern of major phyla, precisely S₁ was predominantly comprised of Bacteroidetes with a lower prevalence of Proteobacteria, Actinobacteria and Acidobacteria (Figure 4A), highlighted the modified microbial ecology. Wada et al. [47] reported the similar change of bacterial floras in the sludge brought ashore by the 2011 East Japan earthquake. Bacteroidetes was more evident than Proteobacteria in the affected coastal water area. Additionally, numerous sulfate-reducing bacteria were evident in the sludge, which corresponded with high concentrations of sulfate ions in the sludge and the affected water area. The latter report was consistent with our finding of the higher sulfur metabolism in S₁ (Figure 7). Further, among the Bacteroidetes, flavobacteria predominated which is generally




Figure 4. Distribution of prokaryotic phyla in S₁ and S₂, without (A) and with (B) individual sample ages categorization.
doi:10.1371/journal.pone.0094236.g004

classified as an environmental bacterium with both commensal and pathogen species of marine animals and humans. Banning et al. [48] discovered several flavobacteria strains in various marine environments could function as predators on other bacteria. These bacteria have minimal growth requirements, only sea salt and the utilization of the lysed bacteria. The marine Flavobacteria thus could have critical consequences on microbial ecology as they could eradicate certain microbial communities [48]. Additionally, *Polynucleobacter* sp. are bacterioplankton that could survive broad ecological niches due to their ability to obtain energy by consuming organic materials from other organisms through nitrogen fixation, nitrification, remineralisation and methanogenesis. *Photobacterium* sp. is also a genus with metabolic versatility, which can degrade chitin and cellulose for carbohydrates [49]. Consequently, the flavobacteria, bacterioplankton and photobacteria activities could partly support the high metabolic subsystems of carbohydrates and protein metabolism in S₁ (Figure 7), albeit the overall poor living condition. Note the increased regulation and cell signalling, and cell wall and capsule subsystems (Figure 7) could in part symbolize the growth activities of these bacteria, given capsule lies outside the bacterial cell wall and considered a virulent factor. Bacterial capsule protects the bacteria against some hostile environment, such as desiccation, and prevents phagocytosis by host immune cells [50]. For examples, *Acinetobacter haemolyticus* in contaminated seafood produces shiga toxin that causes bloody diarrhea [51], and *Flavobacterium* sp. cause cold water disease in salmon and other fish species [52]. *Photobacterium*, a genus in family *Vibrionaceae*, is primarily marine microorganisms that evolved to become pathogenic to marine animals, causing mortality in crabs and fish, and indirect pathogens of humans through contact or consumption [49,53]. Hence residents and workers in these areas were recommended to minimize direct contact with the affected soil,

sludge and water, to prevent their risk of infection, and frequent hand wash [3,8,9,47]. Note the many new uncultured species in S₁ (Figure 4A) further emphasized its environmental change resulting in new identified species. Our 16S rRNA gene analyses supported the high dissimilarity indices between S₁ and S₂ prokaryotic community structures (Figure 6A).

The metabolic potentials in Figure 7 supported the prior results, showing advanced metabolic subsystems of regulation and cell signalling, cell wall and capsule, protein metabolism, sulfur metabolism, and carbohydrates in S₁. In contrast, S₂ microbial communities carried high metabolic potentials for pathways of respiration, photosynthesis, and drug and bioactive compound production. This finding supported the diversified biodiversity in the non-affected terrestrials, and highlighted the more abundant pharmaceutical related microbial producers in the naturally undisturbed environments [15], like S₂.

For eukaryotic phyla and species distribution patterns, while both mollusks and brachiopods predominated in both terrestrials, given both animals were marine animals and were more prominent in tsunami-inundated S₁ site, fungi Basidiomycota and animal Arthropoda were only highly proportionate in the S₂ area (Figure 8A). Like prokaryotic phyla distribution patterns among various terrestrial depths (Figure 4B), the more similar eukaryotic phyla distribution patterns among various terrestrial depths were evident in S₁ (Figure 8B) highlighted the factor that a massive tsunami hit could destroy the biodiversity within microbial ecosystems. Basidiomycota, which were found more evident in S₂, are higher fungi that play important roles as carbon recycler and nutrient decomposer, and posed the chief source of bioactive natural products [54–56].

Since the 2004 tsunami up to our study period, an effect of microbial population mixing and microbial change due to human or animal activity on S₁ and S₂ sites in Phra Thong island should




Figure 5. Distribution of prokaryotic species in S₁ and S₂, categorized by individual sample ages. Different color on the diagram represents a different relative abundance, based on the percent frequency chart on the right.
doi:10.1371/journal.pone.0094236.g005

be minimal. The reasons are because, after the 2004 tsunami, Phra Thong island remains almost no human inhabited and no human activity. The place becomes part of the wildlife sanctuary.

Microbial population mixing through time could only be by rainfall and plant root penetration (mostly grass at both sites); hence it should be minimal and in vertical direction only.



A**B**

Figure 6. UPGMA clustering comparing relatedness among S_1 and S_2 prokaryotic (A) and eukaryotic (B) profiles.

doi:10.1371/journal.pone.0094236.g006

Together, the 16S and 18S rRNA gene profiles indicated both terrestrials marine habitat, corresponding to the fact that the two sites were located on a small island in Andaman Sea of Thailand. Nevertheless, twice the greater prediction for marine habitat for S_1 (Figure 10) highlighted its much higher number of prokaryotic and eukaryotic species that represent marine habitat species-indicators, and tsunami inundation.

Conclusion

During the past decades, tsunamis have occurred more frequently though the correlation between tsunami disturbance and change of terrestrial microbial ecology remain poorly defined. The present study provided a culture-independent prokaryotic and eukaryotic analyses representing 0.22–30 μm metagenomes belonging to Thailand tsunami and non-tsunami affected terrestrials.




Figure 7. Metabolic subsystems of prokaryotic communities in S₁ and S₂.
doi:10.1371/journal.pone.0094236.g007




Figure 8. Distribution of eukaryotic phyla in S₁ and S₂, without (A) and with (B) individual sample ages categorization.
doi:10.1371/journal.pone.0094236.g008




Figure 9. Distribution of eukaryotic species in S₁ and S₂, categorized by individual sample ages. Different color on the diagram represents a different relative abundance, based on the percent frequency chart on the right. doi:10.1371/journal.pone.0094236.g009

The different prokaryotic and eukaryotic profiles highlighted the differences due to tsunami, and helped fulfill our knowledge of diverse terrestrial microbial ecologies. The biodiverse species of S₁ distinguished its microbial communities and metabolic potentials.

For instances, the finding of predator and prey bacterial relationship, and cell wall and capsule subsystem were common for S₁, whereas bioactive compound producers were more common in S₂.




Figure 10. Habitat classification for S₁ and S₂. Prokaryotic and eukaryotic species profiles of S₁ and S₂ were matched against WoRMS database for habitat classification. doi:10.1371/journal.pone.0094236.g010

Further, the marine habitat analysis demonstrating the greater percent of marine prokaryotic and eukaryotic species in S₁ could perhaps help serve as another biomarker for the geological history of a terrestrial site. Nonetheless, more researches are required to utilize these as biomarkers for estimating times and presence of any geological incidences. Presently, identification of historic tsunami was restricted to examination of marine and brackish diatoms while silica shell of diatom could be dissolved through time, especially in hot weather of a tropical country [1,2,57,58].

References

- Jankaew K, Atwater BF, Sawai Y, Choowong M, Charoentitrat T, et al. (2008) Medieval forewarning of the 2004 Indian ocean tsunami in Thailand. *Nature* 455:1228–1231.
- Sawai Y, Jankaew K, Martin ME, Prendergast A, Choowong M, et al. (2009) Diatom assemblages in tsunami deposits associated with the 2004 Indian Ocean tsunami at Phra Thong Island, Thailand. *Mar Micropaleontol* 73:70–79.
- Sri-Aroon P, Chusongsang P, Chusongsang Y, Pornpimol S, Butraporn P, et al. (2010) Snails and trematode infection after Indian ocean tsunami in Phang-Nga province, southern Thailand. *Southeast Asian J Trop Med Public Health* 41: 48–60.
- Prummongkol S, Panasoponkul C, Apiwathnasorn C, Lek-Uthai U (2012) Biology of *Culex sitiens*, a predominant mosquito in Phang Nga, Thailand after a tsunami. *J Insect Sci* 12: 11. doi:10.1673/031.021.1101
- Tharnpoophasiam P, Suthisarnsunton U, Worakhunpiset S, Charoenjai P, Tunyong W, et al. (2006) Preliminary post-tsunami water quality survey in Phang-Nga province, southern Thailand. *Southeast Asian J Trop Med Public Health* 37: 216–220.
- Collivignarelli C, Tharnpoophasiam P, Vaccari M, De Felice V, Di Bella V, et al. (2008) Evaluation of drinking water treatment and quality in Takua Pa, Thailand. *Environ Monit Assess* 142: 345–358.
- Collivignarelli C, Tharnpoophasiam P, Vaccari M, De Felice V, Di Bella V, et al. (2008) Water monitoring and treatment for drinking purposes in 2004 tsunami affected area-Ban Nam Khem, Phang Nga, Thailand. *Environ Monit Assess* 147: 191–198.
- Chastel C (2007) Assessing epidemiological consequences two years after the tsunami of 26 December 2004. *Bull Soc Pathol Exot* 100: 139–142.
- Huansom AJ, Agner T, Backer V, Ebbehoj N, Jacobsen P (2012) Skin and respiratory disorders following the identification of disaster victims in Thailand. *Forensic Sci Med Pathol* 8: 114–117.
- Boszke L, Astel A (2007) Fractionation of mercury in sediments from coastal zone inundated by tsunami and in freshwater sediments from the rivers. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 42: 847–858.
- Lukaszewski Z, Karbowska B, Zembrzusi W, Siepak M (2012) Thallium in fractions of sediments formed during the 2004 tsunami in Thailand. *Ecotoxicol Environ Saf* 80: 184–190.
- Rusch DB, Halpern AL, Sutton G, Heidelberg KB, Williamson S, et al. (2007) The Sorcerer II Global Ocean Sampling expedition: northwest Atlantic through eastern tropical Pacific. *PLoS Biol* 5:e77.
- Yooseph S, Neelson KH, Rusch DB, McCrow JP, Dupont CL, et al. (2010) Genomic and functional adaptation in surface ocean planktonic prokaryotes. *Nature* 468:60–66.
- Zinger L, Ammaral-Zettler LA, Fuhman JA, Horner-Devine MC, Huse SM, et al. (2011) Global patterns of bacterial beta-diversity in seafloor and seawater ecosystems. *PLoS ONE* 6: e24570. doi:10.1371/journal.pone.0024570

Acknowledgments

The authors thank Sarah Hof and Khunluck Kidmoa for general help, Dominik Brill for helping with field work, and Troy Skwor for manuscript editing.

Author Contributions

Conceived and designed the experiments: NS AA. Performed the experiments: NS DS S. Tangphatsornruang. Analyzed the data: NS AW. Contributed reagents/materials/analysis tools: NS KJ S. Tongsimma. Wrote the paper: NS. Revised the manuscript: NS KJ S. Tongsimma.

15. Somboonna N, Assawamakin A, Wilantho A, Tangphatsornruang S, Tongshima S (2012) Metagenomic profiles of free-living archaea, bacteria and small eukaryotes in coastal areas of Sichang island, Thailand. *BMC Genomics* 13:S29. doi:10.1186/1471-2164-13-S7-S29
16. Auld RR, Myre M, Myktyczuk NCS, Leduc LG, Merritt TJS (2013) Characterization of the microbial acid mine drainage microbial community using culturing and direct sequencing techniques. *J Microbiol Methods* 93: 108–115.
17. Redel H, Gao Z, Li H, Alekseyenko AV, Zhou Y, et al. (2013) Quantitation and composition of cutaneous microbiota in diabetic and nondiabetic men. *JID* 207: 1105–1114.
18. Taberlet P, Prud'Homme SM, Campione E, Roy J, Miquel C, et al. (2012) Soil sampling and isolation of extracellular DNA from large amount of starting material suitable for metabarcoding studies. *Mol Ecol* 21:1816–1820.
19. Baker GC, Smith JJ, Cowan DA (2003) Review and re-analysis of domain-specific 16S primers. *J Microbiol Methods* 55:541–555.
20. Humblot C, Guyot J-P (2009) Pyrosequencing of tagged 16S rRNA gene amplicons for rapid deciphering of the microbiomes of fermented foods such as pearl millet slurries. *Appl Environ Microbiol* 75: 4354–4361.
21. Nossa CW, Oberdorf WE, Yang L, Aas JA, Paster BJ, et al. (2010) Design of 16S rRNA gene primers for 454 pyrosequencing of the human foregut microbiome. *WJG* 16:4135–4144.
22. Grant S, Grant WD, Cowan DA, Jones BE, Ma Y, et al. (2006) Identification of eukaryotic open reading frames in metagenomic cDNA libraries made from environmental samples. *Appl Environ Microbiol* 72:135–143.
23. Bailly J, Fraissinet-Tachet L, Verner M-C, Debaud J-C, Lemaire M, et al. (2007) Soil eukaryotic functional diversity, a metatranscriptomic approach. *ISME J* 1:632–642.
24. Meyer M, Stenzel U, Hofreiter M (2008) Parallel tagged sequencing on the 454 platform. *Nat Prot* 3:267–278.
25. Overbeek R, Begley T, Butler RM, Choudhuri JV, Chuang HY, et al. (2005) The subsystems approach to genome annotation and its use in the project to annotate 1000 genomes. *Nucleic Acids Res* 33: 5691–5702.
26. Meyer F, Paarmann D, D'Souza M, Olson R, Glass EM, et al. (2008) The metagenomics RAST server - a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* 9: 386. doi:10.1186/1471-2105-9-386
27. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, et al. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389–3402.
28. Sayers EW, Barrett T, Benson DA, Bolton E, Bryant SH, et al. (2010) Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 38: D5–D16.
29. Maidak BL, Cole JR, Libum TG, Parker CT Jr, Saxman PR, et al. (2001) The RDP-II (ribosomal database project). *Nucleic Acids Res* 29: 173–174.
30. McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, et al. (2012) An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J* 6:610–618.
31. Brunak S, Danchin A, Hattori M, Nakamura H, Shinozaki K, et al. (2002) Nucleotide sequence database policies. *Science* 298: 1333.
32. Leinonen R, Akhtar R, Birney E, Bower L, Cerdeno-Tarraga A, et al. (2011) The European nucleotide archive. *Nucleic Acids Res* 39: D28–D31.
33. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, et al. (2007) SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* 35: 7188–7196.
34. Yue JC, Clayton MK (2005) A similarity measure based on species proportions. *Commun Stat Theor Methods* 34: 2123–2131.
35. Chao A, Chazdon RL, Colwell RK, Shen T-J (2006) Abundance-based similarity indices and their estimation when there are unseen species in samples. *Biometrics* 62: 361–371.
36. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, et al. (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 75: 7537–7541.
37. Aziz RK, Bartels D, Best AA, Dejongh M, Disz T, et al. (2008) The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9: 75. doi:10.1186/1471-2164-9-75
38. Appeltans W, Bouchet P, Boxshall GA, De Broyer C, de Voogd NJ, et al. (eds.) (2012) World Register of Marine Species [http://www.marinespecies.org]
39. Andersen K, Bird KL, Rasmussen M, Haile J, Breuning-Madsen H, et al. (2012) Meta-barcoding of 'dirty' DNA from soil reflects vertebrate biodiversity. *Mol Ecol* 21:1966–1979.
40. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, et al. (2010) Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *PNAS*: doi:10.1073/pnas.1000080107
41. Manichanh C, Chapple CE, Frangeul L, Gloux K, Guigo R, et al. (2008) A comparison of random sequence reads versus 16S rDNA sequences for estimating the biodiversity of a metagenomic library. *Nucleic Acids Res* 36: 5180–5188.
42. Maurice CF, Haiser HJ, Turnbaugh PJ (2012) Xenobiotics shape the physiology and gene expression of the active human gut microbiome. *Cell* 152: 39–50.
43. Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, et al. (2013) Gut microbiota from twins discordant for obesity modulate metabolism in mice. *Science* 341. doi:10.1126/science.1241214
44. Henry EA, Devereux R, Maki JS, Gilmour CC, Woese CR, et al. (1994) Characterization of a new thermophilic sulfate-reducing bacterium *Thermodesulfobacterium yellowstonii*, gen.nov. and sp.nov.: its phylogenetic relationship to *Thermodesulfobacterium commune* and their origins deep within the bacterial domain. *Arch Microbiol* 161: 62–69.
45. Huang LN, Zhou H, Chen YQ, Luo S, Lan CY, et al. (2002) Diversity and structure of the archaeal community in the leachate of a full-scale recirculating landfill as examined by direct 16S rRNA gene sequence retrieval. *FEMS Microbiol Lett* 214:235–240.
46. Kendall MM, Liu Y, Sieprawska-Lupa M, Stetter KO, Whitman WB, et al. (2006) *Methanococcus aeolicus* sp. Nov., a mesophilic, methanogenic aecheon from shallow and deep marine sediments. *Int J Syst Evol Microbiol* 56: 1525–1529.
47. Wada K, Fukuda K, Yoshikawa T, Hirose T, Ikeno T, et al. (2012) Bacterial hazards of sludge brought ashore by the tsunami after the great East Japan earthquake of 2011. *J Occup Health* 54: 255–262.
48. Banning EC, Casciotti KL, Kujawinski EB (2010) Novel strains isolated from a coastal aquifer suggest a predatory role for flavobacteria. *FEMS Microbiol Ecol* 73: 254–270.
49. Vezzi A, Campanaro S, D'Angelo M, Simonato F, Vitulo N, et al. (2005) Life at depth: *Photobacterium profundum* genome sequence and expression analysis. *Science* 307: 1459–1461.
50. Yoshida K, Matsumoto T, Tateda K, Uchida K, Tsujimoto S, et al. (2000) Role of bacterial capsule in local and systemic inflammatory responses of mice during pulmonary infection with *Klebsiella pneumoniae*. *J Med Microbiol* 49: 1003–1010.
51. Grotiuz G, Sirok A, Gadea P, Varela G, Schelotto F (2006) Shiga toxin 2-producing *Acinetobacter haemolyticus* associated with a case of bloody diarrhea. *J Clin Microbiol* 44: 3838–3841.
52. Starliper CE (2011) Bacterial coldwater disease of fishes caused by *Flavobacterium psychrophilum*. *J Adv Res* 2: 97–108.
53. Osorio CR, Toranzo AE, Romalde JL, Barja JL (2000) Multiplex PCR assay for *ureC* and 16S rRNA genes clearly discriminates between both subspecies of *Photobacterium damsela*. *Dis Aquat Org* 40: 177–183.
54. Rosa LH, Machado KM, Rabello AL, Souza-Fagundes EM, Correa-Oliveira R, et al. (2009) Cytotoxic, immunosuppressive, trypanocidal and antileishmanial activities of Basidiomycota fungi present in Atlantic rainforest in Brazil. *Antonie Van Leeuwenhoek* 95: 227–237.
55. Qadri M, Johri S, Shah BA, Khajuria A, Sidiq T, et al. (2013) Identification and bioactive potential of endophytic fungi isolated from selected plants of the Western Himalayas. *SpringerPlus* 2: 8.
56. Jaszek M, Osinska-Jaroszuk M, Janusz G, Matuszewska A, Stefaniuk D, et al. (2013) New bioactive fungal molecules with high antioxidant and antimicrobial capacity isolated from *Cerrena unicolor* idiophasic cultures. *BioMed Res Intl* 2013. doi:10.1155/2013/497492
57. Hemphill-Haley E (1995) Diatoms evidence for earthquake-induced subsidence and tsunamis 300 yr ago in southern coastal Washington. *GSA Bulletin* 107: 367–378.
58. Hemphill-Haley E (1996) Diatoms as an aid in identifying late-Holocene tsunami deposits. *Holocene* 6: 439–448.