



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Detection of human coronaviruses in children with acute gastroenteritis

Minna Risku^{a,*}, Suvi Lappalainen^a, Sirpa Räsänen^{a,b}, Timo Vesikari^{a,b}

^a Department of Virology, Medical School, Biokatu 10, FI-33014 University of Tampere, Tampere, Finland

^b Department of Pediatrics, Tampere University Hospital, Tampere, Finland

ARTICLE INFO

Article history:

Received 1 November 2009

Received in revised form 17 February 2010

Accepted 17 February 2010

Keywords:

Coronavirus

Acute gastroenteritis

RT-PCR

ABSTRACT

Background: Human coronaviruses (HCoVs) are known respiratory pathogens. Moreover, coronavirus-like particles have been seen by electron microscope in stools, and SARS-HCoV has been isolated from intestinal tissue and detected in stool samples.

Objectives: To find out if HCoVs can be found in stools of children with acute gastroenteritis and to assess the significance of HCoVs in the etiology of acute gastroenteritis in children.

Study design: 878 stool specimens from children with acute gastroenteritis and 112 from control children were tested by RT-PCR to detect HCoV groups 1B, 2A and SARS. HCoVs were typed by sequencing all PCR positive samples.

Results: Twenty-two (2.5%) of the 878 stool specimens of children with acute gastroenteritis were positive for HCoVs. The following HCoV types were detected: OC43 (10 cases, 45.5%), HKU1 (6 cases, 27.3%), 229E (2 cases, 9.1%) and NL63 (4 cases, 18.2%). In 4 of the cases a HCoV was the only detected virus; in the remaining cases rotavirus or norovirus was found in the same sample. In control groups there were two HCoV positive samples of 112 tested.

Conclusions: This study shows that all known non-SARS HCoVs can be found in stools of children with acute gastroenteritis. On the basis of this study, the significance of coronaviruses as gastrointestinal pathogens in children appears minor, since most of the coronavirus findings were co-infections with known gastroenteritis viruses.

© 2010 Elsevier B.V. All rights reserved.

1. Background

The first human coronaviruses (HCoVs) 229E and OC43 were identified in the 1960s.^{1–4} These viruses are common causes of upper respiratory tract infections^{5,6} but also have association with lower respiratory tract disease especially in patients with underlying disease.^{7–9} Coronavirus-like particles have also been seen by electron microscope (EM) in stool samples of both diarrheic and healthy patients evoking discussion about human enteric coronaviruses.¹⁰ The clinical significance of these findings has been unclear and there are findings showing that such putative enteric coronaviruses are antigenically unrelated to OC43 and 229E viruses.¹¹

In 2003 interest towards coronaviruses increased when a new coronavirus was found to be a causative agent of SARS (severe acute respiratory syndrome).^{12–14} SARS-HCoV caused a serious lower respiratory tract infection with high mortality.^{15–17} The main symptoms were fever, chills, myalgia, cough and headache, but also diarrhea was common and in one study registered in 38.4%

of patients. In the same study SARS-HCoV was also isolated from intestinal tissue and viral RNA was found in 16% of stool samples, which was comparable to the detection rate in nasopharyngeal aspirates.¹⁸ In another cohort of patients diarrhea was seen in 73% of patients and viral RNA was detected in 97% of stool specimens. In this cohort it was suggested that the outbreak was caused by faulty sewage system whereby the transmission might be fecal–oral rather than respiratory.¹⁹ No association between the presence of diarrhea and mortality in SARS cases has been observed, however.²⁰

A fourth human coronavirus, HCoV-NL63, was identified in 2004 in the Netherlands. It was isolated from a 7-month-old child suffering from bronchiolitis and conjunctivitis, and was categorized to be a new group 1 coronavirus.²¹ Soon after, another group in the Netherlands independently detected the same virus in an 8-month-old child with pneumonia.²² Since the discovery, NL63 has been detected in patients with respiratory tract infection in several countries around the world, including Canada, Australia, Korea and France.^{23–26} NL63 has been associated with severe lower respiratory tract infection and croup.^{24,27} The seroprevalence in 6–12-month-old children for NL63 is 28.6–40.0%.²⁸

Less than a year later a second new coronavirus, HCoV-HKU1, was discovered in Hong Kong. This group 2 coronavirus was detected in a 71-year-old man with pneumonia.²⁹

* Corresponding author. Tel.: +358 40 190 1505; fax: +358 3 3551 8450.

E-mail address: minna.risku@uta.fi (M. Risku).

HKU1 has been found globally^{30–33} and in addition to respiratory samples it has been detected in stools, but no clear connection to enteric disease has been found.³⁴

2. Objectives

Considering the EM findings, presence of gastrointestinal symptoms in SARS, and the findings of HKU1 in stools, we wanted to find out if non-SARS human coronaviruses could be detected in stool samples of children with acute gastroenteritis using RT-PCR assay in order to assess the potential significance of coronaviruses in the etiology of acute gastroenteritis in children.

3. Study design

The clinical material for the study was collected in a prospective study of acute gastroenteritis in children Tampere and Kuopio University Hospital during a 2-year period 2006–2008 (Räsänen et al., unpublished). Healthy children ($N=36$), children with indeterminate fever and vomiting ($N=43$), and children with respiratory tract infection ($N=33$) were used as control groups. Also group A rotavirus, calicivirus (including norovirus genogroups I and II, and sapovirus), aichivirus and human bocavirus were studied from the same material. Adenovirus was not tested systematically but of 101 samples tested 13 were positive for adenovirus (six belonging to subgroup F, types 40 and 41) and none of these were co-infections with coronaviruses including two positive samples for coronavirus. A total of 878 samples from children with acute gastroenteritis, 43 samples from children with indeterminate fever and vomiting, 33 samples from children with respiratory tract infection and 36 samples from healthy children were tested for HCoVs.

4. Laboratory methods

Before RNA extraction the stool samples were diluted in phosphate-buffered saline (PBS) creating 10% stool suspension. Viral RNA was extracted using QIAamp Viral RNA Mini Kit (QIAGEN, Germany) according to the manufacturer's protocol. Extracted RNA was stored at -70°C until used.

Reverse transcription was done using random primers as previously described by Pang et al. (2005)³⁵ except that reaction contained $1\times$ First Strand Buffer (Invitrogen, USA) and the final concentration of dNTPs (Promega, USA) was 375 nM per each nucleotide. The produced cDNA was stored at -20°C .

We used a two-step nested PCR which was set up and optimized in our laboratory. Primers targeted to polymerase gene region were chosen because of regions conserved nature. HCoV strains TC-adapted OC43 (VR-1558) and 229E (VR-740) were purchased from ATCC (USA), and propagated in HCT-8 and MRC-5 cells, respectively. In addition, positive HKU1 and NL63 samples were provided by Professor Tobias Allander (Karolinska University, Sweden) and Professor Lia van der Hoek (University of Amsterdam, Netherlands). The specificity of PCR was confirmed by testing 35 different respiratory or enteric viruses and bacteria, with negative results.

The 1st PCR included a primer pair (fwd-GWTGGGAYTATCCNAARTGTGA and rev-YRTCATCASWNARAAT-CATCAT) universal for all coronaviruses with resulting amplicon of 437 bp in size. $10\ \mu\text{l}$ of the cDNA was added to $40\ \mu\text{l}$ of reaction mixture containing $1\times$ Green GoTaq[®] Flexi Buffer (Promega, USA), 2.5 mM of GoTaq[®] MgCl₂ (Promega, USA), 200 μM of each dNTP (Promega, USA), 2.5 U of GoTaq DNA polymerase (Promega, USA) and 0.5 μM of each primer (Sigma-Genosys Ltd., UK). PCR program was run as follows: 2 min at 94°C , 35 cycles of amplification (30 s at 94°C , 30 s at 54°C , 1 min at 72°C) and final extension at 72°C for 5 min. Program was run in GeneAmp[®] PCR system 9700 (PE

Applied Biosystems, USA) or in Thermal Cycler 2720 (Applied Biosystems, USA). Aqua Sterilisata H₂O was used as negative control and cell cultured 229E or OC43 as a positive control.

Nested PCR contained three primer pairs to distinguish three coronavirus groups: 1B, 2A and SARS. Sizes for the amplicons were 203 bp for group 1B (fwd-GTTGTTTATTCWAATGGTGG and rev-YCTATARCAATTATCATAMAG), 275 bp for group 2A (fwd-WYTRCG-TATTGTTAGTAGTTRGT and rev-CGTATACTWARATCTTCAATCTT) and 230 bp for SARS (fwd-TGCTGTAACCTATCACACCGT and rev-CGGACATACTGTGCTAGCTATCT). $2\ \mu\text{l}$ of 1st PCR-product was added to $48\ \mu\text{l}$ of reaction mixture containing $1\times$ Green GoTaq[®] Flexi Buffer (Promega, USA), 1.5 mM of GoTaq[®] MgCl₂ (Promega, USA), 200 μM of each dNTP (Promega, USA), 2.5 U of GoTaq DNA polymerase (Promega, USA) and 0.5 μM of each primer (Sigma-Genosys Ltd., UK). PCR program was run as follows: 2 min at 94°C , 35 cycles of amplification (30 s at 94°C , 30 s at 53°C , 30 s at 72°C) and final extension at 72°C for 5 min.

PCR-products were separated and recognized by gel electrophoresis, and all positive PCR products were confirmed to be coronaviruses and the specific type defined by sequencing. ABI PRISM[™] 310 Genetic Analyzer (Applied Biosystems, USA) was used in sequencing. Sequences were aligned and confirmed by using Sequencher[™] 4.8 program (Gene Codes Corporation, USA) and confirmed sequences were compared to reference strains by NCBI Blast[®]-program.

5. Results

Twenty-two (2.5%) of the 878 stool specimens of children with acute gastroenteritis were positive for HCoVs. A HCoV as a single pathogen was detected in only four of the samples (18.2% of the positive samples). In the remaining cases either norovirus or rotavirus was detected in the same sample (Table 1). In eleven (50%) of the 22 coronavirus positive cases there were symptoms of respiratory tract infection at the same time with gastroenteritis, or respiratory symptoms had been present before symptoms of gastroenteritis.

Three of the 4 patients with coronavirus as a single pathogen in stool sample had respiratory tract related symptoms. Patient 1 (Table 1) had cough and rhinitis, and had just recovered from otitis media. Patient 7 (Table 1) did not have any respiratory tract symp-

Table 1
Positive detections of human coronaviruses in 878 children with acute gastroenteritis.

Patient	Age (months)	Sex	Sample date	HCoV type	Other virus	Respiratory symptoms ^a
1	17	F	January 2007	OC43	None	Yes
2	27	M	February 2007	OC43	noro	None
3	12	F	February 2007	OC43	noro	Yes
4	41	F	February 2007	OC43	noro	None
5	13	F	March 2007	OC43	noro	Yes
6	27	F	March 2007	NL63	rota	None
7	17	M	April 2007	NL63	None	None
8	25	M	February 2007	NL63	noro	None
9	13	M	April 2007	OC43	noro	Yes
10	9	M	March 2007	OC43	noro	None
11	31	F	June 2007	NL63	None	Yes
12	45	M	November 2007	OC43	rota	Yes
13	27	F	December 2007	OC43	rota	Yes
14	18	M	January 2008	229E	rota	None
15	17	M	January 2008	HKU1	noro	Yes
16	29	M	February 2008	OC43	rota	None
17	17	F	February 2008	HKU1	noro	None
18	10	M	February 2008	HKU1	rota	None
19	75	M	March 2008	HKU1	None	Yes
20	21	M	April 2008	HKU1	rota	Yes
21	30	M	April 2008	229E	rota	Yes
22	16	F	April 2008	HKU1	rota	None

^a Including cough, rhinitis, tonsillitis, otitis media, pneumonia, laryngitis.




Fig. 1. Seasonal distribution of human coronavirus positive stool specimens of children with acute gastroenteritis.

toms. Patient 11 (Table 1) had tonsillitis and patient 19 (Table 1) had headache and dizziness in addition to symptoms of respiratory tract infection.

All non-SARS human coronavirus types were found, members of group 2A; OC43 (10 of the cases, 45.5%) and HKU1 (6 of the cases, 27.3%) were most common, whereas group 1B viruses 229E and NL63 were found only in 2 and 4 cases, respectively. No SARS or SARS-like viruses were found. Still there might be unknown coronaviruses that our PCR method did not detect in spite of the universal primers in the 1st PCR. Most HCoV positive cases were found from January to April (Fig. 1).

The age distribution of the coronavirus infected children was 9–75 months (median, 19.5 months), whereas in the total material the youngest child was 14 days and the oldest 14 years and 4 months (median, 17 months). Of the coronavirus positive children 59% were males.

Within the control groups two (1.8%) of the 112 stool samples were positive for HCoV. One of the cases was a 3-year-old female with pneumonia. OC43 was detected as the only pathogen in her stool samples on February 2007. Two days after sample collection she also developed symptoms of gastroenteritis. The second patient was a healthy female aged 2 years and 11 months tested on July 2007. OC43 was again detected in a stool sample as a single pathogen.

6. Discussion

Our study shows that human coronaviruses OC43, HKU1, 229E and NL63 can be found in stool samples of children with acute gastroenteritis. The significance of coronaviruses as gastrointestinal pathogens seems at most marginal, even though it is also possible that our PCR method did not detect all existing coronaviruses and there still might be unknown coronaviruses related to diarrheal disease. Most of the coronavirus findings were co-infections with well known enteric pathogens, norovirus and rotavirus. Furthermore, half of the patients with coronavirus in stools had symptoms related to respiratory tract infection and, therefore, HCoVs found in the stools could have originated from respiratory tract. Unfortunately, no specimens from respiratory tract were collected to confirm the presence of coronaviruses in the respiratory tract in these patients, and further studies are needed to evaluate simultaneous presence of HCoV in stools and respiratory tract. Even if

coronaviruses were found in respiratory tract in cases of acute gastroenteritis it would be difficult to determine whether they were primarily causing the respiratory or gastrointestinal symptom.

Studies with SARS have shown that RNA of SARS coronavirus can be detected in stool samples for more than 10 weeks after symptom onset.¹⁸ This elicits the question whether also non-SARS coronaviruses might be detected after a prolonged time from the original infection. Previous studies with EM showed that coronavirus-like particles can be seen in stools of both diarrheic and healthy patients.¹⁰ In our study one of the 36 healthy control patients had coronavirus detected in stool specimen and thus, there was no difference in the HCoV detection rate between the cases of acute gastroenteritis and control children. This study was hospital based and did not include mild cases of gastroenteritis treated at home or healthcare centers. Future studies should investigate such mild cases for HCoVs.

In conclusion, non-SARS human coronaviruses can be found in stool samples of children with acute gastroenteritis. However, such findings are rare and occur usually with other well established gastroenteritis viruses. HCoVs may also be found in occasional stool samples of children without gastroenteritis. Taken together, it appears that known HCoVs may at most have a minor etiologic role in the acute gastroenteritis of children.

Ethical approval

The study protocol and consent forms had been approved by the Ethics Committee of the Pirkanmaa Hospital District in 2006.

Funding

None.

Conflict of interest

None declared.

Acknowledgements

We thank Professor Tobias Allander and Professor Lia van der Hoek for providing control samples for setting up the PCR method. We also thank our studynurse Marjo Salonen for hard work, and all

our laboratory technicians and Anna Tiainen for excellent technical assistance in this project.

References

- Hamre D, Procknow JJ. A new virus isolated from the human respiratory tract. *Proc Soc Exp Biol Med* 1966;**121**:190–3.
- Almeida JD, Tyrrell DA. The morphology of three previously uncharacterized human respiratory viruses that grow in organ culture. *J Gen Virol* 1967;**1**:175–8.
- McIntosh K, Dees JH, Becker WB, Kapikian AZ, Chanock RM. Recovery in tracheal organ cultures of novel viruses from patients with respiratory disease. *Proc Natl Acad Sci USA* 1967;**57**:933–40.
- McIntosh K, Becker WB, Chanock RM. Growth in suckling-mouse brain of "IBV-like" viruses from patients with upper respiratory tract disease. *Proc Natl Acad Sci USA* 1967;**58**:2268–73.
- Macnaughton MR. Occurrence and frequency of coronavirus infections in humans as determined by enzyme-linked immunosorbent assay. *Infect Immun* 1982;**38**:419–23.
- Makela MJ, Puhakka T, Ruuskanen O, Leinonen M, Saikku P, Pimpimaki M, et al. Viruses and bacteria in the etiology of the common cold. *J Clin Microbiol* 1998;**36**:539–42.
- Vabret A, Mourez T, Gouarin S, Petitjean J, Freymuth F. An outbreak of coronavirus OC43 respiratory infection in Normandy, France. *Clin Infect Dis* 2003;**36**:985–9.
- Falsey AR, Walsh EE, Hayden FG. Rhinovirus and coronavirus infection-associated hospitalizations among older adults. *J Infect Dis* 2002;**185**:1338–41.
- El-Sahly HM, Atmar RL, Glezen WP, Greenberg SB. Spectrum of clinical illness in hospitalized patients with "common cold" virus infections. *Clin Infect Dis* 2000;**31**:96–100.
- Clarke SK, Caul EO, Egglestone SI. The human enteric coronaviruses. *Postgrad Med J* 1979;**55**:135–42.
- Luby JP, Clinton R, Kurtz S. Adaptation of human enteric coronavirus to growth in cell lines. *J Clin Virol* 1999;**12**:43–51.
- Peiris JS, Lai ST, Poon LL, Guan Y, Yam LY, Lim W, et al. Coronavirus as a possible cause of severe acute respiratory syndrome. *Lancet* 2003;**361**:1319–25.
- Ksiazek TG, Erdman D, Goldsmith CS, Zaki SR, Peret T, Emery S, et al. A novel coronavirus associated with severe acute respiratory syndrome. *N Engl J Med* 2003;**348**:1953–66.
- Drosten C, Gunther S, Preiser W, van der Werf S, Brodt HR, Becker S, et al. Identification of a novel coronavirus in patients with severe acute respiratory syndrome. *N Engl J Med* 2003;**348**:1967–76.
- Tsang KW, Ho PL, Ooi GC, Yee WK, Wang T, Chan-Yeung M, et al. A cluster of cases of severe acute respiratory syndrome in Hong Kong. *N Engl J Med* 2003;**348**:1977–85.
- Lee N, Hui D, Wu A, Chan P, Cameron P, Joynt GM, et al. A major outbreak of severe acute respiratory syndrome in Hong Kong. *N Engl J Med* 2003;**348**:1986–94.
- Poutanen SM, Low DE, Henry B, Finkelstein S, Rose D, Green K, et al. Identification of severe acute respiratory syndrome in Canada. *N Engl J Med* 2003;**348**:1995–2005.
- Leung WK, To KF, Chan PK, Chan HL, Wu AK, Lee N, et al. Enteric involvement of severe acute respiratory syndrome-associated coronavirus infection. *Gastroenterology* 2003;**125**:1011–7.
- Peiris JS, Chu CM, Cheng VC, Chan KS, Hung IF, Poon LL, et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *Lancet* 2003;**361**:1767–72.
- Cheng VC, Hung IF, Tang BS, Chu CM, Wong MM, Chan KH, et al. Viral replication in the nasopharynx is associated with diarrhea in patients with severe acute respiratory syndrome. *Clin Infect Dis* 2004;**38**:467–75.
- van der Hoek L, Pyrc K, Jebbink MF, Vermeulen-Oost W, Berkhout RJ, Wolthers KC, et al. Identification of a new human coronavirus. *Nat Med* 2004;**10**:368–73.
- Fouchier RA, Hartwig NG, Bestebroer TM, Niemeyer B, de Jong JC, Simon JH, et al. A previously undescribed coronavirus associated with respiratory disease in humans. *Proc Natl Acad Sci USA* 2004;**101**:6212–6.
- Bastien N, Anderson K, Hart L, Van Caeseele P, Brandt K, Milley D, et al. Human coronavirus NL63 infection in Canada. *J Infect Dis* 2005;**191**:503–6.
- Arden KE, Nissen MD, Sloots TP, Mackay IM. New human coronavirus, HCoV-NL63, associated with severe lower respiratory tract disease in Australia. *J Med Virol* 2005;**75**:455–62.
- Han TH, Chung JY, Kim SW, Hwang ES. Human Coronavirus-NL63 infections in Korean children, 2004–2006. *J Clin Virol* 2007;**38**:27–31.
- Vabret A, Mourez T, Dina J, van der Hoek L, Gouarin S, Petitjean J, et al. Human coronavirus NL63, France. *Emerg Infect Dis* 2005;**11**:1225–9.
- van der Hoek L, Sure K, Ithorst G, Stang A, Pyrc K, Jebbink MF, et al. Croup is associated with the novel coronavirus NL63. *PLoS Med* 2005;**2**:e240.
- Shao X, Guo X, Esper F, Weibel C, Kahn JS. Seroepidemiology of group I human coronaviruses in children. *J Clin Virol* 2007;**40**:207–13.
- Woo PC, Lau SK, Chu CM, Chan KH, Tsoi HW, Huang Y, et al. Characterization and complete genome sequence of a novel coronavirus, coronavirus HKU1, from patients with pneumonia. *J Virol* 2005;**79**:884–95.
- Gerna G, Percivalle E, Sarasini A, Campanini G, Piralla A, Rovida F, et al. Human respiratory coronavirus HKU1 versus other coronavirus infections in Italian hospitalised patients. *J Clin Virol* 2007;**38**:244–50.
- Garbino J, Crespo S, Aubert JD, Rochat T, Ninet B, Deffernez C, et al. A prospective hospital-based study of the clinical impact of non-severe acute respiratory syndrome (Non-SARS)-related human coronavirus infection. *Clin Infect Dis* 2006;**43**:1009–15.
- Sloots TP, McErlean P, Speicher DJ, Arden KE, Nissen MD, Mackay IM. Evidence of human coronavirus HKU1 and human bocavirus in Australian children. *J Clin Virol* 2006;**35**:99–102.
- Esper F, Weibel C, Ferguson D, Landry ML, Kahn JS. Coronavirus HKU1 infection in the United States. *Emerg Infect Dis* 2006;**12**:775–9.
- Vabret A, Dina J, Gouarin S, Petitjean J, Corbet S, Freymuth F. Detection of the new human coronavirus HKU1: a report of 6 cases. *Clin Infect Dis* 2006;**42**:634–9.
- Pang XL, Preiksaitis JK, Lee B. Multiplex real time RT-PCR for the detection and quantitation of norovirus genogroups I and II in patients with acute gastroenteritis. *J Clin Virol* 2005;**33**:168–71.