

RESEARCH ARTICLE

The *Legionella pneumophila* GIG operon responds to gold and copper in planktonic and biofilm cultures

Kathleen Jwanoswki¹, Christina Wells¹, Terri Bruce², Jennifer Rutt¹, Tabitha Banks¹, Tamara L. McNealy^{1*}

1 Department of Biological Sciences, Clemson University, Clemson, South Carolina, United States of America, **2** Clemson Light Imaging Facility, Clemson University, Clemson, South Carolina, United States of America

* tmcneal@g.clemson.edu



Abstract

Legionella pneumophila contaminates man-made water systems and creates numerous exposure risks for Legionnaires' Disease. Because copper/silver ionization is commonly used to control *L. pneumophila*, its mechanisms of metal response and detoxification are of significant interest. Here we describe an *L. pneumophila* operon with significant similarity to the GIG operon of *Cupriavidus metallidurans*. The *Legionella* GIG operon is present in a subset of strains and has been acquired as part of the ICE- β ox 65-kB integrative conjugative element. We assessed GIG promoter activity following exposure of *L. pneumophila* to multiple concentrations of H₂AuCl₄, CuSO₄ and AgNO₃. At 37°C, control stationary phase cultures exhibited GIG promoter activity. This activity increased significantly in response to 20 and 50uM H₂AuCl₄ and CuSO₄ but not in response to AgNO₃. Conversely, at 26°C, cultures exhibited decreased promoter response to copper. GIG promoter activity was also induced by H₂AuCl₄ or CuSO₄ during early biofilm establishment at both temperatures. When an *L. pneumophila* GIG promoter construct was transformed into *E. coli* DH5 α , cultures showed baseline expression levels that did not increase following metal addition. Analysis of *L. pneumophila* transcriptional regulatory mutants suggested that GIG up-regulation in the presence of metal ions may be influenced by the stationary phase sigma factor, RpoS.

OPEN ACCESS

Citation: Jwanoswki K, Wells C, Bruce T, Rutt J, Banks T, McNealy TL (2017) The *Legionella pneumophila* GIG operon responds to gold and copper in planktonic and biofilm cultures. PLoS ONE 12(5): e0174245. <https://doi.org/10.1371/journal.pone.0174245>

Editor: Zhao-Qing Luo, Purdue University, UNITED STATES

Received: April 24, 2016

Accepted: March 6, 2017

Published: May 2, 2017

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All relevant data are within the paper.

Funding: Clemson University Creative Inquiry Program provided funding through JR and TB participation in the project.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Legionella pneumophila, the etiological agent of Legionnaires' Disease (LD), is the leading cause of bacterial waterborne disease outbreaks in the United States [1]. This Gram-negative bacterium is ubiquitous in both natural and man-made aquatic environments, where it replicates as an intracellular parasite of free-living protozoa [2]. Most cases of LD can be traced back to human-made aquatic systems with above-ambient water temperatures: cooling towers, hot water heaters, fountains, and air conditioning units have all served as sources of outbreaks [2].

Legionella persist as part of the biofilm community in human-made aquatic systems, and these systems are routinely treated to inhibit microbial growth. Chlorine and chloramine are the most common disinfectants used in the US. Other treatments include chlorine dioxide, UV radiation, hyperchlorination, bromine, and copper/silver ionization. While such treatments work well against fecal coliforms and other bacteria that enter the system from outside sources, they are less effective at eliminating pathogens like *Legionella* that exist within resistant biofilms of the system itself. Copper/silver ionization is often used to control *Legionella* in large recirculating water systems, particularly industrial plumbing, but its effectiveness is controversial [3–6]. Some studies indicate that high levels of copper inhibit *Legionella* growth and survival, but others have demonstrated increased persistence of *Legionella* in biofilms formed on copper [7]. *Legionella* also demonstrates increased resistance to copper at lower temperatures [8–9].

Small amounts of copper are required for mitochondrial electron transport and other enzymatic reactions, but high intracellular copper levels are considered toxic to most prokaryotes [10–12]. General mechanisms of bacterial metal resistance include export across the plasma membrane, sequestration by binding proteins, and reduction to a less toxic state. While *Legionella* is sensitive to elevated concentrations of numerous metals [13], few of its metal resistance mechanisms have been described. A *Legionella* copper-translocating PIB-type ATPase (CopA) was shown to confer copper resistance when expressed in a copper-sensitive *E. coli* strain [14], and the *helABC* locus was reported to encode three proteins involved in heavy metal resistance and cytopathogenicity [15]. Additional mechanisms by which *Legionella* sense and respond to metal ions in their environment remain to be characterized.

Here we describe a *Legionella pneumophila* operon (*lpg2105-2108*) with significant homology to the “gold induced genes” (GIG) operon of *Cupriavidus metallidurans* [16]. The operon appears to have been acquired by a subset of *L. pneumophila* strains as part of the horizontally transferred ICE- β ox integrative conjugative element [17]. In planktonic cultures, promoter expression occurs at 37°C in response to gold and copper, but at 26°C it occurs only in response to growth phase. Under biofilm conditions, promoter response to metal ions is seen at both temperatures.

Material and methods

Identification of the *L. pneumophila* GIG operon and homologous operons

The *L. pneumophila* GIG operon was originally identified through a BlastP search of proteins from the *L. pneumophila* subsp. *pneumophila* str. Philadelphia 1 genome (NCBI NC_002942.5), using the four *C. metallidurans* GIG proteins as queries (Table 1). Additional homologous operons in *L. pneumophila* were subsequently identified through BlastP searches of five sequenced *L. pneumophila* subsp. *pneumophila* strains, using the *L. pneumophila* GIG proteins as queries. Hits with a query cover of at least 80% and an E-value of less than $1.0E^{-15}$ were retained in this analysis. Homologous operons were also identified in the genomes of the *Francisella tularensis* (NC_006570.2), *Burkholderia pseudomallei* (NC_012695.1), *Polaramonas* sp. JS666 (NC_007948.1) and *Pseudomonas fluorescens* (NC_007492.2) using similar methods. Predicted protein sequences from each operon were concatenated and aligned with MAFFT v.7 [18]. A maximum likelihood tree was constructed in PhyML using the LG amino acid substitution model and the SPR method of topology estimation [19]. Reliability of the tree was assessed with 500 bootstrap replicates, and branches reproduced in fewer than 50% of the replicates were collapsed.

Table 1. Similarity and identity of gold induced genes (GIG) operon in *L. pneumophila* Philadelphia 1 to *C. metallidurans* GIG operon.

<i>C. metallidurans</i> GIG protein	Length (aa)	<i>L. pneumophila</i> Philadelphia 1 GIG protein	Length (aa)	E-value	Query cover (%)	Identity (%)	Annotation information
Rmet_4682 (ABF11544.1)	156	lpg2105 (YP_096118.1)	165	1.00E-19	85	38	Predicted inner membrane protein with a DoxX domain (PF07681)
Rmet_4683 (ABF11545.1)	258	lpg2106 (YP_096119.1)	259	8.00E-16	89	22	Contains N-terminal DUF2063 domain with putative role in DNA binding and transcriptional regulation
Rmet_4684 (ABF11546.1)	278	lpg2107 (YP_096120.1)	284	7.00E-72	98	40	DUF692 family of uncharacterized bacterial proteins; possibly involved in methanobactin synthesis
Rmet_4685 (ABF11547.1)	94	lpg2108 (YP_096121.1)	97	9.00E-06	81	33	DUF2282 family of putative integral membrane and signal proteins

<https://doi.org/10.1371/journal.pone.0174245.t001>

Strains and media

In all experiments, wild type *L. pneumophila* strain Lp02 was cultured at 37°C or 26°C for three days on buffered charcoal yeast extract agar with 100µg/ml thymidine (BCYE-T). *L. pneumophila* Lp02 is a derivative of *L. pneumophila* Philadelphia 1 and contains the complete ICE-βox region. Lp02 *ΔletA* and *ΔrpoS* mutants were grown on BCYE with 20µg/ml kanamycin. All *Legionella pneumophila* strains were provided by Michelle Swanson (University of Michigan). *Escherichia coli* DH5α was grown at 37°C or 26°C for 24 hours on Tryptic Soy Agar (TSA). Broth cultures consisted of ACES (*N*-(2-Acetamido)-2-aminoethanesulfonic acid)-buffered yeast extract (AYE) for *Legionella* strains (with antibiotics and thymidine as necessary) and Tryptic Soy Broth (TSB) for *E. coli* strains.

Reverse Transcriptase-PCR. Total RNA was isolated from stationary phase cultures using the Promega SV RNA Isolation kit. RNA was DNase treated to ensure removal of contaminating DNA prior to RT-PCR analysis. Three sets of primers spanning the four genes of the operon (2105–2106; 2106–2107; 2107–2108) were used to confirm the single transcript containing the four genes. The Verso 1-step Reverse Transcriptase PCR kit was used following manufacturer’s instructions. Reactions were carried out in 25ul final volume using each primer set and 15ng of RNA per reaction. Amplified fragments were analysed on a 1.0% agarose gel.

Construction of pGIG reporter gene vector

The *flaA* promoter of *pflaA* (a GFP reporter gene vector provided by M. Swanson, [20]), was replaced with 180nt of the upstream *lpg2105-2108* predicted promoter region using standard cloning methods. The 180nt upstream of *lpg2105-2108* represents the intergenic region between *lpg2108* and *lpg2109*. Bacterial promoter regions have been shown to be enriched in the intergenic regions of the genome and depleted from coding regions [21]. The entire intergenic region was cloned in order to capture as many potential promoter binding sites as possible for subsequent experiments. The resulting plasmid, *pGIGgfp*, was transformed into *L. pneumophila* Lp02, *E. coli* DH5α, *L. pneumophila* Lp02 *ΔletA*, and *L. pneumophila* Lp02 *ΔrpoS*.

Reporter gene activity in planktonic cultures

The effect of metal ions on *L. pneumophila* growth kinetics and *pGIGgfp* reporter gene activity were measured by incubating planktonic cultures for 72 hours at 150rpm in the presence or absence of gold, copper or silver ions. Cultures were incubated at 37°C with 20µM or 50µM of

gold chloride (HAuCl₄), 20 μM or 50 μM copper sulfate (CuSO₄), or 50 μM or 150 μM silver nitrate (AgNO₃), or no additional ions (control). Cultures were incubated at 26°C with 5 μM and 10 μM HAuCl₄, 200 μM and 275 μM CuSO₄, or no additional ions (control). Every three hours, absorbance (OD₆₀₀) was measured using a Genesys 6 spectrophotometer, and GFP fluorescence (485 nm excitation/528 nm emission) was measured using a Biotek Synergy H1 plate reader.

The effect of metal ions on *E. coli* growth kinetics and *pGIGgfp* reporter gene activity was measured by incubating planktonic cultures for 24 hours at 150 rpm in the presence or absence of gold or copper. Cultures were incubated at 37°C with 20 μM or 50 μM HAuCl₄, 20 μM or 50 μM CuSO₄, or no additional ions. Absorbance and GFP fluorescence were measured hourly as described above.

All experiments were performed in triplicate. Data were normalized by dividing the GFP relative light units (RLU) by the OD₆₀₀. The effects of metal ion, ion concentration, growth phase, and their interactions on the magnitude of GFP fluorescence were each analyzed separately.

Reporter gene activity in biofilms

L. pneumophila Lp02 *pGIGgfp* biofilms were established as previously described [22–23]. Briefly, bacteria were inoculated into glass petri dishes containing glass slides in 10% AYE solution and incubated for 24 hours, then transferred to 100% AYE for the remainder of the incubation. Cultures were incubated at 26°C or 37°C with 20 μM HAuCl₄, 20 μM CuSO₄, or no additional ions. Biofilms were grown for 120 hours and assessed for GIG activity at 24, 48, 72, 96, and 120 h. At each time point, biofilms were washed twice with sterile ultrapure water (UPW) to remove non-attached bacteria. Slides were aseptically removed, briefly air dried, and fixed in paraformaldehyde for 10 minutes. Slides were rinsed with UPW and dried; then coverslips were mounted using a 50/50 v/v solution of glycerol:phosphate buffered saline (1X PBS). All experiments were performed in triplicate.

Image analysis

Biofilms were imaged using a Leica SPE spectral confocal microscope (63X, oil immersion objective, NA = 1.30; Leica Microsystems, Buffalo Grove, IL) in the Clemson Light Imaging Facility. Three DIC (differential interference contrast) images and three corresponding GFP images were obtained for each slide. For assessment of promoter activity, ImageJ software and DIC images were used to generate regions of interest (ROIs) corresponding to individual bacteria within the biofilm. The ROIs were outlined and numbered, and their areas were measured. A binary image “mask” of the ROIs was used as an overlay on the corresponding GFP image. The signal intensity within each ROI on the GFP image was measured and used as an indication of GFP expression in each bacterium (ROI). All biofilm samples produced some level of green fluorescence. To account for this primary fluorescence, the intensity of control biofilms was determined and gated out (subtracted) from the intensity of treated biofilms to determine GFP expression due solely to the addition of gold or copper. The 37°C control fluorescence intensity value was applied to all samples. Treatment effects were assessed using Z-scores with a 99% confidence interval.

To measure the biofilm biomass that demonstrated GIG activity, confocal images from each time point were analyzed using COMSTAT software [24]. Bio-volume was estimated from calculations of biofilm biomass, which were based on the number of bacteria-containing pixels in all images of a stack, multiplied by voxel size and divided by the stack's substratum

area. A one way analysis of variance (ANOVA) was used to compare bio-volumes among bio-films and time points. A significance level of $p < 0.05$ was used for all tests.

Results

Description of the *L. pneumophila* GIG operon

The *L. pneumophila* GIG operon spans 2421 bases from 2352473 to 2354894 in the *L. pneumophila* subsp. *pneumophila* str. Philadelphia 1 genome (NC_002942.5). It is located within the ICE- β ox, a 65-kB integrative conjugative element found in approximately 18% of surveyed *Legionella* strains and associated with increased virulence and oxidative stress tolerance [17, 25]. The operon contains four open reading frames encoding proteins of unknown function (*lpg2105-lpg2108*).

The first *L. pneumophila* GIG gene, *lpg2105*, encodes a 165-aa predicted inner membrane protein with a DoxX domain (PF07681). DoxX proteins exhibit some similarity to the thiosulphate:quinone oxidoreductase small subunit, DoxD, but their precise function is unknown [26]. The second GIG gene, *lpg2106*, encodes a 259-aa protein with an N-terminal DUF2063 domain that is predicted to function in DNA binding and transcriptional regulation [27]. The third GIG gene, *lpg2107*, encodes a 284-aa protein assigned to the DUF692 family of uncharacterized bacterial proteins. Other members of this family are key enzymes in the biosynthesis of methanobactins, secreted copper-binding and copper-reducing peptides produced by a variety of bacteria [28]. The final gene, *lpg2108*, encodes a small 97-aa protein assigned to the DUF2282 family of putative integral membrane and signal peptide proteins.

Characterization of homologous operons

The *L. pneumophila* str. Philadelphia 1 genome also contains three additional regions with significant sequence similarity to the GIG operon, designated here as homologous operons H2, H3 and H4 (Table 2). The H2 operon (*lpg0665-lpg0669*) is structurally similar to the GIG operon and contains homologs of all four GIG genes, arranged in an identical order and strand orientation. The H3 operon (*lpg2253-lpg2255*) lacks an *lpg2105* homolog and therefore contains only three of the four GIG genes, again arranged in identical order and strand orientation. The H4 operon (*lpg0671-lpg0676*) contains homologs of all four GIG genes, but in a different order and on different strands. The first H4 gene (*lpg0671*) shares a DoxX domain with *lpg2105* but encodes a much larger NADH dehydrogenase transmembrane protein. Two additional genes are also present: an acetoacetate decarboxylase gene (*lpg0672*) and an adenylate cyclase gene (*lpg0674*). Rather than an operon, H4 may be better described as a cluster of genes that includes the four GIG homologs. They are unlikely to be transcribed as a unit, given their differing strand orientations.

We surveyed the genomes of four additional *L. pneumophila* strains for the presence of the GIG, H2, H3 and H4 operons (Table 2). While all strains contained copies of H2, H3 and H4, the GIG operon was present only in *L. pneumophila* Philadelphia 1. This is consistent with its acquisition as part of the ICE- β ox. A maximum likelihood phylogenetic tree indicated that the *L. pneumophila* GIG operon was most similar to the *C. metallidurans* GIG operon, as well as to number of homologous operons from multiple environmental and/or pathogenic bacteria (Fig 1).

lpg2105-2108 operon

The four genes—*lpg2105-2108*—are predicted to be transcribed as a single unit based on genomic analysis. Using reverse transcriptase PCR, we confirmed that the genes are co-transcribed.

Table 2. Homologous operons in *L. pneumophila* genomes to *lpg2105-2108* GIG operon.

Strain	GIG		H2		H3		H4	
	Locus tag	Start position and strand	Locus tag	Start position and strand	Locus tag	Start position and strand	Locus tag	Start position and strand
Philadelphia 1	<i>lpg2105</i>	2352970 (-)	<i>lpg0665</i>	716610 (-)			<i>lpg0671</i>	719987 (+)
NC_002942.5	<i>lpg2106</i>	2353739 (-)	<i>lpg0666</i>	717386 (-)	<i>lpg2253</i>	2557140 (-)	<i>lpg0675</i>	725484 (-)
	<i>lpg2107</i>	2354586 (-)	<i>lpg0667</i>	718248 (-)	<i>lpg2254</i>	2558033 (-)	<i>lpg0676</i>	726280 (-)
	<i>lpg2108</i>	2354894 (-)	<i>lpg0669</i>	718508 (-)	<i>lpg2255</i>	2558333 (-)	<i>lpg0673</i>	723047 (+)
Paris			<i>LPP_RS03625</i>	801134 (-)			<i>LPP_RS03650</i>	804510 (+)
			<i>LPP_RS03630</i>	801910 (-)	<i>LPP_RS11155</i>	2543886 (-)	<i>LPP_RS03670</i>	809982 (-)
			<i>LPP_RS03635</i>	802772 (-)	<i>LPP_RS11160</i>	2544746 (-)	<i>LPP_RS03675</i>	810802 (-)
Corby			<i>LPP_RS03640</i>	803032 (-)	<i>LPP_RS11165</i>	2545073 (-)	<i>LPP_RS03660</i>	807569 (+)
			<i>LPC_RS03790</i>	827443 (-)			<i>LPC_RS03815</i>	830819 (+)
			<i>LPC_RS03795</i>	828003 (-)	<i>LPC_RS11785</i>	2673120 (-)	<i>LPC_RS03835</i>	836292 (-)
NC_009494.1			<i>LPC_RS03800</i>	829080 (-)	<i>LPC_RS11790</i>	2673980 (-)	<i>LPC_RS03840</i>	837112 (-)
			<i>LPC_RS03805</i>	829340 (-)	<i>LPC_RS11795</i>	2674376 (-)	<i>LPC_RS03825</i>	833879 (+)
			<i>LPL_RS03540</i>	790101 (-)			<i>LPL_RS03565</i>	793477 (+)
Lens			<i>LPL_RS03545</i>	790877 (-)	<i>LPL_RS11000</i>	2494966 (-)	<i>LPL_RS03585</i>	798950 (-)
			<i>LPL_RS03550</i>	791739 (-)	<i>LPL_RS11005</i>	2495826 (-)	<i>LPL_RS03590</i>	799770 (-)
			<i>LPL_RS03555</i>	791999 (-)	<i>LPL_RS11010</i>	2496153 (-)	<i>LPL_RS03575</i>	796537 (+)
Alcoy			<i>LPA_RS03750</i>	813482 (-)			<i>LPA_RS03775</i>	816859 (+)
			<i>LPA_RS03755</i>	814258 (-)	<i>LPA_RS11585</i>	2646878 (-)	<i>LPA_RS03795</i>	822332 (-)
			<i>LPA_RS03760</i>	815120 (-)	<i>LPA_RS11590</i>	2647738 (-)	<i>LPA_RS03800</i>	823152 (-)
NC_014125.1		<i>LPA_RS03765</i>	815380 (-)	<i>LPA_RS11595</i>	2648134 (-)	<i>LPA_RS03785</i>	819919 (+)	

<https://doi.org/10.1371/journal.pone.0174245.t002>

Using three sets of primers spanning the end of one gene and the beginning of the next, RT-PCR reactions for gene combinations 2105–2106, 2106–2107 and 2107–2108 were positive (Fig 2).

GIG promoter activity in planktonic cultures

Expression of the *C. metallidurans* GIG operon is influenced by the presence of gold and copper ions in the growth medium [16]. To test whether the *L. pneumophila* GIG operon was similarly affected, we created a GFP reporter construct whose expression was driven by the 180nt upstream promoter region of the *L. pneumophila* GIG operon. We measured GFP fluorescence to monitor expression of the *pGIGgfp* reporter in *L. pneumophila* Lp02 exposed to multiple concentrations of gold, copper, and silver ions. At both 37° and 26°C, we observed an extended lag phase in a concentration dependent manner when H₂AuCl₄ or CuSO₄ was added to the cultures. Cultures grown in the presence of 50µM of gold or copper never reached the same maximum OD₆₀₀ as controls.

To account for the individual growth rates of each culture, mid-exponential (ME) and late stationary (LS) growth phases were identified in individual cultures, and the promoter activity was compared at each growth phase. Late stationary phase was defined as the time point prior to decline of OD₆₀₀ that preceded death of the culture. GFP was normalized to OD₆₀₀ at each time point to accurately compare promoter activity across cultures.

At 37°C, control cultures of *L. pneumophila* Lp02 *pGIGgfp* showed similar levels of *lpg2105-2108* promoter activity across all growth phases. The addition of 50µM AgNO₃ did not affect




Fig 1. Maximum likelihood phylogeny for the GIG operon showing the relationship of *L. pneumophila* lpg2105-2108 to 4 additional *Legionella* species and five other bacterial species. Numbers over branches show bootstrap support values (500 replicates).

<https://doi.org/10.1371/journal.pone.0174245.g001>

culture growth or induce a promoter response. 150µM AgNO₃ inhibited culture growth but did not induce a promoter response (Fig 3). The promoter response to H₂AuCl₄ and CuSO₄ differed between growth phases. At the mid-exponential growth phase, only CuSO₄ treatment induced moderate promoter expression. At the late stationary growth phase, both H₂AuCl₄ and CuSO₄ induced higher levels of promoter expression (Fig 3).

At 26°C, *L. pneumophila* Lp02 growth was inhibited by the 20µM H₂AuCl₄ used at 37°C, and H₂AuCl₄ concentrations were therefore reduced to 5µM and 10µM. *Legionella* demonstrates increased resistance to copper at lower temperatures [9], and 26°C cultures showed no inhibition to the 20µM and 50µM CuSO₄ used at 37°C. CuSO₄ concentrations were therefore increased to 200µM and 275µM at 26°C to assess the promoter response under similar growth kinetics.

The effect of growth phase on promoter expression was more pronounced at 26°C than at 37°C. Control cultures showed a significant increase in promoter activity at LS phase, a result which was not observed at 37°C and that suggested temperature regulation of the operon. 10µM H₂AuCl₄ caused a modest increase in promoter activity at ME but reduced the level of promoter activity at LS. The addition of copper caused a significant decrease in promoter activity at LS (Fig 4).




Fig 2. Reverse Transcriptase PCR using gene spanning primers. M-Marker; 1: positive control 2105F/2106R primers; 2: positive control 2106F/2107R primers; 3: positive control 2107F/2108R primers; 4: no template control; 5: no RT control; 6: RNA 2105F/2106R primers; 7: RNA 2106F/2107R primers; 8: RNA 2107F/2108R primers.

<https://doi.org/10.1371/journal.pone.0174245.g002>




Fig 3. Effect of gold, copper and silver on *pGIGfp* activity in *L. pneumophila* at 37°C. GIG operon activity measured by GFP expression of *pGIGfp* in control media, with 20μM HAuCl₄, 50μM HAuCl₄, 20μM CuSO₄ and 50μM CuSO₄ at mid-exponential (ME) and late stationary (LS) growth phases (a). GFP expression in response to 50μM and 150μM AgNO₃ (b). Data presented from three independent experiments ± SD (* = p < 0.05).

<https://doi.org/10.1371/journal.pone.0174245.g003>




Fig 4. Effect of gold and copper on *pGIGgfp* activity in *L. pneumophila* at 26°C. GIG operon activity measured by GFP expression of *pGIGgfp* in control media, with 20µM HAuCl₄, 50µM HAuCl₄, 20µM CuSO₄ and 50µM CuSO₄ at mid-exponential (ME) and late stationary (LS) growth phases. Data presented from three independent experiments ± SD (* = p< 0.05).

<https://doi.org/10.1371/journal.pone.0174245.g004>

The complex promoter response to temperature, growth phase and metal addition suggests that multiple levels of operon regulation interact in planktonic culture. To further investigate mechanisms of operon regulation, we examined promoter expression in *E. coli* DH5α transformed with *pGIGgfp* as well as in two *L. pneumophila* mutants in the major stationary phase regulation pathways. *E. coli* DH5 α, which does not possess the operon, showed no change in growth kinetics in the presence of HAuCl₄ or CuSO₄. At neither 37°C nor 26°C did *E. coli* show promoter activity to gold or copper (Fig 5). These results suggest a unique regulator in *L. pneumophila* for metal induction of the GIG operon. Significant increase in promoter expression in response to HAuCl₄ and CuSO₄ was seen during stationary phase, and we therefore investigated known regulators of *L. pneumophila* stationary phase gene expression. Mutants for the stationary phase regulators LetA and RpoS were transformed with *pGIGgfp* to assess their potential involvement.

The *Legionella* two-component regulatory system, LetA/S, induces the expression of virulence and transmission traits at stationary phase [29–30]. Lp02 Δ*letA* *pGIGgfp* cultures showed delayed entry into exponential phase when exposed to HAuCl₄ or CuSO₄, similar to that seen in wild type Lp02 *pGIGgfp* cultures. Lp02Δ*letA* cultures with added HAuCl₄ or CuSO₄ exhibited greater promoter activity compared to Lp02 Δ*letA* cultures with no metal addition. The magnitude of metal-induced promoter up-regulation was greater at the LS phase. However, wild type and Δ*letA* cultures responded similarly to metal addition: there were no significant differences in promoter activity between Lp02 Δ*letA* cultures and wild type Lp02 cultures at the same metal concentration.




Fig 5. Effect of gold and copper on *pGIGgfp* activity in *E. coli* at 37°C. GIG operon activity measured by GFP expression of *pGIGgfp* in control media with addition of with 20µM HAuCl₄, 50µM HAuCl₄, 20µM CuSO₄ or 50µM CuSO₄ at mid-exponential (ME) and late stationary (LS) growth phases. Data presented are from three independent experiments ± SD.

<https://doi.org/10.1371/journal.pone.0174245.g005>

Lp02 *ΔrpoS pGIGgfp* cultures exhibited an extended lag phase in response to HAuCl₄ or CuSO₄, and 50µM CuSO₄ cultures showed increased promoter expression at ME phase. The promoter response to metal addition was dampened in Lp02 *ΔrpoS pGIGgfp* stationary phase cultures: mutant cultures showed reduced promoter activity compared to wild type in response to both HAuCl₄ and CuSO₄ addition, suggesting that RpoS may play a role in promoter expression, particularly at stationary phase (Fig 6).

GIG promoter activity in biofilms

L. pneumophila primarily persists in the environment as a biofilm [31–33]. To assess operon activity under more natural growth conditions, Lp02 *pGIGgfp* biofilms were established and grown at two temperatures in the presence and absence of 20µM HAuCl₄ or CuSO₄. Analysis of biofilms grown at 26°C revealed that promoter activity increased significantly in early phase biofilms when HAuCl₄ or CuSO₄ was added (Fig 7a). At 24h, a significant increase in GIG expression in response to both HAuCl₄ and CuSO₄ was observed. At 72h, an additional increase was observed in response to HAuCl₄ exposure but not to CuSO₄. The 26°C biofilms did not exhibit the increased sensitivity to gold as seen in the planktonic cultures.

Similar to planktonic cultures, greater promoter expression was measured in 37°C biofilms than in 26°C (Fig 7b). Increased promoter expression was seen in early phase biofilms in response to gold and copper at 24h and in response to gold at 48h and 72h.

To determine if biomass were responsible for the increase in promoter activity, biofilm biomass was assessed from confocal images using COMSTAT. There were no differences in




Fig 6. Effect of gold and copper on pGIGfp activity *L. pneumophila* Lp02 Δ LetA and Δ RpoS. GIG operon activity measured by GFP expression of pGIGfp in control media with addition of with 20 μ M HAuCl₄, 50 μ M HAuCl₄, 20 μ M CuSO₄ or 50 μ M CuSO₄ at mid-exponential (ME) and late stationary (LS) growth phases. Data presented are from three independent experiments \pm SD. (* = significantly different from WT with same metal addition in same panel; p < 0.05, ‡ = significantly different from mutant with no metal addition in first panel; p < 0.05)

<https://doi.org/10.1371/journal.pone.0174245.g006>

biomass among control, plus gold, and plus copper samples at each temperature. There was a significant decrease in biofilm biomass in all treatments from 48h to 72h, likely due to the dispersal event typically seen at this stage in *Legionella* biofilm development (Fig 8).




Fig 7. Analysis of pGIGfp expression during biofilm establishment and development at a) 26°C in *L. pneumophila* control biofilms and with the addition of 20 μ M HAuCl₄ or 20 μ M CuSO₄ and at b) 37°C in *L. pneumophila* control biofilms and with the addition of 20 μ M HAuCl₄ or 20 μ M CuSO₄. Percent GFP intensity above basal level (determined at 26°C) is shown. Data presented are from three independent experiments \pm SE (p < 0.05)

<https://doi.org/10.1371/journal.pone.0174245.g007>




Fig 8. Biomass of control biofilms and biofilms exposed to 20µM HAuCl₄ or 20µM CuSO₄ and measured at 48 or 72h. Data presented are from three independent experiments ±SE ($p < 0.05$).

<https://doi.org/10.1371/journal.pone.0174245.g008>

Discussion

Based on sequenced strains, the GIG operon appears to be rare in the *Legionella* genus, existing in only 3 sequenced strains—Philadelphia 1, Lp02 and SVir, each of which possess the full operon. Multiple homologs of the operon exist in *L. pneumophila* at genes *lpg0665-0669*, *lpg0671*, *0673*, *0675*, *0676*, and *lpg2253-2255*. Species within the *Legionella* genus appear to contain multiple paralogous copies of the operon, which group into four distinct clades (Fig 1, Table 1). The *lpg2105-2108* operon in *L. pneumophila* Philadelphia 1 is distinct from the others.

The response of the *L. pneumophila* GIG operon to HAuCl₄ and CuSO₄ is similar to that reported for the GIG operon of *C. metallidurans* [16]. At 37°C, planktonic cultures expressed GIG operon promoter activity that increased with the addition of HAuCl₄ or CuSO₄ at stationary phase. Conversely, promoter activity was repressed in response to HAuCl₄ or CuSO₄ at 26°C (Fig 4). Although not well explored, temperature regulation of genes is not unknown in *Legionella*. Piao, et al., [34] showed that biofilms alter their morphology based on temperature and surface material. Biofilms formed at 37°C and 42°C resembled filamentous mycelial mats, while at 26°C, short, rod shaped bacteria made up the majority of the biofilm. Genes involved in type IV pilus biogenesis and type II protein secretion also showed temperature regulation, with increased expression at 30°C compared to 37°C [35]. The increased HAuCl₄ sensitivity and opposite expression patterns between 26°C and 37°C in both biofilm and planktonic cultures suggest that temperature regulation may be involved with operon activity. *Legionellae* growing at low-temperatures have a more unsaturated membrane lipids, suggesting that

changes in membrane fluidics may also be involved in gold sensitivity if membrane proteins are involved in the response to gold [36]. The *lpg2107* gene appears to possess transmembrane domains and may therefore be localized to the bacterial membrane.

Stationary phase changes in promoter activity in both *Legionella* and *E. coli* suggest that known stationary phase regulatory proteins may regulate the operon. *L. pneumophila* exhibits unique biphasic gene expression with tight control over exponential vs. stationary phase proteins. Determinants involved in metal or metalloid resistance that are up-regulated after treatment with gold complexes are often controlled by MerR- or ArsR-type regulators, which usually bind “soft” metals or metalloids. In addition to *C. metallidurans*, gold response systems exist in *E. coli*, controlled through CueR [37] and in *S. typhimurium* controlled by GolS [38]. *C. metallidurans* also possesses a CupR system that responds to both gold and copper [39]. Homologs of CueR and GolS are found in the *L. pneumophila* genome but have not been characterized.

Previous transcriptomic work showed expression of the *Legionella* GIG operon during amoebal infection, suggesting a potential role in virulence [40]. Many *Legionella* virulence genes are regulated by the *rpoS*, *csrA*, and *letA* genes [41–43]. LetA/S activates transmission phase genes, including but not limited to *mip* (macrophage infectivity protein), *dot/icm* T4SS, and *flaA* [29, 43, 44–45]. Deletion of *letA* significantly inhibits virulence [29]. However, we saw no changes in promoter activity in Lp02 Δ *letA* *pGIGgfp* compared to WT Lp02 *pGIGgfp*, suggesting that LetA does not directly regulate expression of the operon. A bioinformatic search for the conserved LetA promoter binding sequence upstream of the GIG operon was also negative, further supporting a lack of direct LetA involvement in operon regulation.

Lp02 Δ *rpoS* cultures showed no response to added H₂AuCl₄ or CuSO₄ when compared to the Δ *rpoS* control. Compared to the Lp02 WT however, promoter activity in Δ *rpoS* strains was significantly repressed (Fig 6). RpoS typically represses motility, infectivity, and cytotoxicity during exponential phase, and up-regulates them during stationary phase [46–47]. It is required for expression of virulence traits and growth within amoebae [41]. RpoS functions in response to stressors in the environment and interacts with other stationary phase regulators such as LetA/S, FliA, *letE*, and LqsR [47–48]. Our data suggest that RpoS may regulate the GIG operon similarly to the way it regulates other stationary phase virulence genes. In fact, regulation may require LetA to release exponential phase repression by RpoS and allow for stationary phase expression, as seen in the regulation of the sRNA molecules RsmYZ [44]. The data support the possibility of multiple regulators for this operon, one for basal expression, and a second for gold/copper response. Investigations into mutants of *fliA* and *letE*, which also coordinate differentiation from replicative phase to transmissible [29,49], as well as *csrA*, a global regulator responsible for the repression of transmission phase genes and activation of replicative phase genes [42], may lead to a better understanding of how this operon is controlled. The recent publication of a *csrA* homolog within the ICE- β ox unit-*csrT*-is also a potential regulatory candidate [50].

Biofilms grown at 26°C with 20 μ M gold or copper showed increased response to the metal ions compared to planktonic cultures at 26°C, with no obvious toxicity. Biomass of biofilms exposed to copper and gold was equivalent to controls (Fig 8). The matrix produced by biofilms that protects bacteria from the effects of toxic metals is not found in planktonic cells. Analyses of operon expression through multiple stages of biofilm development revealed that while baseline expression is present in controls, H₂AuCl₄ or CuSO₄ treatments increase promoter activity, particularly at early developmental stages. Specific up-regulation of the promoter in response to gold occurred at 24h and 72h. Up-regulation early in biofilm development suggests that after the matrix is in place, there is less need for metal response system expression. The up-regulation in response to gold at 72h corresponded with the timing of initial dispersal events in this model (Fig 8). Biofilm dispersal is evident in the difference in

biomass between 48 and 72h, but a significant increase in promoter activity is seen at 72h. The up-regulation at this time point may be from cells newly exposed to the external environment.

The GIG operon is present in several other virulent pathogens, including a category B biological agent *Burkholderia pseudomallei*, the causative agent of melioidosis, and a Category A bioterrorism agent, *Francisella tularensis*, the causative agent of tularemia. The presence of this operon in *F. tularensis* is particularly interesting since its abundance of pseudogenes and disrupted metabolic pathways are indicative of genome reduction [51–52]. The idea of “use it or lose it” can be applied to this process: genes that are nonessential to the survival of the organism that are mutated or lost. That this operon has been retained suggests it may have an essential function in survival or environmental persistence that could be exploited as a target for antimicrobials.

Many studies have been conducted on the efficacy of water treatment methods on *Legionella*, but few have looked at possible genetic response systems to those treatments, specifically those involved with metal ion response. The controversy over the use of copper/silver ionization as an effective means to protect water systems from *Legionella* colonization may be better explained after the metal response systems in the bacteria are more thoroughly characterized. Lack of information on bacterial response to metals hinders development of effective disinfection strategies. Based on the persistence of *Legionella* in the presence of metal-based disinfection treatments, an increased understanding of biofilm ecology, and in particular the environmental ecology of *Legionella*, is necessary. The results of this study lay groundwork for continued investigation of *Legionella* responses to potentially toxic metal ion concentrations.

Acknowledgments

Disclaimer: *This work was prepared while Tamara McNealy was employed at Clemson University. The opinions expressed in this article are the author's own and do not reflect the view of the National Institutes of Health, the Department of Health and Human Services, or the United States government.*

Author Contributions

Conceptualization: CW T. Bruce TM.

Data curation: CW T. Banks TM.

Formal analysis: KJ CW TM.

Funding acquisition: TM.

Investigation: KJ CW T. Banks JR T. Bruce.

Methodology: T. Bruce TM.

Project administration: TM.

Resources: CW T. Bruce TM.

Supervision: T. Bruce TM.

Validation: KJ TM.

Visualization: CW TM.

Writing – original draft: KJ CW JR T. Bruce TM.

Writing – review & editing: KJ CW T. Bruce JR T. Banks TM.

References

1. MMWR. Centers for Disease Control and Prevention. Surveillance for waterborne disease outbreaks associated with drinking water and other non-recreational water—United States, 2009–2010. Morbidity and Mortality Weekly Report. 2013; 62:714–720. PMID: [24005226](#)
2. Fields BS, Benson RF, Besser RE. *Legionella* and Legionnaires' disease: 25 years of investigation. Clin Microbiol Rev. 2002; 15: 506–526. <https://doi.org/10.1128/CMR.15.3.506-526.2002> PMID: [12097254](#)
3. Rohr U, Senger M, Selenka F, Turley R, Wilhelm M. Four years of experience with silver-copper ionization for control of *Legionella* in a German university hospital hot water plumbing system. Clin Infect Dis. 1999; 29: 1507–1511. <https://doi.org/10.1086/313512> PMID: [10585804](#)
4. Mathys W, Curro PH, Junge-Mathys E. Efficacy of copper-silver ionization in controlling Legionella in a hospital hot water distribution system: a German experience. Legionella. Washington, DC: American Society for Microbiology 2002; 419–424.
5. Stout JE and Yu VL. Experiences of the First 16 Hospitals Using Copper—Silver Ionization for Legionella Control: Implications for the Evaluation of Other Disinfection Modalities. Infect Contr Hosp Epidemiol. 2003; 24: 563–568.
6. Lin YE, Stout JE, Yu VL. Controlling Legionella in hospital drinking water: an evidence-based review of disinfection methods. Infect Contr Hosp Epidemiol. 2011; 32: 166–173.
7. Buse HY, Lu J, Struewing IT, Ashbolt NJ. Preferential colonization and release of *Legionella pneumophila* from mature drinking water biofilms grown on copper versus unplasticized polyvinylchloride coupons. International journal of hygiene and environmental health 2014; 217(2–3): 219–225. <https://doi.org/10.1016/j.ijheh.2013.04.005> PMID: [23706882](#)
8. Rohr U, Senger M, Selenka F. Effect of silver and copper ions on survival of Legionella pneumophila in tap water. Zentralbl Hyg Umweltmed. 1996; 198(6): 514–21. PMID: [9409904](#)
9. Landeen LK, Moyasar TY, Gerba CP. Efficacy of copper and silver ions and reduced levels of free chlorine in inactivation of *Legionella pneumophila*. Appl Env Microbiol. 1989; 55(12): 3045–3050.
10. Braymer JJ and Gierdoc DP. Recent developments in copper and zinc homeostasis in bacterial pathogens. Curr Opin Chem Biol. 2014; 19: 59–66. <https://doi.org/10.1016/j.cbpa.2013.12.021> PMID: [24463765](#)
11. Bruins MR, Kapil S, Oehme FW. Microbial resistance to metals in the environment. Ecotox Evn Safety. 2000; 45: 198–207.
12. Nies DH. Microbial heavy metal resistance. Appl Microbiol Biotechnol 1999; 51: 730–750. PMID: [10422221](#)
13. States SJ, Conley LF, Ceraso M, Stephenson TE, Wolford RS, Wadowsky RM, et al. Effects of metals on *Legionella pneumophila* growth in drinking water plumbing systems. Appl Env Microbiol. 1985; 50: 1149–1154.
14. Kim E, Charpentier X, Torres-Urquidí O, McEvoy MM, Rensing C. The metal efflux island of *Legionella pneumophila* is not required for survival in macrophages and amoebas. FEMS Microbiol Lett. 2009; 301: 164–170. <https://doi.org/10.1111/j.1574-6968.2009.01813.x> PMID: [19895645](#)
15. McClain MS, Hurley MC, Brieland JK, Engleberg C. The *Legionella pneumophila* *hel* locus encodes intracellularly induced homologs of heavy-metal ion transporters of *Alcaligenes* spp. Infect Immun. 1996; 64: 1532–1540. PMID: [8613357](#)
16. Reith F, Etschmann B, Grosse C, Moors H, Benotmane M, Monsieurs P, et al. Mechanisms of gold biomineralization in the bacterium *Cupriavidus metallidurans*. PNAS. 2009; 106: 17757–17762. <https://doi.org/10.1073/pnas.0904583106> PMID: [19815503](#)
17. Flynn K and Swanson M. Integrative conjugative element ICE- β ox confers oxidative stress resistance to *Legionella pneumophila* in vitro and in macrophages. mBio. 2014; 5.
18. Katoh K and Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol Biol Evo. 2013; 30(4): 772–80.
19. Guinon S, Dufayrad JF, Lefort V, Anisimova M, Hordijk W and Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. 2010; 59(3): 307–21. Epub 2010 Mar 29.
20. Hammer B, Swanson M. Co-ordination of *Legionella pneumophila* virulence with entry into stationary phase by ppGpp. Mol Microbiol. 1999; 33: 721–731. PMID: [10447882](#)
21. Jacques PE, Rodrigue S, Gaudreau L, Goulet J, Brzezinski R. Detection of prokaryotic promoters from the genomic distribution of hexanucleotide pairs. BMC Bioinformatics. 2006. 7: 423. <https://doi.org/10.1186/1471-2105-7-423> PMID: [17014715](#)

22. Stojak A, Raftery T, Klaine S, McNealy T. Morphological responses of *Legionella pneumophila* biofilm to nanoparticle exposure. *Nanotoxicology*. 2011; 5: 730–742. <https://doi.org/10.3109/17435390.2010.550696> PMID: 21294606
23. Raftery TD, Kerscher P, Hart AE, Saville SL, Qi B, Kitchens CL, et al. Discrete nanoparticles induce loss of *Legionella pneumophila* biofilms from surfaces. *Nanotoxicology*. 2014; 8(5): 477–484. <https://doi.org/10.3109/17435390.2013.796537> PMID: 23586422
24. Heydorn A, Nielsen AT, Hentzer M, Sternberg C, Givskov M, Ersbøll BK, Molin S. Quantification of biofilm structures by the novel computer program COMSTAT. *Microbiology*. 2000 Oct; 146 (Pt 10):2395–407.
25. Brassinga AK, Hiltz M, Sisson G, Morash M, Hill N, Garduno E, et al. A 65-kilobase pathogenicity island is unique to Philadelphia-1 strains of *Legionella pneumophila*. *J Bacteriol*. 2003; 185: 4630–4637. <https://doi.org/10.1128/JB.185.15.4630-4637.2003> PMID: 12867476
26. Muller FH, Bandejas TM, Urlich T, Teixeira M, Gomes CM and Kletzin A. Coupling of the pathway of sulphur oxidation to dioxygen reduction: characterization of a novel membrane-bound thiosulphate:quinone oxidoreductase. *Mol Microbiol*. 2004; 53(4): 1147–1160. <https://doi.org/10.1111/j.1365-2958.2004.04193.x> PMID: 15306018
27. Das D, Grishin NV, Kumar A, Carlton D, Bakolitsa C, Miller D, et al. The structure of the first representative of Pfam family PF09836 reveals a two-domain organization and suggests involvement in transcriptional regulation. *Acta Cryst*. 2010; F66: 1174–1181.
28. Kenney GE and Rosenzweig AC. Genome mining for methanobactins. *BMC Biology*. 2013; 11:17 <http://www.biomedcentral.com/1741-7007/11/17> <https://doi.org/10.1186/1741-7007-11-17> PMID: 23442874
29. Hammer BK, Tateda ES, Swanson MS. A two-component regulator induces the transmission phenotype of stationary-phase *Legionella pneumophila*. *Mol Microbiol*. 2002; 44: 107–118. PMID: 11967072
30. Lynch D, Fieser N, Glöggler K, Forsbach-Birk V, Marre R. The response regulator LetA regulates the stationary-phase stress response in *Legionella pneumophila* and is required for efficient infection of *Acanthamoeba castellanii*. *FEMS Microbiol Lett* 2003; 219(2): 241–248. PMID: 12620627
31. Andreozzi E, Di Cesare A, Sabatini L, Chessa E, Sisti D, Rocchi M, Citterio B. Role of biofilm in protection of the replicative form of *Legionella pneumophila*. *Curr Microbiol*. 2014. 69(6):769–74. Epub 2014 Jul 15. <https://doi.org/10.1007/s00284-014-0648-y> PMID: 25023637
32. Ashbolt NJ. Environmental (Saprozoic) Pathogens of Engineered Water Systems: Understanding Their Ecology for Risk Assessment and Management. *Pathogens*. 2015. 4(2):390–405. <https://doi.org/10.3390/pathogens4020390> PMID: 26102291
33. Berjeaud JM, Chevalier S, Schlusshuber M, Portier E, Loiseau C, Aucher W, et al. *Legionella pneumophila*: The Paradox of a Highly Sensitive Opportunistic Waterborne Pathogen Able to Persist in the Environment. *Front Microbiol*. 2016. 7:486. eCollection 2016. <https://doi.org/10.3389/fmicb.2016.00486> PMID: 27092135
34. Piao Z, Sze CC, Barysheva O, Iida KI, Yoshida SI. Temperature-regulated formation of mycelial mat-like biofilms by *Legionella pneumophila*. *Appl Env Microbiol*. 2006; 72: 1613–1622.
35. Liles MR, Viswanathan VK, Cianciotto NP. Identification and temperature regulation of *Legionella pneumophila* genes involved in Type IV pilus biogenesis and Type II protein secretion. *Infect Immun*. 1998; 66: 1776–1782. PMID: 9529113
36. Mauchline WS, Araujo R, Wait R, Dowsett AB, Dennis PJ, Keevil CW. Physiology and morphology of *Legionella pneumophila* in continuous culture at low oxygen concentration. *J Gen Microbiol*. 1992. 138 (11):2371–80. <https://doi.org/10.1099/00221287-138-11-2371> PMID: 1479356
37. Chen P and He C. A general strategy to convert the MerR family proteins into highly sensitive and selective fluorescent biosensors for metal ions. *J Am Chem Soc*. 2004; 126: 728–729. <https://doi.org/10.1021/ja0383975> PMID: 14733542
38. Checa S, Espariz M, Perez Audero M, Botta P, Spinelli S, Soncini F. Bacterial sensing of and resistance to gold salts. *Mol Microbiol*. 2007; 63: 1307–1318. <https://doi.org/10.1111/j.1365-2958.2007.05590.x> PMID: 17244194
39. Jian X, Wasinger E, Lockard J, Chen L, He C. Highly sensitive and selective gold (I) recognition by a metalloregulator in *Ralstonia metallidurans*. *J Am Chem Soc*. 2009; 131: 10869–10871. <https://doi.org/10.1021/ja904279n> PMID: 19606897
40. Weissenmayer BA, Prendergast JG, Lohan AJ, Loftus BJ. Sequencing illustrates the transcriptional response of *Legionella pneumophila* during infection and identifies seventy novel small non-coding RNAs. *PLoS ONE*. 2011; 6: e17570. <https://doi.org/10.1371/journal.pone.0017570> PMID: 21408607
41. Hales LM and Shuman HA. The *Legionella pneumophila rpoS* gene is required for growth within *Acanthamoeba castellanii*. *J Bacteriol*. 1999; 181: 4879–4889. PMID: 10438758

42. Molofsky AB and Swanson MS. *Legionella pneumophila* CsrA is a pivotal repressor of transmission traits and activator of replication. *Mol Microbiol.* 2003; 50: 445–461. PMID: [14617170](#)
43. Shi C, Forsbach-Birk V, Marre R, McNealy T. The *Legionella pneumophila* global regulatory protein LetA affects DotA and Mip. *Int J Med Microbiol.* 2005; 296: 15–24. <https://doi.org/10.1016/j.ijmm.2005.09.003> PMID: [16423685](#)
44. Rasis M, Segal G. The LetA-RsmYZ-CsrA regulatory cascade, together with RpoS and PmrA, post-transcriptionally regulates stationary phase activation of *Legionella pneumophila* lcm/Dot effectors. *Mol Microbiol.* 2009; 72: 955–1010.
45. Edwards R, Jules M, Sahr T, Buchrieser C, Swanson M. The *Legionella pneumophila* LetA/LetS Two-Component System Exhibits Rheostat-Like Behavior. *Infect Immun.* 2010; 78: 2571–2583. <https://doi.org/10.1128/IAI.01107-09> PMID: [20351136](#)
46. Bachman MA and Swanson MS. RpoS co-operates with other factors to induce *Legionella pneumophila* virulence in the stationary phase. *Mol Microbiol.* 2001; 40: 1201–14. PMID: [11401723](#)
47. Dong T and Schellhorn HE. Role of RpoS in Virulence of Pathogens. *Infect Immun.* 2010; 78: 887–897. <https://doi.org/10.1128/IAI.00882-09> PMID: [19948835](#)
48. Tiaden A, Spirig T, Weber SS, Bruggemann H, Bosshard R, Buchrieser C, Hilbi H. The *Legionella pneumophila* response regulator LqsR promotes host cell interactions as an element of the virulence regulatory network controlled by RpoS and LetA. *Cell Microbiol.* 2007; 9: 2903–2920. <https://doi.org/10.1111/j.1462-5822.2007.01005.x> PMID: [17614967](#)
49. Bachman MA and Swanson MS. Genetic evidence that *Legionella pneumophila* RpoS modulates expression of the transmission phenotype in both the exponential phase and the stationary phase. *Infect Immun.* 2004; 72: 2468–2476. <https://doi.org/10.1128/IAI.72.5.2468-2476.2004> PMID: [15102753](#)
50. Abbott ZD, Flynn KJ, Byrne BG, Mukherjee S, Kearns DB, Swanson MS. *csrT* represents a new class of *csrA*-like regulatory genes associated with integrative conjugative elements of *Legionella pneumophila*. *J Bacteriol.* 2015; 198(3): 553–64. <https://doi.org/10.1128/JB.00732-15> PMID: [26598366](#)
51. Rohmer L, Fong C, Abmayr S, Wasnick M, Larson Freeman TJ, Radey M, et al. Comparison of *Francisella tularensis* genomes reveals evolutionary events associated with the emergence of human pathogenic strains. *Genome biology.* 2007; 8(6): 1–16.
52. Siddaramappa S, Challacombe JF, Petersen JM, Pillai S, Kuske CR. Genetic diversity within the genus *Francisella* as revealed by comparative analyses of the genomes of two North American isolates from environmental sources. *BMC Genomics.* 2013; 13: 422.