

A new species of *Paraonis* and an annotated checklist of polychaetes from mangroves of the Brazilian Amazon Coast (Annelida, Paraonidae)

Rannyele Passos Ribeiro¹, Paulo Ricardo Alves²,
Zafira da Silva de Almeida³, Christine Ruta⁴

1 Universidad Autónoma de Madrid, Cantoblanco, 28049, Madrid, Spain **2** Universidade Federal Fluminense, Programa de Pós-Graduação em Biologia Marinha e Ambientes Costeiros, Departamento de Biologia Marinha. Laboratório de Sistemática e Ecologia de Polychaeta, Niterói, 24020-141, Rio de Janeiro, Brazil **3** Universidade Estadual do Maranhão, Laboratório de Pesca e Ecologia Aquática, São Luís, 65055-000, Maranhão, Brazil **4** Universidade Federal do Rio de Janeiro, Campus UFRJ-Macaé Professor Aloísio Teixeira. Núcleo em Ecologia e Desenvolvimento Sócio-Ambiental de Macaé, Grupo de Sistemática e Ecologia de Organismos Bentônicos. Macaé, 27965-045, Rio de Janeiro, Brazil

Corresponding author: Rannyele Passos Ribeiro (rannyele.passos@uam.es)

Academic editor: G. Rouse | Received 23 June 2017 | Accepted 14 January 2018 | Published 27 February 2018

<http://zoobank.org/2DAF40B3-95FF-46BB-AFB4-86E62F116973>

Citation: Ribeiro RP, Alves PR, Almeida ZS, Ruta C (2018) A new species of *Paraonis* and an annotated checklist of polychaetes from mangroves of the Brazilian Amazon Coast (Annelida, Paraonidae). ZooKeys 740: 1–34. <https://doi.org/10.3897/zookeys.740.14640>

Abstract

The polychaete fauna from the mangroves of the Amazon Coast in Maranhão state, Brazil, is reported in this study. Fourteen species are listed, namely *Alitta succinea* (Leuckart, 1847); *Arabella (Arabella) iricolor* Montagu, 1804; *Capitella capitata* (Fabricius, 1780) complex; *Exogone (Exogone) brevi antennata* Hartmann-Schröder, 1959; *Heteromastus filiformis* (Claparède, 1864); *Isolda pulchella* Müller, 1858; *Mediomastus californiensis* Hartman, 1944; *Namalycastis fauweli* Nageswara Rao, 1981; *Namalycastis geayi* (Gravier, 1901); *Namalycastis senegalensis* (Saint-Joseph, 1901); *Nephtys simoni* Perkins, 1980; *Paraonis amazonica* **sp. n.**; *Sigambra bassi* (Hartman, 1945); and *Sigambra grubii* Müller, 1858. Among them, *Namalycastis fauweli* and *Namalycastis geayi* are recorded for the first time in Brazil. *Paraonis amazonica* **sp. n.** is a new species for science, characterized by a rounded prostomium, 4–8 pairs of foliaceous branchiae, absent eyes, and two types of modified neurochaetae, acicular and hook-shaped.

Keywords

annelid, mangrove, Maranhão, new records, *Paraonis*, taxonomy

Introduction

The two largest rivers that drain South America, the Amazon and the Orinoco, are respectively, the first and third largest rivers in the world in terms of water volume (Degens et al. 1991). The Orinoco and Amazon Rivers are responsible for the discharge of an enormous amount of freshwater and sediment into the ocean, representing nearly 20 % of the total global annual freshwater (Hu et al. 2004; Miloslavich et al. 2011). These rivers have been recognized as zoogeographic barriers to the dispersal of marine fauna between the Caribbean and southwestern Atlantic (Gilbert 1972; Floeter and Gasparini 2000). Accordingly, they influence the Brazilian Northern Coast, also known as the Brazilian Amazon Coast, which extends from the north of the Amapá State to the Gulf of Maranhão and represents 35 % of the entire Brazilian Coast (Silveira 1964; Sousa et al. 2008). This region is characterized by a variety of poorly known estuarine and marine ecosystems with very diverse habitats (Couto et al. 2003).

Most of what is known about the marine biodiversity of the Brazilian Amazon Coast is related to fishing and mangrove habitats. Nevertheless, the REVIZEE Program – Living Resources in the Exclusive Economic Zone provided important information regarding the continental shelf and offshore area (Miloslavich et al. 2011). In general, macrobenthos assemblages are one of the least known of Brazil (Amaral and Jablonski 2005; Miloslavich et al. 2011). The lack of studies in this region represents one of the major gaps in the knowledge of the biodiversity of Brazilian polychaetes (see Lana et al. 2009). The main studies concerning polychaetes in this region include ecological research with records in the coast of the states of Maranhão (see Ribeiro and Almeida 2014) and Pará (Rosa-Filho et al. 2006; Morais and Lee 2014) recording a total of 27 species and 24 families.

In this study, polychaetes collected in the mangrove of São Marcos Bay in the Gulf of Maranhão were examined. The current study contributes to increasing the knowledge of polychaetes in the South Atlantic, particularly in the Amazon coastal zone. This is the first taxonomic study with a focus on the polychaete fauna from Maranhão and includes new records and the description of a new species.

Materials and methods

Mangrove specimens from the Gulf of Maranhão were collected between April of 2010 and June of 2012 from two creeks in São Marcos Bay: Buenos Aires at the São Luís (02°35'56"S, 44°21'11.8"W) and Tronco at the Caranguejos Island (02°49'33.6"S, 44°28'51.1"W) (Fig. 1). Along of a 100 m transect, nine sediment samples were taken using a corer (20 cm long and 10 cm diameter) at the intertidal region of each creek. Samples were washed through a 0.5 mm mesh sieve with filtered freshwater in the




Figure 1. Study area. Buenos Aires and Tronco creeks in Maranhão, Brazilian Amazon Coast.

laboratory; specimens were fixed in 4 % formaldehyde and then transferred to 70 % ethanol for long-term storage.

Polychaetes were identified at the species level using stereo (Olympus SZX-16) and light microscope (Olympus CX31). Specimens were prepared for scanning electron microscopy (SEM) by critical point drying, prior to being mounted on stubs and coated with gold (200 Å thick). These specimens were observed and analyzed in the Jeol JSM-6390LV scanning electron microscope of the Museu Nacional/Universidade Federal do Rio de Janeiro (MNRJ). Light microscopy images were taken with a camera attached to a Leica M205A stereo microscope and a Zeiss Axio Scope microscope. Drawings and pictures were processed using Adobe (San Jose, CA, USA) Photoshop CS6.

The specimens and type material referent to the new species were deposited in the research collection Coleção Científica de Invertebrados Marinhos e Costeiros (NPM) of the Núcleo em Ecologia e Desenvolvimento Sócio-Ambiental de Macaé (NUPEM), Macaé, Brazil, and in the Museo Nacional de Ciencias Naturales (MNCN), Madrid, Spain. Additionally, we elaborated a list of some Brazilian records of the species identified here, taking into account those reported by studies formally published (Suppl. material 1). We designated up to one previous record for each Brazilian state indicated in the species distribution section, plus original description, when applicable. Our criteria to select the studies were, preferably: taxonomic approach, ecological approach providing voucher-specimens, ecological approach without voucher-specimens.

Other abbreviations cited in this study:

BMHN	British Museum (Natural History), London.
IBUFRJ	Coleção de Polychaeta do Museu Nacional, Rio de Janeiro.
MCEM-BPO	Centro de Estudos do Mar, Universidade Federal do Paraná, Pontal do Paraná.
MZUSP	Museu de Zoologia da USP, São Paulo.
MPEG.ANL	Museu Paraense Emílio Goeldi, Belém.
POLY-UFPB	Coleção de Invertebrados Marinhos Paulo Young, Universidade Federal da Paraíba, Paraíba.
UK	United Kingdom.
USA	United States of America.
USNM	United States National Museum, Smithsonian Institution, Washington.
ZMH	Zoologisches Staatsinstitut und Zoologisches Museum Hamburg, Hamburg.
ZUEC	Coleção de Polychaeta do Museu de Zoologia “Prof. Adão José Cardoso”, Universidade Estadual de Campinas, Campinas.

Taxonomy

A total of eight families, eleven genera, and 14 species were identified; new records of *Namalycastis geayi* and *Namalycastis senegalensis* (Nereididae) and a new species of *Paraonis* (Paraonidae) are reported from Brazil.

Phylum Annelida

Subclass Errantia

Order Eunicida

Family Oeonidae Kinberg, 1865

Genus *Arabella* Grube, 1850

Arabella (Arabella) iricolor (Montagu, 1804)

Fig. 2

Type locality. Devonshire, England, United Kingdom (50°34'N, 3°34'W; estimated geolocation).

Material examined. São Luís, 02°35'56"S, 44°211'1.8"W: two specimens, 18 March 2012 (NPM-Pol 115); one specimen, 1 June 2012 (NPM-Pol 090); one specimen, 18 November 2011 (NPM-Pol 886); complete and incomplete specimens.

Distribution. Pacific Ocean: New Zealand, Philippines, USA, Peru. Indian Ocean: Red Sea. Atlantic Ocean: Ireland, UK, France, Mediterranean Sea, Marmara




Figure 2. *Arabella (Arabella) iricolor*, fixed specimen. **A** Whole body, dorsal view, arrow point to maxillae and carriers **B** Anterior end, ventral view, arrow point to mandibles **C** Posterior end, dorsal view. Scale bars: 0.5 mm (**A, B**), 0.3 mm (**C**).

Sea (Turkey), Mauritania, South Africa, USA, Mexico, Caribbean Sea, Brazil (states of Maranhão, Bahia, São Paulo and Paraná, see Suppl. material 1).

Remarks. *Arabella (Arabella) iricolor* was described to the south coast of Devonshire (UK) as *Nereis iricolor* (Montagu, 1804). The description of specimens from the

Caribbean Sea (Augener 1927) closely resembles specimens in this study, which were identified as this species due to the characteristics: ventralmost chaeta tapering gradually to guards in median and posterior chaetigers, the absence of hooded acicular chaetae, maxilla MxI unidentate and posterior post-chaetal lobe shorter than chaetae. Body surface whitish was observed in small fixed individuals, probably juveniles (Fig. 2A–C). (Montagu 1804). The species was recorded in ecological studies of the continental shelf, intertidal zone, coral reefs, estuaries, and mangroves (Paiva 1993, Santa-Isabel et al. 2000), but apparently, the material was not deposited in any collection and was not available for comparison. Previous record from Maranhão reports specimens found in mangroves (Oliveira and Mochel 1999). This species has been described with worldwide distribution and is probably a complex of species (Colbath 1989; Zanol and Ruta 2015). Studies on the variation of the symmetry in maxillae and modified ventral chaetae should be conducted to know the polymorphism in species of the genus *Arabella* (Steiner and Amaral 2009). That would be a challenge to species identification and new descriptions, once Oeonidae species are usually collected in low densities (Zanol 2010, Zanol and Ruta 2015).

Order Phyllodocida

Family Nephtyidae Grube, 1850

Genus *Nephtys* Cuvier, 1817

Nephtys simoni Perkins, 1980

Fig. 3

Type locality. Hutchinson Island, Florida, USA (27°21.6'N, 80°13.2'W; original geolocation).

Material examined. São Luís, 02°35'56"S, 44°21'11.8"W: one specimen, 23 April 2010 (NPM-Pol 868); four specimens, 21 October 2010 (NPM-Pol 869); one specimen, 27 January 2011 (NPM-Pol 870); four specimens, 27 January 2011 (NPM-Pol 871); one specimen, 6 September 2011 (NPM-Pol 872); one specimen, 18 December 2011 (NPM-Pol 873); complete and incomplete specimens. Caranguejos Island, 02°49'33.6"S, 44°28'51.1"W: 31 specimens, 20 October 2010 (NPM-Pol 874); 12 specimens, 17 March 2012 (NPM-Pol 455); six specimens, 5 September 2011 (NPM-Pol 875); complete and incomplete specimens.

Distribution. Atlantic Ocean: Mediterranean Sea, USA, Mexico, Brazil (states of Pará, Maranhão, São Paulo, see Suppl. material 1).

Remarks. First record for Maranhão. The specimens present the characters that define *Nephtys simoni* Perkins, 1980, such as interramal branchiae from the third chaetiger; proboscis with long middorsal and midventral subdistal papilla and 23 conical papilla distal, dorsal cirrus linked to pre-chaetal cirrus, short and finger-like lobes. Individuals that have one pair of eyespots and median reddish pigmentation in the prostomium




Figure 3. *Nephtys simoni*, fixed specimen. **A** Whole body, dorsal view **B** Anterior end, dorsal view **C** Posterior end, dorsal view. Scale bars: 0.5 mm.

were reported for juveniles by Perkins (1980). In this study, some specimens presented eyespots, but not the reddish pigmentation pattern. Specimens of *N. simoni* have been reported in estuarine areas as in the type locality (Perkins 1980) and in Amazon mangroves (Silva et al. 2011). In Brazil, the specimens recorded as *Nephtys simoni* in Paranaguá Bay, Paraná (Lana 1986) are in fact *Nephtys californiensis* Hartman, 1938 (Rizzo and Amaral 2007).

Family Nereididae Blainville, 1818**Genus *Alitta* Kinberg, 1865*****Alitta succinea* (Leuckart, 1847)**

Fig. 4

Type locality. Helgoland and Cuxhaven, Germany (53°53'N, 8°37'E; estimated geolocation).

Material examined. São Luís, 02°35'56"S, 44°21'11.8"W: one specimen, 6 September 2011 (NPM-Pol 083); two specimens, 27 January 2011 (NPM-Pol 876); complete and incomplete specimens. Caranguejos Island, 02°49'33.6"S, 44°28'51.1"W: one specimen, 20 October 2010 (NPM-Pol 877); complete and incomplete specimens.

Distribution. Pacific Ocean: Australia, New Zealand, USA, Mexico. Indian Ocean: Red Sea. Atlantic Ocean: North Sea, Mediterranean Sea, South Africa, Canada, USA, Caribbean Sea, Brazil (state of Pará, Maranhão, Pernambuco, Bahia, Espírito Santo, Rio de Janeiro, São Paulo, Paraná, Santa Catarina and Rio Grande do Sul, see Suppl. material 1).

Remarks. This species was described as *Nereis succinea* (Leuckart 1847), transferred to genus *Neanthes* (Imajima 1972), and later to *Alitta* (Bakken and Wilson 2005). The specimens examined in this study share the features of the specimens re-described by Villalobos-Guerrero and Carrera-Parra (2015), such as paragnaths present in all areas of the pharynx, homogomph spinigerous notochaetae, neurochaetae with heterogomph spinigers and homogomph and heterogomph falcigers and the widely expanded notopodial ligule in posterior parapodia. However they are smaller (major individual measuring 3.5 mm of length from the prostomium to the 25th chaetiger) than those described from the Caribbean Sea (Espinosa et al. 2007) and southern-southeastern Brazil (Amaral et al. 2005). On the other hand, specimens from northeastern Brazil measuring less than 5 mm length from the prostomium to the 25th chaetiger are considered recruits (Sette et al. 2013). Therefore, we suggest all the individuals collected in this study are juveniles. *Alitta succinea* is widely distributed in the world with records in different environments. This species was recorded in mangroves from the Caribbean Sea (Londoño-Mesa et al. 2002) and Brazil, including a record in Maranhão state (Mochel 1997). This species is reported in environments with different salinity levels and has been considered as a euryhaline species (Sato 2013).

Genus *Namalycastis* Hartman, 1959***Namalycastis fauveli* Nageswara Rao, 1981**

Fig. 5

Type locality. Estuary of Tachin River, Thailand (13°44'N, 100°30'E; original geolocation).

Material examined. São Luís, 02°35'56"S, 44°21'11.8"W: one specimen, 6 September 2011 (NPM-Pol 883). Caranguejos Island, 02°49'33.6"S, 44°28'51.1"W: one




Figure 4. *Alitta succinea*, fixed specimen. **A** Anterior end, dorsal view **B** Posterior end, dorsal view. Scale bars: 0.5 mm.

specimen, 26 January 2011 (NPM-Pol 086); three specimens, 28 March 2011 (NPM-Pol 109); one specimen, 22 April 2010 (NPM-Pol 878); three specimens, 17 August 2010 (NPM-Pol 879); two specimens, 2 June 2012 (NPM-Pol 880); four specimens, 5 September 2011 (NPM-Pol 881). Complete and incomplete specimens.

Distribution. Indian Ocean: Thailand, India. Atlantic Ocean: Brazil (Maranhão state).

Diagnosis. Body widest mid-anteriorly. Prostomium anteriorly shallowly cleft or cleft absent. Antennae minute. Notochaeta present. Heterogomph setae with boss extremely prolonged. Supra-neuroacicular falcigers in chaetiger 10 with blades slightly curved (Glasby 1999).

Description. Based on specimens NPM-Pol 878 and 883. Complete specimen with 17.3 mm long, 0.72 mm wide (chaetiger 10), and 79 chaetigers (Fig. 5A). Body widest mid-anteriorly, gradually tapering anteriorly and posteriorly. Dorsum convex. Epidermal pigment absent. Prostomium trapezoidal, some individuals with lateral indentation on pros-




Figure 5. *Namalycastis fauveli*, fixed specimen, NPM-Pol 883. **A** Whole body, dorsal view **B** Anterior end, dorsal view **C** Posterior end, ventral view **D** Neurochaetae supra-acicular, 10th chaetiger **E** Neurochaeta spiniger supra-acicular, 10th chaetiger. Scale bars: 0.5 mm (**A–C**), 0.01 mm (**D, E**).

tomium. Prostomium anterior end smooth or with a shallow cleft (Fig. 5B). Narrow longitudinal groove extending from tip to mid-posterior prostomium. Antennae short, extending short of palpopore anterior end, laterally inserted. Two pair of eyes transversally arranged on prostomium. Four pairs of tentacular cirri with indistinct cirrophores and smooth cir-

rostyles. Posterodorsal pair extending posteriorly to third chaetiger. Pharynx smooth, lacking paragnaths or papillae. Parapodia sesquiramous (sub-biramous). Dorsal cirri increasing in length posteriorly. Neuropodial acicular ligulae bilobed. Notochaeta as sesquigomph spinigers present from third chaetiger. Supra-acicular neurochaeta as sesquigomph spinigers on postacicular fascicle and heterogomph falcigers on preacicular fascicle (Fig. 5D, E). Sub-acicular neurochaetae as heterogomph spinigers on postacicular fascicle and heterogomph falcigers in preacicular fascicle. Supra-acicular sesquigomph spinigers shaft with boss $1.2\times-1.5\times$ length of collar. Shaft of heterogomph chaeta with boss prolonged. Supra-acicular falcigers in chaetiger 10 with blades slightly curved, blades length $8.0\times-9.5\times$ width of shaft head. Sub-acicular falcigers blades in chaetiger 10 length, $8.2\times-11.4\times$ (dorsal-most) and $6.0\times-7.3\times$ (ventral most) width of shaft head. Sub-acicular spinigers in anterior region of body with blades finely serrated. Chaeta pale. Aciculae dark brown. Pigidium buttom-shaped (Fig. 5C). Anus terminal. Anal cirri smooth and subconical, arising ventro-laterally.

Colour. Specimens in alcohol yellow. No pigment visible throughout the body.

Remarks. First species record for the America. These specimens present some differences from the original description (Nageswara Rao 1981), such as a dorsal surface convex, body less arched mid-anteriorly, longer antennae and tentacular cirri (Fig. 2A). However, the projection of heterogomph chaetae with an extremely long boss supports this identification for this species (Fig. 2B, C). The differences found are probably because the specimens in this study are juveniles by the smaller size (around 15 mm long, and 80 chaetigers), compared to type material, 21–45 mm long, 134–282 chaetigers, after Glasby (1999). Some Namareneidinae species, as *Namalycastis abiuma*, can have juveniles with blades longer and up to 80 chaetigers. *Namalycastis fauveli* is recorded in estuarine beaches and coastal lagoons in the type locality (Nageswara Rao 1981) and in mangroves of this study.

Namalycastis geayi (Gravier, 1901)

Fig. 6

Type locality. Ouanary, French Guiana ($4^{\circ}12'N$, $51^{\circ}39'W$; estimated geolocation).

Material examined. Caranguejos Island, $02^{\circ}49'33.6''S$, $44^{\circ}28'51.1''W$: one specimen, 17 December 2011 (NPM-Pol 082); one specimen, 27 March 2011 (NPM-Pol 882); one specimen, 20 October 2010 (NPM-Pol 884); all incomplete specimens.

Distribution. Atlantic Ocean: French Guiana, Brazil (state of Maranhão).

Diagnosis. Prostomium anterior end smooth or with a shallow cleft. Antennae extending short of the palpophore tip or of the prostomium tip. Two pairs of eyes nearly longitudinally arranged. Dorsal cirri short, similar in length throughout the body. Notochaetae present (Glasby 1999).

Description. Based on specimen NPM-Pol 884. Incomplete specimen with 4.93 mm long, 1.1 mm wide and 18 chaetigers. Body widest mid-anteriorly, tapering gradually anteriorly and posteriorly (Fig. 6A). Dorsum and venter convex. Epidermal pigment absent. Prostomium trapezoidal, some individuals with lateral indentation on prostomium (Fig. 6A). Prostomium anterior end smooth. Antennae short and smooth,




Figure 6. *Namalycastis geayi*, fixed specimen, NPM-Pol 882 and 884. **A** Anterior end, dorsal view **B** Parapodia sub-biramous, anterior region, dorsal view **C** Supra-acicular spiniger, arrow point to fine serrations proximally to the base of chaetae blades, parapodium 17, dorsal view of chaetae. Scale bars: 0.2 mm (**A**), 0.05 mm (**B**), 0.01 mm (**C**).

extending short of the anterior end of the prostomium, laterally inserted. Two pairs of eyes, arranged nearly longitudinally on prostomium. Four pairs of tentacular cirri with indistinct cirrophores and smooth cirrostyles. Posterodorsal pair extending posteriorly to third chaetiger. Pharynx smooth, lacking paragnaths or papillae. Parapodia sesquiramous (sub-biramous). Dorsal cirri short, similar in length throughout the body. Neuropodial acicular ligulae bilobed. Notochaeta as sesquigomph spinigers present from third chaetiger. Neurochaetae as heterogomph spinigers in all fascicles. Supra-acicular sesquigomph spinigers (postacicular) shaft with boss $1.9\times-2.0\times$ length of collar (Fig. 6B). Shaft of heterogomph chaeta with boss slightly prolonged. Sub-acicular spinigers in anterior region of body with blades moderately serrated (Fig. 6C). Chaeta pale. Aciculae dark brown.

Colour. Specimens in alcohol yellow. No pigment visible throughout the body.

Remarks. First species record for Brazil. The collected specimens of this study were not complete, but they present the same characters of *Namalycastis geayi* (Gravier, 1901) based on the anterior end (Fig. 6A). The identification of this species is supported by the presence of only heterogomph spinigers in sub- and supra-preacicular fascicle in the parapodia (Fig. 6B, C). In the original description, *N. geayi* has been recorded in freshwater environments, muddy river banks, and in coarse sediments (Gravier 1901). This study recorded *N. geayi* in mangroves and brackish water.




Figure 7. *Namalycastis senegalensis*, fixed specimen, NPM-Pol 105. **A** Anterior end, dorsal view **B** Parapodia of 10th chaetiger, anterior view **C** Sub-acicular neurochaetae spiniger of 10th chaetiger **D** Sub-acicular neurochaeta falciger of 10th chaetiger. Scale bars: 2.3 mm (**A**), 0.2 mm (**B**), 0.02 mm (**C, D**).

***Namalycastis senegalensis* (Saint-Joseph, 1901)**

Fig. 7

Type locality. Marsassoun, Senegal (13°59'N, 16°43'W; estimated geolocation).

Material examined. Caranguejos Island, 02°49'33.6"S, 44°28'51.1"W: one specimen, incomplete, 22 October 2010 (NPM-Pol 105).

Distribution. Atlantic Ocean: Senegal, Nigeria, Congo, Suriname, Brazil (states of Pará and Maranhão, see Suppl. material 1).

Remarks. First record for Maranhão. Complete specimens were not found in this study; however, the features of the anterior body are very similar to the re-description of Glasby (1999). The presence of thick cuticle covering the eyes, supra neuro acicular sesquigomph spinigers in the parapodia of chaetiger 10, with a 1.4 × length of collar or more boss, and distally smooth falciger blades supports the identification of the species. Previous Brazilian records include the Amazon coast, the estuarine beaches of Marajó Island in the mouth of the Amazon River (Glasby 1999), and the delta of the Amazon River (one specimen, ZHM PE405) (Glasby 1999). This species is known to live in brackish water and freshwater environments such as mangroves, creeks, and marshes (Glasby 1999).

Family Pilargidae Saint-Joseph, 1899**Genus *Sigambra* Müller, 1858*****Sigambra bassi* (Hartman, 1945)**

Fig. 8A

Type locality. Lemon Bay, Florida, USA (26°54'N, 82°20'W, estimated geolocation).

Material examined. São Luís, 02°35'56"S, 44°21'11.8"W: one specimen, complete, 6 September 2011 (NPM-Pol 111).

Distribution. Pacific Ocean: USA, Mexico, Chile. Atlantic Ocean: USA, Caribbean Sea, Brazil (state of Maranhão and São Paulo, see Suppl. material 1).

Remarks. First record for Maranhão. The specimens examined in this study present a long medium antenna reaching up to setiger 5–12; a dorsal hook beginning in the posterior chaetigers supports the identification as *Sigambra bassi*. The records in the Caribbean and Brazil include estuaries and beaches (Gillet 1986, Amaral et al. 2003).

***Sigambra grubii* Müller in Grube, 1858**

Fig. 8B

Type locality. Florianópolis, Santa Catarina, Brazil (27°36'30"S, 48°26'30"W; original geolocation).

Material examined. São Luís, 02°35'56"S, 44°21'11.8"W: one specimen, 18 August 2010 (NPM-Pol 110); one specimen, 27 January 2011 (NPM-Pol 887). Caranguejos Island, 02°49'33.6"S, 44°28'51.1"W: one specimen, 22 April 2010 (NPM-Pol 888). Complete and incomplete specimens.

Distribution. Atlantic Ocean: USA, Caribbean Sea, Brazil (states of Pará, Maranhão, Sergipe, Rio de Janeiro, São Paulo, Santa Catarina and Rio Grande do Sul, see Suppl. material 1).

Remarks. First species record for Maranhão. The presence of notopodial hooks distally curved appearing in setiger 20 and a medium antenna reaching the second chaetiger are characteristics that support the identification of the species based on the original description by Müller (1858) and re-description by Salazar-Vallejo (1990). In this study, the hooks appeared among the segments 6–29, in specimens shorter and with reduced number of chaetigers, the hooks appeared before the chaetiger 20. This type of variability in the hooks position related with the body size and number of chaetigers was also reported by Salazar-Vallejo (1990). No other morphological variation was found. This species is widely recorded in the coast of Brazil, mainly in estuarine environments, including mangroves and coastal lagoons as the type locality (Müller 1858). In the Caribbean, the species was recorded in a coastal lagoon (Liñero-Arana and Díaz-Díaz 2005).



Figure 8. Pilargidae species. **A** *Sigambra basi*, whole body, dorsal view **B** *Sigambra grubii*, whole body, dorsal view. Scale bars: 0.5 mm.

Family Syllidae Grube, 1850

Genus *Exogone* Örsted, 1845

Subgenus *Exogone (Exogone)* Örsted, 1845

***Exogone (Exogone) breviantennata* Hartmann-Schröder, 1959**

Fig. 9

Type locality. Estero Jaltepeque, El Salvador (13°18'N, 88°52"W; estimated geolocation).

Material examined. São Luís, 02°35'56"S, 44°21'11.8"W: one specimen, 18 August 2010 (NPM-Pol 889); three specimens, 27 January 2011 (NPM-Pol 890); one specimen, 29 March 2011 (NPM-Pol 112); one specimen, 6 September 2011 (NPM-Pol 891); five specimens, 18 December 2011 (NPM-Pol 892); all complete specimens.

Distribution. Pacific Ocean: Australia, Panama, Ecuador. Indian Ocean: Seychelles, Red Sea, Australia. Atlantic Ocean: Spain (Canary Islands), South Africa, Caribbean Sea, Brazil (states of Maranhão, Paraíba, Pernambuco, Espírito Santo, and São Paulo, see Suppl. material 1).




Figure 9. *Exogone (Exogone) breviantennata*. **A** Whole body, dorsal view **B** Anterior end, dorsal view. Abbreviations: pa, palps; ey, eye; pv, proventricle. Scale bars: 0.15 mm.

Remarks. First species record for the Brazilian Amazon Coast. The features that confirm these specimens as *Exogone (Exogone) breviantennata* Hartmann-Schröder, 1959 are median and lateral antennae of similar size, compound spinigers and falcigers with bidentate blades (subdistal tooth larger than distal tooth) and falcigers in the anterior body with 3–4 relatively thick spines. This species is found worldwide in several habitats such as in seagrass in the intertidal zone, rocky shores, algae assemblages, soft bottoms (San Martín and Bone 2001, Paresque et al. 2014), and others. The type

material of *E. (E.) breviantennata* is from a mangrove (Hartmann-Schröder 1959) as in the present study. However, this species has been recorded in several environments and it presents a circumtropical distribution (Núñez et al. 1992).

Subclass Sedentaria

Order Terebellida

Family Ampharetidae Malmgren, 1866

Genus *Isolda* Müller, 1858

Isolda pulchella Müller, 1858

Fig. 10A–C

Type locality. Florianópolis, Santa Catarina, Brazil (27°36'S, 48°27'W; estimated geolocation).

Material examined. São Luís, 02°35'56" S, 44°21'11.8"W: seven specimens, 6 September 2011 (NPM-Pol 849); two specimens, 18 December 2012 (NPM-Pol 067); 14 specimens, 18 December 2012 (NPM-Pol 848). Caranguejos Island, 02°49'33.6"S, 44°28'51.1"W: one specimen, 17 December 2012 (NPM-Pol 850). Complete and incomplete specimens.

Distribution. Pacific Ocean: Australia, USA. Indian Ocean: Red Sea, Australia. Atlantic Ocean: Portugal, Mediterranean Sea, South Africa, USA, Mexico, Caribbean Sea, Brazil (states of Pará, Maranhão, Sergipe, Rio de Janeiro, São Paulo, Paraná and Santa Catarina, see Suppl. material 1).

Remarks. The presence of two groups of four branchiae, post-branchial notopodium with sharply curved hooks; twelve or thirteen thoracic segments with pectinate uncini with four to seven teeth support the identification of these specimens as *Isolda pulchella* Müller, 1858. This species was described in south Brazil and is found along the coast inhabiting estuarine environments, including mangroves and coastal lagoons. The specimens described by Díaz-Díaz and Liñero-Arana (2012) for Caribbean Sea are also similar to the specimens in this study and are recorded in estuaries.

Infraclass Scolecida

Family Capitellidae Grube, 1862

Genus *Capitella* Blainville, 1828

Capitella capitata (Fabricius, 1780), complex

Fig. 11A, B

Type locality. Uummannaq, West Greenland (71°6.5'N, 51°17'W; original geolocation).

Material examined. São Luís, 02°35'56"S, 44°21'11.8"W: six specimens, 29 March 2011 (NPM-Pol 069); one specimen, 23 September 2012 (NPM-Pol 102); 17 specimens, 18 December 2011 (NPM-Pol 851). Caranguejos Island, 02°49'33.6"S,



Figure 10. *Isolda pulchella*. **A** Whole body, dorsolateral view **B** Anterior end, dorsal view **C** Pygidium, ventral view. Scale bars: 0.5 mm.

44°28'51.1"W: five specimens, 17 October 2010 (NPM-Pol 852); two specimens, 26 January 2011 (NPM-Pol 853); two specimens, 17 December 2011 (NPM-Pol 854). Complete and incomplete specimens.

Distribution. Arctic Ocean: Greenland. Pacific Ocean: China, Japan, Australia, USA, Mexico, Costa Rica. Indian Ocean: Red Sea. Atlantic Ocean: North Sea, Ireland, UK, Germany, Netherlands, France, Spain, Mediterranean Sea, Ukraine, South Africa,

USA, Mexico, Caribbean Sea, Brazil (states of Pará, Maranhão, Ceará, Paraíba, Rio de Janeiro, São Paulo, Paraná, Santa Catarina and Rio Grande do Sul, see Suppl. material 1).

Remarks. First record for Maranhão. The specimens examined in this study are similar to the neotype description of *Capitella capitata* by Blake (2009). *C. capitata* was considered as a globally distributed species, but allozyme analyses have demonstrated that this species is composed of at least six sibling species (Grassle and Grassle 1976). Additionally, Blake (2009) indicates that *C. capitata* may be only distributed in Arctic regions, suggesting that the numerous records from lower latitudes are not this species. Nevertheless, the specimens from warmer waters such as in the Caribbean (Amoureux 1985) and Brazil (Pardo et al. 2010) are also similar to the neotype descriptions by Blake (2009). Recently, four new species of *Capitella* were described from the *Capitella capitata* complex (Silva et al. 2017). Studies including molecular data must to be conducted on these animals from Maranhão, and the specimens should be re-examined.

Genus *Heteromastus* Eisig, 1887

Heteromastus filiformis (Claparède, 1864)

Fig. 11C

Type locality. Port-Vendres, France (42°30'N, 3°07'E; estimated geolocation).

Material examined. São Luís, 02°35'56"S, 44°21'11.8"W: one specimen, incomplete, 6 September 2011 (NPM-Pol 070); two specimens, 27 January 2011 (NPM-Pol 852); four specimens, 18 December 2011 (NPM-Pol 856); one specimen, 18 March 2012 (NPM-Pol 857); complete and incomplete specimens.

Distribution. Pacific Ocean: New Zealand, USA, Costa Rica. Indian Ocean: Red Sea, Mozambique. Atlantic Ocean: Ireland, UK, Belgium, France, Mediterranean Sea, South Africa, USA, Mexico, Caribbean Sea, Brazil (states of Pará, Maranhão, Bahia, Rio de Janeiro, São Paulo, see Suppl. material 1).

Remarks. *Heteromastus filiformis* from São Marcos Bay share the same characters of the specimens described by Day (1967) and Dean (2001) such as thoracic region with 12 segments, the first achaetous; thoracic hooks with long hood and about six denticles above the main tooth; abdominal hooks narrow and three to four denticles above the main tooth, gills in subsequent medial segments. The specimens of *H. filiformis* examined in this study are very similar to *H. similis* Southern, 1921. One of the main differences between those species is the presence of gills processes and the shape of neuropodial hooks in *H. filiformis*. According to Hartman (1947), *Heteromastus similis* is considered an inhabitant of freshwater areas and *H. filiformis* is typical of marine environments. Both species have distribution in estuarine environments such as mangroves from Brazil (Silva et al. 2011). In the Caribbean Sea, the records are also in estuarine areas and especially in the muddy intertidal areas of the Caribbean Sea (Gobin 1990). Both species seems to be distributed worldwide, independent of environmental salinity, but descriptions based on fewer characters can be related to several records around the world.



Figure 11. Capitellidae species. **A** *Capitella capitata* complex, whole body, lateral view and **B** anterior view, arrow point genital spines in 9th chaetiger **C** *Heteromastus filiformis*, thoracic region, lateral view **D** *Mediomastus californiensis*, thoracic region, lateral view. Scale bars: 0.5 mm.

Genus *Mediomastus* Hartman, 1944

Mediomastus californiensis Hartman, 1944

Fig. 11D

Type locality. Tomales Bay, California (38°18'N, 122°56'W; estimated geolocation).

Material examined. São Luís, 02°35'56"S, 44°21'11.8"W: two specimens, 21 October 2010 (NPM-Pol 73); three specimens, 18 August 2010 (NPM-Pol 858); one specimen, 18 March 2012 (NPM-Pol 859). Caranguejos Island, 02°49'33.6"S, 44°28'51.1"W: one specimen, 22 April 2010 (NPM-Pol 860), three specimens, 26 January 2011 (NPM-Pol 861); three specimens, 28 March 2011 (NPM-Pol 862); eight specimens, 28 March 2011 (NPM-Pol 863); one specimen, 17 December 2011 (NPM-Pol 864); four specimens, 2 July 2012 (NPM-Pol 865). Complete and incomplete specimens.

Distribution. Pacific Ocean: Australia, USA, Mexico. Atlantic Ocean: Caribbean Sea, Brazil (states of Pará, Maranhão, Rio de Janeiro, São Paulo, Paraná and Santa Catarina, see Suppl. material 1).

Remarks. First record for Maranhão. The specimens examined in this study have triangular prostomium with cylindrical palpodium, in dorsal view; peristomium devoid of setae with a pair of ocelli; 10 chaetigers in thoracic region; only capillaries in chaetigers 1–4; abdominal chaetigers only with hooded hooks defining them as *Mediomastus californiensis* (Hartman, 1944). In the present study, we found specimens exceeding 100 seg-

ments as observed by Warren et al. (1994). Although *M. californiensis* has been recorded in the Pacific (USA) and Atlantic Ocean (Canada and the USA), Warren et al. (1994) examined specimens from both oceans and did not observe differences among them. This species has been recorded in muddy bottoms of estuarine environments in the Brazilian Amazon Coast (Rosa-Filho et al. 2006) and in the Caribbean Sea (Gobin 1990).

Family Paraonidae Cerruti, 1909

Genus *Paraonis* Cerruti, 1909

Paraonis amazonica sp. n.

<http://zoobank.org/D7449E5D-1126-4135-A4B2-DB76AE4CFCCE>

Figs 12, 13, 14

Type locality. Brazil, Maranhão: São Luís, 02°35'56"S, 44°21'11.8"W, mangrove, 21 October 2010, R.P. Ribeiro.

Material examined. Holotype: São Luís, 02°35'56"S, 44°21'11.8"W, one specimen, complete, 21 October 2010 (NPM-Pol 906). Paratypes: São Luís, 02°35'56"S, 44°21'11.8"W, one specimen, incomplete, 18 August 2010 (NPM-Pol 907); 80 specimens, all incomplete, 27 January 2011 (NPM-Pol 908); two specimens, both complete, 21 October 2010 (NPM-Pol 929); two specimens, both incomplete, 18 March 2012 (MNCN 16.01/17766). Caranguejos Island, 02°49'33.6"S, 44°28'51.1"W, three specimens, all incomplete, 26 January 2011 (NPM-Pol 930); 11 specimens, all incomplete, 28 March 2011 (MNCN 16.01/17765).

Distribution. Only known from the type locality.

Diagnosis. Rounded prostomium, clearly wider than longer. Dorsal brownish pigmentation reaching the beginning of the prostomium. Rounded to foliaceous branchiae (4–8 pairs), from the fourth segment. Neurochaetae of two types: acicular chaeta with lateral spine beginning in pre-branchial segments, and hook-shaped chaeta with terminal spines in post-branchial segments.

Description. Complete holotype, 2.68 mm long, 0.17 mm wide (chaetiger 8), and 46 chaetigers. Three complete paratypes with 2.43–2.94 mm long, 0.18–0.20 mm wide and 36–54 chaetigers. Incomplete paratypes up to 4.607 mm long, 0.283 mm wide, and 16–61 chaetigers. Fixed individuals with brown pigmentation that reaches the distal end of the prostomium and extends along the body. Anteriorly flattened body, wider than longer, cylindrical from the 8th chaetiger and in all middle body region (Fig. 12A, B). Branchial region dorsoventrally flattened. Rounded prostomium, wider than longer (Fig. 12B, D). Absence of antenna, palpode, ciliated bands and eyes in the prostomium (Fig. 12A, B). The anterior segments are short, wider than longer. Long and biannulate segments in the post-branchial region. One pair of nuchal organs located on the posterior edge of the prostomium (Fig. 13A). Notopodial post-chaetal lobes absent in the pre-branchial region, the first notopodial post-chaetal lobe appear in the fifth branchial chaetiger. Notopodial post-chaetal cirriform lobes, longer




Figure 12. *Paraonis amazonica* sp. n., fixed specimen, NPM-Pol 906. **A** Whole body, lateral view **B** Anterior end, dorsal view **C** Posterior end, dorsal view **D** Anterior end, the arrow indicates foliaceous branchiae, dorsolateral view. Scale bars: 0.5 mm (**A**), 0.1 mm (**B**) 0.25 mm (**C, D**).

from the middle and posterior regions. Branchiae from chaetiger 4, rounded to foliaceous, flat, short, 4–8 pairs, first and last pairs are shorter (Fig. 12D). Notopodial capillary chaetae throughout the body. Curved capillary chaetae in the neuropodium and notopodium of the pre-branchial and branchial segments (Fig. 13B). Capillary neurochaetae progressively thinner, longer, and straight in the post-branchial segments. Capillary notochaetae of the posterior segments thicker than those anterior and median segments. Pre-branchial and branchial segments with 3–5 chaetae capil-




Figure 13. *Paraonis amazonica* sp. n., SEM. **A** Anterior end, arrow point to the nuchal organ, dorsal view **B** Anterior parapodium of chaetiger 3, arrow point to the acicular lateral spine, dorsal view **C** Parapodium of setiger 35, acicular chaeta with a lateral spine (**sl**) hook-shaped chaeta with a terminal spine (**st**) **D** Acicular chaeta with a lateral spine enlarged. Scale bars: 0.1 mm (**A**), 20 μ m (**B**), 10 μ m (**C**), 1 μ m (**D**).

laries in the notopodium and 2–5 in the neuropodium. Post-branchial segments with 1–2 chaetae capillary in the notopodium, absent in the neuropodium. First acicular neuropodial chaetae with a lateral spine in chaetiger 2–8, and 2–3 chaetae in the branchial segments (Figs 13B, 14A). Neuropodium in the post-branchial middle segments and posterior end segments with one acicular chaeta with a lateral spine (Figs 13B, C, 14A–C). Hook-shaped neurochaetae with terminal spine beginning in post-branchial chaetigers, 1–2 chaetae. Neuropodium in the post-branchial middle chaetigers with 2–4 hook-shaped chaetae with a terminal spine. Neuropodium in posterior chaetigers with two hook-shaped chaetae with a terminal spine (Figs 13B, C, 14B, C). Pygidium rounded with two anal lobes and three anal cirri: two dorsolateral and one medium-ventral (Figs 12C, 14D).




Figure 14. *Paraonis amazonica* sp. n., chaetae. **A** Acicular chaetae with a lateral spine of anterior setiger 3 **B** Posterior hook-shaped chaeta with terminal spine and acicular chaetae with a lateral spine (left) in setiger 30 **C** Hook-shaped chaeta with a terminal spine in the last setiger, 36 **D** Pygidium with three anal cirri, ventral view. Scale bars 50 μ m (**A–C**), 0.25 mm (**D**).

Colour. Specimens in alcohol show brownish pigment spots all over the body, two pairs of reddish brown lateral spots in the pygidial lobes of some specimens.

Etymology. Named after the Amazon Coast, region where type locality is located.

Remarks. *Paraonis amazonica* sp. n. differs from all other species by the presence of acicular and hook-shaped modified neurochaetae. Currently, there are five valid species named in the genus *Paraonis*: *Paraonis fulgens* (Levinsen, 1884); *Paraonis paucibranchiata* Cerruti, 1909; *Paraonis pycnobranchiata* Fauchald, 1972; *Paraonis pygoenigmatica* Jones, 1968; and *Paraonis strelzovi* Hartmann-Schröder, 1980 (see Table 1). Several species first described as *Paraonis* were established as a synonymy of *Aricidea* (López 2008), *Levinsenia* (Gaston 1984), *Paradoneis* (Mackie 1991), and *Paraonides* (Parapar et al. 2012). *Paraonis tenera* Grube, 1873 is a species considered *nomen oblitum* by Strelzov (1973) because its description was inaccurate, being based on a single specimen and probably referring to a species of *Aricidea*.

Among the five valid species of *Paraonis*, *P. fulgens*, *P. paucibranchiata*, and *P. strelzovi* also have the first pair of branchiae in the fourth chaetiger as seen in *P. amazonica* sp. n. However, *P. fulgens* has more than 25 pairs of branchiae and the first post-chaetal lobe starts in the third chaetiger, whereas *P. amazonica* sp. n. has 4–8 pairs of branchiae and first post-chaetal lobe in the 9th chaetiger. In addition, *P. fulgens* (about 120 chaetigers in total) seems to be longer than *P. amazonica* sp. n. (up to 54 chaetigers in complete individuals). However, longer animals could be found, since incomplete individuals of *P. amazonica* sp. n. showed up to 61 chaetigers. Only four pairs of branchiae are described in *P. paucibranchiata* and *P. strelzovi* whereas *P. amazonica* sp. n. has 4–8 pairs of branchiae. Moreover, *P. paucibranchiata* differs from *P. amazonica* sp. n. by the presence of eyes and longer and straighter branchiae. The other two species mainly differ on the first chaetiger with branchiae and post-chaetal lobe. *Paraonis pygoenigmatica* has approximately 20 pairs of branchiae that begin in the sixth chaetiger, joined to the first dorsal lobes. In *P. pycnobranchiata*, the branchiae (about 19) are present from chaetiger 6–25. *P. amazonica* sp. n. and *P. pycnobranchiata* have the same pigmentation pattern consisting in small pigment spots scattered along the body.

Table 1. Key features of *Paraonis* based on original descriptions and redescriptions. NI: no information.

Features	Species					
	<i>P. amazonica</i> sp. n.	<i>P. fulgens</i> (Levinsen, 1884)	<i>P. paucibranchiata</i> Cerruti, 1909	<i>P. pycnobranchiata</i> Fauchald, 1972	<i>P. pygmaenigmatica</i> Jones, 1968	<i>P. strelzovi</i> Hartmann-Schröder, 1980
Eyes	absent	present	present	present	present	absent
Prostomial ciliated bands	absent	present	absent	absent	present	NI
First chaetiger with branchiae	4	4	4	6	6	4
Number of branchiae pairs	4–8	16–25	4	20	15–19	4
Branchiae shape	foliaceous to rounded	foliaceous to oval	cylindrical	thick and distally blunt	Lanceolate	large, smooth and ciliated
Prostomium	rounded	conical	ovoid	rounded pentagonal	Conical	conical
Number of chaetigers	36–54	110–120	at least 20	at least 48	62–81	> 27
Notochaetae	capillary	capillary	capillary	capillary	capillary or limbate	capillary and capillary fringed
Pre-branchial and branchial neurochaetae	capillary and acicular with lateral spine	capillary and hook-shaped with fringe	capillary and hook-shaped	capillary	capillary or limbate	capillary and capillary fringed and hooded spine
Post-branchial neurochaeta	acicular with lateral spine and hook-shaped with terminal spine	hook-shaped with fringe	hook-shaped	curved and pilose without aristae	capillary or limbate and modified	hooded spine
Number of anal cirri	3	3	3	NI	3 to 8	3
Habitat	estuarine, intertidal	marine, intertidal	marine	deep sea	marine, subtidal	estuarine, subtidal
Bottom	muddy	sand bottom	NI	NI	sand bottom	NI
Type locality	Amazon Coast, Brazil	Denmark	Mediterranean Sea	Gulf of California, USA	Cape Cod Bay, USA	Australia

Species of *Paraonis* are usually reported in marine, inshore and continental shelf environments (Glasby and Wilson 2003). There are some exceptions, such as *P. fulgens*, recorded in the intertidal zone from Caribbean Sea (Helguera et al. 2011), *P. strelzovi* in mangroves from Australia (Hartmann-Schröder 1980), and *P. pygoenigmatica* recorded in estuarine areas from Brazil (Barros et al. 2001). *Paraonis amazonica* sp. n. is the first record of a *Paraonis* species found in muddy bottoms in mangrove vegetated areas.

Conclusion

In total, 14 species belonging to eight families and eleven genera were identified in São Marcos Bay, Maranhão, Brazilian Amazon Coast. Two of them were first recorded to Brazilian Coast (*N. fauveli*, *N. geayi*) and one new species was described (*P. amazonica* sp. n.). Two other species are new records for the Brazilian Amazon Coast (*E. (E.) breviantennata* and *S. bassi*), and five species are new records for the Maranhão Coast (*C. capitata* complex, *M. californiensis*, *N. senegalensis*, *N. simoni*, and *S. grubii*).

This study expands the occurrence of *N. geayi* to the Brazilian Amazon Coast (in estuarine muddy sediments) because the type specimens of *N. geayi* were collected in freshwater and muddy bottoms in the Ouanary Stream in French Guiana (Gravier 1901). In addition, a new species of *Paraonis* is described in Amazon mangroves, although Paraonidae is a family commonly found and highly diversified in deep-sea environments (Aguirrezabalaga and Gil 2009). We encourage further studies on this genus because many species need improved descriptions, given that some features lack information in the original descriptions.

In summary, this checklist increases the number of recorded species in the Brazilian Amazon Coast. Further studies targeting sampling beyond mangroves and soft bottoms, including deep sea, seagrasses, and algal mats, can lead to the discovery of higher diversity of annelids in the Brazilian Amazon Coast. We assume that other new species can be found in this region or described from the worldwide species reported here, since they probably correspond to species complexes.

Acknowledgments

We are grateful to a number of colleagues from LabPEA (UEMA) and LABIN (UFRJ-Macacé) for assistance in the field and lab work, in particular, Alana Leitão, Allana Cutrim, Carlos Rodrigues, Marco Antônio Gomes, and Lorena Sousa. We thank the reviewers and Dr. Cinthya Santos (Universidade Federal Fluminense) for helpful comments and suggestions to improve the manuscript. We also thank Verônica Oliveira for a review of the early draft of our manuscript. We acknowledge the Centro de Microscopia Eletrônica de Varredura (CMEV), Museu Nacional/UFRJ for the SEM service. RPR and PRA thank respectively the research institutions CNPq and CAPES for funding their scholarship. ZSA received support from FAPEMA and MPX-Energia. CR was supported by CNPq (PROTAX-562343/2010-5).

References

- Aguirrezabalaga F, Gil J (2009) Paraonidae (Polychaeta) from the Capbreton Canyon (Bay of Biscay, NE Atlantic) with the description of eight new species. *Scientia Marina* 73(4): 631–666. <https://doi.org/10.3989/scimar.2009.73n4631>
- Almeida TCM, Vivan JM, Pessler BH, Lana PC (2012) Polychaetes of the North-Central Santa Catarina state, Brazil. *Check List* 8(2): 200–206. <https://doi.org/10.15560/8.2.204>
- Amaral ACZ, Jablonski S (2005) Conservação da biodiversidade marinha e costeira no Brasil. *Megadiversidade* 1: 43–51.
- Amaral ACZ, Denadai MR, Turra A, Rizzo AE (2003) Intertidal macrofauna in Brazilian subtropical tide-dominated sandy beaches. *Journal of Coastal Research* SI: 446–455. <https://doi.org/10.1590/s1679-8759200000200005>
- Amaral ACZ, Rizzo AE, Arruda EP (2005) Manual dos invertebrados marinhos da região Sudeste-Sul do Brasil. Editora Universidade de São Paulo, São Paulo, 282 pp.
- Amoureux L (1985) Annélides benthiques récoltées à l'entrée de la lagune de la Manche-à-Eau, Guadeloupe (Antilles). *Bulletin du Muséum national d'histoire naturelle* 7(1): 93–107.
- Andrade JT, Palhano NB, Tagliaro CH, Beasley CR (2014) Spatial and temporal variation in the abundance and taxonomic composition of estuarine and terrestrial macrofauna associated with mangrove logs. *Journal of the Marine Biological Association of the United Kingdom* 94(1): 35–42. <https://doi.org/10.1017/S0025315413001215>
- Assis JE, Alonso C, Brito RJ, Santos AS, Christofferen ML (2012) Polychaetous annelids from the coast of Paraíba state, Brazil. *Revista Nordestina de Biologia* 21(1): 3–45.
- Augener H (1927) Polychaeten von Curaçao. *Bijdragen Tot De Kennis Der Fauna Van Curaçao* 25: 39–82.
- Bakken T, Wilson RS (2005) Phylogeny of nereidids (Polychaeta, Nereididae) with paragnaths. *Zoologica Scripta* 34: 507–547. <https://doi.org/10.1111/j.1463-6409.2005.00200.x>
- Barros F, Borzone CA, Rosso S (2001) Macroinfauna of six beaches near Guaratuba Bay, Southern Brazil. *Brazilian Archives of Biology and Technology* 44(4): 351–364. <https://doi.org/10.1590/s1516-89132001000400005>
- Bemvenuti CE, Angonesi LG, Gandra MS (2005) Effects of dredging operations on soft bottom macrofauna on a harbor in the Patos Lagoon estuarine region, Southern Brazil. *Brazilian Journal of Biology* 65(4): 573–581. <https://doi.org/10.1590/S1519-69842005000400003>
- Blake JA (1996) Family Paraonidae Cerruti, 1909. In: Blake, J.A., Hilbig, B., Scott, P.H. (Eds), *Taxonomic atlas of the benthic fauna of the Santa Maria Basin and Western Santa Barbara Channel – The Annelida part 3 – Polychaeta: Orbiniidae to Cossuridae*. Santa Barbara Museum of Natural History, Santa Barbara, 27–70.
- Blake JA (2009) Redescription of *Capitella capitata* (Fabricius) from West Greenland and designation of a neotype (Polychaeta, Capitellidae). *Zoosymposia* 2: 55–80. <https://doi.org/10.3853/j.0067-1975.34.1982.295>
- Blankensteyn A, Moura RS (2002) Lista preliminar das espécies da macrofauna de fundos consolidados da baía de Guaratuba, Paraná, Brasil. *Revista Brasileira de Zoologia* 19(3): 715–721. <https://doi.org/10.1590/S0101-81752002000300008>
- Bone D, Klein E (2000) Temporal variations in a tropical soft-bottom community, Venezuela. *Journal of Coastal Research* 16(2): 278–286. <https://doi.org/10.2112/06-0790.1>

- Braga CF, Monteiro VF, Rosa-Filho JS, Beasley CR (2011) Benthic macroinfaunal assemblages associated with Amazonian saltmarshes. *Wetlands Ecology and Management* 19(3): 257–272. <https://doi.org/10.1007/s11273-011-9215-5>
- Braga CF, Silva RF, Rosa-Filho JS, Beasley CR (2013) Spatio-temporal changes in macroinfaunal assemblages of tropical saltmarshes, northern Brazil. *Pan-American Journal of Aquatic Sciences* 8(4): 28–298.
- Brasil ACS, Silva SHG (2000) Spatial distribution of Polychaeta in a soft-bottom community at Saco do Céu, Ilha Grande, Rio de Janeiro, Brazil. *Bulletin of Marine Science* 67(1): 103–112.
- Cerruti A (1909) Contributo all'Anatomia, biologia e sistematica delle Paraonidae (Levinseniidae) con particolare riguardo alle specie del golfo di Napoli. *Mitteilungen aus der Zoologischen Station zu Neapel* 19(3): 459–512.
- Claparède E (1864) Glanures zootomiques parmiles annélides de Port-Vendres (Pyrénées Orientales). *Memoires de la Société de physique et d'histoire naturelle de Genève* 17(2): 463–600. <https://doi.org/10.5962/bhl.title.14827>
- Colbath GH (1989) A revision of *Arabella mutans* (Chamberlin, 1919) and related species (Polychaeta: Arbellidae). *Proceedings of the Biological Society of Washington* 102(2): 283–299.
- Colling LA, Capitoli RR, Bemvenuti CE (2007) Epitoquia de *Neanthes succinea* na região estuarina da Lagoa dos Patos. *Atlântica* 29: 61–63.
- Couto E, Silveira FL, Rocha G (2003) Marine biodiversity in Brazil: the current status. *Gayana* 67: 327–340. <https://doi.org/10.4067/S0717-65382003000200014>
- Day JH (1967) A monograph on the Polychaeta of Southern Africa. Trustees of the British London Museum (Natural History), London, 878 pp. <https://doi.org/10.5962/bhl.title.8596>
- Dean HK (2001) Capitellidae (Annelida: Polychaeta) from the Pacific coast of Costa Rica. *Revista de Biología Tropical* 2: 69–84. <https://doi.org/10.11646/zootaxa.3956.2.2>
- Dean HK (2012) A literature review of the Polychaeta of the Caribbean Sea. *Zootaxa* 3596: 1–86. [https://doi.org/10.1016/0198-0254\(87\)90315-3](https://doi.org/10.1016/0198-0254(87)90315-3)
- Degens ET, Kempe S, Richey JE (1991) Summary: biogeochemistry of major world rivers. In: Degens ET, Kempe S, Richey JE (Eds) *Biogeochemistry of major world rivers*. John Wiley and Sons, New York, 323–347. <https://doi.org/10.1002/aqc.3270010209>
- Díaz-Díaz O, Liñero-Arana I (2012) Ampharetidae Malmgren, 1867 (Annelida: Polychaeta) de Venezuela. *Boletín de Investigaciones Marinas y Costeras* 41: 165–177.
- Espinosa VV, Díaz OD, Liñero-Arana I (2007) Nereididae Lamarck, 1818 (Annelida: Polychaeta) de la Costa Occidental de Venezuela. *Boletín del Instituto Oceanográfico de Venezuela* 46: 119–127.
- Fabricius O (1780) *Fauna Groenlandica*. Ioannis Gottlob Rothe, Hafniae et Lipsiae, 466 pp. <https://doi.org/10.5962/bhl.title.13489>
- Fauchald K (1972) Benthic polychaetous annelids from deep water off western Mexico and adjacent areas in the Eastern Pacific Ocean. *Allan Hancock Monographs in Marine Biology* 7: 1–575. <https://doi.org/10.1080/00364827.1972.10411211>
- Fauvel P (1927) *Faune de France*. Polychètes sédentaires addenda aux errantes, archiannelides, mizostomaires. Lechevalier, Paris, 494 pp. <https://doi.org/10.1038/113528b0>
- Floeter SR, Gasparini JL (2000) The southwestern Atlantic reef fish fauna: composition and zoogeographic patterns. *Journal of Fish Biology* 56: 1099–1114. <https://doi.org/10.1111/j.1095-8649.2000.tb02126.x>

- Gaston GR (1984) Paraonidae. In: Uebelacker M, Jonhson PG, Vittor B (Eds) Taxonomic guide to the polychaetes of the northern Gulf of Mexico 1(2). Barry A. Vittor and Associates, Inc., Louisiana, 1–53. <https://doi.org/10.5962/bhl.title.4035>
- Gilbert CR (1972) Characteristics of the western Atlantic reef-fish fauna. Quarterly Journal of the Florida Academy of Sciences 35: 130–144. <https://doi.org/10.2174/1874401x00902010090>
- Gillet P (1986) Contribution à l'étude des Annélides Polychètes des lagunes de la Manche-à-Eau et de Belle-Plaine (Guadeloupe). Description d'un nouveau Capitellidae: *Scyphoprotus guadalupensis* n. sp.. Bulletin du Muséum national d'Histoire naturelle 8: 803–817. <https://doi.org/10.5962/bhl.part.6290>
- Glasby CJ (1999) The Namanereidinae (Polychaeta: Nereididae). Part 1, Taxonomy and Phylogeny. Part 2, Cladistic Biogeography. Records of the Australian Museum 25: 1–129. <https://doi.org/10.3853/j.0812-7387.25.1999.1355>
- Glasby CJ, Wilson RS (2003) Paraonidae (Polychaeta) – A DELTA database of genera, and Australian species. In: Wilson RS, Hutchings PA, Glasby CJ (Eds) Polychaetes: Interactive identification and information retrieval. CSIRO Publishing, Melbourne. <https://doi.org/10.3853/j.0067-1975.40.1988.150>
- Gobin JF (1990) A checklist of marine polychaetous annelids (Polychaeta) for the Gulf of Paria, Trinidad, West Indies. Caribbean Marine Studies 1(1): 37–47. <https://doi.org/10.1017/s1755267213000390>
- Gobin JF (2010) Free-living marine polychaetes (Annelida) inhabiting hard-bottom substrates in Trinidad and Tobago, West Indies. Revista de Biología Tropical 58(1): 147–157. <https://doi.org/10.15517/rbt.v58i1.5200>
- Grassle JP, Grassle JF (1976) Sibling species in the marine pollution indicator *Capitella* (Polychaeta). Science 192: 567–569. <https://doi.org/10.1126/science.1257794>
- Gravier C (1901) Sur deux nouvelles espèces du genre *Lycastis* Savigny, Aud. et Edw. rev., de la Guyane Française. Bulletin du Muséum national d'Histoire naturelle 7(8): 397–402. <https://doi.org/10.5962/bhl.part.80291>
- Grube AE (1873) Über ein paar neue Anneliden aus der Familie der Spiodeen. Jahres-Bericht der Schlesischen Gesellschaft für Vaterländische Cultur 50: 57–58.
- Hartman O (1938) Review of the annelid worms of the family Nephtyidae from the northeast Pacific, with descriptions of five new species. Proceedings of the United States National Museum 8(5): 143–158. <https://doi.org/10.5479/si.00963801.85-3034.143>
- Hartman O (1944) Capitellidae and Nereidae (marine annelids) from the gulf side of Florida, with a review of freshwater Nereididae. Bulletin of Marine Science of the Gulf and Caribbean 9(2): 153–168.
- Hartman O (1945) The marine annelids of North Carolina. Duke University Marine Station Bulletin 2: 1–54. <https://doi.org/10.1086/395019>
- Hartman O (1947) Polychaetous annelids Part VII. Capitellidae. Allan Hancock Pacific Expeditions 10: 391–481.
- Hartmann-Schröder G (1959) Zur Ökologie der Polychaeten des Mangrove-Estero-Gebietes von El Salvador. Beiträge zur Neotropischen Fauna 1(2): 69–183. <https://doi.org/10.1080/01650525909380612>

- Hartmann-Schröder G (1980) Die Polychaeten der tropischen Nordwestküste Australiens (zwischen Port Samson im Norden und Exmouth im Süden). In: Hartmann-Schroeder G, Hartmann G (Eds) Zur Kenntnis des Eulitorals der australischen Küsten unter besonderer Berücksichtigung der Polychaeten und Ostracoden. Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut 77: 41–110.
- Helguera Y, Díaz-Asencio L, Fernández-Garcés R, Gómez-Batista M, Guillén A, Díaz-Asencio M, Armenteros M (2011) Distribution patterns of macrofaunal polychaete assemblages in a polluted semi-enclosed bay: Cienfuegos, Caribbean Sea. *Marine Biology Research* 7: 757–768. <http://dx.doi.org/10.1080/17451000.2011.569552>
- Hu C, Montgomery ET, Schmitt RW, Muller-Karger FE (2004) The dispersal of the Amazon and Orinoco River water in the tropical Atlantic and Caribbean Sea: Observation from space and S-PALACE floats. *Deep-Sea Research Part II* 51: 1151–1171. <http://dx.doi.org/10.1016/j.dsr2.2004.04.001>
- Imajima M (1972) Review of the annelid worms of the family Nereidae of Japan, with descriptions of five new species or subspecies. *Bulletin of the National Science Museum* 15: 37–153.
- Jones ML (1968) *Paraonis pygoenigmatica* new species, a new annelid from Massachusetts (Polychaeta: Paraonidae). *Proceedings of the Biological Society of Washington* 81: 323–334.
- Lana PC (1986) Nephthyidae (Annelida; Polychaeta) do litoral do Estado do Paraná (Brasil). *Nerítica* 1(1): 135–154. <https://doi.org/10.5380/rn.v1i1.41022>
- Lana PC, Amaral ACZ, Souza JRB, Ruta C, Paiva PC, Brasil ACS, Santos CSG, Garraffoni ARS (2009) Polychaeta. In: Rocha RM, Boeger WA (Eds) Estado da arte e perspectivas para a zoologia no Brasil. Editora Universidade Federal do Paraná, Curitiba, 91–100.
- Lana PC, Santos CSG, Garraffoni ARS, Oliveira VM, Radashevsky V (2006) Checklist of polychaete species from Parana State (Southern Brazil). *Check List* 2: 30–64. <https://doi.org/10.15560/2.3.30>
- Leuckart R (1847) Verzeichniss der zur Fauna Helgoland's gehörenden wirbellosen Seethiere. In: Frey H, Leuckart R (Eds) Beiträge zur Kenntniss wirbelloser Thiere, mit besonderer Berücksichtigung der Fauna des Norddeutschen Meeres. Vieweg, Braunschweig, 136–168. <https://doi.org/10.5962/bhl.title.2128>
- Levinsen GMR (1884) Systematisk-geografisk Oversigt over de nordiske Annulata, Gephyrea, Chaetognathi og Balanoglossi. *Videnskabelige Meddelelser fra den naturhistoriske Forening i Kjöbenhavn* 45(1883): 92–350.
- Licher F (1999) Revision der Gattung *Typosyllis* Langerhans, 1879 (Polychaeta: Syllidae). Morphologie, Taxonomie und Phylogenie. *Abhandlungen der Senckenbergischen naturforschenden Gesellschaft* 551: 1–336.
- Liñero-Arana IL, Díaz-Díaz OD (2005) Notes on some pilargid polychaetes (Annelida) from Venezuela. *Boletín del Instituto Oceanográfico de Venezuela* 44(1): 67–70
- Londoño-Mesa M, Polanía MJ, Vélez I (2002) Polychaetes of the mangrove-fouling community at the Colombian Archipelago of San Andrés and Old Providence, Western Caribbean. *Wetlands Ecology and Management* 10: 227–232. <https://doi.org/10.1023/a:1020127814042>
- López E (2008) Redescription of *Aricidea belgicae* (Fauvel, 1936), an Antarctic Paraonidae (Annelida: Polychaeta), with reassessment of its taxonomic status based on new material. *Proceedings of the Biological Society of Washington* 121(3): 297–307. <https://doi.org/10.2988/08-05.1>

- Mackie ASY (1991) *Paradoneis eliasoni* sp. nov. (Polychaeta: Paraonidae) from northern european waters, with a redescription of *Paradoneis lyra* (Southern, 1914). *Ophelia* 5: 147–155. <https://doi.org/10.1080/00785326.1986.10429745>
- Miloslavich P, Klein E, Diaz JM, Hernández CE, Bigatti G, Campos L, Artigas F, Castillo J, Penschazadeh PE, Neil PE, Carranza A, Retana MV, Astarloa JMD, Lewis M, Yorio P, Piriz ML, Rodriguez D, Yoneshigue-Valentin Y, Gambo L, Martin A (2011) Marine Biodiversity in the Atlantic and Pacific Coasts of South America: knowledge and gaps. *PLoS ONE* 6: e14631. <https://doi.org/10.1371/journal.pone.0014631>
- Mochel FR (1997) Mangroves on São Luís Island, Maranhão, Brazil. In: Kjerfve B, Lacerda LD, Diop EHS (Eds) *Mangrove ecosystem studies in Latin America and Africa*, Vol. 1. UNESCO, Paris, 145–154.
- Montagu G (1804) Description of several marine animals found on the south coast of Devonshire. *Transactions of the Linnean Society of London* 7(1): 61–85. <https://doi.org/10.1111/j.1096-3642.1804.tb00282.x>
- Morais GC, Lee JT (2014) Intertidal benthic macrofauna of rare rocky fragments in the Amazon region. *Revista de Biología Tropical* 62(1): 69–86. <https://doi.org/10.15517/rbt.v62i1.8425>
- Müller F (1858) Einiges über die Annelidenfauna der Insel Santa Catharina an der brasilianischen Küste. *Archiv für Naturgeschichte* 24(1): 211–220. http://dx.doi.org/10.1007/978-3-662-40099-9_12
- Nageswara-Rao CA (1981) On two new polychaetes (Nereidae: Annelida) from estuarine waters of India. *Bulletin of the Zoological Survey of India* 3(3): 213–217. <https://doi.org/10.11646/phytotaxa.292.1.8>
- Núñez J, San Martín G, Brito MC (1992) Exogoninae (Polychaeta: Syllidae) from the Canary Islands. *Scientia Marina* 56(1): 43–52. <https://doi.org/10.1007/s12526-009-0026-3>
- Oliveira VM, Mochel FR (1999) Macroendofauna benthica de substratos móveis de um manguezal sob impacto das atividades humanas no sudoeste as ilha de São Luís, Maranhão, Brasil. *Boletim do Laboratório de Hidrobiologia* 12(1): 75–93.
- Oliveira AB, Rizzo AE, Conceição E, Couto G (2011) Benthic macrofauna associated with decomposition of leaves in a mangrove forest in Ilhéus, state of Bahia, Brazil. *Journal of the Marine Biological Association of the United Kingdom* 92(7): 1–9.
- Omena EP, Amaral ACZ (2003) Sandy beach morphodynamic and the polychaete fauna in Southeast Brazil. *Journal of Coastal Research Special Issue* (35): 431–439.
- Omena EP, Lavrado HP, Paranhos R, Silva TA (2012) Spatial distribution of intertidal sandy beach polychaeta along an estuarine and morphodynamic gradient in an eutrophic tropical bay. *Marine Pollution Bulletin* 64(9): 1861–1873.
- Ouvides TMS, Boehs G, Rizzo AE (2011) Composition and spatial distribution of the benthic macrofauna in the Cachoeira River estuary, Ilhéus, Bahia, Brazil. *Revista de Biología Marina y Oceanografía* 46(1): 17–25. <https://doi.org/10.1016/j.marpolbul.2012.06.009>
- Paiva PC (1993) Anelídeos poliquetas da plataforma continental norte do Estado de São Paulo: I - Padrões de densidade e diversidade específica. *Boletim do Instituto Oceanográfico* 41(1/2): 69–80. <https://doi.org/10.1590/s0373-55241993000100006>
- Parapar J, Alós C, Núñez J, Moreira J, López E, Aguirrezabalaga F, Besteiro C, Martínez A (2012) Annelida Polychaeta III. In: Ramos MA, et al. (Eds) *Fauna Ibérica*. Museo Nacional de Ciencias Naturales, CSIC, Madrid, 416 pp. <https://doi.org/10.3989/scimar.2006.70n1159>

- Pardo EV, Teixeira LLS, Amaral ACZ (2010) Morphometric analysis of *Capitella capitata* (Polychaeta, Capitellidae). *Iheringia Série Zoologia* 100(1): 13–18. <https://doi.org/10.1590/s0073-47212010000100002>
- Paresque K, Fukuda MV, Nogueira JMM (2014) The genus *Exogone* (Polychaeta: Syllidae) from the Brazilian coast, with the description of a new species. *Zootaxa* 3790(4): 501–533. <https://doi.org/10.11646/zootaxa.3790.4.1>
- Perkins TH (1980) Review of the species previously referred to *Ceratonereis mirabilis*, and descriptions of new species of *Ceratonereis*, *Nephtys*, and *Goniada* (Polychaeta). *Proceedings of the Biological Society of Washington* 93(1): 1–49. <https://doi.org/10.1007/bf00409279>
- Ribeiro RP, Almeida ZS (2014) Anelídeos Poliquetas do estado do Maranhão, Brasil: síntese do conhecimento. *Bioikos* 28(1): 45–55.
- Rizzo AE, Amaral ACZ (2007) Nephtyidae (Annelida: Polychaeta) from São Paulo State, Brazil, including a new record for the Brazilian coast. *Biota Neotropica* 7(3): 253–263. <https://doi.org/10.1590/S1676-06032007000300028>
- Rosa-Filho JS, Busman DV, Viana AP, Greg AM, Oliveira DM (2006) Macrofauna bentônica de zonas entre-marés não vegetadas do estuário do rio Caeté, Bragança, Pará. *Boletim do Museu Paraense Emílio Goeldi. Ciências Naturais* 1(3): 85–96.
- Rosa-Filho JS, Gomes TP, Almeida MF, Silva RF (2011) Benthic fauna of macrotidal sandy beaches along a small-scale morphodynamic gradient on the Amazon coast (Algodão Island, Brazil). *Journal of Coastal Research Special Issue* (64): 435–439.
- Rullier F, Amoureux L (1979) Annélides polychètes. In: Campagne de la Calypso au large des côtes Atlantiques de l'Amérique du Sud (1961–1962). I. 33. *Annales de l'Institut océanographique* 55: 145–206. <https://doi.org/10.3406/mefr.1979.1196>
- Saint-Joseph A (1901) Sur quelques invertébrés marins des côtes du Sénégal (annelides polychètes, nematoïde endoparasite d'annelide polychète et crustacé décapode parasite). *Annales des sciences naturelles, Zoologie et Paléontologie* 8(12): 217–246. <https://doi.org/10.1163/156854060x00276>
- Salazar-Vallejo SI (1990) Redescription of *Sigambra grubii* Müller, 1858 and *Hermundura tricuspis* Müller, 1858 from Brazil and designation of neotypes (Polychaeta: Pilargidae). *Journal of Natural History* 24: 507–517. <https://doi.org/10.1080/00222939000770351>
- San Martín G, Bone D (2001) Syllidae (Polychaeta) de praderas de *Thalassia testudinum* en el Parque Nacional Morrocoy (Venezuela). *Revista de Biología Tropical* 49: 609–620. <https://doi.org/10.1023/a:1026117503709>
- Santa-Isabel LM, Leão ZMAN, Peso-Aguiar MC (2000) Polychaetes from the Guarajuba coral reefs, Bahia, Brazil. *Bulletin of Marine Science* 67(1): 645–653. https://doi.org/10.1007/978-94-017-1982-7_28
- Santi L, Tavares M (2009) Polychaete assemblage of an impacted estuary, Guanabara Bay, Rio de Janeiro, Brazil. *Brazilian Journal of Oceanography* 57(4): 287–303. <https://doi.org/10.1590/S1679-87592009000400004>
- Santos MA, Santos CSG, Oliveira CMM (1994) Polychaeta in the estuary of the Piauí River, Sergipe, Brazil. *Mémoires du Muséum d'Histoire naturelle* 162: 541–547.
- Santos MFL, Pires-Vanin AMS (2004) Structure and dynamics of the macrobenthic communities of Ubatuba Bay, southeastern Brazilian coast. *Brazilian Journal of Oceanography* 52(1): 59–73. <https://doi.org/10.1590/S1679-87592004000100006>

- Sars M (1851) Beretning om en i Sommeren 1849 foretagen zoologisk Reise i Lofotenog Fimarken. *Nyt Magazin for Naturvidenskaberne* 6: 12–211. <https://doi.org/10.4045/tidsskr.10.1261>
- Sato M (2013) Resurrection of the genus *Nectoneanthes* Imajima, (Nereididae: Polychaeta), with redescription of *Nectoneanthes oxypoda* (Marenzeller, 1879) and description of a new species, comparing them to *Neanthes succinea* (Leuckart, 1847). *Journal of Natural History* 47(1/2): 1–50. <https://doi.org/10.1080/00222933.2012.743609>
- Sette CSC, Shinozaki-Mendes RA, Barros TL, Souza JRB (2013) Age and growth of *Alitta succinea* (Polychaeta : Nereididae) in a tropical estuary of Brazil. *Journal of the Marine Biological Association of the United Kingdom* 93(8): 2123–2128. <https://doi.org/10.1017/S0025315413000854>
- Silva CF, Seixas VC, Barroso R, Di Domenico M, Amaral ACZ, Paiva PC (2017) Demystifying the *Capitella capitata* complex (Annelida, Capitellidae) diversity by morphological and molecular data along the Brazilian coast. *PLOS ONE* 12(5): e0177760. <https://doi.org/10.1371/journal.pone.0177760>
- Silva RF, Rosa-Filho JS, Souza SR, Souza Filho PWM (2011) Spatial and temporal changes in the structure of soft-bottom benthic communities in an Amazon estuary (Caeté estuary, Brazil). *Journal of Coastal Research Special Issue* (64): 440–444. <https://doi.org/10.3989/scimar.03312.16c>
- Silveira JD (1964) Morfologia do litoral. In: Azevedo A (Ed.) *Brasil: a terra e o homem*. Companhia Editora Nacional, São Paulo, 253–305.
- Sousa EB, Costa VB, Pereira LCC, Costa RM (2008) Microfitoplâncton de águas costeiras amazônicas: ilha Canela (Bragança, PA, Brasil). *Acta Botanica Brasilica* 22: 626–636. <https://doi.org/10.1590/s0102-33062008000300004>
- Southern R (1921) Polychaeta of the Chilka Lake and also of fresh and brackish waters in other parts of India. *Memoirs of the Indian Museum* 5: 563–659. <https://doi.org/10.1038/1741131d0>
- Souza-Filho PWM (2005) Costa de manguezais de macromaré da Amazônia: cenários morfológicos, mapeamento e quantificação de áreas usando dados de sensores remotos. *Revista Brasileira de Geofísica* 23(4): 427–435. <https://doi.org/10.1590/s0102-261x2005000400006>
- Steiner TM, Amaral ACZ (2009) *Arabella aracaensis*, a new species with growth rings on its mandibles, and some remarks on the endoparasitic *Labrorostratus prolificus* (Polychaeta: Oeonidae) from southeast Brazil. *Journal of Natural History* 43(41–42): 2537–2551. <https://doi.org/10.1080/00222930903219988>.
- Strelzov VE (1973) Polychaete worms of the family Paraonidae Cerruti, 1909 (Polychaeta, Sedentaria). *Akademiya Nauk, Leningrad*, 170 pp.
- Suárez AM (1981) Poliquetos bentónicos Cubanos II. Sedentaria. *Revista de Investigaciones Marinas* 2: 3–47.
- Uebelacker JM (1984) *Arabellidae* Hartman, 1944. In: Uebelacker JM, Johnson PG, Vittor B (Eds) *Taxonomic guide to the polychaetes of the northern Gulf of Mexico* 6(42). Barry A, Vittor and Associates, Inc., Louisiana, 1–29. <https://doi.org/10.5962/bhl.title.4035>
- Viana MG, Rocha-Barreira CA, Hijo CG (2005) Macrofauna bentônica da faixa entremarés e zona de arrebentação da praia de Paracuru (Ceará-Brasil). *Brazilian Journal of Aquatic Science and Technology* 9(1): 75–82. <https://doi.org/10.14210/bjast.v9n1.p75-82>
- Villalobos-Guerrero T, Carrera-Parra LF (2015) Redescription of *Alitta succinea* (Leuckart, 1847) and reinstatement of *A. acutifolia* (Ehlers, 1901) n. comb. based upon morpho-

- logical and molecular data (Polychaeta: Nereididae). *Zootaxa* 3919(1): 157–178. <https://doi.org/10.11646/zootaxa.3919.1.7>
- Warren LM, Hutchings PA, Doyle S (1994) A revision of the genus *Mediomastus* Hartman, 1944 (Polychaeta: Capitellidae). *Records of the Australian Museum* 46(3): 227–256. <https://doi.org/10.3853/j.0067-1975.46.1994.6>
- Young DK, Young MW (1982) Macrobenthic invertebrates in bare sand and seagrass (*Thalassia testudinum*) at Carrie Bow Cay lagoon. *Smithsonian Contributions to Marine Science* 12: 115–126.
- Zanol J (2010) Homology of prostomial and pharyngeal structures in Eunicida (Annelida) based on innervation and morphological similarities. *Journal of Morphology* 271(9): 1023–1043. <https://doi.org/10.1002/jmor.10843>
- Zanol J, Ruta C (2015) New and previously known species of Oeonidae (Polychaeta: Annelida) from Lizard Island, Great Barrier Reef, Australia. *Zootaxa* 4019(1): 745–772. <https://doi.org/10.11646/zootaxa.4019.1.26>

Supplementary material I

Brazilian records of the species identified from Amazon coast, Maranhão

Authors: Rannyele Passos Ribeiro, Paulo Ricardo Alves, Zafira da Silva de Almeida, Christine Ruta

Data type: species data

Explanation note: Information on deposit number, record, state, coordinates, habitat, substrate and depth and references of the records are provided.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.740.14640.suppl1>