



Randomized phase II trial of survivin 2B peptide vaccination for patients with HLA-A24-positive pancreatic adenocarcinoma

Hiroaki Shima¹ | Giichiro Tsurita² | Satoshi Wada^{3,4} | Yoshihiko Hirohashi⁵  | Hiroshi Yasui⁶ | Hiroshi Hayashi⁷ | Takashi Miyakoshi⁷ | Kazue Watanabe⁵ | Aiko Murai⁵ | Hiroko Asanuma⁸ | Serina Tokita⁵ | Terufumi Kubo⁵ | Munehide Nakatsugawa⁵ | Takayuki Kanaseki⁵ | Tomohide Tsukahara⁵  | Yutaka Nakae⁹ | Osamu Sugita⁷ | Yoichi M. Ito¹⁰ | Yasunori Ota¹¹ | Yasutoshi Kimura¹ | Goro Kutomi¹ | Koichi Hirata¹ | Toru Mizuguchi¹ | Kohzoh Imai⁶ | Ichiro Takemasa¹ | Noriyuki Sato⁵ | Toshihiko Torigoe⁵

¹Department of Surgery, Surgical Oncology and Science, Sapporo Medical University School of Medicine, Sapporo, Japan

²Department of Surgery, Research Hospital, The Institute of Medical Science, The University of Tokyo, Tokyo, Japan

³Department of Clinical Diagnostic Oncology, Showa University, Tokyo, Japan

⁴Cancer Vaccine Center, Kanagawa Cancer Center, Kanagawa, Japan

⁵Department of Pathology, Sapporo Medical University School of Medicine, Sapporo, Japan

⁶The Institute of Medical Science, The University of Tokyo, Tokyo, Japan

⁷Hokkaido University Hospital Clinical Research and Medical Innovation Center, Sapporo, Japan

⁸Department of Surgical Pathology, Sapporo Medical University School of Medicine, Sapporo, Japan

⁹Collaboration Center for Community and Industry, Sapporo Medical University School of Medicine, Sapporo, Japan

¹⁰Department of Biostatistics, Hokkaido University Graduate School of Medicine, Sapporo, Japan

¹¹Department of Pathology, Research Hospital, The Institute of Medical Science, The University of Tokyo, Tokyo, Japan

Correspondence

Yoshihiko Hirohashi or Toshihiko Torigoe, Department of Pathology, Sapporo Medical University School of Medicine, Sapporo, Japan.

Email: hirohash@sapmed.ac.jp (YH); torigoe@sapmed.ac.jp (TT)

Funding information

Japan Agency for Medical Research and Development, Grant/Award Number 16cm0106309h0001; Japan Society for the Promotion of Science, Grant/Award Number 15H04722 17H01540

Abstract

The prognosis of advanced pancreatic adenocarcinoma is still extremely poor. This study sought to determine the efficacy of, and immunological response to, peptide vaccination therapy in patients with this disease. In this multicenter randomized phase II study, patients with advanced pancreatic adenocarcinoma after gemcitabine and/or tegafur/gimeracil/oteracil were randomly assigned to 3 groups that each received a 2-step treatment course. In Step 1, the groups received treatments of: (i) survivin 2B peptide (SVN-2B) plus interferon- β (IFN β); (ii) SVN-2B only; or (iii) placebo until the patients show progression. In Step 2, all patients who consented to participate received 4 treatments with SVN-2B plus IFN β . The primary endpoint was progression-free survival (PFS) after initiation of Step 1 treatment. Secondary endpoints included immunological effects assessed by analysis of PBMCs after Step 1. Eighty-three patients were randomly assigned to receive SVN-2B plus IFN β (n = 30),

Hiroaki Shima, Giichiro Tsurita, and Satoshi Wada equally contributed to this work.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors. *Cancer Science* published by John Wiley & Sons Australia, Ltd on behalf of Japanese Cancer Association.

SVN-2B (n = 34), or placebo (n = 19). No significant improvement in PFS was observed. Survivin 2B-specific CTLs were found to be increased in the SVN-2B plus IFN β group by tetramer assay. Among patients who participated in Step 2, those who had received SVN-2B plus IFN β in Step 1 showed better overall survival compared with those who had received placebo in Step 1. Patients vaccinated with SVN-2B plus IFN β did not have improved PFS, but showed significant immunological reaction after vaccination. Subgroup analysis suggested that a longer SVN-2B plus IFN β vaccination protocol might confer survival benefit. (Clinical trial registration number: UMIN 000012146).

KEYWORDS

immunotherapy, interferon beta, pancreatic cancer, peptide vaccination, survivin

1 | INTRODUCTION

Pancreatic adenocarcinoma is the sixth most common malignancy in the United States; in Japan it also has a high mortality rate, being the fourth most common cause of cancer death¹ (ganjoho.jp, https://ganjoho.jp/reg_stat/statistics/stat/summary.html). Recent advances in chemotherapy have improved overall survival (OS) among patients with pancreatic adenocarcinoma to some extent; however, mortality rates have not improved. Recently, treatment with immune checkpoint inhibitors (ICIs), including anti-programmed death-1 and anti-CTL-associated protein-4, has achieved great success in several malignancies.² However, the effectiveness of ICIs remains unsatisfactory, and severe side-effects have been reported. The ICI treatments induce antitumor immunity by canceling inhibitory effects on immunological effector CTLs, and only antigen-specific activation of CTLs can induce cancer-specific antitumor immunity. In this regard, peptide vaccination using antigenic peptides derived from tumor-associated antigens is an attractive approach for cancer immunotherapy.

Survivin belongs to the inhibitor of apoptosis protein family and is expressed in various types of malignancies, making it an attractive target molecule.³ We previously identified a human leucocyte antigen (HLA)-A24-restricted antigenic peptide (SVN-2B) from the survivin variant survivin 2B, and reported the efficacy of this peptide for several malignancies.⁴ In a phase I trial of peptide vaccination for colorectal cancer, breast cancer, bladder cancer, and oral cancer, we observed both reduction of tumor markers and tumor regression in some cases, indicating that peptide vaccination could be a promising approach for survivin-positive cancer.⁵ Furthermore, we undertook a phase I clinical trial using SVN-2B plus α -interferon (IFN α) for advanced pancreatic cancer.⁶ Although the number of cases was limited (n = 6), we observed good immune reaction for the SVN-2B peptide and a relatively good objective tumor response rate (ORR; 4 stable disease [SD] and 2 progressive disease [PD]), indicating that SVN-2B peptide vaccination with type 1 IFN could be a promising novel approach for pancreatic cancer. We therefore undertook this phase II clinical trial.

2 | MATERIALS AND METHODS

2.1 | Patients

Eligible patients had inoperable histologically diagnosed pancreatic adenocarcinoma and had been treated with at least first-line chemotherapy. Patients were required to have measurable disease, be HLA-A*24:02-positive, have histologically survivin protein-positive immunohistochemical (IHC) staining as described previously,⁷ aged from 20 to 85 years, and have an ECOG performance status of 0 or 1 and adequate bone marrow function and liver function. Patients whose cancer was refractory to gemcitabine or tegafur/gimeracil/oteracil (TS-1) and patients who were intolerant of gemcitabine and TS-1 were eligible.

2.2 | Study design and treatment

This was a multicenter, double-blinded, 3-arm randomized phase II trial of SVN-2B plus IFN β carried out at Sapporo Medical University (Sapporo, Japan), The University of Tokyo (Tokyo, Japan), and Kanagawa Cancer Center (Kanagawa, Japan) among patients with advanced pancreatic adenocarcinoma after chemotherapy (Clinical trial registration number: UMIN 000012146). The study included 2 steps. In Step 1, patients were assigned to 3 arms in a ratio of 2:2:1, as follows: Arm 1, SVN-2B + IFN β ; Arm 2, SVN-2B; Arm 3, placebo. Survivin 2B peptide solution (1 mL; 1 mg/mL) was mixed with 1 mL incomplete Freund's adjuvant (Montanide ISA-51 VG; Seppic) and emulsified, and 1.8 mL solution was injected s.c. every 2 weeks for Arm 1 and Arm 2 patients. One million units of IFN β was dissolved in 1 mL saline and injected s.c. weekly for Arm 1 patients. Survivin 2B placebo (used for Arm 3 patients) was prepared as saline mixed with incomplete Freund's adjuvant and emulsified. Interferon- β placebo (used for Arm 2 and Arm 3 patients) was prepared as 1 mL saline. In Step 1, patients were treated until diagnosed as having PD according to RECIST-PD or clinically apparent disease progression. In Step 2, all patients who consented to participate were treated with SVN-2B plus IFN β until diagnosed with immune-related response





FIGURE 1 A, Flow diagram of a randomized phase II trial of survivin 2B peptide (SVN-2B) vaccination for patients with advanced human leukocyte antigen (HLA)-A24-positive pancreatic adenocarcinoma. B, Immunohistochemistry for pancreatic carcinoma. Survivin protein and HLA class I protein expressions were assessed by immunohistochemical staining. Original magnification, 200 \times . FAS, Full Analysis Set; IFN β , interferon- β

criteria (irRC) or irPD, or for a maximum of 7 months. The primary endpoint was progression-free survival (PFS) in Step 1, defined as the time from the date of first treatment to the date of death or disease progression. Secondary endpoints were the ORR according

to RECIST[®] and immunological reaction to SVN-2B peptide in Step 1. Other study outcomes included survivin expression in tumor tissue, the lymphocyte blast transformation test, PFS and ORR according to the irRC criteria in Step 1 and Step 2,⁹ and safety.

2.3 | Assessment

The expressions of survivin and HLA class I were assessed by IHC staining using tumor tissue specimens as described previously.^{6,10} Tumor size was assessed every 6-8 weeks using computed tomography, with the first assessment occurring 8 weeks after the first treatment. The CTL response was assessed by tetramer assay and ELISpot assay before the trial, after Step 1, and after Step 2, as described previously.⁶

2.4 | Statistical analysis

We calculated that enrollment of 65 patients (in a ratio of 2:2:1) was required to achieve 73.1% power ($\alpha = 0.05$) to estimate median survival of 5 months for the SVN-2B + IFN β group (Arm 1) and 2 months for the placebo group (Arm 3), with a 7-month follow-up period and 24-month registration period, using Lakatos' formula. The sample size for this study was defined as 71, in anticipation of some withdrawals. Unfortunately, 10 patients dropped out after registration but before receiving the first treatment, leaving a total of 94 patients enrolled.

The log rank test at $\alpha = 2.5\%$ was used for PFS and survival post progression (SPP), and progression events included RECIST-PD and clinically apparent PD. Statistical analysis of the tetramer and enzyme-linked immunospot assays was intended to assess the results from patients who underwent immunological testing before treatment and after Steps 1 and 2. However, many patients unexpectedly experienced rapid worsening of performance status, including 11 patients in life-threatening condition, and 30 of the remaining 83 patients were unable to undergo immunological analysis during Step 1 (Figure 1A). Although a substantial number of patients underwent immunological testing as planned in Step 1, a very limited number of

patients completed exploratory immunological testing in Step 2, and only 10 underwent immunological assays (Figure 1A). Unexpectedly, only 1 patient from the SVN-2B monotherapy group was available for immunological analysis in Step 2; this patient was excluded, and the exploratory analysis thus included the SVN-2B + IFN β group (Arm 1; $n = 4$) and the placebo group (Arm 3; $n = 5$).

The *t* test, Dunnett's test and Kruskal-Wallis test were used to evaluate the immunological response secondary endpoint. All statistical analyses were carried out with SAS software, version 9.4.

3 | RESULTS

3.1 | Patient profiles

The 94 patients who showed survivin- and HLA class I-positive by IHC staining (Figure 1B), enrolled in this study were randomly assigned to the study arms as follows: Arm 1, SVN-2B + IFN β ($n = 36$); Arm 2, SVN-2B ($n = 38$); Arm 3, placebo ($n = 20$). Of these, 84 patients were treated between November 22, 2013 and October 27, 2016 (10 patients dropped out because of disease progression and 1 patient withdrew from the study before treatment; Figure 1A). All patients in the study died, with a median follow-up time of 101 days (range, 19-579 days). Patient characteristics are summarized in Table 1. All patients had primary and/or metastatic lesions and had received chemotherapy treatment before enrollment. Other previous treatments included surgery and radiation therapy. Thirty patients in the SVN-2B + IFN β group, 34 in the SVN-2B group, and 19 in the placebo group completed Step 1 treatment; of these, immunological response could be analyzed for 18 patients in the SVN-2B + IFN β group, 16 in the SVN-2B group, and 11 in the placebo group. Ten patients in the SVN-2B + IFN β group, 12 in the SVN-2B group, and 7 in the placebo group consented to Step 2 treatment; exploratory analysis of immunological response was undertaken for 4

TABLE 1 Summary of data for patients with advanced pancreatic adenocarcinoma treated with survivin 2B peptide vaccination, alone or in combination with interferon- β (IFN β), or placebo

Clinical variable		SVN2B + IFN β , n = 30	SVN2B, n = 34	Placebo, n = 19
Gender	Male : female	19:11	21:13	13:6
Age, years	Mean (min-max)	62.0 (39-78)	67.2 (29-82)	63.9 (45-77)
ECOG PS	0:1	21:9	24:10	14:5
Prior surgery	Negative : positive	18:12	20:14	10:9
Prior radiation therapy	Negative : positive	25:5	26:8	12:7
Prior chemotherapy	Negative : positive	0:30	0:34	0:19
Histology	Adenocarcinoma : others	28:2	32:2	17:2
Stage at registration	I/III/IVa/IVb	0:3:3:24	1:1:3:29	0:1:4:14
Target lesion, mm	Mean (min-max)	91.8 (24-226)	86.6 (15-160)	78.0 (10-156)
Primary lesion at registration	Negative : positive	10:20	14:20	9:10
Local lesions at registration	Negative : positive	20:10	26:8	14:5
Metastatic lesions at registration	Negative : positive	3:27	4:30	3:16
Local advanced disease at registration	Negative : positive	13:17	19:15	6:13

max, maximum; min, minimum; PS, performance status.

patients in the SVN-2B + IFN β group and 5 patients in the placebo group before and after administration of SVN-2B + IFN β in Step 2.

3.2 | Efficacy outcome and immune responses

During the study period, none of the enrolled patients received treatment other than peptide vaccinations. There was no statistically significant difference in PFS according to the RECIST criteria across the 3 groups (Figure 2A). In the exploratory analysis, there was also no difference across the 3 groups in OS (for the SVN-2B + IFN β group, SVN-2B group, and placebo group, median [95% confidence interval (CI)] 102 days [77-144], 96.5 days [66-131], and 111 days [63-150], respectively; $P = .4565$).

The disease control rate was evaluated according to the RECIST criteria. Stable disease was observed in 3 of 18 patients in the

SVN-2B + IFN β group, 5 of 15 in the SVN-2B group, and 3 of 11 in the placebo group. There was no significant difference in disease control rate across all groups.

Immune responses were evaluated by tetramer assay and enzyme-linked immunospot (ELISpot) assay using PBMCs from prevaccination and 8 weeks after the first vaccination. SVN-2B peptide-specific tetramer staining was increased after 8 weeks in the SVN-2B + IFN β group; however, the difference did not reach statistical significance ($P = .071$; Figure 2B). There was no significant difference between pre- and posttreatment in the SVN-2B group or the placebo group ($P = .9545$ and $P = .7947$, respectively). The ELISpot assay revealed that SVN-2B peptide-specific IFN γ spots were significantly increased after treatment in the SVN-2B + IFN β group ($P = .0245$), and IFN γ spots were increased in the SVN-2B group, but this did not reach statistical significance





FIGURE 2 A, Kaplan-Meier estimates of progression-free survival among patients with advanced human leukocyte antigen-A24-positive pancreatic adenocarcinoma. Cytotoxic T lymphocyte frequencies pretreatment and posttreatment, by tetramer assay (B) and enzyme-linked immunospot (ELISpot) assay (C). Change in CTL frequency according to time course, by tetramer assay (D) and ELISpot assay (E). IFN β , interferon- β ; SVN-2B, survivin 2B peptide

FIGURE 3 Swimmer plot of patients with advanced human leukocyte antigen-A24-positive pancreatic adenocarcinoma who completed Step 2 treatments, comprising 4 treatments with survivin 2B peptide vaccination (SVN-2B) plus interferon- β (IFN β), IFA, Incomplete Freund's adjuvant; PFS, progression-free survival; SPP, survival post progression



($P = .1323$) (Figures 2C and S1). There was no difference from pretreatment to posttreatment in the placebo group, as shown in Figure 2C ($P = .6184$). The increase in SVN-2B peptide-specific tetramer staining from before treatment to 8 weeks after initiation of Step 1 in the SVN-2B + IFN β group was not significantly larger than that in the placebo group ($P = .6670$). In the SVN-2B group, the increase in tetramer staining also was not significantly higher than that in the placebo group ($P = .5496$). In the ELISpot assay, the increases in SVN-2B peptide-specific IFN γ spots were not significantly larger in the SVN-2B + IFN β group or the SVN-2B group compared with the placebo group ($P = .0849$ and $P = .5892$, respectively).

In an exploratory analysis to investigate immunological effects during the period after Step 2 (the cases are summarized in Table S1), tetramer-positive cells were significantly increased at 8 weeks after Step 2 compared with before Step 1 ($P = .0347$) and before Step 2 ($P = .0190$) in the SVN-2B + IFN β group, whereas there was no difference in the placebo group (Figure 2D). In the ELISpot assay, there was no significant difference between 8 weeks after Step 2 and before Step 1 (Figure 2E). A very small number of SVN-2B peptide-specific IFN γ spots were observed before Step 1 in the SVN-2B + IFN β group, but some patients had a large number of spots at 8 weeks after Step 2. Swimmer plots showing SPP and PFS in each period (Step 1/Step 2) are shown in Figure 3. The median PFS for Step 2 was 69 days (range, 59-74 days). There was no significant difference in PFS between the SVN-2B + IFN β group and the placebo group (median [95% CI], 66 days [63-70] vs 70 days [59-74]; $P = .2617$). Survival post progression was significantly better in the SVN-2B + IFN β group than in the placebo group (median [95% CI], 312 days [43-460] vs 39 days [13-153]; $P = .0318$) (Figure S2).

3.3 | Safety

Forty-one adverse events occurred with SVN-2B use, and 46 adverse events occurred with IFN β use. Treatment-related adverse events, which occurred in over 5% of patients, were injection site reaction and fever. In the SVN-2B + IFN β group, SVN-2B group, and placebo group, injection site reactions were observed in 15/30 patients (50.0%), 14/34 patients (41.2%), and 8/20 patients (40.0%), respectively; fever was observed in 3/30 patients (10.0%) in the SVN-2B + IFN β group, but did not occur in the SVN-2B group or placebo group. Eighty-seven severe adverse events (over grade 3) were observed in the SVN-2B + IFN β group, SVN-2B group, and placebo group in 20 patients (41.7%), 16 patients (33.3%), and 12 patients (25.0%), respectively; events included disseminated intravascular coagulation and multiple organ dysfunction, mainly caused by disease progression and loss of PS. There were no treatment-related severe adverse events.

4 | DISCUSSION

In this study, a 3-armed double-blinded placebo-controlled randomized phase II trial of SVN-2B peptide vaccine was undertaken among patients with HLA-A24-positive pancreatic cancer. In a previous study, we carried out a phase I trial of SVN-2B peptide vaccination with IFN α in patients with HLA-A24-positive advanced pancreatic cancer ($n = 6$), and observed promising results, including a 66.7% ORR rate according to RECIST (4 SD cases and 2 PD cases).⁶ In addition, type 1 IFN (IFN α) was found to enhance the immunological (CTL) reaction to peptide vaccination. We therefore concluded type 1 IFN is beneficial for CTL induction. However,

IFN α requires a relative higher amount for CTL induction compared with IFN β , which brings higher rates of adverse effects including fever, bleeding tendency, and depression. Furthermore, IFN β has been approved for s.c. injection in melanoma cases, as we used in this clinical trial. In this regard, we decided to use IFN β in this study.

In the present study, we aimed to confirm the enhancement of peptide vaccination by type 1 IFN. By examining prevaccination vs postvaccination peripheral blood, we showed that type 1 IFN (IFN β) enhanced the immunological reaction significantly. Thus, type 1 IFN is an appropriate adjuvant for peptide vaccination therapy. In this study, we applied a 2-step peptide vaccination protocol, because the antitumor effects of cancer immunotherapy are frequently delayed after immunization.⁹ The majority of patients in the phase II study showed progression after Step 1 and shorter OS compared with the phase I study result (median, 3.48 months vs 8.08 months), suggesting that patients with very advanced disease were enrolled in this study. However, among the limited number of patients who completed Step 2 treatment, those treated with SVN-2B + IFN β showed longer survival compared with placebo-treated patients, indicating that a longer peptide vaccination protocol might be beneficial for pancreatic adenocarcinoma. The adverse effects experienced by patients in this study were limited and tolerable, including localized rash and fever; thus, SVN-2B + IFN β peptide vaccination therapy appears to be safe and is feasible for further analysis. As this study suggests that longer vaccination might be beneficial for pancreatic cancer patients, peptide vaccination at early stages or adjuvant setting for prevention of recurrence might be a better approach than for patients in advanced stages.

Recently, personalized RNA mutanome vaccines based on gene mutations found in the exome sequences of patients with cancer were shown to be feasible and safe, and to possibly bring survival benefit.^{11,12} Neoantigens developed from gene mutations are expected to be highly immunogenic¹³; however, immunological reactions to neoantigens in patients with cancer have not been high without active immunization.^{11,12} In this study, we used SVN-2B peptide derived from survivin protein. Because survivin is an over-expressed antigen, the immunological reaction to SVN-2B should be low without active immunization. Thus, type 1 interferon could be a promising candidate adjuvant for peptide vaccination therapy.

Recent analyses of patients treated with immune checkpoint inhibitors have suggested that high tumor mutation burden is a predictor of better prognosis,¹⁴ and neoantigen expression is related to long-term survival in pancreatic cancer.¹⁵ Although there is no direct evidence that neoantigens derived from tumor mutations are related to antipancreatic cancer immunity, these observations indicate that gene mutations are related to antitumor immune reactions. Because pancreatic cancers are known to contain relatively few gene mutations,¹⁶ neoantigens for application in pancreatic cancer immunotherapy might be limited. Expression of survivin, which is expressed in pancreatic adenocarcinoma cells at a high rate (88%), is related to proliferation index, and high survivin expression is related to poorer prognosis, indicating that survivin is a reasonable target for pancreatic cancer immunotherapy.¹⁷

In summary, we show that SVN-2B + IFN β peptide vaccination therapy is safe and feasible for patients with pancreatic cancer, and IFN β is a promising adjuvant for peptide vaccination therapy. Longer treatments with SVN-2B + IFN β could be beneficial.

ACKNOWLEDGMENTS

This work was supported by a Grant-in-Aid for Scientific Research (B) from the Japan Society for the Promotion of Science (JSPS) for Y. Hirohashi (Grant Number 15H04722) and the Project for Cancer Research and Therapeutic Evolution (P-CREATE) from the Japan Agency for Medical Research and Development (AMED) for T. Torigoe.

DISCLOSURE

The authors have no conflict of interest in regard to this study.

ORCID

Yoshihiko Hirohashi  <https://orcid.org/0000-0002-0608-3914>

Tomohide Tsukahara  <https://orcid.org/0000-0002-3678-4359>

REFERENCES

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2017. *CA Cancer J Clin.* 2017;67:7-30.
2. Ribas A, Wolchok JD. Cancer immunotherapy using checkpoint blockade. *Science.* 2018;359:1350-1355.
3. Altieri DC. Validating survivin as a cancer therapeutic target. *Nat Rev Cancer.* 2003;3:46-54.
4. Hirohashi Y, Torigoe T, Maeda A, et al. An HLA-A24-restricted cytotoxic T lymphocyte epitope of a tumor-associated protein, survivin. *Clin Cancer Res.* 2002;8:1731-1739.
5. Tsuruma T, Hata F, Torigoe T, et al. Phase I clinical study of anti-apoptosis protein, survivin-derived peptide vaccine therapy for patients with advanced or recurrent colorectal cancer. *J Transl Med.* 2004;2:19.
6. Kameshima H, Tsuruma T, Kutomi G, et al. Immunotherapeutic benefit of alpha-interferon (IFN α) in survivin2B-derived peptide vaccination for advanced pancreatic cancer patients. *Cancer Sci.* 2013;104:124-129.
7. Kameshima H, Tsuruma T, Torigoe T, et al. Immunogenic enhancement and clinical effect by type-I interferon of anti-apoptotic protein, survivin-derived peptide vaccine, in advanced colorectal cancer patients. *Cancer Sci.* 2011;102:1181-1187.
8. Therasse P, Arbuck SG, Eisenhauer EA, et al. New guidelines to evaluate the response to treatment in solid tumors. *J Natl Cancer Inst.* 2000;92:205-216.
9. Wolchok JD, Hoos A, O'Day S, et al. Guidelines for the evaluation of immune therapy activity in solid tumors: immune-related response criteria. *Clin Cancer Res.* 2009;15:7412-7420.
10. Torigoe T, Asanuma H, Nakazawa E, et al. Establishment of a monoclonal anti-pan HLA class I antibody suitable for immunostaining of formalin-fixed tissue: unusually high frequency of down-regulation in breast cancer tissues. *Pathol Int.* 2012;62:303-308.
11. Sahin U, Derhovannessian E, Miller M, et al. Personalized RNA mutanome vaccines mobilize poly-specific therapeutic immunity against cancer. *Nature.* 2017;547:222-226.

12. Ott PA, Hu Z, Keskin DB, et al. An immunogenic personal neoantigen vaccine for patients with melanoma. *Nature*. 2017;547:217-221.
13. Coulie PG, Van den Eynde BJ, van der Bruggen P, Boon T. Tumour antigens recognized by T lymphocytes: at the core of cancer immunotherapy. *Nat Rev Cancer*. 2014;14:135-146.
14. Goodman AM, Kato S, Bazhenova L, et al. Tumor mutational burden as an independent predictor of response to immunotherapy in diverse cancers. *Mol Cancer Ther*. 2017;16:2598-2608.
15. Balachandran VP, Luksza M, Zhao JN, et al. Identification of unique neoantigen qualities in long-term survivors of pancreatic cancer. *Nature*. 2017;551:512-516.
16. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA Jr, Kinzler KW. Cancer genome landscapes. *Science*. 2013;339:1546-1558.
17. Sarela AI, Verbeke CS, Ramsdale J, Davies CL, Markham AF, Guillou PJ. Expression of survivin, a novel inhibitor of apoptosis and cell

cycle regulatory protein, in pancreatic adenocarcinoma. *Br J Cancer*. 2002;86:886-892.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Shima H, Tsurita G, Wada S, et al. Randomized phase II trial of survivin 2B peptide vaccination for patients with HLA-A24-positive pancreatic adenocarcinoma. *Cancer Sci*. 2019;110:2378-2385. <https://doi.org/10.1111/cas.14106>