

SCIENTIFIC REPORTS



OPEN

Susceptible and mCry3A resistant corn rootworm larvae killed by a non-hemolytic *Bacillus thuringiensis* Cyt1Aa mutant

Alejandra Bravo¹ , Jazmin A. López-Díaz¹, Takashi Yamamoto², Kathleen Harding², Jian-Zhou Zhao³, Gretel Mendoza¹, Janette Onofre¹, Mary-Carmen Torres-Quintero¹, Mark E. Nelson³, Gusui Wu², Amit Sethi³ & Mario Soberón¹

The western corn rootworm (WCR) *Diabrotica virgifera virgifera* causes substantial damage in corn. Genetically modified (GM) plants expressing some *Bacillus thuringiensis* (Bt) insecticidal Cry proteins efficiently controlled this pest. However, changes in WCR susceptibility to these Bt traits have evolved and identification of insecticidal proteins with different modes of action against WCR is necessary. We show here for the first time that Cyt1Aa from Bt exhibits toxicity against WCR besides to the dipteran *Aedes aegypti* larvae. Cyt1Aa is a pore-forming toxin that shows no cross-resistance with mosquitoicidal Cry toxins. We characterized different mutations in helix α -A from Cyt1Aa. Two mutants (A61C and A59C) exhibited reduced or absent hemolytic activity but retained toxicity to *A. aegypti* larvae, suggesting that insecticidal and hemolytic activities of Cyt1Aa are independent activities. These mutants were still able to form oligomers in synthetic lipid vesicles and to synergize Cry11Aa toxicity. Remarkably, mutant A61C showed a five-fold increase insecticidal activity against mosquito and almost 11-fold higher activity against WCR. Cyt1Aa A61C mutant was as potent in killing WCR that were selected for resistance to mCry3A as it was against unselected WCR indicating that this toxin could be a useful resistance management option in the control of WCR.

The western corn rootworm (WCR) *Diabrotica virgifera virgifera* is an insect pest that attacks cornfields eating the roots of the plants causing extensive damage. It is principally found in North America and Europe resulting in grower-losses in excess of one billion dollars per year¹. In addition, this pest is highly adaptable and has evolved resistance to chemical insecticides². Genetic modified (GM) plants expressing toxins from *Bacillus thuringiensis* (Bt) such as Cry3Bb1, mCry3A, eCry3.1Ab and Cry34/35Ab1 have been used in the control of this pest in cornfields^{3–5}. However, changes in WCR susceptibility to these traits have evolved in some areas potentially compromising the effectiveness of GM plants^{6,7}. In addition, resistance to Cry3Bb showed cross-resistance to mCry3A and eCry3.1Ab toxins further highlighting the challenges facing Bt-corn technology in the control of this pest^{8,9}.

WCR resistance to Bt-corn has encouraged researchers to look for additional strategies and novel proteins that could be used to control this challenging pest. Recently new proteins that kill WCR have been identified from non-Bt bacteria such as the small IPD072Aa protein from *Pseudomonas chlororaphis* or the PIP-47Aa protein from *P. mosselii*^{10,11}. These novel toxins do not share receptors with Cry3A or Cry34/35Ab1 and neither showed cross-resistance in Cry3A-selected insects^{10,11}, while IPD072Aa also showed no cross-resistance to Cry34/35Ab1-selected insects¹⁰. In addition a binary toxin named AfIP-1A/1B isolated from the bacteria *Alcaligenes faecalis* was shown to be toxic against WCR although it shares a similar mode of action as the Cry34/35Ab1 toxin since WCR population resistant to Cry34/35Ab1 showed cross-resistance to AfIP-1A/1B¹².

Here we show that Cyt1Aa, previously characterized as a mosquitoicidal toxin, display moderate toxicity against WCR. Under sporulation conditions Bt produces two main types of insecticidal toxins named Cry or Cyt, which are toxic to different insect orders¹³. The Cyt toxins are composed of a single α - β domain with seven to eight β -strands wrapped by α -helices^{14–17}. Cyt1Aa toxin is mostly active against dipteran larvae¹⁸ and it is found

¹Instituto de Biotecnología, Universidad Nacional Autónoma de México. Apdo. Postal 510-3, Cuernavaca, 62250, Morelos, Mexico. ²DuPont Pioneer, Hayward, CA, 94545, USA. ³DuPont Pioneer, Johnston, IA, 50131, USA. Correspondence and requests for materials should be addressed to A.B. (email: bravo@ibt.unam.mx)




Figure 1. SDS-PAGE analysis of the Cyt1A wild type and mutant proteins produced in *B. thuringiensis*. Panel A, Prototoxins solubilized from purified crystal inclusions. Panel B, trypsin activated toxins. All samples were boiled 5 min before loading into the SDS-PAGE and the gel was stained with coomassie brilliant blue.

in *Bt* subspecies *israelensis* strain along with the mosquitocidal Cry4Aa, Cry4Ba and Cry11Aa toxins. Besides its insecticidal activity, Cyt1Aa toxin has cytolytic activity against red blood cells¹⁹ and this hemolytic activity could limit its applications in GM products. The Cyt proteins form high molecular weight oligomers, which insert into the membrane of the midgut targeted-cells forming lytic pores^{20–23}. It has been proposed that β 5– β 7 region of Cyt1Aa is likely involved in Cyt1Aa membrane insertion while helices α -A and α -C are involved in Cyt1Aa oligomerization and recognition of membrane lipids^{14,23–26}. It was shown that oligomerization of Cyt1Aa and Cyt2Aa toxins is a key step in membrane binding and pore formation since certain mutations in helix α -C affected oligomerization and membrane insertion, reducing their toxicities against *Aedes aegypti* larvae as well as their hemolytic activities^{23,27}.

One of the most interesting features of Cyt1Aa is its capacity to synergize the toxicity of different Cry toxins such as Cry11Aa or Cry4Ba^{28–30}. Moreover Cyt1Aa is able to overcome resistance of *Culex quinquefasciatus* to Cry4Aa, Cry4Ba and Cry11Aa toxins³¹. It has been proposed that Cyt1Aa functions as a receptor of Cry11Aa since binding of this toxin to Cyt1Aa facilitates oligomer formation of Cry11Aa and its membrane insertion^{29,32}.

To determine the role of Cyt1Aa helix α -A in the mode of action of this toxin, different residues were mutated and these mutants were analyzed for oligomerization, synergism with Cry11Aa, as well for the hemolytic and insecticidal activities against dipteran and coleopteran larvae. Our results show that two mutations located in helix α -A of Cyt1Aa reduced hemolysis of red blood cells, but did not alter oligomerization, synergism with Cry11Aa, or toxicity against *A. aegypti* or WCR larvae. Moreover, one of these mutants showed a significant increase in its insecticidal potency against both mosquitoes and WCR larvae and also displays toxicity against the WCR population resistant to mCry3A. These data indicate that this non-hemolytic Cyt1Aa toxin with increased toxicity to WCR might have the potential for efficient WCR control in transgenic corn.

Results

Insecticidal activity of Cyt1Aa against the coleopteran pest. Previous report showed the potential toxicity of Cyt1Aa against coleopteran species³⁴. The toxicity of Cyt1Aa was evaluated against WCR larvae in bioassays as described in Methods. The evaluation of the mean growth inhibition concentration (IC_{50}) value was performed with solubilized prototoxin proteins (Fig. 1A) that were obtained from crystal inclusions purified by sucrose gradients as reported in Methods. The control included the same volume of solubilization buffer. These bioassays showed that Cyt1Aa has an IC_{50} of 335 μ g per cm^2 of diet against neonate WCR larvae.

Isolation of Cyt1Aa α -A mutants. To determine the potential role of Cyt1Aa helix α -A (⁴⁹PNYILQ AIMLANAFQNAL⁶⁶) in Cyt1Aa mode of action, the amino acid residues L58, A59, A61 and F62, located in the hydrophobic phase of the helix, were mutated as described in Materials and Methods. The plasmids containing the Cyt1Aa mutant genes were transformed into acrySTALLIFEROUS Bt407 strain and protein expression was analyzed under sporulation condition. Mutants A59E and F62R did not produce the corresponding prototoxin, while L58E mutant was produced at much lower levels compared to Cyt1Aa. However, after solubilization of protein

Toxin	LC ₅₀ ^a (ng/ml) against <i>Aedes aegypti</i>	IC ₅₀ ^b (μg/cm ²) against control WCR
Cyt1Aa	1100 (880–1480) ^c	335.6 (284.8–399.2)
A59C	2419 (1861–3653)	63.2 (38.8–96.5)
A61C	212 (131–273)	29.2 (24.8–35.1)
Cry11Aa	669 (476–994)	ND ^d

Table 1. Insecticidal activity of Cyt1Aa mutants against 4th instar *Aedes aegypti* larvae and neonate WCR larvae. ^aMean lethal concentration value. ^bMean growth inhibition concentration value. ^c95% confidential limits calculated by Probit statistical analysis within the parenthesis. ^dNot determined.




Figure 2. Hemolytic activity of Cyt1A wild type and mutant proteins assayed in Rabbit red blood cells. Positive control showing 100 percent hemolysis was defined after incubation of the same volume of rabbit red blood cells with dechlorinated H₂O. Negative controls were red blood cells incubated with buffer A. These assays were performed three times in triplicate each time. Standard deviations are shown in the figure.

crystals by alkaline treatment, L58E protein was not soluble (data not shown). Therefore A59E, F62R and L58E mutants were not further analyzed. In contrast A59C and A61C mutants were produced as 27 kDa proteins upon sporulation (Fig. 1A). These proteins were solubilized and treated with trypsin for toxin activation. Figure 1B shows that, as Cyt1Aa, A59C and A61C yielded a 22 kDa activated protein after trypsin treatment indicating no major structural changes caused by these mutations.

Insecticidal and hemolytic activities of Cyt1Aa α -A mutants. Table 1 shows the mean lethal concentration (LC₅₀) values of the toxicity data against *A. aegypti* and the mean growth inhibition concentration (IC₅₀) values in WCR. The Cyt1Aa A59C mutant display two-fold lower toxicity against *A. aegypti* compared to Cyt1Aa, while A61C showed five-fold higher toxicity to this mosquito larvae (Table 1). Bioassay data against WCR showed that A59C displayed five-fold higher toxicity to WCR while A61C mutant showed 11-fold higher toxicity against WCR. The wild type Cyt1Aa and A59C mutant were not active enough against the mCry3Aa resistant WCR³³ population to evaluate cross-resistance accurately (data not shown). In contrast, Cyt1Aa A61C mutant showed sufficient activity against the mCry3A-resistant WCR (IC₅₀ = 131 μg/cm² diet, with 95% CL of 68.5–1149 μg/cm² diet) to reveal no significant cross-resistance (RR = 2) when compared to the activity against susceptible WCR (IC₅₀ = 65.5 μg/cm² diet, with 95% CL of 41.59–127.5 μg/cm² diet) when tested at the same time.

We determined the hemolytic activity of Cyt1Aa and helix α -A mutants. Rabbit red blood cells were incubated with increasing concentrations of trypsin-activated toxins. Figure 2 shows that both α -A mutants were severely affected in hemolysis since wild type Cyt1Aa toxin showed a fifty percent effective dose (ED₅₀) of 250 ng/ml while mutant A61C lysed only 30% of the red blood cells with 1000 ng/ml. A59C exhibited no hemolytic activity at the highest toxin concentration tested (Fig. 2).

The capacity of Cyt1Aa and α -A mutants to synergize Cry11Aa toxicity to *A. aegypti* larvae was determined. As described in Methods we prepared mixtures of Cyt1Aa and Cry11Aa that would give a toxicity of 20% based on their corresponding LC₅₀ toxicity values. Table 2 shows that A59C and A61C synergize the activity of Cry11Aa since the toxicity of the protein mixtures showed a three- to four-fold higher toxicity than the expected mortality.

Effect of Cyt1Aa A59C and A61C on toxin oligomerization. To determine the effect of A59C and A61C mutations on Cyt1Aa oligomerization, soluble protoxins of Cyt1Aa and of the two α -A mutants were incubated with small unilamellar vesicles (SUV) and trypsin. The vesicles were collected by centrifugation and analyzed by western blot using anti-Cyt1Aa antibody as described in Materials and Methods. As negative control we

Toxin	$S_{(toxin)OBS}^a = \frac{Rep1 + Rep2 + Rep3}{n}$	$S_{(Cyt1Aa, Cry11Aa)EXP}^b = \frac{S_{(Cyt1Aa, Cry11Aa)OBS}}{S_{(Cyt1Aa)OBS} \times S_{(Cry11Aa)OBS}}$	Expected mortality ^c = $(1 - S_{(Cyt1Aa, Cry11Aa)EXP}) \times 100\%$	Observed mortality ^d <i>Cyt1Aa+Cry11Aa</i>
Cyt1Aa	1.00	0.80	20%	90 ± 10%
A59C	1.00	0.80	20%	57 ± 20%
A61C	0.93	0.75	25.3%	83 ± 15%
Cry11Aa	0.80			

Table 2. Analysis of synergism of Cyt1Aa or mutant toxins with Cry11Aa toxin in 4th instar *Aedes aegypti* larvae. ^aObserved survival of individual toxin $S_{(toxin)OBS}$ corresponds to the observed proportion of larvae that survived to the exposure to Cyt1Aa or mutant toxins. Observed mortality was 20% with Cry11Aa at 200 ng per ml and 0% with Cyt1Aa at 75 ng Cyt1Aa per ml. n = 30 larvae for each toxin tested.




Figure 3. Oligomerization of Cyt1Aa and mutant proteins after activation of solubilized protoxin, with trypsin in the presence of SUV liposomes. Samples were heated 3 min at 65 °C before loading into the SDS-PAGE transferred to PVDF and revealed in western blot assay as described in materials and methods using polyclonal anti-Cyt1A antibody and goat anti-rabbit antibody coupled to horseradish peroxidase.

included two Cyt1Aa mutants located in helix α -C that were previously characterized to be affected in oligomerization (V122E and V126E) and two helix α -C mutants that were not affected in oligomerization (L120K and L123K)²³. Figure 3 shows that Cyt1Aa and A59C and A61C mutants produced high molecular weight oligomers after protease activation in the presence of synthetic membranes. This was similar to the previously characterized α -C mutants L120K or L123K that were not affected in oligomer formation²³. In contrast, helix α -C mutants V122E and V126E did not form oligomers as previously reported²³ (Fig. 3A). These results show that Cyt1Aa A59C and A61C mutations did not affect toxin oligomerization.

Discussion

Cyt1Aa toxin shows a specific insecticidal activity mainly to dipteran insects. However, previous reports showed that Cyt1Aa could have coleopteran activity since it was able to kill the cottonwood leaf beetle, *Chrysomela scripta*³⁴. It was also shown that Cyt1Aa exhibits low toxicity against homopteran pests such as the pea aphid *Acyrtosiphon pisum*³⁵. Here we show that Cyt1Aa has insecticidal activity against the agriculturally important coleopteran pest WCR, *Diabrotica virgifera virgifera* (Le Conte). In addition, Cyt1Aa shows cytolytic activity against red blood cells¹⁹.

It was previously suggested that Cyt1Aa helices α -A and α -C were involved in the initial binding of Cyt1Aa to membrane lipids and in toxin oligomerization³⁶. In agreement with this, it was later shown that certain Cyt1Aa helix α -C mutations affected toxin oligomerization that correlated with a severe effect in its insecticidal and hemolytic activities but did not affect its synergism with Cry11Aa toxin²³.

Here we show that Cyt1Aa mutations in helix α -A reveal amino acids (A59C and A61C) that can differentiate the lytic effects of Cyt1Aa on rabbit red blood cells from the insecticidal activity against WCR since cysteine

Oligonucleotide	DNA sequence
A59C	TTG CAA GCA ATT ATG TTA <u>TGT</u> AAT GCC TTT CAA AAT GC
A61C	GCA AGC AAT TAT GTT AGC AAA <u>CTG</u> TTT TCA AAA TGC ATT AGT TCC C
L58E	TAT ATA TTG CAA GCA ATT ATG <u>GAA</u> GCA AAT GCG TTT CAA AAT GC
A59E	TAT ATT GCA AGC AAT TAT GTT <u>AGA</u> AAA TGC GTT TCA AAA TGC
F62R	AGC AAT TAT GTT AGC AAA TGC <u>ACG</u> GCA AAA TGC GTT AGT TCC
IRE1d	TGT GAA TTC ATG GAA AAT TTA AAT CAT TG
IRE4r	CTA CTC GAG GAG GGT TCC ATT AAT AGC

Table 3. Sequence of mutagenic oligonucleotides. The sequence of mutagenic codon is underlined and labeled with bold letters.

replacement of these amino acids resulted in significant reduction of Cyt1Aa hemolytic activity and a significant increase in the toxicity to WCR compared to the wild type Cyt1Aa.

The Cyt1Aa mutations A59C or A61C did not affect oligomerization of the toxin with exposure to synthetic SUV. In the case of the closely related Cyt2Aa toxin, a similar mutation at A61C located also in helix α -A was affected in oligomer formation with exposure to SUV although it showed a similar toxicity to mosquito larvae²⁷. We do not know the reason for this difference but different experimental conditions in the oligomerization assays might explain it. In the case of Cyt2Aa, liposomes were prepared with a lipid mixture of phosphatidylcholine: cholesterol: stearylamine in a molar ratio of 4:3:1²⁷, while we prepared liposomes with the same lipid mixture but in a molar ratio of 10:3:1. Also, Cyt2Aa activated toxin was incubated with the liposomes during the oligomerization assay²⁷, while we co-incubated Cyt1Aa protoxin along with trypsin (for toxin activation) in the presence of the liposomes to induce Cyt1Aa-oligomerization.

Cyt1Aa has a distinct mode of action than Cry toxins since it does not rely on protein receptors for killing mosquitoes. It has been shown that *Culex quinquefasciatus* insects resistant to Cry1Aa showed no cross-resistance to Cyt1Aa³⁷ and that Cyt1Aa can overcome the resistance of the mosquito larvae to Cry4Aa, Cry4Ba and Cry11Aa toxins³¹. These data suggest that Cyt1Aa toxin could represent an interesting alternative for the control of other insect pests.

Our results show that helix α -A is involved in Cyt1Aa toxin specificity. The α -A helix A61C and A59C mutations both resulted in more toxic proteins against WCR while only the A61C mutation was significantly more toxic to *A. aegypti* than Cyt1Aa. The A61C mutant also showed insecticidal activity against the mCry3A resistant population of WCR. Importantly, both Cyt1Aa α -A mutants had reduced hemolytic activity with A59C exhibiting no hemolytic activity, improving its potential for use in transgenic corn for protection from damage caused by WCR. These data indicate that the α -A helix of Cyt1Aa plays an important role in hemolytic activity and that its insecticidal activity is independent of its hemolytic activity. These novel Cyt1Aa mutant toxins offer potential as additional alternatives to currently utilized Bt toxins for the effective control of WCR.

Materials and Methods

Production of Cyt1Aa and Cry11Aa Proteins. Production of Cyt1Aa and Cry11Aa proteins was done as described previously²³. The Cyt1Aa or Cry11Aa protoxins were produced in *B. thuringiensis* 407 acrySTALLIFEROUS strain transformed with plasmid pWF45³⁸ or pCG6³⁹. Cyt1Aa mutants were expressed also in the *B. thuringiensis* 407 acrySTALLIFEROUS strain. These Bt strains were grown in solid nutrient broth sporulation medium⁴⁰ (4 days, 30 °C), these media were supplemented with 10 μ g/ml erythromycin for Cyt1Aa or 25 μ g/ml erythromycin for Cry11Aa. Spores and crystals were washed with 0.3 M NaCl, 0.01 M EDTA, pH 8.0, by centrifugation (10 min, 10,000 rpm, 4 °C), washing steps were done three times. Pellets were stored at -20 °C until used. The crystal/spore pellet was suspended in 0.05% Triton X-100, 300 mM NaCl, 20 mM Tris-HCl pH 7.2, sonicated two times 1 min and the inclusion bodies were purified by sucrose gradient centrifugation as reported⁴¹. Cyt1Aa proteins were solubilized 1 h at 37 °C in 50 mM Na₂CO₃, 10 mM DTT, pH 10.5, agitation at 350 rpm and centrifuged for 10 min at 10,000 rpm 4 °C. The soluble protoxins were recovered in the supernatant. Protein concentrations were determined by the Bradford assay. Finally, Cyt1Aa protoxin was activated with trypsin 1:20 (Trypsin: Cyt1Aa) ratio (Sigma-Aldrich Co., St Louis, MO) w/w for 2 h at 30 °C. Supplementary Figs S1 and S2 show additional images of SDS-PAGE analysis of Cyt1Aa, A59C and A61C proteins purified from *B. thuringiensis*.

Site-Directed Mutagenesis of Cyt1Aa Toxin. Mutagenesis of Cyt1Aa toxin was performed in pWF45 plasmid by using QuikChange XL Site-Directed kit (Stratagene La Jolla, CA) as previously described²³. Table 3 shows the information of mutagenic oligonucleotides that were synthesized by Sigma-Aldrich (St Louis, MO). Mutated plasmids were transformed in *E. coli* X-L1 blue strain and transformants were selected in LB Ampicillin 100 μ g/ml at 25 °C. Plasmid DNA was purified using a DNA extraction kit (Qiagen, Hilden, GER) and sequenced at the Institute of Biotechnology UNAM. Finally plasmids were transformed into Bt 407 strain and transformants were selected in LB erythromycin 10 μ g/ml at 30 °C. We confirmed the sequence of these mutants by PCR amplification using IRE1d-IRE4r oligonucleotides (Table 3).

Insect bioassays. The *Aedes aegypti* mosquitoes used in this work were reared at Instituto de Biología facilities, at 28 °C, 75% humidity and a 12 h: 12 h light: dark photoperiod. Mosquitocidal bioassays were performed as described before²³. We used 10 early 4th-instar larvae in 100 ml of dechlorinated water per doses. We used ten different concentrations from 50 to 10000 ng/ml of spore/crystal suspensions of each Cyt1Aa protein, wild type

or mutants. These samples were previously sonicated (1 min) using an ultrasonic processor (Cole-Palmer) and diluted into the 100 ml water containers that will be used in the bioassay. The negative control was dechlorinated water. Larvae viability was examined after 24 h. Probit analyses were used to determine the mean lethal concentration (LC_{50}) using statistical parameters of the data in triplicate performed in three independent assays (Polo-Plus LeOra Software).

Susceptible WCR and mCry3A resistant WCR (laboratory selected³³) populations were reared at DuPont Pioneer, Johnston, IA headquarters. Bioassays were performed in 96-well plates. Isolated Cyt1Aa crystals were solubilized for 1 h at room temperature in 100 mM sodium bicarbonate buffer supplemented with 20 mM DTT, pH 10. Samples were centrifuged at 13,000 rpm 30 min and the supernatant was collected. The supernatant was concentrated up to 1 ml using Ultra 15 Amicon filters. Serial 2X dilutions of the solubilized protoxins were prepared with sodium bicarbonate buffer and 20 μ l of each protoxin dilution was overlaid on 50 μ l diet in 96-well plates. Three to five neonate susceptible WCR larvae were placed in each well and the assay plates were incubated at 25 °C for four days. For assessing cross-resistance three to five 24 h-born mCry3Aa resistant WCR larvae, that were fed 24 h on diet, were placed in each well and the assay plates were incubated at 25 °C for six days. Sodium bicarbonate buffer was used as a negative control. The growth inhibition concentration (IC_{50}) defined as the concentration of protein that resulted in 50% reduction in larval development when compared to larvae feeding on diet alone was determined using Microsoft Excel from triplicate assays.

Synergism assay. Synergism was evaluated as previously described⁴² by testing for deviation from the null hypothesis of simple independent action of the toxins. This hypothesis assumes that the surviving larvae to the exposure of mixture of toxins are the product of the surviving larvae to each toxin separately. The formula $S_{(ab)EXP} = S_{(a)OBS} \times S_{(b)OBS}$ ⁴² was used where $S_{(ab)EXP}$ is the proportion of larvae expected to survive to the mixture of toxins a and b , $S_{(a)OBS}$ and $S_{(b)OBS}$ are the observed proportion of larvae that survived to toxin a or toxin b . We used 30 larvae per toxin or toxins mixtures. The expected mortality to the mixture of toxins a and b was obtained by using $(1 - S_{(ab)EXP}) \times 100\%$ and the expected numbers of dead and live larvae were obtained by multiplying the expected mortality and survival rates by the sample size used with each toxin. All synergism assays were done in triplicate and Fisher's exact test was used to determine the significant differences among observed and expected mortality data.

Hemolysis assays. Hemolytic assays were done as previously described²². We used red blood cells from Rabbits which were washed using buffer A (0.1 M dextrose, 0.07 M NaCl, 0.02 M sodium citrate, 0.002 M citrate, pH 7.4) three times and then diluted to 2×10^8 cells/ml in the same buffer. The reaction was performed in a final volume of 0.2 ml in 96 wells microtiter plates, which contains 20 μ l of red blood cells and different concentrations of Cyt1Aa toxin (20–1000 ng) in the same buffer. Samples were incubated at 37 °C for 30 min, after that time were centrifuged (2,500-x g, 5 min, 4 °C), the supernatants were collected and absorbance at 405 nm was determined. Positive control was rabbit red blood cells with dechlorinated H₂O showing 100% hemolysis. Negative control was red blood cells with buffer A. All assays were done three times in triplicate and t -test analysis were performed using GraphPad Prism statistical program.

Preparation of Small Unilamellar Vesicles (SUV). SUV were prepared as previously reported²³. A mixture of lipids egg-yolk phosphatidyl choline (PC), cholesterol (Ch) (Avanti Polar Lipids, Alabaster, AL) and stearylamine (S) (Sigma-Aldrich, St Louis, MO) at 10:3:1 proportion, was used. The final concentration of the total lipid mixture was 0.65 μ mol. The mixture was dried by nitrogen flow, stored overnight under vacuum to remove residual chloroform and finally hydrated in 0.65 ml of 10 mM CHES, 150 mM KCl pH 9 by a 30 min incubation followed by vortex. The lipid suspension was sonicated three to five times during 20 sec each in a Branson-1200 bath sonicator (Danbury, CT). SUV were not stored, they were used the same day upon their preparation.

Oligomerization of Cyt1Aa and mutant toxins. Oligomerization of Cyt1Aa and mutants was performed as previously described²³. We incubate 200 ng of Cyt1Aa or Cyt1Aa mutants solubilized protoxin with 90 μ l SUV and 10 ng of trypsin in a final volume of 100 μ l (2 h, 30 °C, 350 rpm). One mM PMSF (final concentration) was added to stop the reaction. The membrane pellet was separated from the supernatant by centrifugation (30 min, 55,000 rpm). Samples were heated at 65 °C for 3 min, and loaded in SDS-PAGE gels. Proteins were transferred to PVDF Immobilon-P Millipore membranes using a wet chamber (12 h, 150 mA, 4 °C) and revealed by western blot assay as described²³, using 5% skimmed milk in PBS for blocking (1 h, room temperature), and two washes (5 min each) with PBS containing 0.1% Tween 20 (PBS-Tween). The PVDF was incubated 1 h, room temperature with polyclonal anti-Cyt1A antibody (1:30,000 dilution in PBS-Tween) washed twice with PBS-Tween (5 min each) and incubated with goat anti-rabbit antibody coupled to horseradish peroxidase (Santa Cruz Biotechnology, Dallas, TX) (1:10000 dilution in PBS-Tween). The peroxidase signal was visualized with SuperSignal chemiluminescent substrate (ECL; Amersham Pharmacia Biotech). Oligomerization assays were done three times using different samples of the Cyt1Aa or mutant toxins. Molecular weight markers were Precision Plus Protein Standards All Blue (Bio-Rad) and molecular masses are indicated in kDa. Supplementary Fig. S3 shows additional images of the oligomerization analysis of CytAa, A59C and A61C proteins.

Data Availability

All data generated during this study are included in this published article.

References

- Gray, M. E. *et al.* Adaptation and Invasiveness of western corn rootworm: intensifying research on a worsening pest. *Annu. Rev. Entomol.* **54**, 303–321 (2009).
- Zhu, K. Y. *et al.* Evidence of evolving carbamyl resistance in western corn rootworm (*Coleoptera:Chrysomelidae*) in area wide-managed cornfields in north central Kansas. *J. Econ. Entomol.* **94**, 929–934 (2001).
- Moellenbeck, D. J. *et al.* Insecticidal proteins from *Bacillus thuringiensis* protect corn from corn rootworms. *Nat. Biotechnol.* **19**, 668–672 (2001).
- Rice, M. E. Transgenic rootworm corn: Assessing potential agronomic, economic, and environmental benefits. *Plant Health Prog.* **10**, 1094/PHP-2004-0301-01-RV (2004).
- Narva, K. E., Siegfried, B. D. & Storer, N. P. Transgenic approaches to western corn rootworm control. *Adv. Biochem. Eng. Biotechnol.* **136**, 135–162 (2013).
- Gassmann, A. J., Petzold-Maxwell, J. L., Keweshan, R. S. & Dunbar, M. W. Field-evolved resistance to Bt maize by western corn rootworm. *PLoS ONE* **6**, e22629 (2011).
- Gassmann, A. J. *et al.* Evidence of resistance to Cry34/35Ab1 corn by western worn rootworm (*Coleoptera: Chrysomelidae*): Root injury in the field and larval survival in plant-based bioassays. *J. Econ. Entomol.* **109**, 1872–1880 (2016).
- Gassmann, A. J. *et al.* Field-evolved resistance by western corn rootworm to multiple *Bacillus thuringiensis* toxins in transgenic maize. *Proc. Natl. Acad. Sci. USA* **111**, 5141–5146 (2014).
- Jakka, S. R. K., Shrestha, R. B. & Gassmann, A. J. Broad-spectrum resistance to *Bacillus thuringiensis* toxins by western corn rootworm (*Diabrotica virgifera virgifera*). *Sci. Rep.* **6**, 27860 (2016).
- Schellenberger, U. *et al.* A selective insecticidal protein from *Pseudomonas* for controlling corn rootworms. *Science* **354**, 634–637 (2016).
- Wei, J.-Z. *et al.* A selective insecticidal protein from *Pseudomonas mosselii* for corn rootworm control. *Plant Biotech. J.* Accepted Manuscript, <https://doi.org/10.1111/pbi.12806> (2017).
- Yalpani, N. *et al.* An *Alicigenes* strain emulates *Bacillus thuringiensis* producing a binary protein that kills corn rootworm through a mechanism similar to Cry34Ab1/35AB1. *Sci. Rep.* **7**, 3063 (2017).
- Pardo-López, L., Soberón, M. & Bravo, A. *Bacillus thuringiensis* insecticidal 3-domain Cry toxins: Mode of action, insect resistance and consequences for crop protection. *FEMS Microbiol. Rev.* **37**, 3–22 (2013).
- Cohen, S. *et al.* Cyt1Aa toxin: crystal structure reveals implications for its membrane-perforation function. *J. Mol. Biol.* **413**, 804–814 (2011).
- Li, J., Koni, P. A. & Ellar, D. J. Structure of the mosquitocidal delta-endotoxin CytB from *Bacillus thuringiensis* sp. *kyushuensis* and implications for membrane pore formation. *J. Mol. Biol.* **257**, 129–152 (1996).
- Cohen, S. *et al.* High-resolution crystal structure of activated Cyt2Ba monomer from *Bacillus thuringiensis* subsp. *israelensis*. *J. Mol. Biol.* **380**, 820–827 (2008).
- Soberón, M., López-Díaz, J. A. & Bravo, A. Cyt toxins produced by *Bacillus thuringiensis*: A protein fold conserved in several pathogenic microorganisms. *Peptides* **41**, 87–93 (2013).
- Koni, P. A. & Ellar, D. J. Biochemical characterization of *Bacillus thuringiensis* cytolytic delta-endotoxins. *Microbiology* **140**, 1869–1880 (1994).
- Knowles, B. H., White, P. J., Nicholls, C. N. & Ellar, D. J. A broad-spectrum cytolytic toxin from *Bacillus thuringiensis* var. *kyushuensis*. *Proc. R. Soc. Lon.* **248**, 1–7 (1992).
- Knowles, B. H. *et al.* A cytolytic delta-endotoxin from *Bacillus thuringiensis* var. *israelensis* forms cation-selective channels in planar lipid bilayers. *FEBS Lett.* **244**, 259–262 (1989).
- Promdonkoy, B. & Ellar, D. J. Investigation of the pore-forming mechanism of a cytolytic delta-endotoxin from *Bacillus thuringiensis*. *Biochem. J.* **374**, 255–259 (2003).
- Rodríguez-Almazán, C. *et al.* The amino- and carboxyl-terminal fragments of the *Bacillus thuringiensis* Cyt1Aa toxin have differential roles on toxin oligomerization and pore formation. *Biochemistry* **50**, 388–396 (2011).
- López-Díaz, J. A., Cantón, P. A., Gill, S. S., Soberón, M. & Bravo, A. Oligomerization is a key step in Cyt1Aa membrane insertion and toxicity but not necessary to synergize Cry11Aa toxicity in *Aedes aegypti* larvae. *Environ. Microbiol.* **15**, 3030–3039 (2013).
- Promdonkoy, B. & Ellar, D. J. Membrane pore architecture of a cytolytic toxin from *Bacillus thuringiensis*. *Biochem. J.* **350**, 275–282 (2000).
- Du, J., Knowles, B. H., Li, J. & Ellar, D. J. Biochemical characterization of *Bacillus thuringiensis* cytolytic toxins in association with a phospholipid bilayer. *Biochem. J.* **338**, 185–193 (1999).
- Li, J., Derbyshire, D. J., Promdonkoy, B. & Ellar, D. J. Structural implications for the transformation of the *Bacillus thuringiensis* delta-endotoxins from water-soluble to membrane-inserted forms. *Biochem. Soc. Trans.* **29**, 571–577 (2001).
- Promdonkoy, B. *et al.* Amino acid substitutions in α A and α C of Cyt2Aa2 alter hemolytic activity and mosquito larvicidal specificity. *J. Biotechnol.* **133**, 287–293 (2008).
- Crickmore, N., Bone, E. J., Williams, J. A. & Ellar, D. J. Contribution of the individual components of the δ -endotoxin crystal to the mosquitocidal activity of *Bacillus thuringiensis* subsp. *israelensis*. *FEMS Microbiol. Lett.* **131**, 249–254 (1995).
- Pérez, C. *et al.* Bti Cry11Aa and Cyt1Aa toxins interactions support the synergism-model that Cyt1Aa functions as membrane-bound receptor. *Proc. Natl. Acad. Sci. USA* **102**, 18303–18308 (2005).
- Canton, P. E., Reyes, E. Z., Ruiz deEscudero, I., Bravo, A. & Soberón, M. Binding of *Bacillus thuringiensis* subsp. *israelensis* Cry4Ba to Cyt1Aa has an important role in synergism. *Peptides* **32**, 595–600 (2011).
- Wirth, M., Georghiou, G. P. & Federici, B. A. CytA enables CryIV endotoxins of *Bacillus thuringiensis* to overcome high levels of CryIV resistance in the mosquito *Culex quinquefasciatus*. *Proc. Natl. Acad. Sci. USA* **9**, 10536–10540 (1997).
- Pérez, C. *et al.* *Bacillus thuringiensis* subsp. *israelensis* Cyt1Aa enhances activity of Cry11Aa toxin by facilitating the formation of a pre-pore oligomeric structure. *Cell. Microbiol.* **9**, 2931–2937 (2007).
- Zhao, J.-Z. *et al.* mCry3A-selected western corn rootworm (*Coleoptera:Chrysomelidae*) colony exhibits high resistance and has reduced binding of mCry3A to midgut tissue. *J. Econm. Entomol.* **109**, 1369–1377 (2016).
- Federici, B. A. & Bauer, L. S. Cyt1Aa protein of *Bacillus thuringiensis* is toxic to the cottonwood leaf beetle, *Chrysomela scripta*, and suppresses high levels of resistance to Cry3Aa. *Appl. Environ. Microbiol.* **64**, 4368–4371 (1998).
- Porcar, M., Grenier, A.-M., Federici, B. & Rahbe, Y. Effects of *Bacillus thuringiensis* δ -endotoxins on the pea aphid (*Acyrtosiphon pisum*). *Appl. Environ. Microbiol.* **78**, 4897–4900 (2009).
- Gazit, E. & Shai, Y. Structural characterization, membrane interaction, and specific assembly within phospholipid membranes of hydrophobic segments from *Bacillus thuringiensis* var. *israelensis* cytolytic toxin. *Biochemistry* **32**, 12363–12371 (1993).
- Wirth, M. C., Park, H.-W., Walton, W. E. & Federici, B. A. Cyt1A of *Bacillus thuringiensis* delays evolution of resistance to Cry11A in the mosquito *Culex quinquefasciatus*. *Appl. Environ. Microbiol.* **71**, 185–189 (2005).
- Wu, D., Johnson, J. J. & Federici, B. A. Synergism of mosquitocidal toxicity between CytA and CryIV proteins using inclusions produced from cloned genes of *Bacillus thuringiensis* subsp. *israelensis*. *Mol. Microbiol.* **13**, 965–972 (1994).
- Chang, C., Yu, Y. M., Dai, S. M., Law, S. K. & Gill, S. S. High-level cryIVD and cytA gene expression in *Bacillus thuringiensis* does not require the 20-kilodalton protein, and the coexpressed gene products are synergistic in their toxicity to mosquitoes. *Appl. Environ. Microbiol.* **59**, 815–821 (1993).

40. Lereclus, D., Agaisse, H., Gominet, M. & Chaufaux, J. Overproduction of encapsulated insecticidal crystal proteins in a *Bacillus thuringiensis* spoOA mutant. *Bio/Technology* **13**, 67–71 (1995).
41. Thomas, W. E. & Ellar, D. J. *Bacillus thuringiensis* var. *israelensis* crystal delta-endotoxin: effects on insect and mammalian cells *in vitro* and *in vivo*. *J. Cell Sci.* **60**, 181–197 (1983).
42. Fernandez-Luna, M. T. *et al.* Single-concentration tests show synergism among *Bacillus thuringiensis* subsp. *israelensis* toxins against the malaria vector mosquito *Anopheles albimanus*. *J. Invertebr. Pathol.* **104**, 231–233 (2010).

Acknowledgements

We thank Lizbeth Cabrera, Blanca Ines García-Gómez and Jorge F. Sánchez-Quintana for technical assistance. This work was supported in part by CONACyT 221200, and PAPIIT IN202718.

Author Contributions

A.B. and M.S. designed experiments; A.B. analyzed data and wrote the paper; J.A.L.D. built initial constructs and validated them; T.Y., J.Z.Z., A.S. and K.H. perform bioassays; G.M. perform oligomerization assays; M.C.T.Q. and J.O. perform bioassays and hemolysis assays; M.E.N., G.W., A.S. and M.S. edited the manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-36205-6>.

Competing Interests: A.B. and M.S. participate as co-authors on a US Provisional 62/337656 “*Bacillus thuringiensis* Cyt1a Mutants”.

Publisher’s note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018