


## Research Article

# Zingiber officinale and Vernonia amygdalina Infusions Improve Redox Status in Rat Brain

**Damilare Emmanuel Rotimi** <sup>1,2</sup>, **Goodnews Mavoghenegbero Ben-Goru**,<sup>1,2</sup>  
**Ikponmwoa Owen Evbuomwan**,<sup>3</sup> **Tobiloba Christiana Elebiyo**,<sup>1,2</sup> **Mohammed Alorabi**,<sup>4</sup>  
**Abdullah Farasani**,<sup>5</sup> **Gaber El-Saber Batiha** <sup>6</sup>, and **Oluyomi Stephen Adeyemi** <sup>1,2,7</sup>

<sup>1</sup>SDG 03 Group-Good Health & Well-Being, Landmark University, Omu-Aran 251101, Kwara State, Nigeria

<sup>2</sup>Department of Biochemistry, Medicinal Biochemistry, Nanomedicine & Toxicology Laboratory, Landmark University, PMB 1001, Omu-Aran 251101, Nigeria

<sup>3</sup>Department of Microbiology, Cellular Parasitology Unit, College of Pure and Applied Sciences, Landmark University, PMB 1001, Omu-Aran 251101, Nigeria

<sup>4</sup>Department of Biotechnology, College of Sciences, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia

<sup>5</sup>Department of Medical Laboratory Technology, Biomedical Research Unit, Medical Research Center, College of Applied Medical Sciences, Jazan University, Jazan 45142, Saudi Arabia

<sup>6</sup>Department of Pharmacology and Therapeutics, Faculty of Veterinary Medicine, Damanshour University, Damanshour 22511, AlBeheira, Egypt

<sup>7</sup>Laboratory of Sustainable Animal Environment, Graduate School of Agricultural Science, Tohoku University, 232-3 Yomogida, Naruko-Onsen, Osaki, Miyagi 989-6711, Sendai, Japan

Correspondence should be addressed to Oluyomi Stephen Adeyemi; yomibowa@yahoo.com

Received 14 February 2022; Accepted 6 September 2022; Published 26 September 2022

Academic Editor: Ruchika Garg

Copyright © 2022 Damilare Emmanuel Rotimi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The study investigated the effects of *Zingiber officinale* root and *Vernonia amygdalina* leaf on the brain redox status of Wistar rats. Twenty-four (24) rats weighing  $160 \pm 20$  g were randomly assigned into four (4) groups, each with six (6) rats. Animals in Group 1 (control) were orally administered distilled water (1 mL), while the test groups were orally administered 5 mg/mL of either *Z. officinale*, *V. amygdalina* infusion, or a combination of both, respectively, for 7 days. The rats were sacrificed at the end of treatments and blood and tissue were harvested and prepared for biochemical assays. Results showed that administration of *V. amygdalina* and *Z. officinale*, as well as their coadministration, reduced the levels of malondialdehyde (MDA), nitric oxide (NO), acetylcholinesterase (AChE), and myeloperoxidase (MPO) in rat brain tissue compared with the control group. Conversely, coadministration of *V. amygdalina* and *Z. officinale* increased the levels of reduced glutathione (GSH) in rat brain tissue compared with the control group. However, the administration of the infusions singly, as well as the combination of both infusions, did not have any effect on the rat brain levels of glutathione peroxidase (GPx) and catalase (CAT) antioxidant enzymes compared to the control. Taken together, the findings indicate that the *V. amygdalina* and *Z. officinale* tea infusions have favorable antioxidant properties in the rat brain. The findings are confirmatory and contribute to deepening our understanding of the health-promoting effects of *V. amygdalina* and *Z. officinale* tea infusions.

## 1. Introduction

Over time, man has gradually gained knowledge and application of plants as sources of food and medicine through trial and error, and he has become able to source for his

needs from his immediate environment. The application of these plants is being refined bit by bit due to technological and scientific advances. In many regions of the world, medicinal plants contribute to primary health care delivery especially in their use as vital sources of pharmaceutical and

therapeutic products, with approximately 80% of the human population globally depending on natural products for primary health care [1, 2]. Medicinal plants are used singly or in combination with other substances (to increase treatment efficacy for several therapies) in the preparation of traditional or alternative medicines [3, 4].

Ginger (*Zingiber officinale*), belonging to the family Zingiberaceae, is a popular spice and herb used as a delicacy. *Z. officinale* has long been in use therapeutically and is currently validated as a potent medicinal spice to manage numerous diseases such as diabetes, hypertension, cancer, ulcer, diarrhea, cold, cough, spasms, and vomiting [5]. It is an effective spasmolytic, antipyretic, antiemetic, antioxidant, antiulcer, analgesic, hypotensive, antidiabetic, and anti-inflammatory agent containing scented essential oils and spicy oleoresins. The phenolic compounds (gingerols, shogaols, paradol, and gingerdiones) in *Z. officinale* have been discovered to be the source of their pharmacological benefits [5]. On the other hand, bitter leaf (*Vernonia amygdalina*) is a shrub or small tree belonging to the family Asteraceae that grows predominantly in tropical regions in Africa and Asia with the leaves having a characteristic bitter taste. For hundreds of years, bitter leaf serves as fodder, food, and herbal medicine [6]. *V. amygdalina* is rich in minerals such as potassium, magnesium, zinc, iron, and vitamins A, C, and E [7]. The shrub has been known to exhibit profound pharmacological properties such as antimalarial, antioxidant, antimicrobial anticancer, neuroprotective, anti-inflammatory, and anthelmintic properties [8–12]. The pharmacological properties of the bitter leaf have been reported to be connected to its phytochemical constituents which include flavonoids, steroidal saponins, alkaloids, sesquiterpene lactones, anthraquinones, polyphenolics, and coumarins [3, 13]. Interestingly, *Z. officinale* and *V. amygdalina* are useful components in traditional medical systems like Ayurveda to manage neurodegenerative conditions [14, 15].

Oxidative phosphorylation leads to the production of oxidants and free radicals including reactive oxygen species (ROS) and reactive nitrogen species (RNS), the two most important signaling molecules that help to sustain cellular proliferation and differentiation, trigger stress-responsive survival mechanisms, and maintain cellular homeostasis [16–18]. Under normal physiological conditions at redox equilibrium, harmful effects of the production of ROS and other free radicals during aerobic metabolism are counterbalanced by the antioxidant system, and in this way, the organs in the body including the brain efficiently regulate their oxygen consumption and redox generation capacity [19, 20]. However, when the production of ROS, other free radicals, and radical reactive species surpasses the scavenging ability of the antioxidant response system, a situation referred to as redox imbalance ensues [21]. The redox imbalance could cause oxidative cellular damage because of lipid peroxidation as well as protein and nucleic acid oxidation. The resulting oxidative stress could mediate the pathogenesis and pathophysiology of several diseases including neurodegenerative disorders and cardiovascular diseases [22–26].

The brain is one of the most metabolically active organs in the body and its oxygen consumption and redox generation capacity is under strict control. Although the brain of an adult human being weighs about 1.4 kg, it consumes approximately 20% of the total basal oxygen to power its high metabolic activity [27]. In addition, the low endogenous antioxidant defense mechanism of the brain makes it highly sensitive to oxidative damage. Therefore, ischemic/hypoxic or hyperoxic condition raises the levels of free radicals and disrupts redox homeostasis in the brain which could lead to cellular death [28, 29]. Various types of antioxidants play significant roles in redox homeostasis. These antioxidants comprise vitamins A, C, and E; glutathione, coenzyme Q, bilirubin, and ferritin; and endogenous antioxidant enzymes such as glutathione peroxidase (GPx), catalase (CAT), superoxide dismutase (SOD), glutathione reductase (GR), and heme oxygenase (HO) [30, 31]. Furthermore, many medicinal plants and their products have been shown to possess antioxidant defense capacity that helps protect against oxidative damage and maintain redox homeostasis [32–34]. Therefore, the present study aimed to determine the effect of *Z. officinale* and *V. amygdalina* infusions on rat brain redox status.

## 2. Materials and Methods

### 2.1. Identification and Authentication of the Sample.

*Z. officinale* roots were obtained from Omu-Aran market, Kwara state, while *V. amygdalina* leaves were acquired from Landmark University, Omu-Aran. Both plants were identified and authenticated at the Department of Plant Biology, University of Ilorin, Kwara State with voucher numbers; UILH|001|1023|2021 for *Z. officinale* and UILH|002|1083|2021 for *V. amygdalina*.

#### 2.1.1. Preparation of *Z. officinale* and *V. amygdalina* Infusion.

The *Z. officinale* root was washed with water, peeled, sun-dried, and pulverized into powder. The *V. amygdalina* leaf was washed, sun-dried, and the stalk was removed from the leaf before it was pulverized into powder. To make the infusion, the plant powder in the tea bag was steeped in hot water (5 mg/mL, 100°C) for 3–5 minutes [35]. The infusion was allowed to cool at room temperature (25°C) before use.

### 2.2. Phytochemical Analysis

#### 2.2.1. Qualitative Phytochemical Screening.

Qualitative screening for phytoconstituents of ginger and bitter leaf tea infusions was performed as previously described [13]. The phytochemicals assayed included saponin, phenolic compounds, water-soluble phenol, water-insoluble phenol, free and combined anthraquinones, flavonoids, poly steroid, cardiac glycosides, terpenoids, alkaloids, and tannins.

#### 2.2.2. Animal Grouping and Treatments.

The animals (24 rats) were obtained from the Department of Biochemistry, University of Ilorin, Nigeria. They were housed in a clean cage, allowed to acclimatize for 14 days, and administered

pelletized feed and water. The rats were assigned randomly into four (4) groups, consisting of six (6) rats each. Animals in Group 1 were orally administered distilled water (1 mL). Groups 2, 3, and 4 were orally administered 5 mg/mL of *Z. officinale*, *V. amygdalina*, and an oral combination of *Z. officinale* and *V. amygdalina* infusion, respectively. The treatments lasted for 7 days.

The animals received humane care in compliance with the institution's guidelines and criteria as outlined in the National Institute of Health (NIH) Guidelines for the care and use of laboratory animals, and this work had approval from the ethical committee of the Department of Biochemistry at Landmark University, reference number: LUAC/2021/003A.

After the last treatments, the animals were fasted overnight and sacrificed under mild anesthesia of diethyl ether. The brain was harvested, weighed, and homogenized in an isotonic medium (0.25 M sucrose solution). The homogenate was centrifuged at 5000 rpm for 10 min to obtain supernatants, which were used for biochemical analysis. A section of the brain was also preserved in 10% buffered neutral formalin (BNF) for a histopathology examination.

**2.2.3. Biochemical Assays.** The brain homogenates collected were used in the biochemical determination of redox markers. Total protein content in the rat brain homogenate was determined according to Gornall et al. [36]. Superoxide dismutase (SOD) and glutathione peroxidase (GPx) activities were assayed as previously reported by Misra and Fridovich [37] and Rotruck et al. [38], respectively. The procedures described by Beers and Sizer [39] and Pulli et al. [40] were used to assay for catalase (CAT) and myeloperoxidase (MPO), respectively. For the determination of reduced glutathione (GSH) and malondialdehyde (MDA) levels, the methods described by Beutler and Yeh [41] and Varshney and Kale [42] were used, respectively. Nitric oxide (NO) level and acetylcholinesterase (AChE) activity were determined using Adeyemi et al. [43] and Ellman et al. [44], respectively. DNA fragmentation was carried out following the procedure described by Perandones et al. [45].

**2.2.4. Histopathology Examination.** After the rat brain was excised from the animal, it was fixed in 10% BNF and subsequently processed for histopathology examinations as described by Adeyemi and Akanji [46]. Photomicrograph capture and scoring for morphological changes were done at the Pathology Unit, University of Ilorin Teaching Hospital, Ilorin, Nigeria.

**2.2.5. Statistical Evaluation.** The data were analyzed on GraphPad Prism Software (San Diego, CA, USA), using one-way ANOVA. Data are presented as the mean value of six replicates plus or minus the standard error of the mean (SEM). At *p* value of 0.05, mean values were considered significant.

### 3. Results

**3.1. Qualitative Phytochemical Analysis.** The qualitative phytochemical screening revealed the presence of saponin,

TABLE 1: Qualitative Phytochemical tests of the infusion of ginger, bitter leaf, and a combination of Ginger + Bitterleaf.

Phytochemical compounds	Ginger	Bitter leaf	Ginger + bitter leaf
Saponin	+	+	+
Phenolic compounds	+	+	+
Water insoluble phenol	+	+	+
Flavonoids	+	+	+
Free anthraquinones	+	+	+
Tannins	-	+	+
Alkaloids	+	+	+

+: Detected -: Not detected.

phenolics, flavonoids, alkaloids, and anthraquinones in the tea infusions of ginger and bitter leaf. However, tannins were not detected in ginger infusion but were present in bitter leaf (Table 1).

The administration of *Z. officinale* and *V. amygdalina* infusions separately had no significant effect on the rat brain superoxide dismutase (SOD) activity relative to control. In contrast, the coadministration of both infusions significantly increased the rat brain SOD activity in comparison with control (Figure 1(a)). As shown in Figure 1(d), the separate administration of *Z. officinale* and *V. amygdalina* infusions as well as the coadministration of both infusions did not significantly change the rat brain catalase (CAT) activity in relation to control (Figure 1(b)). As represented in Figure 1(c), there were no significant changes in the rat brain glutathione peroxidase activity in relation to control when *Z. officinale* and *V. amygdalina* infusions were administered separately. In addition, the coadministration of both infusions had a negligible effect on the rat brain glutathione peroxidase activity in relation to control. Glutathione level in the rat brain was reduced when *Z. officinale* infusion only was administered, as well as the coadministration of both infusions, when compared with the control (Figure 1(d)).

An appreciable decrease in the rat brain MDA level in comparison with the control was observed when the infusions were administered separately as well as coadministration of the infusions together (Figure 2(a)). In the same vein, the administration of the tea infusions separately or as a combination significantly decreased the level of the rat brain DNA fragmentation when compared with the control (Figure 2(b)).

Furthermore, the administration of the infusions separately, as well as in a combination, did not significantly affect the rat brain protein concentration when compared to control (Figure 3(a)). However, the administration of the infusions separately or as a combination significantly decreased the level of the rat brain nitric oxide (NO) in relation to control (Figure 3(b)). Figure 3(c) shows a substantial decrease in rat brain AChE activity in comparison with the control when the infusions were administered singly or in combination. Similarly, the rat brain MPO activity decreased when the infusions were administered separately or as a combination (Figure 3(d)).

For all the treatment groups, the histopathology examination revealed normal brain tissue composed of




FIGURE 1: The effect of *Zingiber officinale* and *Vernonia amygdalina* on the levels of rat brain antioxidant molecules: (a) superoxide dismutase, (b) Catalase, (c) glutathione peroxidase, and (d) reduced glutathione. Each value is a mean of six replicates  $\pm$  SEM. Values bearing the asterisk are significantly different compared to the control at  $p < 0.05$ .




FIGURE 2: The effect of *Zingiber officinale* and *Vernonia amygdalina* on the level of rat brain oxidative stress markers: (a) malondialdehyde and (b) DNA fragmentation. Values are the mean of six replicates  $\pm$  SEM. Bars with the asterisk are significantly different from the control at  $p < 0.05$ .




FIGURE 3: The effect of *Zingiber officinale* and *Vernonia amygdalina* on the level of rat brain biochemical parameters: (a) protein concentration, (b) nitric oxide, (c) acetylcholine esterase, and (d) myeloperoxidase. Values are the mean of six replicates ± SEM. Bars with the asterisk are significantly different from the control at  $p < 0.05$ .




FIGURE 4: The effect of *Zingiber officinale* and *Vernonia amygdalina* on rat brain morphology: (a) control, (b) *Vernonia amygdalina*, (c) *Zingiber officinale*, and (d) combination of *Zingiber officinale* and *Vernonia amygdalina*. H&E staining (x100). The sections show normal brain tissue composed of preserved neuronal bodies surrounded by a fibrillary glial matrix (cerebral and hippocampal).

preserved neuronal bodies surrounded by a fibrillary glial matrix (cerebral and hippocampal). No observation of degenerative changes or significant inflammation (Figure 4).

#### 4. Discussion

This work investigated the effect of *Z. officinale* root and *V. amygdalina* leaf infusion on the redox status of the brain of rats to provide more understanding of the health-promoting effects of these commonly consumed medicinal plants. The phytochemical screening showed the presence of saponin, phenolics, flavonoids, free anthraquinones, and alkaloids. However, tannins were not detected in ginger but only in the tea infusions of bitter leaf and that of the combination of bitter leaf and ginger.

We found that the combined mixture of *V. amygdalina* and *Z. officinale* infusions led to a significant increase in SOD activity and an insignificant increase in GPx activity across all experimental groups when compared with the control. This finding could mean that the infusions enhanced the rat brain's oxidative defense status by improving the activities of SOD [47]. In addition, our study revealed that the combined administration of the infusions caused an insignificant increase in CAT activity but a significant decrease in reduced glutathione (GSH). GSH is an antioxidant molecule that quenches  $O_2^{\bullet-}$ ,  $HO^{\bullet}$ , and peroxy nitrite anions (ONOO<sup>-</sup>) by the action of its thiol group, which donates a reducing equivalent to unstable radicals and anions that may have arisen due to respiratory processes and other metabolic activities in the body cells. In contrast to the increased antioxidant activity of SOD, CAT, and GPx, there was a significant decrease in the level of malondialdehyde (MDA), an end-product of lipid peroxidation during oxidative degradation of lipids usually caused by free radicals [48]. The coordinated function of enzymatic antioxidant systems including CAT, SOD, and GPx, and nonenzymatic antioxidants such as glutathione (GSH) and vitamin E play a significant role in the prevention of oxidative damage by ROS such as  $O_2^{\bullet-}$ ,  $HO^{\bullet}$  and  $H_2O_2$  [49, 50]. A significant decrease in the level of nitric oxide (NO), a proinflammatory agent, may indicate the anti-inflammatory action of the tea infusions. Nitric oxide (NO) participates in several physiological processes and is considered a free radical as it could accumulate and react with superoxide anion to form poisonous nitrite anion, thereby leading to nitrosative stress [51]. The decrease in the levels of both MDA and NO across all experimental groups in comparison with the control may be due to the antioxidant properties of *V. amygdalina* and *Z. officinale*. This implies that the tea infusions can suppress oxidative stress by inhibiting the production of MDA and NO in support of earlier reports [14, 52–54]. The tea infusions can limit the production of MDA and NO, and thereby prevent oxidative stress probably by enhancing the activities of the inherent enzymatic antioxidants in the rat brain. Myeloperoxidase (MPO) is a heme-containing peroxidase mainly expressed in neutrophils and, to a lesser extent, in monocytes. In the presence of  $H_2O_2$  and halides (HOCl), MPO catalyzes the formation of reactive oxygen intermediates such as hypochlorous acid. Microbial death by

neutrophils relies heavily on the MPO/HOCl system. In addition, MPO acts as a local mediator of tissue injury through inflammation [55]. The observed decrease in MPO activity due to the separate or combined administration of the tea infusions is consistent with the findings of Hussein et al. [56] and Erukainure et al. [57]. A decrease in MPO activity impedes neutrophil infiltration [58], supporting the anti-inflammatory effect of tea infusions. Furthermore, results showed a decrease in DNA fragmentation following the separate or combined administration of the tea infusions, corroborating an earlier report [57]. This fact further reinforces not only the antioxidant but also the anti-inflammatory potential of the tea infusions of *Z. officinale* and *V. amygdalina* and their capacity to protect against oxidative-induced DNA damage.

Acetylcholinesterase (AChE) is a serine hydrolase primarily responsible for the termination of signal transmission in the cholinergic system. Its substrate, acetylcholine (ACh), is a cholinergic neurotransmitter that has a strong effect on motor neurons involved in memory formation [59]. Therefore, the observed decrease in the activity of this AChE after a separate and combined administration of the tea infusions is similar to the findings reported in earlier studies [11, 59–61]. The present finding suggests that the tea infusions possess bioactive compounds that could decrease neurodegeneration or treat neurodegenerative diseases by modulating the activity of AChE. Protein is an important structural component of cells as well as a source of energy and one of the building blocks of human tissues. Extracellular and intracellular enzymes, as well as other proteins, contribute to the total protein concentrations [62]. The separate and the combined administration of the tea infusions of *V. amygdalina* and *Z. officinale* caused a negligible rise in total protein concentration, attributable to their capacity to boost the production of antioxidant proteins as reflected in the elevated levels of SOD and CAT. Furthermore, morphological examination of the rat brain revealed normal histological architectures of the cerebral and hippocampal in all the treatment groups. There are no features of degenerative changes or inflammation, thus corroborating the biochemical data and further underscoring the medicinal value of *V. amygdalina* and *Z. officinale* tea infusions in rat brains.

#### 5. Conclusions

The data from this work showed that *V. amygdalina* and *Z. officinale* infusions improved brain antioxidants and reduced oxidative stress after a 7-day separate or combined oral administration. The results support the efficacy of the tea infusions in maintaining redox status/homeostasis in rat brains. Taken together, the findings support the health-promoting effects of *V. amygdalina* and *Z. officinale*. Future studies may focus on the long-term effects of the tea infusions separately or as a combination to gain a better understanding of the therapeutic benefits of the medicinal plants.

#### Data Availability

The raw data are available from the corresponding author upon request.

## Additional Points

*Institutional Review Board Statement:* The protocol used in handling the animals was approved by the ethical committee of the Department of Biochemistry at Landmark University and assigned the reference number: LUAC/2021/003A.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Authors' Contributions

O.S.A. and A.T.A. conceptualized the data; O.S.A., H.F.H., and S.A.-R. curated the data; O.S.A., A.T.A., H.F.H., S.A.-R., D.R., and G.E.-S.B. formally analyzed the data; O.S.A. investigated the data; O.S.A., D.R., and G.E.-S.B. created methodology; H.H. and S.A.-R. found resources; O.S.A., A.T.A., D.R., and G.E.-S.B. wrote the original draft; O.S.A., A.T.A., H.F.H., S.A.-R., D.R., T.C.E., and G.E.-S.B. wrote, reviewed, and edited the draft. All authors have read and agreed to the published version of the manuscript.

## Acknowledgments

The authors appreciate the support of Taif University Researchers Supporting Project (number: TURSP-2020/310), Taif University, Taif, Saudi Arabia. In addition, the authors appreciation goes to Landmark University Nigeria.

## References

- [1] F. Jamshidi-Kia, Z. Lorigooini, and H. Amini-Khoei, "Medicinal plants: past history and future perspective," *Journal of Herbméd Pharmacology*, vol. 7, no. 1, p. 1, 2018.
- [2] S. Mlilo and S. Sibanda, "An ethnobotanical survey of the medicinal plants used in the treatment of cancer in some parts of Matebeleland, Zimbabwe," *South African Journal of Botany*, vol. 146, pp. 401–408, 2022, <https://doi.org/10.1016/j.sajb.2021.11.022>.
- [3] X. Liu, W. Tian, G. Wang et al., "Stigmastane-type steroids with unique conjugated  $\Delta^7$ , 9(11) diene and highly oxygenated side chains from the twigs of *Vernonia amygdalina*," *Phytochemistry*, vol. 158, pp. 67–76, 2019.
- [4] M. M. Matotoka and P. Masoko, "Phytochemical screening and pharmacological evaluation of herbal concoctions sold at Ga Maja Limpopo Province," *South African Journal of Botany*, vol. 117, pp. 1–10, 2018, <https://doi.org/10.1016/j.sajb.2018.04.013>.
- [5] F. O. Balogun, E. T. AdeyeOluwa, and A. O. T. Ashafa, "Pharmacological potentials of ginger," *Ginger Cultivation and its Antimicrobial and Pharmacological Potentials*, 2019.
- [6] O. Alara, N. Abdurahman, and O. Olalere, "Ethanol extraction of flavonoids, phenolics and antioxidants from *Vernonia amygdalina* leaf using two-level factorial design," *Journal of King Saud University-Science*, vol. 32, no. 1, pp. 7–16, 2020.
- [7] M. Johnson, O. S. Kolawole, and L. A. Olufunmilayo, "Phytochemical analysis, in vitro evaluation of antioxidant and antimicrobial activity of methanolic leaf extract of *Vernonia amygdalina* (bitter leaf) against *Staphylococcus aureus* and *Pseudomonas aeruginosa*," *Int. J. Curr. Microbiol. App. Sci*, vol. 4, pp. 411–426, 2015.
- [8] O. B. Akpor, J. Ndakotsu, I. O. Evbuomwan, T. D. Olaolu, and O. O. Osemwegie, "Bacterial growth inhibition and antioxidant potentials of leaf infusions of (*Moringa oleifera*), locust beans (*Parkia biglobosa*) and bitter leaf (*Vernonia amygdalina*)," *Scientific African*, vol. 14, 2021.
- [9] G. Oboh, B. C. Adedayo, M. B. Adetola, I. S. Oyeleye, and O. B. Ogunsuyi, "Characterization and neuroprotective properties of alkaloid extract of *Vernonia amygdalina* Delile in experimental models of Alzheimer's disease," *Drug and Chemical Toxicology*, vol. 45, no. 2, pp. 731–740, 2020.
- [10] J. O. Oladele, O. M. Oyeleke, O. T. Oladele, and A. T. Oladiji, "Covid-19 treatment: Investigation on the phytochemical constituents of *Vernonia amygdalina* as potential Coronavirus-2 inhibitors," *Computational Toxicology*, vol. 18, p. 100161, 2021.
- [11] J. O. Oladele, O. M. Oyeleke, O. T. Oladele, and M. Olaniyan, "Neuroprotective mechanism of *Vernonia amygdalina* in a rat model of neurodegenerative diseases," *Toxicology Reports*, vol. 7, pp. 1223–1232, 2020, <https://doi.org/10.1016/j.toxrep.2020.09.005>.
- [12] C. G. Yedjou, A. T. Mbemi, F. Noubissi et al., "Prostate cancer disparity, chemoprevention, and treatment by specific medicinal plants," *Nutrients*, vol. 11, no. 2, p. 336, 2019.
- [13] G. El-Saber Batiha, A. Magdy Beshbishy, O. Stephen Adeyemi et al., "Phytochemical screening and antiprotozoal effects of the methanolic berberis vulgaris and acetonetic rhus coriaria extracts," *Molecules*, vol. 25, no. 3, p. 550, 2020, <https://doi.org/10.3390/molecules25030550>.
- [14] R. Sharma, K. Kuca, E. Nepovimova, A. Kabra, M. M. Rao, and P. K. Prajapati, "Traditional Ayurvedic and herbal remedies for Alzheimer's disease: from bench to bedside," *Expert Review of Neurotherapeutics*, vol. 19, no. 5, pp. 359–374, 2019, <https://doi.org/10.1080/14737175.2019.1596803>.
- [15] R. Sharma, A. Kabra, M. M. Rao, and P. K. Prajapati, "Herbal and holistic solutions for neurodegenerative and depressive disorders: leads from Ayurveda," *Current Pharmaceutical Design*, vol. 24, no. 22, pp. 2597–2608, 2018.
- [16] M. Jelinek, M. Jurajda, and K. Duris, "Oxidative stress in the brain: basic concepts and treatment strategies in stroke," *Antioxidants*, vol. 10, no. 12, p. 1886, 2021.
- [17] A. Nugud, D. Sandeep, and A. T. El-Serafi, "Two faces of the coin: minireview for dissecting the role of reactive oxygen species in stem cell potency and lineage commitment," *Journal of Advanced Research*, vol. 14, pp. 73–79, 2018, <https://doi.org/10.1016/j.jare.2018.05.012>.
- [18] R. Radi, "Oxygen radicals, nitric oxide, and peroxynitrite: redox pathways in molecular medicine," *Proc Natl Acad Sci U S A*, vol. 115, no. 23, pp. 5839–5848, 2018.
- [19] P. Poprac, K. Jomova, M. Simunkova, V. Kollar, C. J. Rhodes, and M. Valko, "Targeting free radicals in oxidative stress-related human diseases," *Trends in Pharmacological Sciences*, vol. 38, no. 7, pp. 592–607, 2017.
- [20] S. Salim, "Oxidative stress and the central nervous system," *J Pharmacol Exp Ther*, vol. 360, no. 1, pp. 201–205, 2017.
- [21] H. Sies, "Biochemie des oxidativen Stress," *Angewandte Chemie*, vol. 98, no. 12, pp. 1061–1075, 1986.
- [22] X. Ba and I. Boldogh, "8-Oxoguanine DNA glycosylase 1: beyond repair of the oxidatively modified base lesions," *Redox Biology*, vol. 14, pp. 669–678, 2018.
- [23] J. Banerjee, A. Das, M. Sinha, and S. Saha, *Biological efficacy of medicinal plant extracts in preventing oxidative damage*, vol. 2018, 2018.

- [24] D. A. Butterfield and B. Halliwell, "Oxidative stress, dysfunctional glucose metabolism and Alzheimer disease," *Nature Reviews Neuroscience*, vol. 20, no. 3, pp. 148–160, 2019.
- [25] B. Matter, C. L. Seiler, K. Murphy et al., "Mapping three guanine oxidation products along DNA following exposure to three types of reactive oxygen species," *Free Radical Biology and Medicine*, vol. 121, pp. 180–189, 2018.
- [26] C. Peña-Bautista, M. Baquero, M. Vento, and C. Cháfer-Pericás, "Free radicals in Alzheimer's disease: lipid peroxidation biomarkers," *Clinica Chimica Acta*, vol. 491, pp. 85–90, 2019.
- [27] P. J. Magistretti and I. Allaman, "A cellular perspective on brain energy metabolism and functional imaging," *Neuron*, vol. 86, no. 4, pp. 883–901, 2015.
- [28] J. N. Copley, M. L. Fiorello, and D. M. Bailey, "13 reasons why the brain is susceptible to oxidative stress," *Redox Biology*, vol. 15, pp. 490–503, 2018.
- [29] I. Torres-Cuevas, M. Corral-Debrinski, and P. Gressens, "Brain oxidative damage in murine models of neonatal hypoxia/ischemia and reoxygenation," *Free Radical Biology and Medicine*, vol. 142, pp. 3–15, 2019, <https://doi.org/10.1016/j.freeradbiomed.2019.06.011>.
- [30] B. Marengo, M. Nitti, A. L. Furfaro et al., "Redox homeostasis and cellular antioxidant systems: crucial players in cancer growth and therapy," *Oxidative Medicine and Cellular Longevity*, vol. 2016, pp. 6235641–16, 2016.
- [31] J. Taile, A. Arcambal, P. Clerc, A. Gauvin-Bialecki, and M. P. Gonthier, "Medicinal plant polyphenols attenuate oxidative stress and improve inflammatory and vasoactive markers in cerebral endothelial cells during hyperglycemic condition," *Antioxidants (Basel)*, vol. 9, no. 7, p. 573, 2020.
- [32] J. Namukobe, P. Sekandi, R. Byamukama et al., "Antibacterial, antioxidant, and sun protection potential of selected ethno medicinal plants used for skin infections in Uganda," *Trop Med Health*, vol. 49, no. 1, p. 49, 2021.
- [33] C. O. Nwonuma, O. O. Osemwegie, O. O. Alejlowo et al., "Antioxidant and the ameliorating effect of *Allium cepa* (Onion) fortified feed against potassium bromate induced oxidative damage in Wistar rats," *Toxicology Reports*, vol. 8, pp. 759–766, 2021.
- [34] A. Septembre-Malaterre, G. Stanislas, E. Douraguia, and M. P. Gonthier, "Evaluation of nutritional and antioxidant properties of the tropical fruits banana, litchi, mango, papaya, passion fruit and pineapple cultivated in Reunion French Island," *Food Chemistry*, vol. 212, pp. 225–233, 2016.
- [35] A. Zachara, D. Gałkowska, and L. Juszczak, "Contamination of tea and tea infusion with polycyclic aromatic hydrocarbons," *International Journal of Environmental Research and Public Health*, vol. 15, no. 1, p. 45, 2017.
- [36] A. G. Gornall, C. J. Bardawill, and M. M. David, "Determination of serum proteins by means of the biuret reaction," *Journal of Biological Chemistry*, vol. 177, no. 2, pp. 751–766, 1949.
- [37] H. P. Misra and I. Fridovich, "The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase," *Journal of Biological Chemistry*, vol. 247, no. 10, pp. 3170–3175, 1972.
- [38] J. T. Rotruck, A. L. Pope, H. E. Ganther, A. B. Swanson, D. G. Hafeman, and W. G. Hoekstra, "Selenium: biochemical role as a component of glutathione peroxidase," *Science*, vol. 179, no. 4073, pp. 588–590, 1973.
- [39] R. F. Beers and I. W. Sizer, "A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase," *Journal of Biological Chemistry*, vol. 195, no. 1, pp. 133–140, 1952.
- [40] B. Pulli, M. Ali, R. Forghani et al., "Measuring myeloperoxidase activity in biological samples," *PLoS One*, vol. 8, no. 7, p. e67976, 2013.
- [41] E. Beutler and M. K. Y. Yeh, "Erythrocyte glutathione reductase," *Blood*, vol. 21, no. 5, pp. 573–585, 1963.
- [42] R. Varshney and R. K. Kale, "Effects of calmodulin antagonists on radiation-induced lipid peroxidation in microsomes," *International Journal of Radiation Biology*, vol. 58, no. 5, pp. 733–743, 2009.
- [43] O. S. Adeyemi, O. Atolani, P. Banerjee et al., "Computational and experimental validation of antioxidant properties of synthesized bioactive ferulic acid derivatives," *International Journal of Food Properties*, vol. 21, no. 1, pp. 86–98, 2018.
- [44] G. L. Ellman, K. D. Courtney, V. Andres Jr, and R. M. Featherstone, "A new and rapid colorimetric determination of acetylcholinesterase activity," *Biochemical Pharmacology*, vol. 7, no. 2, pp. 88–95, 1961.
- [45] C. E. Perandones, V. A. Illera, D. Peckham, L. L. Stunz, and R. F. Ashman, "Regulation of apoptosis in vitro in mature murine spleen T cells," *Journal of Immunology (Baltimore, Md, 1950)*, vol. 151, no. 7, pp. 3521–3529, 1993.
- [46] O. S. Adeyemi and M. A. Akanji, "Psidium guajava leaf extract: effects on rat serum homeostasis and tissue morphology," *Comparative Clinical Pathology*, vol. 21, no. 4, pp. 401–407, 2010.
- [47] K. Ji, L. Fang, H. Zhao et al., "Ginger oleoresin alleviated  $\gamma$ -ray irradiation-induced reactive oxygen species via the Nrf2 protective response in human mesenchymal stem cells," *Oxidative Medicine and Cellular Longevity*, vol. 2017, pp. 1–12, 2017.
- [48] S. B. Nimse and D. Pal, "Free radicals, natural antioxidants, and their reaction mechanisms," *RSC Advances*, vol. 5, no. 35, pp. 27986–28006, 2015.
- [49] S. Demirci-Çekiç, G. Özkan, A. N. Avan, S. Uzunboy, E. Çapanoğlu, and R. Apak, "Biomarkers of oxidative stress and antioxidant defense," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 209, 2022.
- [50] K. Ferro, D. Ferro, F. Corrà, R. Bakiu, G. Santovito, and J. Kurtz, "Cu, Zn superoxide dismutase genes in *Tribolium castaneum*: evolution, molecular characterisation, and gene expression during immune priming," *Frontiers in Immunology*, vol. 8, p. 1811, 2017.
- [51] J. C. Toledo Jr. and O. Augusto, "Connecting the chemical and biological properties of nitric oxide," *Chem Res Toxicol*, vol. 25, no. 5, pp. 975–989, 2012.
- [52] K. Danwilai, J. Konmun, B.-o. Sripanidkulchai, and S. Subongkot, "Antioxidant activity of ginger extract as a daily supplement in cancer patients receiving adjuvant chemotherapy: a pilot study," *Cancer Management and Research*, vol. 9, pp. 11–18, 2017.
- [53] I. Mustafa, N. L. Chin, S. Fakurazi, and A. Palanisamy, "Comparison of phytochemicals, antioxidant and anti-inflammatory properties of sun-oven- and freeze-dried ginger extracts," *Foods*, vol. 8, no. 10, p. 456, 2019.
- [54] Z. Naderi, H. Mozaffari-Khosravi, A. Dehghan, A. Nadjarzadeh, and H. F. Huseini, "Effect of ginger powder supplementation on nitric oxide and C-reactive protein in elderly knee osteoarthritis patients: a 12-week double-blind randomized placebo-controlled clinical trial," *Journal of Traditional and Complementary Medicine*, vol. 6, no. 3, pp. 199–203, 2016.



- [55] Y. Aratani, "Myeloperoxidase: its role for host defense, inflammation, and neutrophil function," *Archives of Biochemistry and Biophysics*, vol. 640, pp. 47–52, 2018.
- [56] S. A. Hussein, Y. A. El Senosi, M. K. Mansour, and M. F. Hassan, "Role of antioxidant and anti-inflammatory of Ginger (*Zingiber officinale* Roscoe) against metalaxyl induced oxidative stress in rats," *Benha Veterinary Medical Journal*, vol. 33, no. 2, pp. 504–516, 2017.
- [57] O. L. Erukainure, C. I. Chukwuma, O. Sanni, M. G. Matsabisa, and M. S. Islam, "Histochemistry, phenolic content, antioxidant, and anti-diabetic activities of *Vernonia amygdalina* leaf extract," *Journal of Food Biochemistry*, vol. 43, no. 2, 2019.
- [58] W. Q. Liu, Y. Z. Zhang, Y. Wu et al., "Myeloperoxidase-derived hypochlorous acid promotes ox-LDL-induced senescence of endothelial cells through a mechanism involving beta-catenin signaling in hyperlipidemia," *Biochemical and Biophysical Research Communications*, vol. 467, no. 4, pp. 859–865, 2015.
- [59] H. Khan, S. Amin, M. A. Kamal, and S. Patel, "Flavonoids as acetylcholinesterase inhibitors: current therapeutic standing and future prospects," *Biomedicine & Pharmacotherapy*, vol. 101, pp. 860–870, 2018.
- [60] L. Czernicka, A. Ludwiczuk, E. Rój, Z. Marzec, A. Jarzab, and W. Kukula-Koch, "Acetylcholinesterase inhibitors among *Zingiber officinale* terpenes—extraction conditions and thin layer chromatography-based bioautography studies," *Molecules*, vol. 25, no. 7, p. 1643, 2020.
- [61] B. T. Tung, D. K. Thu, N. T. K. Thu, and N. T. Hai, "Antioxidant and acetylcholinesterase inhibitory activities of ginger root (*Zingiber officinale* Roscoe) extract," *Journal of Complementary & Integrative Medicine*, vol. 14, no. 4, 2017.
- [62] F. Laddomada, M. M. Miyachiro, and A. Dessen, "Structural insights into protein-protein interactions involved in bacterial cell wall biogenesis," *Antibiotics (Basel)*, vol. 5, no. 2, p. 14, 2016.