



Silicosis Incidence and Mortality after Occupational Exposure with Silica Dust: A Systematic Review and Dose-Response Meta-Analysis

Fatemeh Shahbazi^{1, 2*} , Mohammad Javad Assari^{2, 3}, Yahya Shadi^{1, 4}, Marzieh Fattahi-Darghlou^{1, 4*} 

Received: 17 May 2024

Published: 23 Nov 2024

Abstract

Background: We conducted a systematic review of all published epidemiological research related to the relationship between occupational silica exposure and the rates of silicosis incidence and mortality.

Methods: We searched Scopus, PubMed, and Web of Sciences up to 11/07/2023, for original in any language. The search start date was not limited. Observational studies, including cohort, case-control, and cross-sectional that have reported risk estimates for the association between silica exposure and silicosis mortality and incidence rates were considered. The methodological quality of the included articles was assessed using the Newcastle-Ottawa scale. Pooled estimates were calculated using the random effects model. Dose-response relations were explored through a two-stage random-effects model with “drmeta” command in Stata software version 14.

Results: Nineteen observational studies were included in the present systematic review and meta-analysis. Based on the linear dose-response analysis, with each mg/m³ increase in daily occupational exposure to silica, the mortality risk of silicosis, the odds and risk of silicosis occurrence significantly increased by 10.19%, 360.02%, and $4.43 \times 108\%$, respectively.

Conclusion: This review revealed that there is a linear dose-response relationship between occupational exposure to silica and incidence and mortality from silicosis. Our findings could have practical applications for occupational and public health. Considering the direct relationship between occupational silica exposure and high incidence and mortality rates of silicosis, the level of silica dust should be decreased in different industries.

Keywords: Silicon Dioxide, Silica; Silicosis, Systematic-Review, Meta-Analysis, Dose-Response Relationship.

Conflicts of Interest: None declared

Funding: This review was supported financially by the Research Vice-Chancellor of Hamedan University of Medical Sciences (Research ID: 140306275259).

*This work has been published under CC BY-NC-SA 4.0 license.

Copyright© Iran University of Medical Sciences

Cite this article as: Shahbazi F, Assari MJ, Shadi Y, Fattahi-Darghlou M. Silicosis Incidence and Mortality after Occupational Exposure with Silica Dust: A Systematic Review and Dose-Response Meta-Analysis. *Med J Islam Repub Iran*. 2024 (23 Nov);38:135. <https://doi.org/10.47176/mjiri.38.135>

Introduction

Silicosis is a potentially fatal, irreversible, fibrotic pulmonary disease that may develop subsequent to the inhalation of large amounts of silica dust over time (1). This health outcome is an occupational disease (2) mainly that occurs in almost all countries and in many industries and occupations, including sandblasting, mining, rock drilling,

quarrying, brick cutting, glass manufacturing, tunneling, foundry work, stone working, ceramic manufacturing and construction activities (3, 4). Globally, silicosis accounts for 90% of all pneumoconiosis cases (5). The prevalence of silicosis in 1990 and 2019 was 33.13 and 31.60 per 100,000 people, respectively (5). According to the global burden of

Corresponding authors: Dr Fatemeh Shahbazi, f.shahbazi@umsha.ac.ir
Marzieh Fattahi-Darghlou, m.fattahid@edu.umsha.ac.ir

¹ Department of Epidemiology, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran

² Occupational Health and Safety Research Center, Health Sciences & Technology Research Institute, Hamadan University of Medical Sciences, Hamadan, Iran

³ Department of Occupational Health, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran

⁴ Students Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran

↑What is “already known” in this topic:

Although the relationship between Silicosis incidence and mortality after occupational exposure to silica been established in previous literature, its quantity is still not well defined. The purpose of this meta-analysis is to explore potential links between exposure to silica in the workplace and the occurrence and death rates related to silicosis.

→What this article adds:

The current study highlights there is a linear dose-response relationship between occupational exposure to silica and incidence and mortality from silicosis. With more details, each mg/m³ increase in daily occupational exposure to silica, the mortality risk of silicosis, and the odds and risk of silicosis occurrence significantly increased by 10.19%, 360.02% and $4.43 \times 108\%$, respectively.

diseases study, the death number of silicosis between 1990 and 2019 has decreased from 9973 to 10043 cases, and the disability-adjusted life years (DALYs) rates have decreased from 5.05 cases per 100,000 people to 3.33 cases in 2019 (6). The highest disease burden is carried by the middle and high-middle socio-demographic index regions and elderly people (7). According to previous studies, occupational exposure to silica has an excessive mortality rate due to silicosis, although the quantity of this relationship is different in various research (8-10).

Although the relationship between exposure to silica and the risk of silicosis has been well established in previous literature (4, 11-14), its quantity is still not well defined. The purpose of this meta-analysis is to explore potential links between occupational silica exposure and the incidence and mortality rates of silicosis. While recent literature has examined this association through epidemiological studies, we believe this is the first systematic review and dose-response meta-analysis conducted so far.

Methods

Eligibility criteria (PICOS)

Population: The study population comprised people who were exposed to silica due to their jobs.

Exposure: The exposure group included people who had different levels of exposure to silica.

Control: The control group consisted of people who were not exposed to silica or were exposed to it at the lowest level.

Outcome: The primary outcome of interest was the occurrence of silicosis, as confirmed through radiological examination, or death from silicosis.

Studies: Observational studies, including cross-sectional, retrospective cohort, prospective cohort, and case-control studies, were included in the analysis regardless of their publication time, publication status or language.

Information sources and search

A systematic search was conducted in Scopus, PubMed, and Web of Science databases until 11/07/2023. The search start date was not limited. Additionally, the references of the included studies were reviewed to identify any additional studies that met the inclusion criteria. The search strategy was as follows: (((silica [Title/Abstract]) OR (Silicon Dioxide [MeSH Terms])) AND ((Silicosis [MeSH Terms]) OR (Silicosis [Title/Abstract])) AND (((Case-Control Studies [MeSH Terms]) OR (Cohort Studies [MeSH Terms])) OR (Observational Study [Publication Type]))).

Study selection

The search results from the three mentioned databases were combined using EndNote software, and duplicate records were removed. Two authors (Y.S. and M.FD.) independently reviewed the titles and abstracts of the preliminary articles to exclude those that did not meet the inclusion criteria. Then the full text of the articles whose title and abstract were in accordance with the purpose of the study was examined to determine their eligibility for inclusion in the analysis. Disagreements between authors were resolved by

negotiation. Also, the Kappa statistic was calculated to measure the agreement between the two authors.

Data extraction

The data from the included studies were extracted by two authors (F.SH. and M.FD.) and recorded in Stata software. The extracted data included the first author's name, publication year, country, language, age mean/ range, study design, the number of cases at each level of exposure to silica, type of effect size with confidence interval or standard error of the outcome at each exposure level, comorbidity with tuberculosis, and daily exposure to silica (mg/m^3).

Methodological quality

We used the Newcastle-Ottawa scale (NOS) to evaluate the methodological quality of the studies included in this systematic review and meta-analysis (15). This scale was developed to assess the quality of nonrandomized studies such as case-control, cohort, and cross-sectional studies. In the NOS, a set of items about the selection of study participants, comparability of groups, and evaluation of exposure/ outcome were assessed. This scale assigns a maximum of nine stars to each study (4 stars for selection, 2 stars for comparability, and 3 stars for outcome or exposure). Studies that obtained seven or more stars were classified as high-quality, while those with fewer than seven stars were classified as low-quality.

Assessment of heterogeneity and publication bias

We used I^2 statistic and chi-square test to investigate the existence of heterogeneity between studies (16). We also performed meta-regression to identify any potential factors that could explain the observed heterogeneity in the results. We used the Begg test (17) and funnel plot (18) to check the possibility of publication bias.

Statistical analysis

The "metan" command was performed to summarize data from studies (19) using Stata software version 14 (StataCorp, College Station, TX, USA) and the random effects model. In order to investigate the dose-response relationship between silica exposure and the incidence and mortality rates from silicosis, the two-stage random-effects dose-response model using linear, quadratic, and cubic splines was used (20). Different models were fitted to the data and the best model was chosen according to Bayesian information criterion.

Results

Study selection & Characteristics

The primary search found 513 articles. Following excluding 128 duplicates, 385 articles were entered for the title and abstract reviewing. Next, 51 articles were included for the full-text screening. Finally, the data from 19 studies (6 studies for mortality and 13 studies for incidence) were included in the systematic review and dose-response meta-analysis (Figure 1). These studies included 288336 participants from six different countries (Australia, Brazil, China, Finland, Germany, and the USA). The mean of exposure to silica in incidence studies was $0.54 \text{ mg}/\text{m}^3$ per day and in




Figure 1. Flow of information through the various phases of the systematic review

the mortality studies, it was 9.50 mg/m³ per day. The studies were published between 1994 and 2021. All studies were published in English. The percentage of agreement between data extractors was 81.5%.

Overall systematic review and meta-analysis results

We found a significant positive association between occupational silica exposure and mortality rate from silicosis (RR = 4.54, 95% CI: 2.41, 6.67) (Figure 2). We performed a subgroup analysis by measuring the association in the relationship between occupational silica exposure and silicosis incidence. The pooled OR, RR, and HR for silicosis incidence were 1.13 (95% CI: 1.03, 1.22), 2.22 (95% CI: 1.32, 3.12), and 2.41 (95% CI: -0.15, 4.97), respectively (Figure 3). The meta-regression showed no significant association between country, years of publication, study quality and risk of silicosis mortality and incidence ($P = 0.530$).

Publication bias and quality assessment

The Begg test indicates no significant publication bias in the association between occupational silica exposure and

mortality from silicosis ($P = 0.535$) and silicosis incidence ($P = 0.617$). Also, the symmetry distribution of studies around the null line in all funnel plots indicates the absence of publication bias (Figure 4). In terms of quality assessment based on the NOS eight of the studies had low quality and eleven had high quality (Table 1).

Dose-response meta-analysis

We included 11 studies in a dose-response meta-analysis between occupational inhalation of silica dust and silicosis mortality and incidence. In this analysis, the studies that had not reported the number of silicosis patients or the number of deaths due to silicosis in each level of exposure to silica were excluded. According to linear dose-response analysis, the risk of mortality from silicosis increased by 10.19% for every mg/m³ increase in daily occupational silica inhalation ($P < 0.001$). Also, the odds and risk of silicosis occurrence increased by 360.02% and $4.43 \times 10^8\%$ with an increase in mg/m³ per day of silica exposure, respectively (P for both effect sizes were less than 0.001) (Figure 5). Given that more than one study is needed for the two-stage dose-response meta-analysis and only Mundt's study remained for HR, we could not perform a dose-response meta-analysis for this effect size.

Discussion

Although the relationship between exposure to silica and silicosis is straightforward, its magnitude has not been quantitatively investigated using a systematic review and meta-analysis. This study is the first attempt to quantify this association by a dose-response meta-analysis. According to this systematic review, there is a positive relationship between occupational silica exposure and silicosis mortality and incidence. Based on the dose-response analysis, with each mg/m³ increase in daily exposure to silica, the mortality and incidence rates of silicosis increase linearly.

The mechanism for the risk of silicosis among various occupational industries and factories is unknown. Experimental studies have suggested that the respirable form of silica is small enough to reach the terminal bronchioles and alveoli of the respiratory system. Standard immune responses are unable to remove these particles from the lungs,




Figure 2. Forrest plot of the association between occupational silica exposure and mortality from silicosis (a measure of association is Rate Ratio)




Figure 3. Forrest plot of the association between occupational silica exposure and silicosis incidence




Figure 4. Funnel plots of included studies by incidence and mortality studies

setting off a harmful cycle of inflammation and tissue damage that eventually results in silicosis (8, 21). On the other hand, increasing exposure to silica increases the silicosis mortality with high intensity and steep slope. According to the Centers for Disease Control and Prevention report, despite the significant progress and global efforts, silicosis death still occurs and is a public and occupational health problem. These deaths reflect exposure to silica particles that cause silicosis-related deaths (22). So, in order to reduce the terrible deaths attributed to silicosis in workplaces,

efforts should be made in cases where silicosis cannot be prevented, and the silicosis death risk factors should be identified to prevent deaths in patients with silicosis (23).

The findings of this study have important applications for occupational and public health because occupational silica exposure has long been a main universal occupational health problem. The findings of this study can be a basis to help direct public policies related to protective measures to keep workers in good health and safe working environments and away from other work activities responsible for




Figure 5. Dose-response analysis for the association between occupational silica exposure and silicosis mortality and incidence rates.

Table 1. Characteristic of the included studies

| ID | First author | Year | Country | Study design | Sample size | Age mean/range | Silica mean exposure | NOS Score | Quality |
|----|-----------------------|------|-------------------------|-----------------|-------------|----------------|----------------------|-----------|--------------|
| 1 | Carneiro, A. P (27) | 2006 | Brazil | Case-Control | 82 | 39.57 | N.R [©] | ★★★★★ | Low Quality |
| 2 | W. Chen (28) | 2005 | China | Cohort | 23002 | 42.18 | 0.395 | ★★★★★ | Low Quality |
| 3 | De Klerk (29) | 1998 | Australia | Cohort | 2297 | N.R | N.R | ★★★★★★★ | High Quality |
| 4 | M Dosemeci (30) | 1994 | USA | Case-Control | 1638 | N.R | 0.191 | ★★★★★★★ | High Quality |
| 5 | Kathleen Kreiss (31) | 1996 | USA | Cross-Sectional | 149 | 59.03 | N.R | ★★★★★ | Low Quality |
| 6 | Kenneth Mundt (32) | 2011 | Germany | Cohort | 17644 | N.R | 0.115 | ★★★★★★★ | High Quality |
| 7 | Kenneth Rosenman (25) | 1996 | USA | Cohort | 1072 | 58.90 | 0.171 | ★★★★★★★ | High Quality |
| 8 | Kyle Steenland (24) | 1995 | USA | Cohort | 3330 | N.R | 1.800 | ★★★★★★★ | High Quality |
| 9 | Yi Sun (33) | 2011 | China | Cohort | 3250 | 60.30 | 0.196 | ★★★★★ | Low Quality |
| 10 | Lap Ah Tse (34) | 2015 | China | Cohort | 3499 | 63.13 | 0.087 | ★★★★★ | Low Quality |
| 11 | Vacek PM (26) | 2019 | USA | Case-Control | 234 | 43.50 | 1.143 | ★★★★★★★ | High Quality |
| 12 | Dongming Wang (10) | 2020 | China | Cohort | 39808 | 45.30 | 2.734 | ★★★★★★★ | High Quality |
| 13 | Min Zhang (35) | 2010 | China | Cohort | 1300 | N.R | N.R | ★★★★★★★ | High Quality |
| 14 | G M Calvert (36) | 2003 | USA | Cross-Sectional | 7422 | N.R | 1.858 | ★★★★★★★ | High Quality |
| 15 | Michaela Kreuzer (37) | 2013 | Germany | Cohort | 58982 | N.R | 16.200 | ★★★★★★★ | Low Quality |
| 16 | Michaela Kreuzer (38) | 2021 | Germany | Cohort | 35204 | N.R | 22.625 | ★★★★★★★ | Low Quality |
| 17 | A 't Mannetje (39) | 2002 | USA, Finland, Australia | Cohort | 18364 | 68.70 | 10.632 | ★★★★★★★ | High Quality |
| 18 | Kyle Steenland (40) | 2001 | USA | Cohort | 5086 | N.R | 0.827 | ★★★★★ | Low Quality |
| 19 | Kyle Steenland (41) | 2005 | USA | Cohort | 65980 | N.R | 10.632 | ★★★★★★★ | High Quality |

©: Not reported.

lung disease (4).

The dose-response association between silica and silicosis mortality and incidence has been confirmed in several previous studies (10, 24-26); however, most previous studies focus on a single industry and evidence on different work settings and industries is limited. In this systematic review, various job environments from the metal mines, iron ore, and gold industry to ceramics and pottery workers, were examined and pooled.

Our study has some limitations. First, we did not calculate a dose-response analysis of silicosis incidence for the hazard ratio effect size because some studies did not report

silicosis cases at each level of exposure to silica dust. Finally, only one study remained for HR so we could not perform a dose-response meta-analysis for this measure. Second, high heterogeneity between studies is another limitation of this study, which can be attributed to the difference in study design, small number of included articles, different working environments, or differences in the mechanism of measuring exposure to silica dust in different studies. Although we performed meta-regression analysis to find the potential sources of heterogeneity, due to the small sample number of included studies, the heterogeneities remained high. The most important strength of this study was measuring the dose-response relationship of exposure to silica

on the incidence and mortality rates of silicosis with a systematic review study for the first time.

Conclusion

The results of this systematic review and meta-analysis propose that the risk of silicosis mortality and incidence increases as the level of silica exposure increases. This result confirms the linear association between occupational silica exposure and incidence and mortality rates from silicosis. The existence of this dose-response relationship is one of Hill's causal criteria and can strengthen the scientific background for strenuous occupational and industrial health interventions for the control of exposure to this dust.

Authors' Contributions

FSH and MJA conceived and designed the study. MFD and YS conducted the literature search and screening. FSH and MFD gathered the data. FSH conducted the statistical analysis. FSH and YS contributed to data interpretation. MFD and MJA drafted the manuscript, which FSH critically reviewed. All authors reviewed the final version before publication. FSH assumes full accountability for the accuracy and reliability of the data analysis and has unrestricted access to the study's data.

Ethical Considerations

This study has been approved by the Research Ethics Committee of Hamadan University of Medical Sciences with ethical ID: IR.UMSHA.REC.1403.401 and research ID: 140306275259.

Acknowledgment

We would like to appreciate the Vice-Chancellor for Research and Technology of the Hamadan University of Medical Sciences for supporting this work.

Conflict of Interests

The authors declare that they have no competing interests.

References

- Vanka KS, Shukla S, Gomez HM, James C, Palanisami T, Williams K, et al. Understanding the pathogenesis of occupational coal and silica dust-associated lung disease. *Eur Respir Rev*. 2022;31(165).
- Popescu FG, Stoia M, Morariu SI. A cross-national perspective on silicosis: incidence, occupational settings, and trends. *Romanian J. Occup Med*. 2023;74(1):6-11.
- Farooq S, Nawaz R, Nasim I, Irshad MA. Silicosis and Health Effects among Industrial Workers. *J Environ Agric Sci (JEAS)* Farooq et al. 2020;22(4):8-15.
- Rees D, Murray J. Silica, silicosis and tuberculosis. *Occup Health South Afr*. 2020;26(5):266-76.
- Liu X, Jiang Q, Wu P, Han L, Zhou P. Global incidence, prevalence and disease burden of silicosis: 30 years' overview and forecasted trends. *BMC Public Health*. 2023;23(1):1366.
- Global Burden of Diseases (GBD). Global Burden of Disease Study 2019 (GBD 2019) Data Resources 2024 [Available from: <https://vizhub.healthdata.org/gbd-results/>]
- Chen S, Liu M, Xie F. Global and national burden and trends of mortality and disability-adjusted life years for silicosis, from 1990 to 2019: results from the Global Burden of Disease study 2019. *BMC Pulm Med*. 2022;22(1):240.
- Adamcakova J, Mokra D. New insights into pathomechanisms and treatment possibilities for lung silicosis. *Int J Mol Sci*. 2021;22(8):4162.
- Kreuzer M, Deffner V, Schnelzer M, Fenske N. Mortality in Underground Miners in a Former Uranium Ore Mine: Results of a Cohort Study Among Former Employees of Wismut AG in Saxony and Thuringia. *Dtsch Arztebl Int*. 2021;118(4):41.
- Wang D, Zhou M, Liu Y, Ma J, Yang M, Shi T, et al. Comparison of Risk of Silicosis in Metal Mines and Pottery Factories: A 44-Year Cohort Study. *Chest*. 2020;158(3):1050-9.
- Ehrlich R, Akugizibwe P, Siegfried N, Rees D. The association between silica exposure, silicosis and tuberculosis: a systematic review and meta-analysis. *BMC Public Health*. 2021;21(1):953.
- Salum KCR, Castro MCS, Nani ÂSF, Kohlrausch FB. Is individual genetic susceptibility a link between silica exposure and development or severity of silicosis? A systematic review. *Inhal Toxicol*. 2020;32(9-10):375-87.
- Requena-Mullor M, Alarcón-Rodríguez R, Parrón-Carreño T, Martínez-López JJ, Lozano-Paniagua D, Hernández AF. Association between crystalline silica dust exposure and silicosis development in artificial stone workers. *Int J Environ Res Public Health*. 2021;18(11):5625.
- Shahbazi F, Morsali M, Poorolajal J. The effect of silica exposure on the risk of lung cancer: A dose-response meta-analysis. *Cancer Epidemiol*. 2021;75:102024.
- Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Ontario: Ottawa Hospital Research Institute; 2009 [cited 12 November 2018].
- Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Br Med J*. 2003;327(7414):557-60.
- Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics*. 1994;1088-101.
- Sterne J, RMJTSj H. Funnel plots in meta-analysis. *Stata J*. 2004;4(2):127-41.
- Harris RJ, Deeks JJ, Altman DG, Bradburn MJ, Harbord RM, Sterne JA. Metan: fixed-and random-effects meta-analysis. *Stata J*. 2008;8(1):3-28.
- Orsini N, Larsson SC, Salanti G. Dose-Response Meta-Analysis. *Systematic Reviews in Health Research: Meta-Analysis in Context*. 2022:258-69.
- Marrocco A, Ortiz LA. Role of metabolic reprogramming in pro-inflammatory cytokine secretion from LPS or silica-activated macrophages. *Front Immunol*. 2022;13:936167.
- Dhooria S, Sehgal IS, Agarwal R. Silica-associated lung disease in developing countries. *Curr Opin Pulm Med*. 2023;29(2):65-75.
- Hatman EA, Karagül DA, Oyman EK, Tüzün B, Şimşek KO, Kılıçaslan Z. Premature Deaths Due to Silicosis in Turkey, 2006-2017: A Twelve-Year Longitudinal Study. *Balkan Med J*. 2021;38(6):374.
- Steenland K, Brown D. Silicosis among gold miners: Exposure-response analyses and risk assessment. *Am J Public Health*. 1995;85(10):1372-7.
- Rosenman KD, Reilly MJ, Rice C, Hertzberg V, Tseng CY, Anderson HA. Silicosis among foundry workers. Implication for the need to revise the OSHA standard. *Am J Epidemiol*. 1996;144(9):890-900.
- Vacek PM, Glenn RE, Rando RJ, Parker JE, Kanne JP, Henry DA, et al. Exposure-response relationships for silicosis and its progression in industrial sand workers. *Scand J Work Environ Health*. 2019;45(3):280-8.
- Carneiro AP, Barreto SM, Siqueira AL, Cavariani F, Forastiere F. Continued exposure to silica after diagnosis of silicosis in Brazilian gold miners. *Am J Ind Med*. 2006;49(10):811-8.
- Chen W, Hnizdo E, Chen JQ, Attfield MD, Gao P, Hearl F, et al. Risk of silicosis in cohorts of Chinese tin and tungsten miners, and pottery workers (I): An epidemiological study. *Am J Ind Med*. 2005;48(1):1-9.
- de Klerk NH, Musk AW. Silica, compensated silicosis, and lung cancer in Western Australian goldminers. *Occup Environ Med*. 1998;55(4):243-8.
- Dosemeci M, McLaughlin JK, Chen JQ, Hearl F, McCawley M, Wu Z, et al. Indirect validation of a retrospective method of exposure assessment used in a nested case-control study of lung cancer and silica exposure. *Occup Environ Med*. 1994;51(2):136-8.
- Kreiss K, Zhen B. Risk of silicosis in a Colorado mining community. *Am J Ind Med*. 1996;30(5):529-39.
- Mundt KA, Birk T, Parsons W, Borsch-Galetke E, Siegmund K, Heavner K, et al. Respirable crystalline silica exposure-response evaluation of silicosis morbidity and lung cancer mortality in the German porcelain industry cohort. *J Occup Environ Med*.

- 2011;53(3):282-9.
33. Sun Y, Bochmann F, Morfeld P, Ulm K, Liu Y, Wang H, et al. Change of exposure response over time and long-term risk of silicosis among a cohort of Chinese pottery workers. *Int J Environ Res Public Health*. 2011;8(7):2923-36.
 34. Tse LA, Dai J, Chen M, Liu Y, Zhang H, Wong TW, et al. Prediction models and risk assessment for silicosis using a retrospective cohort study among workers exposed to silica in China. *Sci Rep*. 2015;5:11059.
 35. Zhang M, Zheng YD, Du XY, Lu Y, Li WJ, Qi C, et al. Silicosis in automobile foundry workers: a 29-year cohort study. *Biomed Environ Sci*. 2010;23(2):121-9.
 36. Calvert GM, Rice FL, Boiano JM, Sheehy JW, Sanderson WT. Occupational silica exposure and risk of various diseases: an analysis using death certificates from 27 states of the United States. *Occup Environ Med*. 2003;60(2):122-9.
 37. Kreuzer M, Sogl M, Bröske I, Möhner M, Nowak D, Schnelzer M, et al. Silica dust, radon and death from non-malignant respiratory diseases in German uranium miners. *Occup Environ Med*. 2013;70(12):869-75.
 38. Kreuzer M, Deffner V, Schnelzer M, Fenske N. Mortality in Underground Miners in a Former Uranium Ore Mine—Results of a Cohort Study Among Former Employees of Wismut AG in Saxony and Thuringia. *Dtsch Arztebl Int*. 2021;118(4):41-8.
 39. Mannetje At, Steenland K, Attfield M, Boffetta P, Checkoway H, DeKlerk N, et al. Exposure-response analysis and risk assessment for silica and silicosis mortality in a pooled analysis of six cohorts. *Occup Environ Med*. 2002;59(11):723-8.
 40. Steenland K, Sanderson W. Lung cancer among industrial sand workers exposed to crystalline silica. *Am J Epidemiol*. 2001;153(7):695-703.
 41. Steenland K. One agent, many diseases: Exposure-response data and comparative risks of different outcomes following silica exposure. *Am J Ind Med*. 2005;48(1):16-23.