

Activation of the Akt Survival Pathway Contributes to TRAIL Resistance in Cancer Cells

Jing Xu¹, Jun-Ying Zhou¹, Wei-Zen Wei², Gen Sheng Wu^{1*}

1 Program in Molecular Biology & Genetics, Karmanos Cancer Institute, Department of Pathology, Wayne State University School of Medicine, Detroit, Michigan, United States of America, **2** Program in Breast Cancer, Karmanos Cancer Institute, Department of Immunology and Microbiology, Wayne State University School of Medicine, Detroit, Michigan, United States of America

Abstract

The mechanism of tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) resistance in cancer cells is not fully understood. Here, we show that the Akt survival pathway plays an important role in TRAIL resistance in human cancer cells. Specifically, we found that TRAIL treatment activates the Akt survival pathway and that inhibition of this pathway by the PI3K inhibitor LY294002 or knockdown of Akt sensitizes resistant cancer cells to TRAIL. Since Akt is negatively regulated by the tumor suppressor PTEN, we examined the TRAIL sensitivity in PTEN knockdown mouse prostate epithelial cells and found that PTEN^{-/-} cells are more resistant than PTEN^{+/+} cells while the sensitivity of PTEN^{+/-} cells fell in between. Further, we showed that overexpression of a mutant PTEN confers TRAIL resistance in PTEN^{+/+} cells, supporting a role of PTEN in TRAIL sensitivity. In TRAIL resistant breast T47D cells, overexpression of the mutant PTEN further increased their resistance to TRAIL. Taken together, our data indicate that inactivation of functional PTEN and the consequent activation of the Akt pathway prevents TRAIL-induced apoptosis, leading to TRAIL resistance. Therefore, our results suggest that TRAIL resistance can be overcome by targeting PTEN or the Akt survival pathway in cancer cells.

Citation: Xu J, Zhou J-Y, Wei W-Z, Wu GS (2010) Activation of the Akt Survival Pathway Contributes to TRAIL Resistance in Cancer Cells. PLoS ONE 5(4): e10226. doi:10.1371/journal.pone.0010226

Editor: Dong-Yan Jin, University of Hong Kong, Hong Kong

Received: February 22, 2010; **Accepted:** March 19, 2010; **Published:** April 19, 2010

Copyright: © 2010 Xu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported in part by NIH/NCI grant R01 CA100073 and Department of Defense Breast Cancer Program W81XWH-07-1-05-21. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: wug@karmanos.org

Introduction

TRAIL (or Apo2L) is a member of the tumor necrosis factor superfamily that selectively induces apoptosis in cancer but not normal cells, making it an attractive agent for cancer therapy. TRAIL induces apoptosis through activation of its death receptors DR4 (TRAIL-R2) and/or DR5 (KILLER, TRAIL-R1, TRICK2) [1,2,3,4]. When TRAIL binds to DR4 and/or DR5 receptors, the death receptors become trimerized and then recruit the adaptor protein FADD (Fas-associated protein with death domain) and procaspase 8 or procaspase 10 to form the DISC, leading to caspase 8 activation [1,2]. Activated caspase 8 can directly activate the effector caspases 3, 6 and 7 to trigger apoptosis or indirectly induces apoptosis through the tBid-induced caspase 9 mediated mitochondrial apoptosis pathway [2,4].

Although TRAIL is an attractive therapeutic agent many cancer cells are resistant to this agent [2]. The mechanisms of TRAIL resistance are not fully understood, but may occur at multiple levels from its receptors to executor caspases within its signaling pathway [2,4]. At the receptor level, mutations in DR4 and DR5 have been found in some tumors including breast and lung cancers [4]. Increased expression of decoy receptors and decreased DR4/DR5 expression have been correlated with TRAIL resistance, although other studies indicated that the expression levels of TRAIL receptors do not correlate with TRAIL sensitivity. At the death-inducing signaling complex (DISC) level, c-FLIP is believed to be a major inhibitor for TRAIL signaling [5,6]. A recent study

showed that TRAIL sensitive non-small cell lung cancer (NSCLC) cells assemble DISC in lipid rafts of the plasma membrane that leads to caspase 8 activation, whereas non-raft DISC assembly leads to the recruitment of c-FLIP and RIP and the activation of the pro-survival signaling pathways such as NF-κB and ERK1/2 pathways [7]. In addition, it has been shown that anti-apoptotic members of the Bcl-2 family, including Bcl-2 and Mcl-1, are also involved in TRAIL resistance [8,9]. The activation of other survival pathways including the Akt and NF-κB pathways leads to cell survival and chemoresistance [10]. Consistently, it has been shown that the activation of the Akt pathway is closely associated with drug resistance including TRAIL resistance [11,12]. However, the exact mechanism by which the Akt pathway causes TRAIL resistance is not fully understood.

The PI3K/Akt signaling pathway is a prototypic survival pathway that plays a central role in diverse cellular functions, including proliferation, growth, survival, and metabolism [13]. Upon growth factor stimulation, PI3K is activated to phosphorylate phosphatidylinositol-3, 4-bisphosphate (PIP₂), generating phosphatidylinositol-3, 4, 5-triphosphate (PIP₃). PIP₃ binds to Akt, then translocates to the plasma membrane where Akt is activated by sequential phosphorylation at T308 and S473 residues. Phosphorylation at T308 is mediated by PDK-1 (3'-phosphoinositide-dependent kinase 1) while phosphorylation at S473 can be mediated by several kinases including PDK-1 and mTORC2 [13]. Once activated, Akt can phosphorylate many substrates to exert its functions. The best-characterized substrate of Akt is the serine/

threonine kinase mTOR (mammalian target of rapamycin). Akt can directly phosphorylate and activate mTOR and indirectly activate mTOR by phosphorylating and inactivating tuberous sclerosis complex 2 (TSC2). Activated mTOR forms a complex with Raptor to activate ribosomal protein S6 kinases (S6K) and inhibit 4E-BP, leading to increased protein translation [13]. It is known that activation of the PI3K/Akt/mTOR pathway can increase cell survival and apoptosis resistance induced by a number of agents [12,13]. For example, overexpression of Akt increases TRAIL resistance [11,14]. On the other hand, the tumor suppressor gene PTEN (phosphatase and tensin homolog deleted on chromosome ten) can dephosphorylate PIP3 to function as a negative regulator of PI3K/Akt/mTOR signaling. PTEN inactivation leads to the activation of the PI3K/Akt/mTOR signaling pathway. However, the exact mechanism by which the PI3K/Akt/mTOR pathway confers TRAIL resistance is not fully understood.

In this study, we demonstrated that increased activation of the Akt pathway plays an important role in TRAIL resistance. Specifically, we showed that TRAIL activates the Akt pathway and that blockade of Akt activation by the PI3K inhibitor LY294002 or knockdown of Akt expression by siRNA sensitizes resistant cells to TRAIL. We also found that reintroduction of wild-type PTEN into PTEN knockout cells sensitizes PTEN^{-/-} cells to TRAIL whereas inactivation of PTEN in PTEN^{+/+} and PTEN^{+/-} cells leads to TRAIL resistance. These results suggest that increased activation of the Akt pathway via inactivation of PTEN plays an important role in TRAIL resistance.

Materials and Methods

Reagents

LY294002 was purchased from Alexis Biochemicals (San Diego, CA). Recombinant TRAIL was purchased from PeproTech (Rocky Hill, NJ). Rabbit polyclonal antibodies against phosphorylated Akt (Ser473), phosphorylated mTOR, phosphorylated p70S6K, phosphorylated eIF4E, PTEN, total Akt, and poly (ADP-ribose) polymerase (PARP) were purchased from Cell Signaling Technology (Beverly, MA). Anti-actin antibody was purchased from Sigma. The adenoviruses expressing wild-type and dominant negative PTEN, as well as control adenoviruses expressing galactosidase (Ad-LacZ), were kindly provided by Dr. Mengjer Lee (University of Louisville).

Cell lines, culture conditions, and treatment

The human breast cancer T47D cells were obtained from American Type Culture Collection and maintained in DMEM with 10% fetal bovine serum (FBS). Human ovarian cancer SKOV3 cells were cultured in RPMI1640 with 10% FBS, OVCA432 cells were cultured in MCDB105/M199 medium with 10% FBS as described [15]. PTEN-knockout mouse prostate epithelial cells (PTEN^{-/-} and PTEN^{+/-}) and normal cells (PTEN^{+/+}) were obtained from Dr. Young Chen (Wake Forest University) and maintained in DMEM with 10% FBS, as described previously [16]. All cell lines were cultured at 37°C in a humidified atmosphere consisting of 5% CO₂ and 95% air.

siRNA transfection for knockdown of Akt

Signal Silence siRNA for Akt and corresponding control siRNA were purchased from Cell Signaling. The transfections were performed as suggested by the manufacturer with slight modifications, as described previously [17]. Briefly, T47D cells were plated at 6×10⁵ per well in six-well plates. The next day, cells were transfected with Akt or non-target control siRNAs using Oligofect-

mine (Invitrogen). After 24 h, transfected cells were trypsinized and plated at 6×10⁵ per well in six-well plates. 48 h later, transfected cells were left untreated or treated with TRAIL (100 ng/ml) for 24 h and then harvested for assessing the expression of Akt and cleaved PARP by Western blot analysis. To determine chemosensitivity, cells with or without transfection were placed at 8,000 per well in 96-well plates and then treated with TRAIL for 24 h, and cell viability was determined by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assays.

MTT assays

MTT assays were described previously [18].

Western blot analysis

The procedures for preparation of whole-cell protein lysates and Western blot analysis were described previously [18].

Adenovirus infections

PTEN^{+/+}, PTEN^{+/-} and PTEN^{-/-} mouse prostate epithelial cells and human breast cancer T47D cells were plated in 60 mm dishes at 8×10⁵, 6×10⁵, 5×10⁵ and 8×10⁵ cells per dish, respectively. The next day, cells were washed with normal medium and infected with adenoviruses expressing wild-type PTEN, dominant negative PTEN, or LacZ. During infection, cells were gently shaken every 10 min for 1 h, and then maintained in adenovirus containing medium at 37°C in a humidified atmosphere containing 5% CO₂. 24 h later, infected cells were switched to normal medium without adenoviruses and then continuously cultured for another 48 h. At 72 h after infection, cells were trypsinized and plated in 60 mm dish at 5×10⁵ cells. The next day, cells were left untreated or treated with TRAIL (100 ng/ml) for 48 h. Cells were harvested for Western blot analysis and the MTT assay, respectively.

Statistical analysis

Statistical analysis was done using Student's *t* test. The data were presented as the mean ± SD, and *P*≤0.05 was considered significant.

Results

Characterization of TRAIL sensitivity in a panel of breast and ovarian cancer cell lines

To understand the mechanisms of TRAIL resistance, we first investigated TRAIL sensitivity in 14 cancer cell lines including 3 breast and 11 ovarian cancer cell lines. These cell lines were treated with different doses of TRAIL for 24 h, and cell proliferation was measured by the MTT assay. Fig. 1 shows that these cell lines exhibited the differential TRAIL sensitivity; MDA-231, DOV13, OV433 and OVCA420 cells were sensitive to TRAIL whereas the rest of lines were resistant to TRAIL except MCF-7 cells that showed modest resistance. These data show that many ovarian and breast cancer cell lines are resistant to TRAIL.

The Akt/mTOR pathway is activated in TRAIL resistant cell lines

Since many breast and ovarian cancer cell lines are resistant to TRAIL, it is important to understand the underlying mechanism of TRAIL resistance. It has been reported that TRAIL can activate several pro-survival pathways, including Akt, NF-κB and ERK pathways [19,20,21,22]. Because activation of these pathways modulates cell survival and chemoresistance, it is conceivable that activation of these pathways contributes to




Figure 1. Effect of TRAIL on cell proliferation. Human breast cancer cell lines T47D, MDA231, and MCF7 and ovarian cancer cell lines DOV13, E52, OV90, OV433, OVCA432, RMG-1, TOV-21G, TOV-112D, CAOV3, OVCA420, and SKOV3 were treated with various doses of TRAIL for 24 h. Cell proliferation was determined by MTT assays. Cell proliferation data are expressed as percentage of untreated cells. Representative of three independent experiments. doi:10.1371/journal.pone.0010226.g001

TRAIL resistance. To test this possibility, we treated four resistant lines T47D, SKOV3, OVCA432 and OV90 with 100 ng/ml TRAIL for different periods of time and examined the activation of the Akt, NF- κ B and ERK pathways, as well as expression of the Bcl-2 family proteins. As shown in Fig. 2A–D, the levels of both phosphorylated Akt and mTOR protein were increased as early as 2 h and lasted up to 6 h after TRAIL treatment. We also showed that such activation was not due to an increase in the total Akt and mTOR proteins, which remained constant (Fig. 2A–D). Moreover, we showed that p70S6K and eIF4E, both are downstream targets of mTOR, are activated (Fig. 2A and D). These data indicate that TRAIL can activate the Akt/mTOR pathway in TRAIL resistant cell lines. Importantly, we showed that Akt and mTOR are not activated in TRAIL sensitive MCF-7 and OVCA420 cells (Fig. 2E and F). Thus, our results suggest that activation of Akt/mTOR pathway is a common event in TRAIL resistant cancer cell lines and that this pathway may be important in TRAIL resistance. In addition, we found that TRAIL could activate the MEK/ERK and NF- κ B pathways in OV90 cells, but not in other three cell lines (data not shown). Furthermore, inhibition of the ERK and NF- κ B pathways by their corresponding pharmacological inhibitors did not affect TRAIL sensitivity in OV90 cells (data not shown), indicating that the activation of these two pathways may not be important in TRAIL resistance in this cell line.




Figure 2. Activation of Akt/mTOR pathway by TRAIL in resistant cell lines but not in sensitive cell lines. T47D (A), SKOV3 (B), OVCA432 (C), OV90 (D), MCF7 (E) and OVCA420 (F) cells were treated with 100 ng/ml TRAIL for 0, 2, 4 and 8 h (in T47D cells, 6 h treatment was included), and total protein was extracted. The levels of total and phosphorylated Akt (*p-Akt*), and phosphorylated mTOR (*p-mTOR*) were determined by Western blot analysis. In T47D (A) and OV90 (D) cells, the levels of phosphorylated p-p70S6K and phosphorylated eIF4E (*p-eIF4E*) were also examined. β -actin was used as a loading control. Of note, there were two bands for Akt (Akt1 and Akt2) in both OVCA432 and OV90 cells while only one band for Akt was detected in T47D and SKOV3 cells. doi:10.1371/journal.pone.0010226.g002

Inhibition of Akt activation sensitizes resistant cells to TRAIL

Since the Akt pathway was activated in all resistant cell lines tested, we asked whether inhibition of Akt activity could increase their TRAIL sensitivity in resistant cells. To this end, we treated T47D cells with the PI3K inhibitor LY294002 and examined the effect of Akt inhibition on TRAIL-induced apoptosis. As shown in Fig. 3, TRAIL treatment increased Akt phosphorylation in T47D cells, which was abolished by LY294002, showing that Akt activation is PI3K dependent. Similarly, we also showed that activation of Akt is blocked by LY294002 in OVCA432 cells (Fig. 3C). Importantly, inhibition of Akt phosphorylation by LY294002 significantly increased TRAIL-induced PARP cleavage as compared to cells treated with TRAIL, or LY294002 alone (Fig. 3A and C). Furthermore, we showed that combined treatment of TRAIL and LY294002 significantly inhibits cell proliferation as compared to either agent alone (Fig. 3B and D). These results strongly suggest that the activation of Akt plays an important role in TRAIL resistance in cancer cells.

Knockdown of Akt sensitizes breast cancer T47D cells to TRAIL

Although treatment with the PI3K inhibitor LY294002 increased TRAIL-induced apoptosis, the results may not com-

pletely reflect the exclusive role of Akt in TRAIL resistance because LY294002 may inhibit other kinases. To investigate the direct role of Akt activation in TRAIL resistance, we used siRNA to knockdown Akt expression and then determine the effect of Akt knockdown on TRAIL sensitivity. We transfected T47D cells with siRNA against Akt (Akt 1, 2 and 3) or control siRNA and showed that Akt was efficiently knock downed by Akt siRNA in cells with or without TRAIL treatment, as compared to cells transfected with control siRNA (Fig. 4B). Importantly, we showed that knockdown of Akt sensitized TRAIL induced PARP cleavage as compared to the same cells transfected with control siRNA (Fig. 4B). Further, knockdown of Akt increased TRAIL-induced growth inhibition while such changes were minimal in cells transfected with control siRNA (Fig. 4A). Thus, our results indicate that while TRAIL can cause apoptosis, it also activates the Akt survival pathway to counteract TRAIL-induced apoptosis.

Loss of PTEN confers TRAIL resistance

It has been known that PTEN is a negative regulator of Akt and that increased Akt activity has been associated with inactivation of PTEN in several cancers. To understand the mechanism of Akt activation-mediated TRAIL resistance, we treated PTEN^{+/+}, PTEN^{+/-} and PTEN^{-/-} mouse prostate epithelial cells with TRAIL for 48 h and measured cell proliferation by MTT assays.




Figure 3. Inhibition of Akt activity sensitizes resistant cancer cells to TRAIL. A and C, effect of the PI3K inhibitor LY294002 on TRAIL-induced apoptosis. Breast cancer cell T47D (A) and ovarian cancer cell OVCA432 (C) were left untreated or pretreated with 10 μ M LY294002 (LY) for 30 min, and then treated with or without 100 ng/ml TRAIL for 6 h. Total protein was extracted, and cleaved PARP and total and phosphorylated Akt were determined by Western blotting. B and D, effect of LY294002 on TRAIL-induced growth inhibition. T47D (B) and OVCA432 cells (D) were left untreated or pretreated with 10 μ M LY294002 for 30 min, and then treated with or without 100 ng/ml TRAIL for 24 h. Cell proliferation was determined by MTT assays and expressed as percentage of untreated cells. Representative of three independent experiments. **, $P < 0.001$, statistical significance; *, $P < 0.01$, statistical significance.

doi:10.1371/journal.pone.0010226.g003




Figure 4. Knockdown of Akt sensitizes breast cancer T47D cells to TRAIL. *A*, effect of Akt knockdown on cell proliferation. T47D cells were plated at 6×10^5 per well in six-well plates. The next day, cells were transfected with Akt or control siRNAs using Oligofectamine. After 2 d, cells were left untreated or treated with TRAIL (100 ng/ml) for 24 h, and cell proliferation was determined by MTT assays. Cell proliferation data are expressed as percentage of untreated cells. Representative of three independent experiments. *, $P < 0.01$, statistical significance. *B*, effect of Akt knockdown on TRAIL-induced apoptosis. T47D cells were transfected with Akt or control siRNAs and then treated with or without TRAIL (100 ng/ml) for 24 h as described in (A). Total protein was extracted for assaying total and phosphorylated Akt (p-Akt) and PARP by Western blot analysis. β -actin was used as a loading control.

doi:10.1371/journal.pone.0010226.g004

Fig. 5A shows that TRAIL-induced about 32% growth inhibition in $PTEN^{+/+}$ cells whereas TRAIL did not affect cell proliferation in $PTEN^{-/-}$ cells, while $PTEN^{+/-}$ cells showed about 13% growth inhibition after TRAIL treatment. Furthermore, we showed that Akt is activated in $PTEN^{-/-}$ cells but not in $PTEN^{+/+}$, while minimal activation in $PTEN^{+/-}$ cells. Importantly, TRAIL treatment caused significant PARP cleavage in $PTEN^{+/+}$ cells, moderate cleavage in $PTEN^{+/-}$ cells and complete absence of cleavage in $PTEN^{-/-}$ cells (Fig. 5B). These results suggest that PTEN is required for TRAIL-induced apoptosis in mouse prostate epithelial cells.

Since $PTEN^{-/-}$ cells are resistant to TRAIL, we asked whether restoration of PTEN expression affects TRAIL sensitivity in $PTEN^{-/-}$ cells. $PTEN^{-/-}$ cells were infected with adenoviruses expressing wild-type PTEN or LacZ. As shown in Fig. 5C, PTEN expression was restored in cells infected with adenoviruses expressing PTEN but not in cells infected with adenoviruses expressing LacZ. Importantly, we were able to show that

restoration of PTEN expression renders $PTEN^{-/-}$ cells sensitive to TRAIL induced apoptosis and growth inhibition whereas such effects were not observed in cells infected with control viruses (Fig. 5C). Since $PTEN^{+/+}$ cells are sensitive to TRAIL, we asked whether inhibition of PTEN function affects TRAIL-induced apoptosis in $PTEN^{+/+}$ cells. To this end, we infected $PTEN^{+/+}$ and $PTEN^{+/-}$ cells with adenoviruses expressing a dominant negative form of PTEN or LacZ, and treated infected cells with TRAIL for 48 h. Fig. 5D shows that introduction of dominant negative form of PTEN increased p-Akt level in $PTEN^{+/+}$ cells. Importantly, introduction of a dominative negative form of PTEN renders $PTEN^{+/+}$ and $PTEN^{+/-}$ cells more resistant to TRAIL (Fig. 5D). Consistent with the role of Akt in inhibiting apoptosis, TRAIL induced PARP cleavage was decreased dramatically in $PTEN^{+/+}$ and $PTEN^{+/-}$ cells after they were infected with viruses expressing a dominant negative form of PTEN. Furthermore, we showed that TRAIL-induced growth inhibition in $PTEN^{+/+}$ cells infected with




Figure 5. The role of PTEN in TRAIL sensitivity. *A*, effect of loss of PTEN in TRAIL-induced growth inhibition. Mouse prostate epithelial cells with wild type PTEN (PTEN^{+/+}), PTEN knockout (PTEN^{-/-}) and heterozygous (PTEN^{+/-}) were plated in 96-well plates, and then treated with 100 ng/ml TRAIL for 48 h. Cells proliferation was determined by MTT assays and expressed as percentage of untreated cells. Representative of three independent experiments. *, $P < 0.01$, statistical significance. *B*, effect of PTEN loss on TRAIL-induced apoptosis. PTEN^{+/+}, PTEN^{+/-}, and PTEN^{-/-} cells were treated with 100 ng/ml TRAIL for 48 h as described in (A). Total protein was extracted for assaying total and phosphorylated Akt (p-AKT), PTEN and PARP by Western blot analysis. β -actin was used as a loading control. *C*, restored PTEN expression sensitizes PTEN^{-/-} cells to TRAIL. PTEN^{-/-} cells (5×10^5) were infected with adenoviruses expressing wild-type PTEN (Ad-PTEN wt) or expressing LacZ (Ad-lacZ). At 72 h after infection, cells were trypsinized and plated in 60 mm dish at 5×10^5 cells. The next day, cells were left untreated or treated with TRAIL (100 ng/ml) for 48 h. Cells were harvested for MTT assays (Upper panel) and Western blot analysis was employed for detecting the levels of PARP and PTEN proteins (lower panel), respectively. Cell proliferation data were expressed as percentage of untreated cells. Representative of three independent experiments. **, $P < 0.001$, statistical significance. *D*, inhibition of PTEN function by a dominant negative form of PTEN confers TRAIL resistance. PTEN^{+/+} and PTEN^{+/-} cells were infected with adenoviruses expressing a dominant negative form of PTEN for 72 h and then treated with TRAIL (100 ng/ml) for 48 h. Cells were either used for growth inhibition by MTT assays (upper panel) or harvested for detecting the levels of PARP, PTEN, and total and phosphorylated Akt by Western blot analysis (lower panel). Cell proliferation data were expressed as percentage of untreated cells. Representative of three independent experiments. *, $P < 0.01$, statistical significance. **, $P < 0.001$, statistical significance. doi:10.1371/journal.pone.0010226.g005

an adenovirus expressing a dominant negative form of PTEN was about 5% but about 27% in the same cells infected with control adenoviruses (Fig. 5D). We also showed that in PTEN^{+/-} cells infected with dominant negative form of PTEN, cell proliferation was about 102% as compared to the same cells infected with control viruses in which cell proliferation was about 81% upon TRAIL treatment (Fig. 5D, upper panel).

The role of PTEN in TRAIL sensitivity in breast cancer T47D cells

Since breast cancer T47D cells with a wild-type PTEN were resistant to TRAIL (Fig. 1), we asked whether modulation of

PTEN function would affect TRAIL sensitivity in T47D cells. To this end, we infected T47D cells with adenoviruses expressing a dominant negative form of PTEN or control viruses and then treated infected cells with TRAIL (100 ng/ml) for 48 h, and the effect of such treatment on TRAIL sensitivity was determined. Fig. 6 shows that cells infected with adenovirus expressing a dominant negative form of PTEN express a higher level of p-Akt while such an increase was not observed in cells infected with adenoviruses expressing LacZ, which suggests that PTEN function was inhibited by a dominant negative form of PTEN (Fig. 6B). Consistent with the role of Akt in inhibiting apoptosis, cleaved PARP was significantly reduced in cells expressing a dominant

negative form of PTEN as compared to cells expressing control viruses (Fig. 6B). Furthermore, overexpression of a dominant negative form of PTEN renders T47D cells more resistant to TRAIL as compared to the same cells infected with viruses expressing LacZ (Fig. 6). Therefore, these data suggest that increased Akt activation due to inactivation of PTEN could confer TRAIL resistance in cancer cells.

Discussion

In this study, we show that TRAIL activates the Akt survival pathway in TRAIL-resistant cancer cell lines. We also show that the underlying mechanism may be due to loss of functional PTEN because PTEN knockout mouse prostate epithelial cells are resistant to TRAIL while cells with intact PTEN are sensitive to TRAIL. Restoring PTEN expression rendered PTEN^{-/-} cells

sensitive to TRAIL. Further, inactivation of PTEN enhances the level of TRAIL resistance in T47D cells. These findings suggest that activation of the Akt survival pathway plays a critical role in TRAIL resistance and that the Akt pathway is a potential therapeutic target to enhance TRAIL-based therapy.

The activation of the Akt pathway has been implicated in protecting cells by many death stimuli, thereby conferring cell survival [10,13]. Previous studies suggested that overexpression of Akt and its upstream regulator PI3K increased TRAIL resistance in breast and ovarian cancer cells [23,24]. However, the underlying mechanism of its resistance is not fully understood. In this study, we showed that while TRAIL induces apoptosis in cancer cells, it also activates several survival pathways, which may counteract TRAIL-induced apoptosis, leading to resistance. In many TRAIL resistant breast and ovarian cancer cell lines but not sensitive cell lines, the Akt pathway was activated (Fig. 2), suggesting that the activation of the Akt pathway may be a common event in TRAIL resistant cells. Consistent with this notion, we found that inhibition of Akt activation by its pharmacological inhibitor LY294002 or knockdown of its expression by siRNA sensitizes TRAIL-resistant cells to TRAIL. Collectively, these data strongly suggest that activation of the Akt survival pathway plays a critical role in TRAIL resistance in ovarian and breast cancer cells and imply that inhibition of this survival pathway might overcome TRAIL resistance.

We also showed that inhibition of PI3K activity blocked Akt activation in TRAIL resistant cell lines. Since PI3K can serve as an upstream positive regulator of Akt, these results suggest that increased Akt activation is at least in part through the increased activation of PI3K activity. At the downstream level, it is well known that Akt exerts its biological functions by phosphorylating its downstream substrates including mTOR. Activation of mTOR can increase protein translation, resulting in cell survival [13]. Consistent with this, we found that TRAIL treatment can cause mTOR phosphorylation and subsequent activation. Our data suggest that TRAIL may activate the PI3K/Akt/mTOR pathway to counteract TRAIL-induced apoptosis, leading to TRAIL resistance.

In cancer cells, the Akt survival pathway can be activated by the several mechanisms including inactivation of its upstream negative regulator PTEN. By dephosphorylating and converting PIP3 to PIP2, PTEN inhibits Akt activation and subsequently sensitizes cells to death [10,13]. Consistent with the role of the PI3K/Akt pathway in TRAIL resistance, we found that loss of PTEN in mouse prostate epithelial cells confers TRAIL resistance whereas the same cells with intact PTEN are sensitive to TRAIL while PTEN^{+/-} cells show intermediate resistance. These data suggest that PTEN indeed plays an important role in TRAIL sensitivity. Furthermore, we showed that reintroduction of a mutant form of PTEN into TRAIL resistant T47D cells further increases TRAIL resistance. Collectively, these data indicate that in TRAIL resistant cells, loss of PTEN contributes to TRAIL resistance. It has been known that decreased or loss of PTEN expression occurs in at least 50% of breast and ovarian cancers [25]. It is possible that TRAIL resistant breast and ovarian cancer cells have decreased levels of PTEN, which need further investigation. Further, a recent study showed that overexpression of miR-221&222 down regulates PTEN, leading to increased activation of Akt and subsequent TRAIL resistance [26]. Additionally, TRAIL resistant leukemia cells expressed a higher level of phosphorylated inactive form of PTEN, which resulted in an increase in Akt activation and TRAIL resistance [27]. In addition to the role of PTEN in regulating Akt activation, several studies have shown that Akt activation by TRAIL treatment can be due to the other mechanisms. For




Figure 6. Inhibition of PTEN function renders breast cancer T47D cells more resistant to TRAIL. A, effect of inhibition of PTEN function on cell proliferation. T47D cells (8×10^5) were infected with adenovirus expressing a dominant negative form of PTEN or adenoviruses expressing LacZ for 72 h and then treated with TRAIL (100 ng/ml) for 48 h. Cell proliferation was determined by MTT assays. Cell proliferation data are expressed as percentage of untreated cells. Representative of three independent experiments. *, $P < 0.01$, statistical significance. B, effect of inhibition of PTEN function on apoptosis. T47D cells infected with adenoviruses as described in A were treated with TRAIL (100 ng/ml) for 48 h and then harvested for analyzing the levels of PARP, PTEN, total and phosphorylated Akt proteins by Western blot analysis (lower panel). β -actin was used as a loading control. doi:10.1371/journal.pone.0010226.g006

example, TRAIL was shown to activate Src, leading to Akt activation and TRAIL resistance [28]. Another mechanism involves DNA-PKcs-mediated Akt activation [29]. Nevertheless, the mechanisms by which TRAIL activates Akt are still not fully understood.

Although TRAIL selectively kills cancer cells but not normal cells, many cancer cells including breast and ovarian cancer cells are resistant to TRAIL [30,31]. Accumulating evidence indicates that TRAIL resistance involves many molecules and pathways including PI3K/Akt pathway [4]. Although previous studies showed that overexpression or constitutive expression of Akt confers TRAIL resistance in cancer cells [11,14], this study is unique because our work indicates that inducible activation of the Akt/mTOR survival pathway counteracts TRAIL-induced apoptosis, leading to TRAIL resistance, as opposed to previous studies in which overexpression or constitutive activation of Akt caused TRAIL resistance [11,14]. Our conclusion is based on the fact that Akt and its downstream target mTOR are activated in all TRAIL resistant cell lines tested. Further, we were able to increase TRAIL sensitivity by inhibiting the Akt pathway using its pharmacological inhibitor LY294002 or siRNA against Akt.

References

- Almasan A, Ashkenazi A (2003) Apo2L/TRAIL: apoptosis signaling, biology, and potential for cancer therapy. *Cytokine Growth Factor Rev* 14: 337–348.
- Jin Z, El-Deiry WS (2005) Overview of cell death signaling pathways. *Cancer Biol Ther* 4: 139–163.
- Wu GS, Burns TF, McDonald ER, Jiang W, Meng R, et al. (1997) KILLER/DR5 is a DNA damage-inducible p53-regulated death receptor gene. *Nat Genet* 17: 141–143.
- Wu GS (2009) TRAIL as a target in anti-cancer therapy. *Cancer Letters* 285: 1–5.
- Burns TF, El-Deiry WS (2001) Identification of inhibitors of TRAIL-induced death (ITIDs) in the TRAIL-sensitive colon carcinoma cell line SW480 using a genetic approach. *J Biol Chem* 276: 37879–37886.
- Hietakangas V, Poukkula M, Heiskanen KM, Karvinen JT, Sistonen L, et al. (2003) Erythroid differentiation sensitizes K562 leukemia cells to TRAIL-induced apoptosis by downregulation of c-FLIP. *Mol Cell Biol* 23: 1278–1291.
- Song JH, Tse MC, Bellail A, Phuphanich S, Khuri F, et al. (2007) Lipid rafts and nonrafts mediate tumor necrosis factor related apoptosis-inducing ligand induced apoptotic and nonapoptotic signals in non small cell lung carcinoma cells. *Cancer Res* 67: 6946–6955.
- Sun SY, Yue P, Zhou J-Y, Wang Y, Kim HC, et al. (2001) Overexpression of bcl2 blocks TNF-related apoptosis-inducing ligand (TRAIL)-induced apoptosis in human lung cancer cells. *BBRC* 280: 788–797.
- Ricci MS, Kim SH, Ogi K, Plastaras JP, Ling J, et al. (2007) Reduction of TRAIL-induced Mcl-1 and cIAP2 by c-Myc or sorafenib sensitizes resistant human cancer cells to TRAIL-induced death. *Cancer Cell* 12: 66–80.
- LoPiccolo J, Blumenthal GM, Bernstein WB, Dennis PA (2008) Targeting the PI3K/Akt/mTOR pathway: effective combinations and clinical considerations. *Drug Resist Updat* 11: 32–50.
- Kandasamy K, Srivastava RK (2002) Role of the phosphatidylinositol 3'-kinase/PTEN/Akt kinase pathway in tumor necrosis factor-related apoptosis-inducing ligand-induced apoptosis in non-small cell lung cancer cells. *Cancer Res* 62: 4929–4937.
- Plastaras JP, Dorsey JF, Carroll K, Kim SH, Birnbaum MJ, et al. (2008) Role of PI3K/Akt signaling in TRAIL- and radiation-induced gastrointestinal apoptosis. *Cancer Biol Ther* 7: 2047–2053.
- Hay N (2005) The Akt-mTOR tango and its relevance to cancer. *Cancer Cell* 8: 179–183.
- Nesterov A, Lu X, Johnson M, Miller GJ, Ivashchenko Y, et al. (2001) Elevated AKT activity protects the prostate cancer cell line LNCaP from TRAIL-induced apoptosis. *J Biol Chem* 276: 10767–10774.
- Wang J, Zhou JY, Zhang L, Wu GS (2009) Involvement of MKP-1 and Bcl-2 in acquired cisplatin resistance in ovarian cancer cells. *Cell Cycle* 8: 3191–3198.
- Berquin IM, Min Y, Wu R, Wu J, Perry D, et al. (2007) Modulation of prostate cancer genetic risk by omega-3 and omega-6 fatty acids. *J Clin Invest* 117: 1866–1875.
- Xu J, Zhou JY, Tainsky MA, Wu GS (2007) Evidence that tumor necrosis factor-related apoptosis-inducing ligand induction by 5-aza-2'-deoxycytidine sensitizes human breast cancer cells to adriamycin. *Cancer Res* 67: 1203–1211.
- Wang Z, Xu J, Zhou JY, Liu Y, Wu GS (2006) Mitogen-activated protein kinase phosphatase-1 is required for cisplatin resistance. *Cancer Res* 66: 8870–8877.
- Trauzold A, Wermann H, Arlt A, Schutze S, Schafer H, et al. (2001) CD95 and TRAIL receptor-mediated activation of protein kinase C and NF-kappaB contributes to apoptosis resistance in ductal pancreatic adenocarcinoma cells. *Oncogene* 20: 4258–4269.
- Ehrhardt H, Fulda S, Schmid I, Hiscott J, Debatin KM, et al. (2003) TRAIL-induced survival and proliferation in cancer cells resistant towards TRAIL-induced apoptosis mediated by NF-kappaB. *Oncogene* 22: 3842–3852.
- Secchiero P, Gonelli A, Carnevale E, Milani D, Pandolfi A, et al. (2003) TRAIL promotes the survival and proliferation of primary human vascular endothelial cells by activating the Akt and ERK pathways. *Circulation* 107: 2250–2256.
- Secchiero P, Zerbinati C, Rimondi E, Corallini F, Milani D, et al. (2004) TRAIL promotes the survival, migration and proliferation of vascular smooth muscle cells. *Cell Mol Life Sci* 61: 1965–1974.
- Cuello M, Ettenberg SA, Clark AS, Keane MM, Posner RH, et al. (2001) Down-regulation of the erbB-2 receptor by trastuzumab (herceptin) enhances tumor necrosis factor-related apoptosis-inducing ligand-mediated apoptosis in breast and ovarian cancer cell lines that overexpress erbB-2. *Cancer Res* 61: 4892–4900.
- Lane D, Robert V, Grondin R, Rancourt C, Piche A (2007) Malignant ascites protect against TRAIL-induced apoptosis by activating the PI3K/Akt pathway in human ovarian carcinoma cells. *Int J Cancer* 121: 1227–1237.
- Brugge J, Hung MC, Mills GB (2007) A new mutational AKTivation in the PI3K pathway. *Cancer Cell* 12: 104–107.
- Garofalo M, Di Leva G, Romano G, Nuovo G, Suh SS, et al. (2009) miR-221&222 regulate TRAIL resistance and enhance tumorigenicity through PTEN and TIMP3 downregulation. *Cancer Cell* 16: 498–509.
- Dida F, Li Y, Iwao A, Deguchi T, Azuma E, et al. (2008) Resistance to TRAIL-induced apoptosis caused by constitutational phosphorylation of Akt and PTEN in acute lymphoblastic leukemia cells. *Exp Hematol* 36: 1343–1353.
- Song JJ, Kim JH, Sun BK, Alcalá MA, Jr., Bartlett DL, et al. (2010) c-Cbl acts as a mediator of Src-induced activation of the PI3K-Akt signal transduction pathway during TRAIL treatment. *Cell Signal* 22: 377–385.
- Kim MJ, Kim HB, Bac JH, Lee JW, Park SJ, et al. (2009) Sensitization of human K562 leukemic cells to TRAIL-induced apoptosis by inhibiting the DNA-PKcs/Akt-mediated cell survival pathway. *Biochem Pharmacol* 78: 573–582.
- Keane MM, Ettenberg SA, Nau MM, Russell EK, Lipkowitz S (1999) Chemotherapy augments TRAIL-induced apoptosis in breast cell lines. *Cancer Res* 59: 734–741.
- Cuello M, Ettenberg SA, Nau MM, Lipkowitz S (2001) Synergistic induction of apoptosis by the combination of trail and chemotherapy in chemoresistant ovarian cancer cells. *Gynecol Oncol* 81: 380–390.

Acknowledgments

We thank Dr. Yong Chen for providing PTEN knockout mouse prostate epithelial cells and Dr. Mengjer Lee for providing adenoviruses expressing PTEN.

Author Contributions

Conceived and designed the experiments: JX GSW. Performed the experiments: JX JYZ. Analyzed the data: JX GSW. Contributed reagents/materials/analysis tools: JX JYZ WZW GSW. Wrote the paper: JX WZW GSW.