

Morpho-molecular analyses reveal two novel species and two new records of Dictyosporiaceae (Pleosporales) from Dulongjiang River in northwestern Yunnan Province, China

Ying Wang¹, D. Jayarama Bhat^{2,3}, Dan-Feng Bao⁴, Hong-Wei Shen¹, Yuan Feng¹, Zong-Long Luo¹

¹ College of Agriculture and Biological Science, Dali University, Dali 671003, Yunnan, China

² Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

³ Vishnugupta Vishwavidyapeetam, Ashoke, Gokarna 581326, India

⁴ Engineering and Research Center for Southwest Biopharmaceutical Resource of National Education Ministry of China, Guizhou University, Guiyang, 550025, Guizhou, China

Corresponding author: Zong-Long Luo (luozonglongfungi@163.com)



This article is part of:

**Exploring the Hidden Fungal Diversity:
Biodiversity, Taxonomy, and Phylogeny of
Saprobic Fungi**

Edited by Samantha C. Karunarathna,
Danushka Sandaruwan Tennakoon,
Ajay Kumar Gautam

Academic editor:

Danushka Sandaruwan Tennakoon

Received: 30 December 2024

Accepted: 28 March 2025

Published: 30 April 2025

Citation: Wang Y, Bhat DJ, Bao D-F, Shen H-W, Feng Y, Luo Z-L (2025) Morpho-molecular analyses reveal two novel species and two new records of Dictyosporiaceae (Pleosporales) from Dulongjiang River in northwestern Yunnan Province, China. MycoKeys 117: 1–27. <https://doi.org/10.3897/mycokeys.117.145587>

Copyright: © Ying Wang et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Abstract

Yunnan Province is rich in freshwater resources, providing a favorable environment for freshwater fungi. As a result, numerous new species have been described in recent years, further supporting the extensive diversity of freshwater fungi in this region. During an investigation of lignicolous freshwater fungi from the Dulongjiang River, six fresh isolates of Dictyosporiaceae were obtained. The new collection includes two new species (*Dictyocheirospora splendida* and *Jalapriya guttulata*), two new records (*D. nabanheensis* and *Dictyosporium hughesii*), and one known species (*Dictyosporium duliujiangense*). They are introduced based on morphological evidence and molecular phylogenetic analysis of combined ITS, LSU, and *tef1-α* sequence data. Detailed descriptions and illustrations of these species are provided, and morphological comparison with similar taxa is discussed.

Key words: Dothideomycetes, lignicolous freshwater fungi, new species, phylogeny, taxonomy

Introduction

Lignicolous freshwater fungi are key decomposers in freshwater ecosystems (Rebeca et al. 2021; Tamam et al. 2021). Recent studies have shown that freshwater fungi globally encompass 13 phyla in the Kingdom Fungi (Calabon et al. 2022). Ascomycota as the dominant group, and Sorsariomycetes leading, followed by Dothideomycetes, is a trend observed in China, especially in Yunnan (Calabon et al. 2022; Shen et al. 2022a). Dothideomycetes represents a major taxonomic group, characterized by bitunicate asci, typically with fissitunicate dehiscence, and many species have been collected from freshwater habitats (Dong et al. 2020). Recent data indicated that freshwater dothideomycetous species reside in six orders, 43 families, and 145 genera, with 46 genera exclusively documented from freshwater environments (Dong et al. 2020; Calabon et al. 2022, 2023).

Boonmee et al. (2016) established Dictyosporiaceae to accommodate most cheirosporous hyphomycetous genera, with *Dictyosporium* as the type genus. Tennakoon et al. (2023) documented that the Dictyosporiaceae comprises 154 species. However, *Dictyocheiropora himachalensis* was not included in the revised version (Sushma et al. 2022). Presently, the species number in Dictyosporiaceae has increased to 178 (Phookamsak et al. 2019; Dong et al. 2023; Liu et al. 2023; Zhang et al. 2023; Feng et al. 2024; Han et al. 2024; Liu et al. 2024; Phookamsak et al. 2024; Shu et al. 2024; Sun et al. 2024; Wang et al. 2024; Wu et al. 2024; Zhang et al. 2024a; Zhang et al. 2024b; Du et al. 2025). These taxa subsist as saprobes on decaying wood and plant debris in aquatic and terrestrial habitats worldwide (Boonmee et al. 2016; Fu et al. 2021; Shen et al. 2022b). Currently, Dictyosporiaceae comprises 23 genera, with most of the asexual morphic members of this family being hyphomycetous and characterized by cheiroid, digitate, palmate, and/or dictyosporous, and pale brown to dark brown conidia. These include the following: *Aquadictyospora*, *Aquaticheirospora*, *Cheirosporum*, *Dendryphiella*, *Dictyocheiropora*, *Dictyopalmispora*, *Dictyosporium*, *Digitodesmium*, *Jalapriya*, *Kamatia*, *Neodendryphiella*, *Neodigitodesmium*, *Neogregarithecium*, *Paradictyocheiropora*, *Pseudodictyosporium*, and *Vikalpa*, and six coelomycetous genera, viz., *Immotthia*, *Pseudocoleophoma*, *Pseudoconiothyrium*, *Pseudocyclothyriella*, *Sajamaea*, and *Verrucocum*. Whereas five genera are known to have sexual morphs, viz., *Dictyosporium*, *Gregarithecium*, *Immotthia*, *Pseudocoleophoma*, and *Verrucocum*. Notably, *Dictyosporium* exhibits both sexual and asexual hyphomycetous morphs, while *Immotthia*, *Pseudocoleophoma*, and *Verrucocum* display sexual and asexual coelomycetous morphs (<https://www.outlineoffungi.org/>; Boonmee et al. 2016; Jiang et al. 2021; Shen et al. 2022b; Wijayawardene et al. 2022; Tennakoon et al. 2023; Shu et al. 2024).

Dictyocheiropora was established by Boonmee et al. (2016) with *D. rotunda* as the type species. This genus is characterized by dark sporodochial colonies that produce aeroaquatic cheiroid, doliiform to cylindrical conidiogenous cells, with or without hooked apices, with or without appendages, and non-complanate or cylindrical conidia (Boonmee et al. 2016). The genus comprises 32 species (Hyde et al. 2017; Jayasiri et al. 2019; Shen et al. 2022b; Sushma et al. 2022; Liu et al. 2023; Shu et al. 2024; Sun et al. 2024; Wang et al. 2024; Wu et al. 2024), with eight being transferred from *Dictyosporium* (Boonmee et al. 2016; Yang et al. 2018).

Corda (1836) established *Dictyosporium* with *Di. elegans* as the type species. This genus is characterized by dark brown, subglobose ascomata, bitunicate cylindrical asci, and hyaline, fusiform uniseptate ascospores with or without a sheath, and sporodochial or effuse conidiomata, micronematous and mononematous conidiophores (or absent), and cheiroid, digitate complanate conidia with several parallel rows of cells (Prasher et al. 2015; Boonmee et al. 2016; Silva et al. 2016; Yang et al. 2018; Shen et al. 2022b). Dubey (2022) documented 13 species of *Dictyosporium*, which have already been reassigned to other genera (Boonmee et al. 2016; Yang et al. 2018). The genus comprises 58 species that are mainly found on dead wood and decaying leaves in terrestrial and freshwater environments (Dubey 2022; Shen et al. 2022b; Liu et al. 2023; Shu et al. 2024; Zhang et al. 2024b).

Jalapriya was introduced by Boonmee et al. (2016) with *J. pulchra* as the type species. The members of this genus are characterized by dark brown to black

colonies, acrogenous, solitary, and cheiroid conidia (Boonmee et al. 2016). Presently, six species are accepted in the genus, viz., *J. apicalivaginat*um, *J. aquaticum*, *J. cheirospora*, *J. inflata*, *J. pulchra*, and *J. toruloides* (Du et al. 2025).

Dulongjiang River is one of the six major river systems in Yunnan Province, China. Originating from Xizang Province, it flows through the northwestern Yunnan Province, China, and then enters into Myanmar. We are carrying out a survey of species diversity of lignicolous freshwater fungi in the Dulongjiang River basin, and this is the first in a series of papers to appear on these fungi. In this study, six fresh isolates of Dictyosporiaceae were obtained from submerged wood in Dulongjiang River and are identified as five species, two of which are new. Descriptions and illustrations are provided for these species. In addition, an updated backbone tree of Dictyosporiaceae is performed to clarify species relationships and support the establishment of novel taxa.

Materials and methods

Sample collection, specimen examination, and isolation

Submerged decaying woods were collected from the Dulongjiang River in Yunnan Province, China, in May 2023 and taken to the laboratory in Ziploc plastic bags. The sample incubation, examination, and morphological studies were carried out following the methods described by Luo et al. (2018a). Macromorphological characteristics were observed using an Optec SZ 760 compound stereomicroscope (Chongqing Optec Instrument Co., Ltd., Chongqing, China). The morphologies of colonies on original substrates were photographed with a Nikon SMZ1000 stereo zoom microscope (Nikon Corporation, Tokyo, Japan). Semi-permanent slides were observed using a Nikon ECLIPSE Ni-U compound stereomicroscope (Nikon Corporation, Tokyo, Japan) and photographed micro-morphological characteristics.

Single spore isolations were carried out based on the method described by Luo et al. (2018a) and Senanayake et al. (2020). Using a sterilized toothpick, a small amount of conidia was picked and streaked on potato dextrose agar (PDA) or water agar (WA) plates to spread the conidia. The plates were incubated at room temperature for approximately 12 hours to facilitate the formation of germ tubes. Germinated conidia were aseptically transferred to new PDA plates and incubated in an incubator room at 22 °C.

Specimens (dry wood with fungal material) were deposited in the Herbarium of Cryptogams Kunming Institute of Botany, Chinese Academy of Sciences (KUN-HKAS), Kunming, China. Living cultures were deposited in the Kunming Institute of Botany Culture Collection Center, Kunming, China (KUNCC). Fungal Names number (FN) (<https://nmdc.cn/fungalnames/>) for the new species was registered. New species were established following the recommendations outlined by Chethana et al. (2021).

DNA extraction, PCR amplification, and sequencing

Fungal mycelium was scraped from the surface of colonies grown on PDA and transferred to a 1.5 mL centrifuge tube. The Trelief TM Plant Genomic DNA Kit (TSP101-50) was used to extract DNA from the ground mycelium

according to the manufacturer's instructions. The ITS, LSU, and *tef1*- α gene regions were amplified using the primer pairs ITS5/ITS4, LR0R/LR7, and 983F/2218R, respectively (Vilgalys and Hester 1990; Liu et al. 1999). The amplification was performed in a 25 μ L reaction volume containing 9.5 μ L deionized water, 12.5 μ L 2 $\times$ Taq PCR Master Mix with blue dye (Sangon Biotech, Shanghai, China), 1 μ L of DNA template, and 1 μ L of each primer (10 μ M). The amplification condition for ITS, LSU, and *tef1*- α followed Bao et al. (2022) and Shen et al. (2021). PCR amplification was confirmed on 1% agarose electrophoresis gels. Purification and sequencing of PCR products were sent for sequencing at Tsingke Biological Engineering Technology and Services Company, Yunnan, China.

Molecular phylogenetic analyses

Sequence alignment

Sequences were assembled using BioEdit v. 7.2.5.0 (Dagona 1999). A BLAST search in the National Center of Biotechnology Information (NCBI) was performed on sequences with high similarity indices to find the closest matches with taxa. The ITS, LSU, and *tef1*- α used for phylogenetic analysis are selected based on the preliminary identification results and the related publications (Shen et al. 2022b; Tennakoon et al. 2023; Shu et al. 2024). Closely relevant sequences were downloaded from the National Center of Biotechnology Information (<https://www.ncbi.nlm.nih.gov/>). All consensus sequences and the reference sequences were automatically aligned with MAFFT v. 7.0 (<http://mafft.cbrc.jp/alignment/server/index.html>, accessed on 11 December 2024; Katoh et al. 2019). The sequence dataset was combined using SequenceMatrix v.1.7.8 (Vaidya et al. 2011). The alignment formats were converted to Phylip and Nexus formats by AliView (Larsson 2014).

Maximum likelihood (ML) analysis was performed by setting RaxML-HPC2 on XSEDE (8.2.12) (Stamatakis 2006; Stamatakis et al. 2008) in the CIPRES Science Gateway (<https://www.phylo.org/portal2/login!input.action>). The final RAXML search was conducted using the GTRGAMMA+I model (Nylander 2004; Ronquist et al. 2012). Maximum likelihood bootstrap support was calculated from 1000 bootstrap replicates (<https://www.phylo.org/portal2>).

Bayesian analysis was performed using MrBayes v. 3.2 (Ronquist et al. 2012). The evolution model was estimated using MrModeltest 2.3 (Nylander 2004), and the GTR+I+G model is the best-fit model of ITS, LSU, and *tef1*- α . Posterior Probabilities (PP) (Rannala and Yang 1996) were performed by Markov Chain Monte Carlo sampling (MCMC) in MrBayes v.3.1.2 (Liu et al. 2012). Six simultaneous Markov chains were run for 100 million generations, and trees were sampled every 10000th. The first 25% trees representing the burn-in phase of the analyses were discarded, and the remaining 75% trees (post-burning) were used for calculating posterior probabilities (PP) in the majority rule consensus tree. Phylogenetic trees were printed with FigTree v.1.4.4. (<http://tree.bio.ed.ac.uk/software/figtree/>), while editing and typesetting were achieved using Adobe Illustrator (AI) (Adobe Systems Inc., California, USA). Newly generated sequences in this study were submitted to GenBank (Table 1).

Table 1. GenBank and culture collection accession numbers of species used in the phylogenetic analyses. The newly generated sequences are indicated in red, and the type strains are indicated in bold.

Species	Voucher/Culture	GenBank accession number		
		ITS	LSU	tef1- α
<i>Aquadictyospora clematidis</i>	MFLUCC 17-2080	MT310592	MT214545	MT394727
<i>Aquadictyospora lignicola</i>	MFLUCC 17-1318	MF948621	MF948629	MF953164
<i>Aquaticheirosora lignicola</i>	HKUCC 10304	AY864770	AY736378	–
<i>Cheirosporium triseriale</i>	HMAS 180703	EU413953	EU413954	–
<i>Dendryphiella eucalyptorum</i>	CBS 137987	KJ869139	KJ869196	–
<i>Dendryphiella fasciculata</i>	MFLUCC 17-1074	MF399213	MF399214	–
<i>Dendryphiella paravinosae</i>	CBS 141286	KX228258	KX228309	–
<i>Dendryphiella phitsanulokensis</i>	MFLUCC 17-2513	MG754400	MG754401	–
<i>Dendryphiella variabilis</i>	CBS 584.96	LT963453	LT963454	–
<i>Dictyocheirosora acaciae</i>	SDBR-CMU454	OP965332	OP965372	OQ000838
<i>Dictyocheirosora acaciae</i>	SDBR-CMU455	OP965333	OP965373	OQ000839
<i>Dictyocheirosora aquadulcis</i>	MFLUCC 17-2571	MK634545	MK634542	–
<i>Dictyocheirosora aquadulcis</i>	MFLUCC 22-0095	OP526634	OP526644	OP542236
<i>Dictyocheirosora aquatica</i>	KUMCC 15-0305	KY320508	KY320513	–
<i>Dictyocheirosora bannica</i>	KH 332	LC014543	AB807513	AB808489
<i>Dictyocheirosora bannica</i>	MFLUCC 16-0874	MH381765	MH381774	–
<i>Dictyocheirosora cheirosora</i>	KUMCC 17-0035	MF177035	MF177036	–
<i>Dictyocheirosora Chiangmaiensis</i>	MFLUCC 22-0097	OP526630	OP526640	OP542232
<i>Dictyocheirosora clematidis</i>	MFLUCC 17-2089	MT310593	MT214546	MT394728
<i>Dictyocheirosora Garethjonesii</i>	MFLUCC 16-0909	KY320509	KY320514	–
<i>Dictyocheirosora Garethjonesii</i>	DLUCC 0848	MF948623	MF948631	MF953166
<i>Dictyocheirosora gigantica</i>	BCC 11346	DQ018095	–	–
<i>Dictyocheirosora heptaspora</i>	MFLUCC 22-0096	OP526635	OP526645	OP542237
<i>Dictyocheirosora heptaspora</i>	CBS 396.59	DQ018090	–	–
<i>Dictyocheirosora indica</i>	MFLUCC 15-0056	MH381763	MH381772	MH388817
<i>Dictyocheirosora lithocarpi</i>	MFLUCC 17-2537	MK347781	MK347999	–
<i>Dictyocheirosora metroxylonis</i>	MFLUCC 15-0028a	MH742321	MH742313	–
<i>Dictyocheirosora metroxylonis</i>	MFLUCC 15-0028b	MH742322	MH742314	MH764301
<i>Dictyocheirosora multiappendiculata</i>	KUNCC 22-10736	OP526633	OP526643	OP542235
<i>Dictyocheirosora multiappendiculata</i>	KUNCC 22-10734	OP526632	OP526642	OP542234
<i>Dictyocheirosora nabanheensis</i>	MFLUCC 17-0562	MH388340	MH376712	MH388375
<i>Dictyocheirosora nabanheensis</i>	MFLUCC 22-0094	OP526637	OP526647	OP542239
<i>Dictyocheirosora nabanheensis</i>	KUNCC 23-15886	PQ309051	PQ309042	PQ346446
<i>Dictyocheirosora pandanicola</i>	MFLUCC 16-0365	MH388341	MH376713	MH388376
<i>Dictyocheirosora pseudomusae</i>	yone 234	LC014550	AB807520	AB808496
<i>Dictyocheirosora rotunda</i>	MFLUCC 14-0293	KU179099	KU179100	–
<i>Dictyocheirosora rotunda</i>	MFLUCC 17-0222	MH381764	MH381773	MH388818
<i>Dictyocheirosora rotunda</i>	GZCC 19-0436	OQ842723	MW133811	OQ850744
<i>Dictyocheirosora rotunda</i>	GZCC 19-0429	OQ842724	MW133814	OQ850745
<i>Dictyocheirosora rotunda</i>	MFLUCC 17-1313	MF948625	MF948633	MF953168
<i>Dictyocheirosora splendida</i>	KUNCC 23-15971	PQ309052	PQ309043	PQ346447
<i>Dictyocheirosora suae</i>	KUNCC 22-12424	OP526631	OP526641	OP542233
<i>Dictyocheirosora submersa</i>	ZHKUCC 24-0001	PP326193	PP326216	PP333113
<i>Dictyocheirosora subramaniana</i>	BCC 3503	DQ018094	–	–

Species	Voucher/Culture	GenBank accession number		
		ITS	LSU	<i>tef1</i> -α
<i>Dictyocheiropora taiwanense</i>	MFLUCC 17-2654	MK495821	MK495820	–
<i>Dictyocheiropora thailandica</i>	MFLUCC 18-0987	MT627734	MN913743	–
<i>Dictyocheiropora vinaya</i>	MFLUCC 14–0294	KU179102	KU179103	–
<i>Dictyocheiropora xishuangbannaensis</i>	KUMCC 17-0181	MH388342	MH376714	MH388377
<i>Dictyosporium alatum</i>	ACC 34953	NR077171	DQ018101	–
<i>Dictyosporium appendiculatum</i>	MFLUCC 17-2259	MH388343	MH376715	–
<i>Dictyosporium aquaticum</i>	MF1318	KM610236	–	–
<i>Dictyosporium bulbosum</i>	yone 221	LC014544	AB807511	AB808487
<i>Dictyosporium digitatum</i>	KH 401	LC014545	AB807515	AB808491
<i>Dictyosporium digitatum</i>	yone 280	LC014547	AB807512	AB808488
<i>Dictyosporium duliuijiangense</i>	GZCC 19-0426	OQ842725	MW133815	OQ850746
<i>Dictyosporium duliuijiangense</i>	KUNCC 23-15949	PQ309055	PQ309046	PQ346450
<i>Dictyosporium elegans</i>	NBRC 32502	DQ018087	DQ018100	–
<i>Dictyosporium guangdongense</i>	ZHKUCC 24-0002	PP326190	PP326213	–
<i>Dictyosporium guttulatatum</i>	MFLUCC 16-0258	MH388345	MH376717	MH388379
<i>Dictyosporium hongkongensis</i>	KUMCC 17-0268	MH388346	MH376718	MH388380
<i>Dictyosporium hughesii</i>	K 1847	LC014548	AB807517	AB808493
<i>Dictyosporium hughesii</i>	KUNCC 23-15923	PQ309056	PQ309047	–
<i>Dictyosporium krabiense</i>	MFLU 16-1890	–	MH376719	MH388381
<i>Dictyosporium meiosporum</i>	MFLUCC 10-0131	KP710944	KP710945	–
<i>Dictyosporium muriformis</i>	GZCC 20-0006	MT002304	MN897834	MT023011
<i>Dictyosporium nigroapice</i>	MFLUCC 17-2053	MH381768	MH381777	MH388821
<i>Dictyosporium nigroapice</i>	BCC 3555	DQ018085	–	–
<i>Dictyosporium olivaceosporum</i>	KH 375	LC014542	AB807514	AB808490
<i>Dictyosporium palmae</i>	CBS-H 22129	–	KX555648	–
<i>Dictyosporium pandanicola</i>	MFLU 16-1886	MH388347	MH376720	MH388382
<i>Dictyosporium sexualis</i>	MFLUCC 10-0127	KU179105	KU179106	–
<i>Dictyosporium</i> sp.	MFLUCC 15-0629	MH381766	MH381775	MH388819
<i>Dictyosporium stellatum</i>	CCFC 241241	NR 154608	JF951177	–
<i>Dictyosporium strelitziae</i>	CBS 123359	NR 156216	FJ839653	–
<i>Dictyosporium tetrasporum</i>	K 2865	LC014551	AB807519	AB808495
<i>Dictyosporium thailandicum</i>	MFLUCC 13-0773	KP716706	KP716707	–
<i>Dictyosporium tratense</i>	MFLUCC 17-2052	MH381767	MH381776	MH388820
<i>Dictyosporium tubulatum</i>	MFLUCC 15-0631	MH381769	MH381778	MH388822
<i>Dictyosporium tubulatum</i>	MFLUCC 17-2056	MH381770	MH381779	–
<i>Dictyosporium variabilisporum</i>	ZHKUCC 24-0003	PP326192	PP326215	PP333112
<i>Dictyosporium wuyiense</i>	CGMCC 3.18703	KY072977	–	–
<i>Dictyosporium zhejiangense</i>	MW–2009a	FJ456893	–	–
<i>Digitodesmium aquaticum</i>	MFLU 22-0203	OP749872	OP749877	OP756064
<i>Digitodesmium bambusicola</i>	CBS 110279	DQ018091	DQ018103	–
<i>Digitodesmium chiangmaiense</i>	HKAS 102163	–	MK571766	–
<i>Digitodesmium chishuiense</i>	GZCC 20-0510	OP377808	OP377907	OP472990
<i>Digitodesmium polybrachiatum</i>	COAD 3174	MW879318	MW879316	MW890262
<i>Digitodesmium polybrachiatum</i>	COAD 3175	MW879319	MW879317	MW890263
<i>Digitodesmium</i> sp.	BRC 10038	MK405235	MK405233	MK405231
<i>Digitodesmium</i> sp.	BRC 10037	MK405234	MK405232	MK405230
<i>Digitodesmium tectonae</i>	NFCCI 4878	MW854646	MW854647	MW854832
<i>Gregarithecium curvisporum</i>	MFLUCC 13-0853	KX364281	KX364282	–

Species	Voucher/Culture	GenBank accession number		
		ITS	LSU	tef1- α
<i>Gregarithecium curvisporum</i>	K 922	AB809644	AB807547	–
<i>Immotthia atrograna</i>	Z-Myc-64283	MW489540	–	–
<i>Immotthia bambusae</i>	HKAS 112012AI	MW489455	MW489450	MW504646
<i>Immotthia bambusae</i>	HKAS 112012C	MW489458	MW489453	MW504648
<i>Jalapriya apicalivaginatum</i>	HKAS 115801	MZ621167	MZ621168	–
<i>Jalapriya aquaticum</i>	DLUCC 2351	MZ621151	MZ621165	–
<i>Jalapriya aquaticum</i>	HKAS 115807	MZ621152	MZ621169	MZ851995
<i>Jalapriya guttulata</i>	KUNCC 23-15861	PQ309053	PQ309044	PQ346448
<i>Jalapriya guttulata</i>	KUNCC 23-14576	PQ309054	PQ309045	PQ346449
<i>Jalapriya inflata</i>	NOU 3855	JQ267362	JQ267363	–
<i>Jalapriya pulchra</i>	MFLUCC 15-0348	KU179108	KU179109	–
<i>Jalapriya pulchra</i>	MFLUCC 17-1683	MF948628	MF948636	MF953171
<i>Jalapriya toruloides</i>	CBS 209.65	DQ018093	DQ018104	–
<i>Neodendryphiella mali</i>	FMR 16561	LT906655	LT906657	–
<i>Neodendryphiella michoacanensis</i>	FMR 16098	LT906660	LT906658	–
<i>Neodendryphiella tarraconensis</i>	FMR 16234	LT906659	LT906656	–
<i>Neodigitodesmium cheirosporum</i>	UESCC 22-0020	ON595714	ON595713	ON595700
<i>Nigrograna mycophila</i>	CBS 141478	NR 147654	–	–
<i>Nigrograna mycophila</i>	MF6	KX650554	–	KX650526
<i>Periconia igniaria</i>	CBS 379.86	LC014585	AB807566	AB808542
<i>Periconia igniaria</i>	CBS 845.96	LC014586	AB807567	AB808543
<i>Pseudocoleophoma bauhiniae</i>	MFLUCC 17-2586	MK347736	MK347953	MK360076
<i>Pseudocoleophoma bauhiniae</i>	MFLUCC 17-2280	MK347735	MK347952	MK360075
<i>Pseudocoleophoma calamagrostidis</i>	K 3284	LC014592	LC014609	LC014614
<i>Pseudocoleophoma flavescen</i>	CBS 178.93	–	GU238075	–
<i>Pseudocoleophoma polygonicola</i>	K 731	AB809634	AB807546	AB808522
<i>Pseudocoleophoma puerensis</i>	ZHKUCC 22 0204	NR 184995	NG154047	OP321568
<i>Pseudocoleophoma puerensis</i>	ZHKUCC 22 0205	OP297800	OP297770	OP321569
<i>Pseudocoleophoma rhapsidis</i>	ZHKUCC 21 0124	ON244664	ON244661	–
<i>Pseudocoleophoma rhapsidis</i>	ZHKUCC 22 0004	ON244665	ON244662	–
<i>Pseudocoleophoma rusci</i>	MFLUCC 16-1444	MT185549	MT183514	–
<i>Pseudocoleophoma typhicola</i>	MFLUCC 16-0123	KX576655	KX576656	–
<i>Pseudocoleophoma zingiberacearum</i>	NCYUCC 19-0052	MN615939	MN616753	MN629281
<i>Pseudocoleophoma zingiberacearum</i>	NCYUCC 19-0053	MN615940	MN616754	MN629282
<i>Pseudoconiothyrium broussonetiae</i>	CBS 145036	NR 163377	NG066331	–
<i>Pseudocyclothyriella clematidis</i>	MFLU 16-0280	MT310596	MT214549	–
<i>Pseudocyclothyriella clematidis</i>	MFLUCC 17-2177A	MT310595	MT214548	MT394730
<i>Pseudodictyosporium elegans</i>	CBS 688.93	DQ018099	DQ018106	–
<i>Pseudodictyosporium indicum</i>	CBS 471.95	DQ018097	–	–
<i>Pseudodictyosporium thailandica</i>	MFLUCC 16-0029	KX259520	KX259522	KX259526
<i>Pseudodictyosporium wauense</i>	NBRC 30078	DQ018098	DQ018105	–
<i>Pseudodictyosporium wauense</i>	DLUCC 0801	MF948622	MF948630	MF953165
<i>Verrucocum coppinsii</i>	E00814291	MT918784	MT918770	–
<i>Verrucocum spribillei</i>	SPO 2343	MT918780	MT918765	–
<i>Vikalpa australiensis</i>	HKUCC 8797	DQ018092	–	–
<i>Vikalpa grandispora</i>	KUNCC 22-12425	OP526638	OP526648	OP542240
<i>Vikalpa sphaerica</i>	CGMCC 3.20682	OP526639	OP526649	OP542241




Figure 1. The maximum likelihood (ML) tree is based on the combined ITS, LSU, and *tef1*-α sequence data. Bootstrap support values with ML greater than 75% and Bayesian posterior probabilities (PP) greater than 0.95 are given above the nodes, shown as “ML/PP”. The tree was rooted to *Periconia igniaria* (CBS 379.86 and CBS 845.96). New species are indicated in red, and type strains are in bold.




Figure 1. Continued.

Results

Phylogenetic analyses

The phylogram generated from maximum likelihood analysis based on combined ITS, LSU, and *tef1*- α sequence data represents the relationships of all genera of the Dictyosporiaceae. One hundred thirty-nine strains are included in the combined analyses, which comprise 2180 characters, including gaps (ITS: 1–470 bp, LSU: 471–1304 bp, *tef1*- α : 1305–2180 bp) after aligning, including *Periconia igniaria* (CBS 379.86 and CBS 845.96) as the outgroup taxon. RAxML and Bayesian analyses were conducted and resulted in generally congruent topologies. The best RAxML tree with a final likelihood value of -21822.800488 is presented. The matrix had 996 distinct alignment patterns, with 25.73% undetermined characters or gaps. Estimated base frequencies were as follows: A = 0.233883, C = 0.255161, G = 0.273222, T = 0.237734; substitution rates AC = 1.700360, AG = 4.088659, AT = 2.394708, CG = 0.842455, CT = 9.208380, GT = 1.000000; gamma distribution shape parameter α = 0.226697. Statistical values for maximum likelihood (ML) above 75% and Bayesian posterior probabilities (PP) greater than 0.95 are given at the nodes.

Phylogenetic analyses showed that the new isolates were nested within Dictyosporiaceae and distributed in three genera, viz., *Dictyocheiropora*, *Dictyosporium*, and *Jalapriya*. *Dictyocheiropora nabanheensis* (KUNCC 23-15886) clustered with its type strain with 79% ML and 0.96 PP support. *Dictyocheiropora splendida* formed a distinct lineage with *D. garethjonesii* and *D. nabanheensis* with 83% ML and 0.98 PP support. Two newly obtained strains of *Jalapriya guttulata* (KUNCC 23-14576 and KUNCC 23-15861) clustered sister to *J. toruloides* with 99% ML and 1.00 PP support. *Dictyosporium hughesii* (KUNCC 23-15923) and *Di. duliuijiangense* (KUNCC 23-15949) clustered within the *Dictyosporium*. *Dictyosporium duliuijiangense* clustered sister to its type strain (GZCC 19-0426) with 100% ML and 1.00 PP support. The new isolate of *Di. hughesii* (KUNCC 23-15923) clustered with its type strain (KT 1847) with 97% ML and 1.00 PP support.

Taxonomy

Dictyocheiropora nabanheensis Tibpromma & K.D. Hyde, Fungal Diversity 93: 10 (2018)

Index Fungorum: IF554474

Facesoffungi Number: FoF04483

Fig. 2

Description. *Saprobic* on submerged decaying wood in the Dulongjiang River. **Sexual morph** Undetermined. **Asexual morph:** Hyphomycetous. **Colonies** on natural substrate in small, scattered clusters, dark brown, velvety. **Mycelium** composed of immersed or partly superficial, hyaline to pale brown, septate, branched hyphae. **Conidiophores** micronematous, mononematous, septate, smooth, thin-walled, cylindrical, reduced to conidiogenous cells. **Conidiogenous cells** 8–13 $\times$ 3–5 μm ($\bar{x}$ = 13.8 $\times$ 4.2 μm , n = 10), holoblastic, monoblastic, integrated, cylindrical, hyaline to pale brown, smooth-walled. **Conidia** 31–36 $\times$ 14–16 μm ($\bar{x}$ = 33 $\times$ 15 μm , n = 36), solitary, ellipsoid to cylindrical, cheiroid, with a basal connecting cell, brown to yellow–brown, smooth-walled, euseptate,




Figure 2. *Dictyocheirospora nabanheensis* (KUN-HKAS 135951) **a** colonies on submerged decaying wood **b, c** conidio-phores and conidia **d–e** conidia **f** germinated conidia; Culture on PDA from front (**g**) and reverse (**h**). Scale bars: 30 μ m (**b–c**); 20 μ m (**d, e, f**).

with discoid individual cells arranged in 6 compact rows ($\bar{x} = 34.8 \times 5.8 \mu\text{m}$, $n = 30$) closely clustered at the apex, each row with 6–8 cells compactly and linearly clustered, with 1–2 rounded to cylindrical hyaline, $6\text{--}10 \times 5\text{--}7 \mu\text{m}$, appendages arising from near the middle of conidial rows.

Culture characteristics. Conidia germinating on WA within 12 h and germ tubes were produced at the basal region (Fig. 2f). Colonies on PDA at 22 °C, circular, white at margin around the edge and light gray in the center, raised on surface, with white, pale orange concentric rings, yellow–orange in the middle on the reverse side.

Material examined. CHINA • Yunnan Province, Dulongjiang River, on submerged decaying wood, 2 May 2023 (Altitude: 1422 m, 27.793592°N, 98.330416°E), Ying Wang, S4719 (KUN-HKAS 135951), living culture, KUNCC 23-15886.

Notes. The newly collected fungus has closely clustered terminal cells at the apex of the conidia, which is similar to those of the genus *Dictyocheiropora*. Phylogenetic analysis of the combined ITS, LSU, and *tef1*-α sequence data showed that the new strain KUNCC 23-15886 clustered sister to the ex-type strain of *D. nabanheensis* (KUMCC 16-0152) with 79% ML and 0.96 PP statistical support (Fig. 1). Morphologically, KUNCC 23-15886 resembles *D. nabanheensis* in having conidia with hyaline, globose to subglobose appendages in the middle region. *Dictyocheiropora nabanheensis* (KUMCC 16-0152) differs from KUNCC 23-15886 in having slightly larger conidia ($38 \times 20 \mu\text{m}$ vs. $33 \times 15 \mu\text{m}$) (Tibpromma et al. 2018a; Shen et al. 2022b) and with more cells in each row (6–10 cells vs. 6–8 cells) (Tibpromma et al. 2018a; Shen et al. 2022b). However, there are only 0.83% (4/480) differences between ITS sequence data. Therefore, we identified our collection as *D. nabanheensis*. The type of *D. nabanheensis* was found on dead leaves of *Pandanus* sp. in a terrestrial habitat in Yunnan, China, and the strain of MFLUCC 22-0094 was found on submerged decaying wood in a freshwater habitat in Thailand. Our study isolated this species for the first time on submerged decaying wood in Yunnan province, China.

Dictyocheiropora splendida Y. Wang & Z.L. Luo, sp. nov.

Fungal Names: FN 572104

Fig. 3

Etymology. Referring to the glistening colony on natural substrates.

Holotype. KUN-HKAS 135954.

Description. *Saprobic* on submerged decaying wood. **Sexual morph:** Undetermined. **Asexual morph:** Hyphomycetous. **Colonies** glistening on natural substrate, in small pulvinate groups, scattered on surface, dark brown, velvety. **Mycelium** immersed on substrate, composed of brown, smooth, septate, branched hyphae. **Conidiophores** micronematous, mononematous, hyaline to pale brown, $4\text{--}5 \mu\text{m}$ wide, mostly reduced to conidiogenous cells. **Conidiogenous cells** $6\text{--}11 \times 3\text{--}6 \mu\text{m}$ ($\bar{x} = 8.5 \times 4.7 \mu\text{m}$, $n = 10$), holoblastic, integrated, terminal, cylindrical, hyaline to pale brown, smooth-walled. **Conidia** $37\text{--}43 \times 14\text{--}17 \mu\text{m}$ ($\bar{x} = 40 \times 16 \mu\text{m}$, $n = 30$) solitary, acrogenous, cheiroid, with a basal connecting cell, brown to yellow-brown, arranged in 6 compact rows closely clustered at the apex, with each row composed of 6–9 cells, euseptate, with one appendage at the center of conidia.



Figure 3. *Dictyocheirospora splendida* (KUN-HKAS 135954, holotype) **a, b** colonies on submerged decaying wood **c–e** conidiophores with conidia **f** crushed single conidia **g** rows of conidial cells **h, i** conidia **j** germinated conidium; Culture on PDA from front (**k**) and reverse (**l**). Scale bars: 30 μ m (**d, e**); 20 μ m (**c, f–j**).

Culture characteristics. Conidia germinating on PDA within 12 h and germ tubes produced at the base (Fig. 3k). Colonies grew on PDA at 22 °C, circular, white at margin and light gray in the center, slightly raised on surface cottony, with mycelium partially infiltrating into the culture and white on the edge, grayish-white in the middle on the reverse side.

Material examined. CHINA • Yunnan Province, Dulongjiang River, on submerged decaying wood, 2 May 2023 (Altitude: 1418 m, 27.783341°N, 98.330741°E), Ying Wang, S4957 (KUN-HKAS 135954, holotype); ex-type culture, KUNCC 23-15971.

Notes. In the phylogenetic analyses, *Dictyocheiropora splendida* was basal to *D. garethjonesii* and *D. nabanheensis*, with 83% ML and 0.98 PP support (Fig. 1). The new strain KUNCC 23-15971 has 2.29% (11/480 bp) ITS and 3.14% (26/829 bp) *tef1*-α base pair difference from *D. nabanheensis* (MFLUCC 17-0562) as well as 2.55% (13/509 bp) ITS and 2.55% (31/874 bp) *tef1*-α base pair difference from *D. garethjonesii* (DLUCC 0848). However, LSU sequences of these three species are identical. Morphologically, *D. splendida* has sporodochial conidiomata, darkly pigmented, micronematous or semi-macronematous conidiophores, and acrogenous, cheiroid, and brown-colored conidia, which match the characteristics of *Dictyocheiropora* (Boonmee et al. 2016). However, *D. nabanheensis* exhibits distinct conidial appendages (Tibpromma et al. 2018b), which were inconspicuous in *D. splendida*. In addition, *D. splendida* exhibits a significant difference in conidial size compared to *D. garethjonesii* (40 × 16 μm vs. 50 × 20 μm) (Boonmee et al. 2016). Based on morphological and phylogenetic analysis, we introduce *Dictyocheiropora splendida* as a new species.

***Dictyosporium duliuijiangense* L.L. Liu & Z.Y. Liu, Phytotaxa 606: 266 (2023)**

Index Fungorum: IF900403

Facesoffungi Number: FoF14119

Fig. 4

Description. **Saprobic** on submerged decaying wood. **Sexual morph:** Undetermined. **Asexual morph:** Hyphomycetous. **Colonies** on natural substrate punctiform, sporodochial, scattered, dark brown to black, velvety, glistening. **Mycelium** mostly immersed, composed of septate, branched, hyaline to pale brown hyphae. **Conidiophores** macronematous, branched, septate, cylindrical, hyaline to pale yellow, rough-walled, 3–5 μm wide, sometimes reduced to conidiogenous cells. **Conidiogenous cells** 11–18 × 3–5 μm ($\bar{x}$ = 15 × 4 μm, n = 17) monoblastic, integrated, terminal, determinate, pale yellow to yellow. **Conidia** 38–45 × 22–26 μm ($\bar{x}$ = 41.5 × 24 μm, n = 30), acrogenous, solitary, cheiroid, smooth-walled, complanate, yellowish-brown to pale brown, range from moderate orange at the base to pale yellow at the apex when immature, consisting of five closely adpressed rows of cells 33–38 × 4–6 μm ($\bar{x}$ = 35.5 × 5.5 μm, n = 30), side rows shorter than middle rows, 4–10-euseptate in each row of cells, constricted at septa, with one or two hyaline, tubular, elongated appendages, narrower in the middle, 42–59 × 6–9 μm ($\bar{x}$ = 50.2 × 7.2 μm, n = 28) containing small guttules, suspended from the apical part of the outer rows of cells.

Cultural characteristics. Conidia germinating on PDA within 24 h and germ tubes produced from the apex. Colonies grew on PDA at 22 °C, circular, with



Figure 4. *Dictyosporium duliuijiangense* (KUN-HKAS 135956) **a** colonies on submerged decaying wood **b–d** conidiophores with conidia **e–i** conidia and conidiogenous cells **j** germinated conidium; Culture on PDA from front (**k**) and reverse (**l**). Scale bars: 50 μ m (**b–c**); 30 μ m (**d, i–j**); 20 μ m (**e–h**).

fluffy, dense, white mycelium. The reverse side has a pale-yellow edge and an orange center.

Material examined. CHINA • Yunnan Province, Dulongjiang River, on submerged decaying wood, 2 May 2023 (Altitude: 1516 m, 27.686450°N, 98.349933°E), Ying Wang, S4923 (KUN-HKAS 135956), living culture, KUNCC 23-15949.

Notes. The conidia in *Dictyosporium* are complanate and not closely clustered at the apex (Boonmee et al. 2016; Yang et al. 2018). The morphological characteristics of the newly collected strain (KUNCC 23-15949) match with *Dictyosporium* species (Boonmee et al. 2016, Yang et al. 2018). Based on the phylogenetic analyses of the combined ITS, LSU, and *tef1*-α sequence data, our new strain (KUNCC 23-15949) shows a close affinity to *Di. duliuijiangense* (GZCC 19-0426). There is a 1.56% (8/512) difference in ITS sequence data between KUNCC 23-15949 and GZCC 19-0426, but the LSU and *tef1*-α sequences of these two species are identical. Morphologically, our collection differs from the type material in the size of conidia (41.5 × 24 μm vs. 33 × 20 μm) and the length of the appendages (50.2 × 7.2 μm vs. 27 × 6 μm) (Liu et al. 2023). *Di. duliuijiangense* was first reported on a decaying branch submerged in a freshwater river in Dushan County, Guizhou Province, China (Liu et al. 2023). Based on these findings, we identified our collection as *Di. duliuijiangense*.

***Dictyosporium hughesii* McKenzie, Mycotaxon 111: 156 (2010)**

MycoBank No: 515243

Fig. 5

Description. *Saprobic* on submerged decaying wood. **Sexual morph:** Undetermined. **Asexual morph:** Hyphomycetous. **Colonies** on natural substrate, punctiform, sporodochia, scattered but coalescing, black, irregular. **Mycelium** immersed, pale to brown. **Conidiophores** micronematous, mononematous, subhyaline, thin-walled, smooth. **Conidiogenous cells** holoblastic, determinate, cylindrical, 3.5–5 μm wide. **Conidia** 56–68 × 23–29 μm ($\bar{x}$ = 62 × 26.3 μm, *n* = 40), solitary, acrogenous, medium brown, complanate, ellipsoidal or cylindrical, cheiroid, arranged in 7 closely adpressed rows of cells, with outer rows of cells arising from a basal cell, the side rows lower than middle rows, outer rows extending about two-thirds of the way along the conidium, each row containing 6–12 cells, constricted at septa, slightly thickened walls and septa, with or without appendages.

Cultural characteristics. Conidia germinating on PDA within 24 h and germ tubes produced from the basal cells and the apex of the row of cells. Colonies grew on PDA at 22 °C, circular, with fluffy mycelium; the mycelium is dense in the center and sparse at the edges, white at the outer margin and light yellow at the center, light yellow on the reverse side.

Material examined. CHINA • Yunnan Province, Dulongjiang River, on submerged decaying wood, 2 May 2023 (Altitude: 1418 m, 27.783341°N, 98.330741°E), Ying Wang, S4812 (KUN-HKAS 135963), living culture, KUNCC 23-15923.

Notes. This collection was identified as *Dictyosporium hughesii* based on the phylogenetic analysis and morphological characteristics. Phylogenetic analysis of the combined ITS, LSU, and *tef1*-α sequence data showed that our new strain (KUNCC 23-15923) clustered with the ex-type strain of *Di. hughesii* (KT 1847)




Figure 5. *Dictyosporium hughesii* (KUN-HKAS 135963) **a** colonies on submerged decaying wood **b** conidia **c** conidiophores and Conidiogenous cells **d** conidia with appendages **e–j** conidia **k** germinated conidium; Culture on PDA from front (**l**) and reverse (**m**). Scale bars: 30 µm (**b, d, f–k**); 20 µm (**c, e**).

with 97% ML and 1.00 PP support (Fig. 1). Morphologically, the conidia of our new collection are slightly larger (56–68 μm vs. 40–47 μm), and the appendages are not prominent. *Di. hughesii* has been found on dead leaves of *Rhopalostylis sapida* in New Zealand (McKenzie 2010) and on dead branches of *Stewartia monadelphica* in Japan (Ghosh et al. 2015; Tanaka et al. 2015), and herein it is reported from submerged decaying wood in a freshwater habitat in China.

***Jalapriya guttulata* Y. Wang & Z.L. Luo, sp. nov.**

Fungal Names: FN 572106

Fig. 6

Etymology. Referring to the guttulate cells of the conidia.

Holotype. KUN-HKAS 135959.

Description. *Saprobic* on submerged decaying wood. **Sexual morph:** Undetermined. **Asexual morph:** Hyphomycetous. **Colonies** on natural substrate effuse, scattered, dark brown to black. **Mycelium** mostly immersed, partly superficial, composed of smooth, septate, branched, pale brown to brown, or pale orange. **Conidiophores** micronematous, unbranched, thin-walled, smooth, 3–4 μm wide. **Conidiogenous cells** holoblastic, integrated, terminal, pale brown to brown, or pale brown, thin-walled. **Conidia** 31–43 $\times$ 25–38 μm ($\bar{x}$ = 37.3 $\times$ 31.5 μm , n = 30) (KUNCC 23-15861) or 34–41 $\times$ 30–37 μm ($\bar{x}$ = 37.5 $\times$ 33.7 μm , n = 30) (KUNCC 23-14576), acrogenous, septate, distinctly constricted at septa, holoblastic, solitary or paired (Fig. 6c), palmate, light olive-brown when immature and olive-brown to reddish-brown or pale brown, with some cells slightly darker in the center and lighter at the edges when mature. with 5–11 overlapping rows of unequal length and cells arranged in a plane, 1–2 outer rows arising from a basal cell, with additional rows arising from the base of the previous row, these rows are slightly curved inward, with each row consisting of 2–9 cells, with number of cells gradually decreasing from base to upward except for basal row of cells, with some cells in the uppermost row enlarged.

Culture characteristics. Conidia germinating on PDA within 24 h and germ tubes produced at the basal cell and the apex of the conidium. Colonies grew on PDA at 22 °C, circular, mycelium was loose, flocculent, on the obverse, at first white, later becoming light yellow. The reverse side is orange, with the center being the darkest.

Material examined. CHINA • Yunnan Province, Dulongjiang River, on submerged decaying wood, 2 May 2023, Fig. 6a–f: (Altitude: 1397 m, 27.702847°N, 98.349787°E), Ying Wang, S4558 (KUN-HKAS 135959, holotype); ex-type culture, KUNCC 23-15861. Fig. 6i–l: (Altitude: 1702 m, 28.033444°N, 98.317358°E), Ying Wang, S4587 (KUN-HKAS 135948), living culture, KUNCC 23-14576.

Notes. Six species are currently accepted in *Jalapriya*, among which *J. inflata* and *J. toruloides* have been reported from wood in marine habitats (Goh et al. 1999; Kirschner et al. 2013; Tibell et al. 2020; Du et al. 2025). These two species were originally classified under *Dictyosporium* and were transferred to the genus *Jalapriya* by Boonmee et al. (Boonmee et al. 2016). *J. apicalivaginatam*, *J. aquaticum*, and *J. pulchra* were found on submerged decaying wood in freshwater habitats, the first one from Henan province and the latter two from Yunnan province (Boonmee et al. 2016; Fu et al. 2021). While *J. cheirospora* was found on dead stems of the




Figure 6. *Jalapriya guttulata* (a–h KUN-HKAS 135959, holotype; i–l KUN-HKAS 135948) a, i colonies on submerged decaying wood b–e, j–l conidiophores and conidia f germinated conidia; Culture on PDA from front (g) and reverse (h). Scale bars: 20 µm (b–f, j–l).

medicinal plant *Disporum longistylum* in Kunming City, Yunnan Province, China. In this study, we collected two morphologically similar strains (KUNCC 23-15861 and KUNCC 23-14576) from submerged decaying wood. Despite the differences in conidial color (olive-brown to reddish-brown vs. pale brown to moderate brown) and size ($37.3 \times 31.5 \mu\text{m}$ vs. $37.5 \times 33.7 \mu\text{m}$), both strains share similar characteristics in culture and natural colonies, have guttulate conidia, and relatively few cells in the terminal cell row. Notably, no differences were found in the sequences of ITS and *tef1*- α gene regions between KUNCC 23-15861 and KUNCC 23-14576. Which indicated that the two strains belong to the same species.

Based on the phylogenetic analyses of the combined ITS, LSU, and *tef1*- α sequence data, our new strains (KUNCC 23-15861 and KUNCC 23-14576) clustered with *J. toruloides* (CBS 209.65) with 99% ML and 1.00 PP support (Fig. 1). The ITS sequence of *J. guttulata* has a 2.7% (14/518 bp) difference from *J. toruloides* (CBS 209.65). But the LSU sequence has only one base pair (1/494 bp) difference from *J. toruloides*. Morphologically, *J. guttulata* has more conidial rows of cells (5–11 rows vs. 6–8 rows); the conidia of *J. guttulata* are shorter (37.3 – $37.5 \mu\text{m}$ vs. 38 – $56 \mu\text{m}$) and wider (31.5 – $33.7 \mu\text{m}$ vs. 25 – $24 \mu\text{m}$) than *J. toruloides*; the rows of these two species are of unequal length, and the apical rows of the conidia in most *J. guttulata* have fewer cells (Henningsson 1974; Goh et al. 1999). Notably, solitary or paired conidia of *J. guttulata* were observed for the first time in the genus *Jalapriya*. Based on morphological characteristics and phylogenetic analysis, we identify our isolates as a new species.

Discussion

Dulongjiang River is rich in lignicolous freshwater fungi. To date, 18 lignicolous freshwater fungi have been described and illustrated from this river, including 11 new species (Su et al. 2016; Wang et al. 2016; Luo et al. 2017, 2018b, 2019; Bao et al. 2018; Li et al. 2020). It is worth continuing the investigation on the diversity of microfungi in the Dulongjiang river basin. Dulongjiang River is particularly rich in species of Dictyosporiaceae. Wang et al. (2016) introduced a new species, *Dictyocheiropora aquatica*, from the Dulongjiang river basin. In the present study, we reported five Dictyosporiaceae species.

The conidia of *Jalapriya* are typically solitary (Boonmee et al. 2016; Tibell et al. 2020; Fu et al. 2021), whereas in this study paired conidia were observed for the first time. Notably, the overall morphologies of KUNCC 23-15861 and KUNCC 23-14576 differ significantly. However, their sequence data showed no differences which indicates that it is difficult to accurately identify species based solely on morphology, and combining both morphological and phylogenetic analyses is essential when identifying species.

Since the early 2000s, mycologists have been working on the diversity of Dictyosporiaceae. However, early studies relied heavily on the morphological characters; only a few species were associated with molecular data, resulting in the taxonomic status of numerous species remaining inaccurate (Luo et al. 2004). Over the past decade, the family has been studied intensively based on both morphology and multi-gene (ITS, LSU, and *tef1*- α) phylogenetic analyses; as a result, the taxonomy of several species has been revised (Boonmee et al. 2016; Yang et al. 2018). ITS and *tef1*- α sequences play a crucial role in the identification of species within the Dictyosporiaceae. The comparison revealed

significant nucleotide differences between the ITS and *tef1*- α sequences among different species. However, many species lack protein-coding genes (*tef1*- α) sequences. As the number of species increases, it becomes more difficult to distinguish different species based on ITS and LSU sequences alone. For instance, *Dictyosporium aquaticum* and *Di. stellatum* only have ribosomal sequences (ITS and LSU) available in GenBank; although there are significant morphological differences, they cannot be distinguished on the phylogenetic tree (Crous et al. 2011; Liu et al. 2015; Shen et al. 2022b; Shu et al. 2024). Therefore, it is necessary to add protein-coding genes for identifying species in future studies.

Dictyosporiaceae has been accommodated as a pleomorphic group and is recently being well-studied based on the combination of morphological and phylogenetic analysis. However, there are still some unsettled issues within the family (Tennakoon et al. 2023; Shu et al. 2024). The phylogenetic distinctions between *Immotthia*, *Pseudocoleophoma*, *Pseudoconiothyrium*, *Pseudocyclothyriella*, and *Verrucocum* are still ambiguous (Shen et al. 2022b; Liu et al. 2023). Whereas *Pseudocoleophoma* is polyphyletic, forming three clades within the family (Feng et al. 2024). Therefore, integrated further research on this family is urgently necessary.

Acknowledgements

Ying Wang is grateful to Shaun Pennycook for checking the nomenclature and to Zheng-Quan Zhang, Guang-Yun Zhu, Qiong Ma, Wen-Peng Wang, and Ting-Xiang Liu for their assistance with sample collection. Li-Qun Wen and Liang Zhang are acknowledged for help with DNA extraction and PCR amplification.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

We would like to thank the National Natural Science Foundation of China (Project ID: 32060005) and the Yunnan Fundamental Research Project (202201AW070001) for financial support. This work was also supported by the High-Level Talents Program of Yunnan Province (YNQR-QNRC-2020-113) and the Foundation of Yunnan Province Science and Technology Department (202305AM070003). D. Jayarama Bhat gratefully acknowledges the financial support provided under the Distinguished Scientist Fellowship Programme (DSFP) at King Saud University, Riyadh, Saudi Arabia. Danfeng Bao would like to thank the Postdoctoral Fellowship Program of CPSF under Grant Number GZC20240346.

Author contributions

Conceptualization: ZLL, DFB. Formal analysis: YW. Funding acquisition: ZLL, JDB. Investigation: YW, DFB, HWS. Methodology: ZLL, JDB. Resources: YW, ZLL, DFB. Software: DFB. Supervision: ZLL, YF. Writing – original draft: YW, ZLL, JDB. Writing – review and editing: JDB, DFB, ZLL, HWS, YF, YW.

Author ORCIDs

Ying Wang  <https://orcid.org/0009-0004-1995-1671>
D. Jayarama Bhat  <https://orcid.org/0000-0002-3800-5910>
Dan-Feng Bao  <https://orcid.org/0000-0002-5697-4280>
Hong-Wei Shen  <https://orcid.org/0000-0003-2508-1970>
Zong-Long Luo  <https://orcid.org/0000-0001-7307-4885>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bao DF, Luo ZL, Liu JK, Bhat DJ, Sarunya N, Li WL, Su HY, Hyde KD (2018) Lignicolous freshwater fungi in China III: New species and record of *Kirschsteiniothelia* from northwestern Yunnan Province. *Mycosphere: Journal of Fungal Biology* 9: 755–768. <https://doi.org/10.5943/mycosphere/9/4/4>
- Bao DF, Hongsanan S, Hyde KD, Luo ZL, Nalumpang S (2022) *Pseudomonodictys aquatica* sp. nov., the sexual morph of *Pseudomonodictys* from freshwater habitats. *Phytotaxa* 567: 222–232. <https://doi.org/10.11646/phytotaxa.567.3.2>
- Boonmee S, D'souza MJ, Luo ZL, Pinruan U, Tanaka K, Su HY, Bhat DJ, McKenzie EHC, Gareth Jones EB, Taylor JE, Phillips AJL, Hirayama K, Eungwanichayapant PD, Hyde KD (2016) Dictyosporiaceae fam. nov. *Fungal Diversity* 80: 457–482. <https://doi.org/10.1007/s13225-016-0363-z>
- Calabon MS, Hyde KD, Gareth Jones EB, Luo ZL, Dong W, Hurdeal VG, Gentekaki E, Rossi W, Leonardi M, Thiagaraja V, Lestari AS, Shen HW, Bao DF, Boonyuen N, Zeng M (2022) Freshwater fungal numbers. *Fungal Diversity* 114: 3–235. <https://doi.org/10.1007/s13225-022-00503-2>
- Calabon MS, Gareth JEB, Pang KL, Abdel-Wahab MA, Jin J, Devadatha B, Sadaba RB, Apurillo CC, Hyde KD (2023) Updates on the classification and numbers of marine fungi. *Botanica Marina* 66: 213–238. <https://doi.org/10.1515/bot-2023-0032>
- Chethana KWT, Manawasinghe IS, Hurdeal VG, Bhunjun CS, Appadoo MA, Gentekaki E (2021) What are fungal species and how to delineate them? *Fungal Diversity* 109: 1–25. <https://doi.org/10.1007/s13225-021-00483-9>
- Corda ACI (1836) Mykologische Beobachtungen. In Weitenweber, Beitrage zur gesammten Natur- und Heilwissenschaften.
- Crous PW, Groenewald JZ, Shivas RG, Edwards J, Seifert KA, Alfenas AC, Alfenas RF, Burgess TI, Carnegie AJ, Hardy GESJ, Hiscock N, Hüberli D, Jung T, Louis-Seize G, Okada G, Pereira OL, Stukely MJC, Wang W, White GP, Young AJ, McTaggart AR, Pascoe IG, Porter IJ, Quaedvlieg W (2011) Fungal Planet Description Sheets: 69–91. *Persoonia* 26: 108–156. <https://doi.org/10.3767/003158511X581723>
- Dagona AG (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic acids symposium series*.
- Dong W, Wang B, Hyde KD, McKenzie EHC, Bhat DJ, Raja HA, Tanaka K, Abdel-Wahab MA, Abdel-Aziz FA, Doilom M, Phookamsak R, Hongsanan S, Wanasinghe DN, Yu XD, Wang GN, Yang H, Yang J, Thambugala AN, Tian Q, Luo ZL, Yang JB, Miller AN, Fournier J, Boonmee S, Hu DM, Nalumpang S, Zhang H (2020) Freshwater Dothideomycetes. *Fungal Diversity* 105: 319–575. <https://doi.org/10.1007/s13225-020-00463-5>
- Dong W, Hyde KD, Jeewon R, Liao CF, Zhao HJ, Kularathnage ND, Li H, Yang YH, Pem D, Shu YX, Gafforov Y, Manawasinghe IS, Doilom M (2023) *Mycosphere notes* 449–468:

- Saprobic and endophytic fungi in China, Thailand, and Uzbekistan. *Mycosphere: Journal of Fungal Biology* 14: 2208–2262. <https://doi.org/10.5943/mycosphere/14/1/26>
- Du HZ, Liu NG, Wu N, Cheewangkoon R, Liu JK (2025) Morpho-phylogenetic evidence reveals novel hyphomycetous fungi on medicinal plants in Southwestern China. *Mycology*. <https://doi.org/10.1080/21501203.2024.2444436>
- Dubey R (2022) *Dictyosporium matherense* sp. nov.: A new-fangled cheirosporous fungal species described from the Western Ghats of India. *Asian Journal of Forestry* 6: 1–8. <https://doi.org/10.13057/asianjfor/r060101>
- Feng Y, Liu ZY, Chen XF, Yang ML, Zhang ZY, Chen YY (2024) Phylogeny and taxonomy of two new species in Dictyosporiaceae (Pleosporales, Dothideomycetes) from Guizhou, China. *MycKeys* 106: 251–264. <https://doi.org/10.3897/mycokeys.106.125693>
- Fu X, Bao DF, Luo ZL, He X, Su HY (2021) Two new species of *Jalapriya* and a new record, *Dictyocheirospora vinaya* from freshwater habitats in China. *Biodiversity Data Journal* 9: e74295. <https://doi.org/10.3897/BDJ.9.e74295>
- Ghosh AK, Kar R, Chatterjee R (2015) Leaf galls on *Dicrodium hughesii* (Feistmantel) Lele from the Triassic of India – A new record. *Alcheringa. An Australasian Journal of Palaeontology* 39: 92–98. <https://doi.org/10.1080/03115518.2015.958285>
- Goh TK, Hyde KD, Ho WH, Yan N (1999) A revision of the genus *Dictyosporium*, with descriptions of three new species. *Fungal Diversity* 2: 65–100.
- Han M, Lu L, Karunarathna SC, Zheng D, Elgorban AM, Tibpromma S (2024) Morpho-molecular analyses reveal *Aquadictyospora yunnanensis* sp. nov. from coffee (*Coffea arabica*) in Yunnan Province, China. *Phytotaxa* 674(3): 239–251. <https://doi.org/10.11646/phytotaxa.674.3.1>
- Henningsson M (1974) Aquatic lignicolous fungi in the Baltic and along the west coast of Sweden. *Svensk Botanisk Tidskrift* 68: 401–425.
- Hyde KD, Norphanphoun C, Abreu VP, Bazzicalupo A, Chethana KWT, Clericuzio M, Dayarathne D, Dissanayake MC, Ekanayaka AJ, He AH, Hongsanan MQ, Huang S, Jayasiri SK, Jayawardena SC, Karunarathna RS, Konta A, Kušan S, Lee I, Li H, Lin J, Liu CG, Lu NG, Luo YZ, Manawasinghe ZL, Mapook IS, Perera A, Phookamsak RH, Phukhamsakda R, Siedlecki C, Soares I, Tennakoon AM, Tian DS, Tibpromma Q, Wanasinghe S, Xiao DN, Yang YP, Zeng J, Abdel-Aziz XY, Li FA, Senanayake WJ, Shang IC, Daranagama QJ, de Silva DA, Thambugala NI, Abdel-Wahab KM, Bahkali MA, Berbee AH, Boonmee ML, Bhat S, Bulgakov DJ, Buyck TS, Camporesi B, Castañeda-Ruiz E, Chomnunti RF, Doilom P, Dovana M, Gibertoni F, Jadan TB, Jeewon M, Jones R, Kang EBG, Karunarathna JC, Lim SC, Liu YW, Liu JK, Plautz ZY, Jr HL, Lumyong S, Maharachchikumbura SSN, Matočec N, McKenzie EHC, Mešić A, Miller D, Pawłowska J, Pereira OL, Promputtha I, Romero AI, Ryvarden L, Su HY, Suetrong S, Tkáčec Z, Vizzini A, Wen TC, Wisitrassameewong K, Wrzosek M, Xu JC, Zhao Q, Zhao RL, Mortimer PE (2017) Fungal diversity notes 603–708: Taxonomic and phylogenetic notes on genera and species. *Fungal Diversity* 87: 1–235. <https://doi.org/10.1007/s13225-017-0391-3>
- Jayasiri SC, Hyde KD, Jones EBG, McKenzie EHC, Jeewon R, Phillips AJL, Bhat DJ, Wanasinghe DN, Liu JK, Lu YZ, Kang JC, Xu J, Karunarathna SC (2019) Diversity, morphology and molecular phylogeny of Dothideomycetes on decaying wild seed pods and fruits. *Mycosphere: Journal of Fungal Biology* 10: 1–186. <https://doi.org/10.5943/mycosphere/10/1/1>
- Jiang HB, Jeewon R, Karunarathna SC, Phukhamsakda C, Doilom M, Kakumyan P, Suwannarach N, Phookamsak R, Lumyong S (2021) Reappraisal of *Immotthia* in Dictyosporiaceae, Pleosporales: Introducing *Immotthia bambusae* sp. nov. and *Pseudocyclothyriella clematidis* comb. et gen. nov. Based on Morphology and Phylogeny. *Frontiers in Microbiology* 12: 656235. <https://doi.org/10.3389/fmicb.2021.656235>

- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20: 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Kirschner R, Pang KL, Gareth Jones EB (2013) Two cheirosporous hyphomycetes reassessed based on morphological and molecular examination. *Mycological Progress* 12: 29–36. <https://doi.org/10.1007/s11557-012-0812-3>
- Larsson A (2014) AliView: Afastandlightweight alignment viewer and editor for large datasets. *Bioinformatics (Oxford, England)* 30(22): 3276–3278. <https://doi.org/10.1093/bioinformatics/btu531>
- Li WL, Bao DF, Bhat DJ, Su HY (2020) *Tetraploa aquatica* (Tetraplosphaeriaceae), a new freshwater fungal species from Yunnan Province, China. *Phytotaxa* 459: 181–189. <https://doi.org/10.11646/phytotaxa.459.2.8>
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16: 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Liu JK, Phookamsak R, Doilom M, Wikee S, Li YM, Ariyawansa H, Boonmee S, Chomnunti P, Dai DQ, Bhat JD, Romero AI, Zhuang WY, Monkai J, Jones EBG, Chukeatirote E, Ko Ko TW, Zhao YC, Wang Y, Hyde KD (2012) Towards a natural classification of Botryosphaeriales. *Fungal Diversity* 57: 149–210. <https://doi.org/10.1007/s13225-012-0207-4>
- Liu JK, Hyde KD, Jones EBG, Ariyawansa HA, Bhat DJ, Boonmee S, Maharachchikumbura SSN, McKenzie EHC, Phookamsak R, Phukhamsakda C, et al. (2015) Fungal diversity notes 1–110: Taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity* 72: 1–197. <https://doi.org/10.1007/s13225-015-0324-y>
- Liu LL, Song LC, Gu XF, Wei QQ, Zhang M, Liu ZY, Gou JL (2023) *Dictyosporium duliujian-gense* sp. nov. (Dictyosporiaceae, Pleosporales) from freshwater habitat in Guizhou Province, China. *Phytotaxa* 606: 259–272. <https://doi.org/10.11646/phytotaxa.606.4.2>
- Liu SL, Zhao P, Cai L, Shen S, Wei HW, Na Q, Han M, Wei R, Ge Y, Ma H, Karunarathna SC, Tibprommab S, Zhang B, Dai D, Lin L, Fan XL, Luo ZL, Shen HW, Lu L, Lu W, Xu RF, Tohtir-jap A, Wu F, Zhou LW (2024) Catalogue of fungi in China 1. New taxa of plant-inhabiting fungi. *Mycology* 16(1): 1–58. <https://doi.org/10.1080/21501203.2024.2316066>
- Luo J, Yin JF, Cai L, Hyde KD (2004) Freshwater fungi in Lake Dianchi, a heavily polluted lake in Yunnan, China[J]. *Fungal Diversity* 16: 93–112.
- Luo ZL, Bhat DJ, Jeewon R, Boonmee S, Bao DF, Zhao YC, Chai HM, Su XJ, Hyde KD (2017) Molecular phylogeny and morphological characterization of asexual fungi (Tubeufiaceae) from freshwater habitats in Yunnan, China. *Mycologie* 38: 27–53. <https://doi.org/10.7872/crym/v38.iss1.2017.27>
- Luo ZL, Hyde KD, Bhat DJ, Jeewon R, Maharachchikumbura SSN, Bao DF, Li WL, Su HY, Yang XY (2018a) Morphological and molecular taxonomy of novel species Pleurotheciaceae from freshwater habitats in Yunnan, China. *Mycological Progress* 17: 511–530. <https://doi.org/10.1007/s11557-018-1377-6>
- Luo ZL, Hyde KD, Liu JK, Bhat DJ, Bao DF, Li WL, Su HY (2018b) Lignicolous freshwater fungi from China II: Novel *Distoseptispora* (Distoseptisporaceae) species from north-western Yunnan Province and a suggested unified method for studying lignicolous freshwater fungi. *Mycosphere: Journal of Fungal Biology* 9: 444–461. <https://doi.org/10.5943/mycosphere/9/3/2>
- Luo ZL, Hyde KD, Liu JK, Maharachchikumbura SSN, Jeewon R, Bao DF, Bhat DJ, Lin CG, Li WL, Yang J, Liu NG, Lu YZ, Jayawardena RS, Li JF, Su HY (2019) Freshwater Sordariomycetes. *Fungal Diversity* 99: 451–660. <https://doi.org/10.1007/s13225-019-00438-1>

- McKenzie EHC (2010) Two new dictyosporous hyphomycetes on *Rhopalostylis sapida* (Arecaceae) in New Zealand. *Mycotaxon* 111: 155–160. <https://doi.org/10.5248/111.155>
- Nylander JAA (2004) MrModeltest 2.0. Program distributed by the author. Evolutionary Biology Centre, Uppsala University, Sweden.
- Phookamsak R, Hyde KD, Jeewon R, Bhat DJ, Jones EBG, Maharachchikumbura SSN, Raspé O, Karunarathna SC, Wanasinghe DN, Hongsanan S, Doilom M, Tennakoon DS, Machado AR, Firmino AL, Ghosh A, Karunarathna A, Mešić A, Dutta AK, Thongbai B, Devadatha B, Norphanphoun C, Senwanna C, Wei D, Pem D, Ackah FK, Wang GN, Jiang HB, Madrid H, Lee HB, Goonasekara ID, Manawasinghe IS, Kušan I, Cano J, Gené J, Li J, Das K, Acharya K, Anil Raj KN, Deepna Latha KP, Chethana KWT, He MQ, Dueñas M, Jadan M, Martín MP, Samarakoon MC, Dayarathne MC, Raza M, Park MS, Telleria MT, Chaiwan N, Matočec N, de Silva NI, Pereira OL, Singh PN, Manimohan P, Uniyal P, Shang QJ, Bhatt RP, Perera RH, Alvarenga RLM, Nogal-Prata S, Singh SK, Vadthanarat S, Oh SY, Huang SK, Rana S, Konta S, Paloi S, Jayasiri SC, Jeon SJ, Mehmood T, Gibertoni TB, Nguyen TTT, Singh U, Thiyagaraja V, Sarma VV, Dong W, Yu XD, Lu YZ, Lim YW, Chen Y, Tkalčec Z, Zhang ZF, Luo ZL, Daranagama DA, Thambugala KM, Tibpromma S, Camporesi E, Bulgakov TS, Dissanayake AJ, Senanayake IC, Dai DQ, Tang LZ, Khan S, Zhang H, Promputtha I, Cai L, Chomnunti P, Zhao RL, Lumyong S, Boonmee S, Wen TC, Mortimer PE, Xu J (2019) Fungal diversity notes 929–1035: Taxonomic and phylogenetic contributions on genera and species of fungi. *Fungal Diversity* 95: 1–273. <https://doi.org/10.1007/s13225-019-00421-w>
- Phookamsak R, Hongsanan S, Jayarama BD, Wanasinghe DN, Promputtha I, Suwanarach N, Kumla J, Xie N, Dawoud TM, Mortimer PE, Xu J, Lumyong S (2024) Exploring ascomycete diversity in Yunnan II: Introducing three novel species in the suborder Massarineae (Dothideomycetes, Pleosporales) from fern and grasses. *MycKeys* 104: 9–50. <https://doi.org/10.3897/mycokeys.104.112149>
- Prasher IB, Verma RK (2015) Two new species of *Dictyosporium* from India. *Phytotaxa* 204(3): 193–202. <http://doi.org/10.11646/phytotaxa.204.3.2>
- Rannala B, Yang Z (1996) Probability distribution of molecular evolutionary trees: A new method of phylogenetic inference. *Journal of Molecular Evolution* 43: 304–311. <https://doi.org/10.1007/BF02338839>
- Rebeca AR, Cayetano GC, Isabel M, Cláudia P, Margarita M (2021) Fungal Biodiversity Mediates the Effects of Drying on Freshwater Ecosystem Functioning. *Ecosystems* (New York, N.Y.) 25: 780–794. <https://doi.org/10.1007/s10021-021-00683-z>
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systems Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Senanayake IC, Rathnayaka AR, Marasinghe DS, Calabon MS, Gentekaki E, Lee HB, Hurdeal VG, Pem D, Dissanayake LS, Wijesinghe SN, Bundhun D, Nguyen TTT, Goonasekara ID, Abeywickrama PD, Bhunjun CS, Jayawardena RS, Wanasinghe DN, Jeewon R, Bhat DJ, Xiang MM (2020) Morphological approaches in studying fungi: Collection, examination, isolation, sporulation and preservation. *Mycosphere: Journal of Fungal Biology* 11: 2678–2754. <https://doi.org/10.5943/mycosphere/11/1/20>
- Shen HW, Bao DF, Hyde KD, Su HY, Bhat DJ, Luo ZL (2021) Two novel species and two new records of *Distoseptispora* from freshwater habitats in China and Thailand. *MycKeys* 84: 79–101. <https://doi.org/10.3897/mycokeys.84.71905>

- Shen HW, Bao DF, Bhat DJ, Su HY, Luo ZL (2022a) Lignicolous freshwater fungi in Yunnan Province, China: An overview. *Mycology* 13: 119–132. <https://doi.org/10.1080/21501203.2022.2058638>
- Shen HW, Bao DF, Wanasinghe DN, Boonmee S, Liu JK, Luo ZL (2022b) Novel species and records of Dictyosporiaceae from freshwater habitats in China and Thailand. *Journal of Fungi* (Basel, Switzerland) 8: 1200. <https://doi.org/10.3390/jof8111200>
- Shu YX, Doilom M, Boonmee S, Xu B, Dong W (2024) Three Novel Cheiroid Hyphomycetes in *Dictyocheiropora* and *Dictyosporium* (Dictyosporiaceae) from Freshwater Habitats in Guangdong and Guizhou Provinces, China. *Journal of Fungi* (Basel, Switzerland) 10: 259. <https://doi.org/10.3390/jof10040259>
- Silva CR, Gusmão LFP, Castañeda-Ruiz RF (2016) *Dictyosporium amoenum* sp. nov. from Chapada, Diamantina, Bahia, Brazil. *Mycotaxon* 130: 1125–1133. <https://doi.org/10.5248/130.1125>
- Stamatakis, A (2006) RAXML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688–2690. <http://doi.org/10.1093/bioinformatics/btl446>
- Stamatakis A, Hoover P, Rougemont J (2008) A rapid bootstrap algorithm for the RAXML web servers. *Systematic Biology* 57: 758–771. <http://doi.org/10.1080/10635150802429642>
- Su HY, Hyde KD, Maharachchikumbura SSN, Ariyawansa HA, Luo ZL, Promputtha I, Tian Q, Lin CG, Shang QJ, Zhao YC (2016) The families Distoseptisporaceae fam. nov., Kirschsteinioteliaceae, Sporormiaceae and Torulaceae, with new species from freshwater in Yunnan Province, China. *Fungal Diversity* 80: 375–409. <https://doi.org/10.1007/s13225-016-0362-0>
- Sun YR, Hyde KD, Liu NG, Jayawardena RS, Wijayawardene NN, Ma J, Zhang Q, Al-Otibi F, Wang Y (2024) Micro-fungi on medicinal plants in southern China and northern Thailand. *Research Square*. <https://doi.org/10.21203/rs.3.rs-4813665/v1>
- Sushma, Verma RK, Prasher IB, Gautam AK, Rajeshkumar KC, Castañeda-Ruiz RF (2022) *Dictyocheiropora himachalensis* sp. nov. from Himachal Pradesh, India. *Mycotaxon* 137(3): 455–463. <https://doi.org/10.5248/137.455>
- Tamam EE, Huzefa A, Raja MF, Ahmed HAS, Rick LB, Nicholas HO (2021) Freshwater fungi as a source of chemical diversity: A review. *Journal of Natural Products* 84(3): 898–916. <https://doi.org/10.1021/acs.jnatprod.0c01340>
- Tanaka K, Hirayama K, Yonezawa H, Sato G, Toriyabe A, Kudo H, Hashimoto A, Matsumura M, Harada Y, Kurihara Y, Shirouzu T, Hosoya T (2015) Revision of the Massariaceae (Pleosporales, Dothideomycetes). *Studies in Mycology* 82: 75–136. <https://doi.org/10.1016/j.simyco.2015.10.002>
- Tennakoon DS, de Silva NI, Maharachchikumbura SN, Bhat DJ, Kumla J, Suwannarach N, Lumyong S (2023) Exploring more on Dictyosporiaceae: The species geographical distribution and intriguing novel additions from plant litter. *Diversity* 15(3): 410. <https://doi.org/10.3390/d15030410>
- Tibell S, Tibell L, Pang KL, Calabon M, Gareth Jones EB (2020) Marine fungi of the Baltic Sea. *Mycology* 11: 195–213. <https://doi.org/10.1080/21501203.2020.1729886>
- Tibpromma S, Hyde KD, Bhat DJ, Mortimer PE, Xu J, Promputtha I, Doilom M, Yang JB, Tang AM, Karunaratna SC (2018a) Identification of endophytic fungi from leaves of Pandanaceae based on their morphotypes and DNA sequence data from southern Thailand. *MycKeys* 33: 25–67. <https://doi.org/10.3897/mycokeys.33.23670>
- Tibpromma S, Hyde KD, McKenzie EHC, Jayarama BD, Phillips AJD, Wanasinghe DN, Samarakoon MC, Jayawardena RS, Dissanayake AJ, Tennakoon DS, Doilom M, Phookamsak R, Tang AMC, Xu JC, Mortimer PE, Promputtha I, Maharachchikumbura SSN, Khan S,

- Karunarathna SC (2018b) Fungal diversity notes 840–928: Micro-fungi associated with Pandanaceae. *Fungal Diversity* 93: 1–160. <https://doi.org/10.1007/s13225-018-0408-6>
- Vaidya G, Lohman DJ, Meier R (2011) SequenceMatrix: Concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* 27: 171–180. <https://doi.org/10.1111/j.1096-0031.2010.00329.x>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Wang RX, Luo ZL, Hyde KD, Bhat DJ, Su XJ, Su HY (2016) New species and records of *Dictyocheiropora* from submerged wood in north-western Yunnan, China. *Mycosphere: Journal of Fungal Biology* 7(9): 1357–1367. <https://doi.org/10.5943/mycosphere/7/9/9>
- Wang WP, Hyde KD, Bao DF, Wanasinghe DN, Lin CG, Shen HW, Lu YZ, Zhang ZQ, Su XJ, Li YX, Al-Otibi F, Yang LQ, Luo ZL (2024) Lignicolous freshwater fungi from karst in Yunnan Province, China. *Mycosphere: Journal of Fungal Biology* 15(1): 6525–6640. <https://doi.org/10.5943/mycosphere/15/1/29>
- Wijayawardene NN, Hyde KD, Dai D, Sánchez-García M, Goto BT, Saxena RK, Erdoğan M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszczowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunarathna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkálčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalaseana SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang LZ, Stadler M, Yurkov A, Thines M (2022) Outline of Fungi and fungus-like taxa – 2021. *Mycosphere: Journal of Fungal Biology* 13: 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Wu N, Du HZ, Chethana KWT, Khongphinitbunjong K, Maharachchikumbura SSN, Hyde KD, Liu JK (2024) Additions to Dictyosporiaceae: *Neoxylochrysis typhicola* comb. et gen. nov., two new species and four new host records from medicinal plants in Southwestern China. *Journal of Fungi* (Basel, Switzerland) 10: 872. <https://doi.org/10.3390/jof10120872>
- Yang J, Liu JK, Hyde KD, Gareth Jones EB, Liu ZY (2018) New species in *Dictyosporium*, new combinations in *Dictyocheiropora* and an updated backbone tree for Dictyosporiaceae. *MycKeys* 36: 83–105. <https://doi.org/10.3897/mycokeys.36.27051>
- Zhang JF, Liu JK, Hyde KD, Chen YY, Ran HY, Liu ZY (2023) Ascomycetes from karst landscapes of Guizhou Province, China. *Fungal Diversity* 122: 1–160. <https://doi.org/10.1007/s13225-023-00524-5>
- Zhang L, Bao DF, Shen HW, Luo ZL (2024b) Diversity of Lignicolous Freshwater Fungi from Yuanjiang River in Yunnan (China), with the Description of Four New Species. *Journal of Fungi* (Basel, Switzerland) 10: 881. <https://doi.org/10.3390/jof10120881>
- Zhang WJ, Xu GP, Liu Y, Gao Y, Song HY, Hu HJ, Zhou JP, Chen MH, Fan DM, Hu DM, Zhai ZJ (2024a) Multi-gene phylogenetic analyses revealed two novel species and one new record of *Trichobotrys* (Pleosporales, Dictyosporiaceae) from China. *MycKeys* 106: 117–132. <https://doi.org/10.3897/mycokeys.106.123279>