



# Scientific landscape and trend analysis of keloid research: a 30-year bibliometric review

Yuming Chong<sup>1</sup>, Xiao Long<sup>1</sup>, Yuh-Shan Ho<sup>2</sup>

<sup>1</sup>Department of Plastic and Aesthetic Surgery, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, China; <sup>2</sup>Trend Research Centre, Asia University, Taichung 41354

**Contributions:** (I) Conception and design: All authors; (II) Administrative support: X Long; (III) Provision of study materials or patients: YS Ho; (IV) Collection and assembly of data: YS Ho; (V) Data analysis and interpretation: Y Chong, YS Ho; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

**Correspondence to:** Xiao Long. Peking Union Medical College Hospital, No. 1, Shuaifuyuan, Dongcheng District, Beijing, China. Email: longxiao@pumch.cams.cn; xiao.longpumch@outlook.com; Yuh-Shan Ho. Trend Research Centre, Asia University, No. 500, Lioufeng Road, Wufeng, Taichung 41354. Email: ysho@asia.edu.tw.

**Background:** Keloids remain troublesome for clinicians because of the lack of standard therapy, and the underlying mechanisms are still unclear. Bibliometric analysis could be a powerful tool to comprehensively review research trends, evaluate publication performances and provide future perspectives. There is no bibliometric analysis focusing on keloid research.

**Methods:** Keloid related publications were searched in Web of Science Core Collection from the publication year 1990 to 2019. Bibliometric data were provided on document type, annual publication number, most productive journals, publication geography, top potential authors, and highly cited articles. The distribution of single words in article titles was analyzed to evaluate the main research focuses and determine their development trends. Word cluster analysis was further performed to detect emerging trends in keloid research.

**Results:** The number of annual articles increased from 24 in 1990 to 63 in 2006 and then increased sharply, reaching 139 in 2019. *Dermatologic Surgery* published the highest number of articles followed by *Plastic and Reconstructive Surgery*. The USA ranked top with six publication indicators and China had a dramatic increase in the annual number of articles since 2012. Recent research hotspots include underlying pathogenetic mechanisms, keloid treatment, and therapeutic effect evaluation.

**Discussion:** Keloids remain a research focus. Efforts will be continuously made to understand the underlying mechanism of keloid formation. Despite many treatment modalities, there is no gold standard for keloid treatment, and many efforts are being made in the exploration of new therapies. Moreover, it is foreseeable that objective measurement tools will have a higher status in the assessment of keloids and scars.

**Keywords:** Keloid; bibliometric; citation analysis; Web of Science Core Collection; research trends

Submitted Feb 02, 2021. Accepted for publication Apr 23, 2021.

doi: 10.21037/atm-21-508

View this article at: <http://dx.doi.org/10.21037/atm-21-508>

## Introduction

Keloids are characterized by aberrant production of collagen fibers that are longer, thicker, and more randomly oriented than normal (1,2). Other biochemical features include a high fibroblast proliferation rate, increased

enzyme concentrations, and growth factors stimulation (3-5). Keloids can lead to pain, pruritus, and cosmetic problems (6). Many efforts have been made to establish a gold standard in keloid treatment, but to date, a consensus has not been reached for standard therapy (7).

Bibliometric analysis provides information on the annual

number of publications, citation life of articles, topic-related journals, publication performances of authors and institutions, and research trends. These bibliometric profiles illustrate important studies, the authority, journals that favor related publications, and the focus of future studies. Experienced researchers who are eager to know the frontier or individuals who intend to start research will benefit from a bibliometric study.

Bibliometrics has been applied to other medical topics including pain, Ebola, dengue, HIV, and pneumonia (8-12). It is surprising that there is no bibliometric analysis focusing on such an old disease which has been intensively studied because of unsettled problems concerning not only pathogenesis but also treatment. Bibliometric analysis could be a powerful tool to comprehensively review research trends, evaluate publication performances and provide future perspectives. This bibliometric study aims to map the research landscape of keloids and, more importantly, identify research trends to assist researchers in future directions.

We present the following article in accordance with the PRISMA reporting checklist (available at <http://dx.doi.org/10.21037/atm-21-508>).

## Methods

It has been pointed out that it is not appropriate to use all databases which are not in the same level in Web of Science for bibliometric studies (13,14). The data in this study was through the Science Citation Index Expanded (SCI-EXPANDED) database of the Clarivate Analytics Web of Science Core Collection (Data last updated on March 10, 2021). After our pre-study, the search keywords: “keloid”, “keloids”, “keloider”, “keloides”, “keloide”, “keloidi”, “keloidentwicklung”, “keloiddisposition”, “keloidsden”, “keloidlike”, “keloidogenesis”, “keloidosis”, “keloidiformis”, “keloidectomy”, “keloidal”, “intrakeloid”, “narbenkeloid”, and “antikeloid” were searched in the Topic field, including title, abstract, author keywords, and *KeyWords Plus*, in the Web of Science Core Collection within the publication year limitation from 1990 to 2019. The retrieving time of a bibliometric analysis usually starts from 1991 because many publications in SCI-EXPANDED before that are lack of abstracts. However, in this study, the publication year was limited from 1990 because in this way the research focuses and development trends could be observed in the past three decades (1990s, 2000s, and 2010s). *KeyWords Plus* supplies additional search terms extracted from the titles of articles cited by authors in their bibliographies and footnotes in the

ISI (now Clarivate Analytics) database, and substantially augments title-word and author-keyword indexing (15). This resulted 3,997 documents (not including the document type of early access) as keloid related publications. Those documents only found by *KeyWords Plus* are more likely to be unrelated to the “keloid” (16). Ho’s group firstly proposed the “front page” filter (17-19)—which covers only documents with searching keywords in their “front page”, including only the title, abstract, and author keywords; this filter might avoid introducing unrelated publications for analysis (17). Finally, 3,394 documents (85% of 3,997 documents) were defined as keloid research publications. The full record of SCI-EXPANDED and the number of citations in each year for each document were downloaded into Microsoft Excel 2016 and checked. Additional coding was manually performed (19,20). The journal impact factors ( $IF_{2019}$ ) were taken from the Journal Citation Reports (JCR) published in 2019.

In the SCI-EXPANDED database, the corresponding author is designated as the “reprint” author; “corresponding author” will remain as the primary terminology instead of reprint author (21). In a single-author article where authorship is unspecified, the single author is considered both the first author and the corresponding author (22). Similarly, in an individually institutional article, the institution is both classified as the first institution and the corresponding institution (22). In cases of multiple corresponding authors, only the last corresponding author and their affiliation were considered (23). In cases of single-author articles, only the first affiliation and first country were considered.

Affiliations in England, Scotland, Northern Ireland, and Wales were reclassified as in the United Kingdom (UK) (24). Affiliations of authors in Hong Kong prior to 1997 were included under the heading of China (25). Affiliations in the Union of Soviet Socialist Republics (USSR) were checked and reclassified as in Russia and Ukraine (19). Similarly, USSR Academy of Medical Sciences (Acad Med Sci USSR) was also checked and re-classified under the heading of Russian Academy of Medical Sciences (Russian Acad Med Sci).

To investigate the citations received by the publications, we used four citation indicators:

- ❖  $C_0$ : the the number of citations from the Web of Science Core Collection in publication year (26);
- ❖  $C_{year}$ : the number of citations from the Web of Science Core Collection in a particular year.  $C_{2019}$  means the number of citations in 2019 (21);

**Table 1** Citations and authors according to document type

Document type	<i>TP</i>	%	<i>AU</i>	<i>APP</i>	<i>TC</i> <sub>2019</sub>	<i>CPP</i> <sub>2019</sub>
Article	2,253	66	11,288	5.0	48,426	21
Meeting abstract	489	14	2,185	4.5	190	0.39
Review	264	7.8	1,025	3.9	11,208	42
Letter	257	7.6	826	3.2	1,315	5.1
Proceedings' paper	97	2.9	418	4.3	3,361	35
Editorial material	88	2.6	238	2.7	343	3.9
Correction	21	0.62	84	4.0	7	0.33
Note	19	0.56	60	3.2	447	24
Book chapter	7	0.21	23	3.3	423	60
Discussion	1	0.029	1	1.0	4	4.0
News item	1	0.029	0	0	0	0
Reprint	1	0.029	3	3.0	1	1.0

*TP*, number of publications; *AU*, number of authors; *APP*, number of authors per publication; *TC*<sub>2019</sub>, the total number of citations from Web of Science Core Collection since publication year to the end of 2019; *CPP*<sub>2019</sub>, number of citations (*TC*<sub>2019</sub>) per publication (*TP*).

- ❖ *TC*<sub>year</sub>: the total number citations from the Web of Science Core Collection since publication to the end of the most recent year. In this study this is 2019 (*TC*<sub>2019</sub>) (8,27);
- ❖ *CPP*<sub>year</sub>: citations per publication (*CPP*<sub>2019</sub> = *TC*<sub>2019</sub>/*TP*) (21). *TP* indicates the total number of publications.

## Results and discussion

### Characteristics of document types


Document type characteristics with their citations per publication ( $CPP_{\text{year}} = TC_{\text{year}}/TP$ ) and number of authors per publication ( $APP = AU/TP$ ) was recently suggested. The advantage of using *TC*<sub>2019</sub> for *CPP*<sub>2019</sub> is that they are invariable and ensure repeatability compared with the index of citation from Web of Science Core Collection (28). A total of 3,394 keloid related documents published in SCI-EXPANDED were found among 12 document types. *Table 1* shows characteristics of the 12 document types, including 2,253 articles (66% of 3,394 documents) with number of authors per publication (*APP*) of 5.0 which was higher than other document types. The percentage of articles (66%) was lower than other medical related topics, for example 79% in asthma in children (29), 75% human papillomavirus (30), and 70% in Ebola (9) but similar to cisplatin-containing

chemotherapy for small cell lung cancer (68%) (31), acupuncture (66%) (32).

A total of 489 meeting abstracts were published in 60 different journals but mainly focused on three journals: *Journal of Investigative Dermatology* (115 meeting abstracts; 24% of 489 meeting abstracts) with a *CPP*<sub>2019</sub> of 0.30, *Wound Repair and Regeneration* (95; 19%) with a *CPP*<sub>2019</sub> of 1.3, and *Journal of the American Academy of Dermatology* (48; 10%) with a *CPP*<sub>2019</sub> of 0.10.

The document type of book chapters with seven documents had the greatest *CPP*<sub>2019</sub> value of 60 followed by reviews with a *CPP*<sub>2019</sub> of 42. The *CPP*<sub>2019</sub> of document type of reviews was found to be double of articles. It should be pointed out that documents could be classified in two document types in Web of Science, for example, the highly cited document entitled “Basic components of connective tissues and extracellular matrix: Elastin, fibrillin, fibulins, fibrinogen, fibronectin, laminin, tenascins and thrombospondins” (33) was classified as “article” and “book chapter”; thus, the sum of percentages was higher than 100%.

Only the 2,253 articles were selected for further analysis on the basis of containing whole research ideas and results (34). There were nine languages in use and one article was unspecified in SCI-EXPANDED. English, as the most popular language, comprised 94% of the 2,253 keloid articles. Other non-English languages



**Figure 1** Number of keloid articles and citations per publication by year.

that were less used were listed as follows, German (41 articles), French (34), Chinese (6), Russian (5), Portuguese (3), Japanese (2), and one in each of Polish and Spanish, respectively.

### Characteristics of publication outputs

The use of  $TC_{2019}$  for determination of  $CPP_{2019}$  is preferred as compared with the citation index from Web of Science Core Collection directly because they are consistent and ensure repeatability (35). In order to understand publications and their impact trends in a research field, Ho proposed a relationship between total number of articles ( $TP$ ) in a year and their citations per publication ( $CPP_{year}$ ) by the decades (21) and years (31). In recent years, it has been applied in medical related topics, for example pain (8), Ebola (9), and dengue (10). A total of the 2,253 keloid related articles were published between 1990 and 2019. The mean value of  $TC_{2019}$  was 21 with 492 as the maximal value for an article. *Figure 1* demonstrated the distribution of the annual number of articles ( $TP$ ) and their citations per publication ( $CPP_{2019}$ ) by year, which was expressed as  $TC_{2019}/TP$  (22), where  $TP$  is number of articles published in that particular year. From 1990 through 2006, the annual number of articles increased from 24 in

1990 to 63 in 2006 and then sharply increasing to reach 139 in 2019.

### Web of science category and journal

JCR indexed 9,381 journals with citation references across 178 Web of Science categories in SCI-EXPANDED in 2019. In order to know development among research fields and their interactions, a relationship between the number of articles in categories and publication years was reported (34). The keloid related articles were published by 635 journals among the 98 Web of Science categories in SCI-EXPANDED. A total of 1,348 articles (60% of 2,253 articles) were published in the top two Web of Science categories: dermatology with 850 articles (38% of 2,253 articles) and surgery with 722 articles (32%). In 2019, a total 68 journals and 210 journals were classified to categories of dermatology and surgery, respectively. Other top ten productive categories were research and experimental medicine (183 articles; 8.1% of 2,253 articles), cell biology (149; 6.6%), pathology (128; 5.7%), oncology (104; 4.6%), biochemistry and molecular biology (98; 4.3%), general and internal medicine (85; 3.8%), otorhinolaryngology (74; 3.3%), and pharmacology and pharmacy (67; 3.0%). It has been noticed that journals

**Table 2** The top 11 most productive journals

Journal	TP (%)	IF <sub>2019</sub>	APP	CPP <sub>2019</sub>	Web of Science category
<i>Dermatologic Surgery</i>	83 (3.7)	2.567	4.0	33	Dermatology; surgery
<i>Plastic and Reconstructive Surgery</i>	81 (3.6)	4.235	5.0	41	Surgery
<i>Annals of Plastic Surgery</i>	58 (2.6)	1.354	4.9	23	Surgery
<i>Journal of Investigative Dermatology</i>	56 (2.5)	7.143	7.1	50	Dermatology
<i>British Journal of Dermatology</i>	50 (2.2)	7.000	5.8	35	Dermatology
<i>Wound Repair and Regeneration</i>	48 (2.1)	2.471	5.8	30	Cell biology; dermatology; research and experimental medicine; surgery
<i>International Journal of Dermatology</i>	40 (1.8)	2.067	4.3	16	Dermatology
<i>Archives of Dermatological Research</i>	39 (1.7)	2.339	5.6	22	Dermatology
<i>Experimental Dermatology</i>	32 (1.4)	3.368	6.1	19	Dermatology
<i>Aesthetic Plastic Surgery</i>	30 (1.3)	1.798	4.2	18	Surgery
<i>Burns</i>	30 (1.3)	2.066	5.2	15	Critical care medicine; dermatology; surgery

TP, number of articles; %, the percentage of articles in total publications; IF<sub>2019</sub>, journal impact factor in 2019; APP, number of authors per publication; CPP<sub>2019</sub>, number of citations (TC<sub>2019</sub>) per publication (TP).

could be classified in two or more categories in Web of Science, for example *Wound Repair and Regeneration* was classified in the categories of cell biology, dermatology, research and experimental medicine, and surgery thus the sum of percentages was higher than 100% (22).

In total 2,253 keloid related articles were published in 635 journals including 565 were listed in SCI-EXPANDED in 2019. The top 11 productive journals with 30 articles or more were listed in Table 2. *Dermatologic Surgery* (IF<sub>2019</sub> =2.567) published the most articles with 83 articles (3.7% of 2,253 articles) followed by *Plastic and Reconstructive Surgery* (IF<sub>2019</sub> =4.235) with 81 articles (3.6%). Compare the top 11 journals, keloid articles published in *Journal of Investigative Dermatology* (IF<sub>2019</sub> =7.143) had the highest CPP<sub>2019</sub> of 50 while articles in *Burns* (IF<sub>2019</sub> =2.066) had a CPP<sub>2019</sub> of 15. Articles published in *Journal of Investigative Dermatology* also had the highest APP of 7.1 while articles in *Dermatologic Surgery* (IF<sub>2019</sub> =2.567) had an APP of 4.0. Eight of the top 11 journals were classified in the Web of Science category of dermatology, six in surgery, and one in each of cell biology, critical care medicine, and research and experimental medicine, respectively. *Plastic and Reconstructive Surgery* (81 articles; ranked 2<sup>nd</sup>) published the most six keloid articles in 2019. The journal with the highest IF<sub>2019</sub> of 60.390 was *Lancet* followed by *Science* (IF<sub>2019</sub> =41.846) and *Nature Genetics* (IF<sub>2019</sub> =27.605) with one article, respectively.

### Publication performances: countries and institutions

To compare publication performance of countries and institutions, Ho's group proposed six publication indicators such as total number of articles (TP), single-country articles (IP), internationally collaborative articles (CP), first-author articles (FP), corresponding-author articles (RP), and single-author articles (SP) (24,34,36). First author and corresponding author are the two most contributed authors in an article (37). At the institutional level, the determined institution of the corresponding-author might be a home base of the study or origin of the paper (21). There were six articles without affiliations information in SCI-EXPANDED. Of 2,247 articles with author affiliations from 86 countries, 1,941 (86% of 2,247 articles) were single-country articles from 69 countries and 306 (14%) were internationally collaborative articles from 61 countries. The top 15 productive countries/regions are listed in Table 3 with six publication indicators (36) and a citation indicator (CPP<sub>2019</sub>). The advantage of using CPP<sub>2019</sub> is that it is invariant in comparison with the index of citations from the Web of Science Core Collection which has to be updated from time to time (17). Seven Asian countries, five European countries, and three American countries were ranked on the top 15 of publications. USA ranked top in the six publication indicators with a TP of 608 articles (27% of 2,247 articles), an IP of 470 articles


**Table 3** Top 15 productive countries/regions with six publication indicators and their  $CPP_{2019}$ 

Countries/regions	<i>TP</i>	<i>TP R</i> (%)	<i>IP R</i> (%)	<i>CP R</i> (%)	<i>FP R</i> (%)	<i>RP R</i> (%)	<i>SP R</i> (%)	$CPP_{2019}$
USA	608	1 (27)	1 (24)	1 (45)	1 (24)	1 (24)	1 (45)	34
China	341	2 (15)	2 (15)	2 (19)	2 (14)	2 (14)	12 (1.6)	10
Japan	202	3 (9.0)	3 (8.4)	5 (13)	3 (8.1)	3 (8.1)	5 (3.9)	29
Germany	177	4 (7.9)	4 (6.1)	4 (19)	4 (6.1)	4 (6.1)	4 (4.7)	29
UK	162	5 (7.2)	5 (5.3)	2 (19)	5 (5.9)	5 (5.6)	2 (5.5)	28
South Korea	128	6 (5.7)	6 (5.3)	8 (8.5)	6 (5.0)	6 (5.2)	12 (1.6)	13
Italy	95	7 (4.2)	7 (3.2)	6 (10)	7 (3.4)	7 (3.3)	12 (1.6)	22
France	70	8 (3.1)	9 (2.8)	12 (4.9)	10 (2.6)	9 (2.7)	8 (2.3)	24
Brazil	66	9 (2.9)	10 (2.8)	14 (3.9)	9 (2.6)	10 (2.6)	19 (0.78)	16
India	66	9 (2.9)	8 (3.1)	21 (2.0)	8 (2.7)	8 (2.8)	8 (2.3)	16
Canada	61	11 (2.7)	14 (1.8)	7 (8.8)	12 (2.0)	15 (2.1)	6 (3.1)	27
Singapore	56	12 (2.5)	15 (1.5)	8 (8.5)	12 (2.0)	14 (2.1)	12 (1.6)	32
Netherlands	52	13 (2.3)	13 (1.9)	11 (5.2)	12 (2.0)	11 (2.1)	N/A	22
Taiwan	50	14 (2.2)	12 (2.1)	17 (3.3)	11 (2.2)	11 (2.1)	19 (0.78)	21
Turkey	48	15 (2.1)	11 (2.2)	21 (2.0)	12 (2.0)	11 (2.1)	2 (5.5)	13

*TP*, number of total articles; *IP*, single-country articles; *CP*, internationally collaborative articles; *FP*, first-author articles; *RP*, corresponding-author articles, *SP*, single-author articles; *R*, rank;  $CPP_{2019}$ , number of citations ( $TC_{2019}$ ) per publication (*TP*); N/A, not available.

(24% of 1,941 single-country articles), a *CP* of 138 articles (45% of 306 internationally collaborative articles), an *FP* of 537 articles (24% of 2,247 first-author articles), an *RP* of 516 articles (24% of 2,193 corresponding-author articles), and an *SP* of 57 articles (45% of 128 single-author articles). Domination in keloid articles by the USA was not surprising since this pattern occurs in other medical topics: stem cells (38), obstetrics and gynecology (39), and orthopaedic surgery (40). USA also had the highest  $CPP_{2019}$  of 34. China ranked second in the four publication indicators such as *TP*, *IP*, *CP*, *FP*, and *RP* but had lower  $CPP_{2019}$  of 10. Among the 86 countries, 17 countries (20% of 86 countries) had no single-country articles while 25 countries (29%) had no internationally collaborative articles. In addition, 9 (10%), 13 (15%), and 56 (65%) countries have no first-author, corresponding-author, and single-author articles, respectively. *Figure 2* shows publication trends of the top six productive countries with 128 articles or more. Only China had a dramatic increase in annual number of articles since 2012. As for institutions, 925 keloid related articles (41% of 2,247 articles) were of single-institution and 1,322 articles (59%) were institutional collaborations. *Table 4* demonstrates the

characteristics of the top 15 productive institutions. Six of them were in USA, four in China, two in South Korea, and one in each of Japan, Singapore, and the UK, respectively. The University of Manchester in the UK took the leading position for four of the six publication indicators with a *TP* of 56 articles (2.5% of 2,247 articles), a *CP* of 45 articles (3.4% of 1,322 inter-institutionally collaborative articles), an *FP* of 44 articles (2.0% of 2,247 first-author articles), and an *RP* of 42 articles (1.9% of 2,193 corresponding-author articles) while the University of Miami in the USA ranked top in single-institution articles with an *IP* of 17 articles (1.8% of 925 single-institution articles). The St. Luke's-Roosevelt Hospital Center in the USA published 13 keloid articles (ranked 27th) including the most single-author articles with an *SP* of five articles (3.9% of 128 single-author articles). Compare the top 15 productive institutes, the Thomas Jefferson University in the USA had the highest  $CPP_{2019}$  of 55. Articles published by the institutes in the USA had higher  $CPP_{2019}$ . However, four Chinese institutions including the Fourth Military Medical University, the Shanghai Jiao Tong University, the Peking Union Medical College, and the Chinese Academy of Medical Sciences had lower  $CPP_{2019}$  of 16, 7.5, 7.4, and 6.8, respectively.



**Figure 2** Comparison of the growth trends of the top six productive countries ( $TP \geq 128$ ).

**Table 4** Top 15 productive institutions with six publication indicators and their  $CPP_{2019}$

Institute	TP	TPR (%)	IPR (%)	CPR (%)	FPR (%)	RPR (%)	SPR (%)	CPP <sub>2019</sub>
University of Manchester, UK	56	1 (2.5)	4 (1.2)	1 (3.4)	1 (2.0)	1 (1.9)	7 (0.78)	32
Shanghai Jiao Tong University, China	45	2 (2.0)	2 (1.3)	2 (2.5)	2 (1.5)	2 (1.6)	N/A	7.5
National University of Singapore, Singapore	41	3 (1.8)	2 (1.3)	5 (2.2)	5 (1.0)	4 (1.0)	N/A	37
Stanford University, USA	37	4 (1.6)	19 (0.54)	3 (2.4)	7 (0.89)	7 (0.73)	7 (0.78)	34
Yonsei University, South Korea	34	5 (1.5)	7 (0.86)	6 (2.0)	4 (1.2)	5 (0.87)	N/A	8.8
University of Miami, USA	31	6 (1.4)	1 (1.8)	14 (1.1)	3 (1.2)	3 (1.0)	7 (0.78)	51
Harvard University, USA	31	6 (1.4)	153 (0.11)	4 (2.3)	23 (0.4)	19 (0.41)	2 (1.6)	36
Nippon Medical School, Japan	28	8 (1.2)	6 (1.0)	8 (1.4)	5 (1.0)	5 (0.87)	2 (1.6)	22
Fourth Military Medical University, China	24	9 (1.1)	12 (0.76)	11 (1.3)	9 (0.62)	8 (0.64)	N/A	16
Chinese Academy of Medical Sciences, China	23	10 (1.0)	33 (0.32)	7 (1.5)	8 (0.76)	9 (0.59)	N/A	6.8
Sungkyunkwan University, South Korea	21	11 (0.93)	12 (0.76)	14 (1.1)	14 (0.49)	9 (0.59)	N/A	12
University of Southern California, USA	20	12 (0.89)	33 (0.32)	11 (1.3)	18 (0.45)	23 (0.36)	N/A	43
University of California Los Angeles, USA	19	13 (0.85)	153 (0.11)	9 (1.4)	30 (0.36)	30 (0.32)	7 (0.78)	36
Peking Union Medical College, China	18	14 (0.80)	N/A	9 (1.4)	N/A	36 (0.27)	N/A	7.4
Thomas Jefferson University, USA	17	15 (0.76)	7 (0.86)	35 (0.68)	11 (0.58)	23 (0.36)	N/A	55

TP, total number of highly cited articles; TPR (%), IPR (%), CPR (%), FPR (%), RPR (%), and SPR (%), the rank and percentage of total articles, single-institution articles, inter-institutionally collaborative articles, first-author articles, corresponding-author articles, single-author articles in their total articles; CPP<sub>2019</sub>, number of citations (TC<sub>2019</sub>) per publication (TP); N/A, not available.




Figure 3 Top 17 authors with Y-index ( $j \geq 10$ ).

**Publication performances: authors**

First and corresponding authors has the most contribution to the overall paper (41,42). The Y-index was proposed in recent years (21,22,26) to evaluate publication potential and characterize the scientific contributions by authors, institutions, and countries in relations to the numbers of first-author articles (FP) and corresponding-author articles (RP). The Y-index with two parameters ( $j, b$ ) can aid in visualization and comparison among different authors' contributions and is expressed as:

$$j = FP + RP \tag{1}$$

$$h = \tan^{-1}\left(\frac{RP}{FP}\right) \tag{2}$$


where  $j$  is the publication potential, a constant related to publication quantity; and  $b$  is publication characteristics which can describe the proportion of RP to FP. The greater the value of  $j$ , the greater the number of first- and corresponding-author articles by the author. Different values of  $b$  represent different proportions of corresponding-author articles from first-author articles.

- ❖  $b = \pi/2, j$  is the number of corresponding-author articles;

- ❖  $\pi/2 > b > 0.7854$  indicates more corresponding-author articles;
- ❖  $b = 0.7854$  indicates the same number of first- and corresponding-author articles;
- ❖  $0.7854 > b > 0$  indicates more first-author articles;
- ❖  $b = 0, j$  is the number of first-author articles.

In total of 2,163 keloid articles (96% of 2,253 articles) with both first- and corresponding-authors information in SCI-EXPANDED were extensively analysed by using the Y-index. Figure 3 demonstrated the distribution of the Y-index ( $j, b$ ) of the top 17 potential authors with  $j \geq 10$ . Each point has a coordinate ( $j, b$ ) that could symbolize a single author or multiple authors (22). A. Bayat (50, 1.435) published 52 articles including six first-author articles and 44 corresponding-author with  $j$  value of 50 which is far away from the original (not in Figure 3). Bayat had the highest publication potential in keloid research (43). Some authors are located on the same curve (for example,  $j = 11$ ), indicating that they had the same publication potential but different publication characteristics (35). It has been pointed out that with these data has a potential for bias in the analysis of authorship; it might attribute to different authors having the same name, or the same





**Figure 4** The citation histories of the 11 highly cited keloid articles with  $TC_{2019} \geq 216$ .

author using different names over time (28). Another potential confounder arises when an author moves from one affiliation to another (44).

#### **Citation histories of the 11 most frequently cited articles**


Information of total citations from Web of Science Core Collection was updated weekly. To improve bibliometric study, the total number of citations from Web of Science Core Collection since publication year to the end of 2019 ( $TC_{2019}$ ) was applied to improve the bias using data from Web of Science directly. The advantage of  $TC_{2019}$  is that they are invariable and ensure repeatability compared with the citation index from Web of Science Core Collection (17). The citation histories of the top 11 most frequently cited keloid articles ( $TC_{2019} \geq 216$ ) are shown in *Figure 4*. Eight of the top 11 articles published in the 1990s, which had a longer impact history, but less impact in the recent years.

“Green Giants” are articles have sharply increasing citations after publication for some years compared with others in the same field, and they quickly become high impact publications in recent year with a high  $C_{year}$ , which is more than double compare with others (45).

Article entitled “Hypertrophic scarring and keloids: Pathomechanisms and current and emerging treatment strategies” (46) was the only “Green Giant” in keloid studies, which ranked top in  $C_{2019}$  of 98 which is much higher than any others. *Figure 4* shows the “Green Giant” in keloid studies had sharply increasing citation rates for eight full years after its publication. “Green Giants” can be also found in other research fields, for example category of environmental sciences (47), pain research (8), and publications in Taiwan (45).

#### **Citation histories of the ten most impact articles in 2019**

The amount of citation received in the year 2019 ( $C_{2019}$ ) and in the publication year ( $C_0$ ) might provide additional information for readers to understand the impact of a highly cited article today versus its immediate effect post publication (21). The 2,253 keloid related articles ranked differently if sorted by  $TC_{2019}$ ,  $C_{2019}$  than sorted by  $C_0$ . A total of 836 articles (37% of 2,253 articles) had no citation in the most recent year of 2019 ( $C_{2019} = 0$ ) and 1,793 (80%) articles had no citation in the initial publication year ( $C_0 = 0$ ). Furthermore, among the top 100  $C_0$  articles, only 9.4% and 23% of the articles were



**Figure 5** The citation histories of the 10 most impact keloid articles in 2019 with  $C_{2019} \geq 22$ .

among the top 100  $TC_{2019}$  and  $C_{2019}$  articles, respectively. The percentages were found different from research topics: occupational therapy (12% and 13%) (48) and metal-organic frameworks (22% and 24%) (19). Top ten most impact keloid articles in 2019 with  $C_{2019} \geq 22$  and their citation history trends are outlined in *Figure 5*. Articles by Gauglitz *et al.* (46), Wolfram *et al.* (49), and Bock *et al.* (50) ranked not only the top ten in  $C_{2019}$  but also the top ten in  $TC_{2019}$ . Since the recent articles had a shorter article life and a shorter time to accumulate citations, the high impact articles in 2019 published by Cheng *et al.* (51), Miao *et al.* (52), and Berman *et al.* (53) had a  $TC_{2019}$  of 22 (rank 639<sup>th</sup>), 40 (rank 341<sup>th</sup>), and 79 (rank 118<sup>th</sup>), respectively. Article entitled “Regeneration of fat cells from myofibroblasts during wound healing” published in *Science* had a sharply increasing trend after publication in 2017 (54).

#### **Research focuses and development trends from 1990 to 2019**

Ho’s group proposed the distribution of words in article titles, abstracts, author keywords, and *KeyWords Plus* in different periods as information to evaluate the main research focuses and determine their development trends in research topics (27,55). In this study, the distribution of

single words in article titles was analyzed because the title of an article always best summarizes the main idea of the study. After eliminating empty words such as “a” and “the”, the 20 most frequently used single words in titles are presented in *Table 5*. Their occurrence times were calculated in the 30-year period and three decades (1990–1999, 2000–2009, and 2010–2019).

Further, word cluster analysis with noun phrases in title, abstract, author keywords, and *KeyWords Plus* compensates for the weakness of single word distribution analysis (27). Noun phrases were generated from the sources above using the LLR algorithm and imported to MapEquation to generate an alluvial diagram that vividly demonstrated how research hotspots changed over time (56). The alluvial diagram shows the progression of research hotspots in keloids (*Figure 6*). Each column represents a cluster. The height of the column indicates the sum of the page rank value of all noun phrases in a cluster. The page rank values of each noun phrase indicates its importance.

First, there is an emerging trend in the underlying pathogenetic mechanisms of keloids. Supporting words in title included “fibroblasts”, “expression”, “collagen”, “formation”, and “proliferation”. Particularly, the frequency of “fibroblasts” and “expression” increased from rank 6 and 7 in the first decade to 4 and 6 in the last decade,

**Table 5** Top 20 most frequently used substantives in title during the past three decades


Words in title	TP	Rank (%)			
		1990–2019	1990–1999	2000–2009	2010–2019
Keloid	768	1 (34)	1 (27)	1 (30)	1 (38)
Keloids	465	2 (21)	2 (22)	2 (22)	2 (19)
Treatment	347	3 (15)	4 (16)	4 (13)	3 (16)
Fibroblasts	307	4 (14)	6 (13)	3 (14)	4 (14)
Scars	300	5 (13)	3 (16)	5 (12)	5 (13)
Hypertrophic	228	6 (10)	4 (16)	6 (12)	9 (7.7)
Expression	200	7 (8.9)	11 (6.4)	7 (10)	6 (9.2)
Skin	181	8 (8)	8 (7.6)	8 (7.8)	8 (8.3)
Scar	163	9 (7.2)	9 (7)	12 (4.9)	7 (8.5)
Collagen	148	10 (6.6)	7 (8.2)	10 (6.6)	11 (6.1)
Human	138	11 (6.1)	10 (6.7)	11 (5.4)	10 (6.4)
Growth	122	12 (5.4)	17 (3.2)	9 (6.9)	12 (5.2)
Therapy	97	13 (4.3)	28 (2.3)	21 (3.6)	12 (5.2)
Laser	88	14 (3.9)	33 (2)	21 (3.6)	14 (4.6)
Intralesional	87	15 (3.9)	20 (2.9)	16 (3.7)	15 (4.2)
Surgical	86	16 (3.8)	33 (2)	13 (4.2)	16 (4.1)
Clinical	83	17 (3.7)	33 (2)	15 (4)	18 (3.9)
Patients	83	17 (3.7)	14 (3.5)	16 (3.7)	22 (3.7)
Formation	81	19 (3.6)	20 (2.9)	16 (3.7)	22 (3.7)
Proliferation	78	20 (3.5)	45 (1.7)	24 (3.1)	16 (4.1)

TP, occurrence times of substantives in title during 1990 to 2019.

respectively. The word “proliferation” soared from rank 45 to 16 (*Table 5*). Thus, researchers tend to focus more on the pathogenetic mechanism in the molecular level, which can be cross-verified by cluster “keloid fibroblasts”, “molecular mechanism”, “protein expression”, and “connective tissue growth factor” in *Figure 6*. It has been demonstrated that keloid fibroblasts and myofibroblasts originate from activated resident skin fibroblasts that interact with other cells including multiple types of immune cells and keratinocytes via signaling molecules such as matrix metalloproteinases, interleukins and cytokines (57). One important difference between keloid and normal fibroblasts is the overexpression of the TGF- $\beta$  signaling cascade (46). The TGF- $\beta$  signaling pathway not only plays a critical role in the fibroproliferative process leading to extracellular matrix accumulation (58) but also has a crosstalk with the

mitogen-activated protein kinase (MAPK) pathway (59). The crosstalk is assumed to be associated with the invasiveness of keloid fibroblasts and, the MAPK pathway might be a specific target of keloid therapy (60). Other signaling pathways, including insulin-like growth factor-I and integrin pathways, have also been found to be activated in keloid fibroblasts (61,62).

Second, increasingly more research attention has focused on keloid treatment. For example, the word “treatment” ranks top 4 in every decade, and the word “therapy” shifted dramatically from rank 28 in the first decade to 12 in the third decade. Other words related to treatment modalities including “laser”, “intralesional” and “surgical” all soared during the 30-year period (*Table 5*). Keloid treatment has always been the focus of dermatologists and plastic surgeons. As single treatment also fails to result in a consistently low



**Figure 6** The progression of research hotspots in the recent three decades.

recurrence rate, increasingly more studies have focused on combination therapy, and it has been reported as the most efficacious and safe management for keloids (63). One common combination is laser therapy and triamcinolone acetonide (TAC) administration. Carbon dioxide laser (64,65), pulsed dye laser with 5-fluorouracil (66), and neodymium-doped yttrium garnet laser (67) yielded good results in combination with TAC. In addition to laser therapy with TAC administration, a low recurrence rate is achieved when surgical excision is combined with other treatment modalities, such as TAC, 5-Fluorouracil, pressure therapy, silicone gel, electron beam, and radiation therapy (68). Although there are already many treatment modalities, no gold standard for keloid treatment has been established, and many efforts are being made in the exploration of new therapies (69).

Lastly, researchers tend to be more comprehensive and objective in the evaluation of therapeutic effects. Supporting evidence includes cluster “recurrence rate”, “Vancouver Scar Scale”, “patient satisfaction”, and “Observer Scar

Assessment Scale” (Figure 6). The Vancouver Scar Scale and Patient and Observer Scar Assessment Scale are two commonly used scales in patients with keloids (70,71). These scales help clinicians and patients gain a comprehensive understanding of treatment responses and guide future treatment. Standardized scales also make the evaluation more comparable, allowing analysis across different studies. However, it should be acknowledged that scales provide quantitative but subjective assessments of therapeutic effects. The scale scores are subjected to a number of human factors, and it is also unclear whether the scale scores correlate with the underlying molecular profile of scars. Thus, objective scar assessment tools should be advocated in practicing evidence-based medicine. Verhaegen *et al.* reviewed noninvasive objective measurement tools to assess different scar features, including color, thickness, relief, pliability, and surface area (72). It is foreseeable that objective measurement tools will have a higher status in the assessment of keloids and scars.

### Limitations

Several limitations of this study should be addressed. First, the main body of this study only includes peer-reviewed original articles, which are the only document type that provides new findings and innovative insights supported by scientific proofs. The absence of other document types might result in the omission of some information. Second, it should be mentioned that citation numbers do not necessarily link with research quality. The citation number is determined by many factors, including research quality, self-citation, and citations from one's coworkers or the same institution. Finally, publication is a complex and time-consuming process. Bibliometric analysis based on publications is more like a summary of the past and is unable to identify the most up-to-date findings and treatment modalities.

### Conclusions

This bibliometric analysis quantitatively studies the scientific landscape and research trends in keloid publications. We presented bibliometric data on document type, annual publication number, most productive journals, publication geography, top potential authors, and highly cited articles. Additionally, research trends were analyzed through single word analysis in title and noun-phrase co-occurrence analysis. Keloids have remained a research focus. Recent research hotspots include underlying pathogenetic mechanisms, keloid treatment, and therapeutic effect evaluation.

### Acknowledgments

*Funding:* None.

### Footnote

*Reporting Checklist:* The authors have completed the PRISMA reporting checklist. Available at <http://dx.doi.org/10.21037/atm-21-508>

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/atm-21-508>). The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are

appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

### References

1. Ehrlich HP, Desmoulière A, Diegelmann RF, et al. Morphological and immunochemical differences between keloid and hypertrophic scar. *Am J Pathol* 1994;145:105-13.
2. Blackburn WR, Cosman B. Histologic basis of keloid and hypertrophic scar differentiation: Clinicopathologic correlation. *Arch Pathol* 1966;82:65-71.
3. Nakaoka H, Miyauchi S, Miki Y. Proliferating activity of dermal fibroblasts in keloids and hypertrophic scars. *Acta Derm Venereol* 1995;75:102-4.
4. Ueda K, Furuya E, Yasuda Y, et al. Keloids have continuous high metabolic activity. *Plast Reconstr Surg* 1999;104:694-8.
5. Younai S, Venters G, Vu S, et al. Role of Growth Factors in Scar Contraction. *Ann Plast Surg* 1996;36:495-501.
6. Murray JC. Keloids and hypertrophic scars. *Clin Dermatol* 1994;12:27-37.
7. Durani P, Bayat A. Levels of evidence for the treatment of keloid disease. *J Plast Reconstr Aesthet Surg* 2008;61:4-17.
8. Chuang KY, Ho YS. A bibliometric analysis on top-cited articles in pain research. *Pain Med* 2014;15:732-44.
9. Pouris A, Ho YS. A bibliometric analysis of research on Ebola in Science Citation Index Expanded. *S Afr J Sci* 2016;112:83-8.
10. Ho YS, Siu E, Chuang KY. A bibliometric analysis of dengue-related publications in the Science Citation Index Expanded. *Future Virol* 2016;11:631-48.
11. Rosas SR, Kagan JM, Schouten JT, et al. Evaluating research and impact: a bibliometric analysis of research by the NIH/NIAID HIV/AIDS clinical trials networks. *PLoS One* 2011;6:e17428.
12. Head MG, Fitchett JR, Newell ML, et al. Mapping pneumonia research: A systematic analysis of UK investments and published outputs 1997-2013. *EBioMedicine* 2015;2:1193-9.
13. Ho YS. Rebuttal to: Su et al. "The neurotoxicity of nanoparticles: A bibliometric analysis". *Toxicol Ind Health*

- 2019;35:399-402.
14. Ho YS. Critical comment on: Zhu, Jin, and He 'On evolutionary economic geography: a literature review using bibliometric analysis', *European Planning Studies* vol. 27, pp 639-660. *European Planning Studies* 2019;27:1235-7.
  15. Garfield E. KeyWords Plus: ISI's breakthrough retrieval method. Part 1. Expanding your searching power on Current Contents on Diskette. *Curr Contents* 1990;32:5-9.
  16. Fu HZ, Ho YS. Top cited articles in thermodynamic research. *Journal of Engineering Thermophysics* 2015;24:68-85.
  17. Fu HZ, Wang MH, Ho YS. The most frequently cited adsorption research articles in the Science Citation Index (Expanded). *J Colloid Interface Sci* 2012;379:148-56.
  18. Fu HZ, Ho YS. Top cited articles in adsorption research using Y-index. *Res Evaluat* 2014;23:12-20.
  19. Ho YS, Fu HZ. Mapping of metal-organic frameworks publications: A bibliometric analysis. *Inorg Chem Commun* 2016;73:174-82.
  20. Li Z, Ho YS. Use of citation per publication as an indicator to evaluate contingent valuation research. *Scientometrics* 2008;75:97-110.
  21. Ho YS. Top-cited Articles in Chemical Engineering in Science Citation Index Expanded: A Bibliometric Analysis. *Chinese J Chem Eng* 2012;20:478-88.
  22. Ho YS. Classic articles on social work field in Social Science Citation Index: A bibliometric analysis. *Scientometrics* 2014;98:137-55.
  23. Ho YS. Bibliometric Analysis of the Journal of Orthopaedic Research from 1991 To 2018. *Ortho Res Online J* 2019;6. doi: 10.31031/OPROJ.2019.06.000632.
  24. Chiu WT, Ho YS. Bibliometric analysis of homeopathy research during the period of 1991 to 2003. *Scientometrics* 2005;63:3-23.
  25. Fu HZ, Ho YS. Comparison of independent research of China's top universities using bibliometric indicators. *Scientometrics* 2013;96:259-76.
  26. Ho YS. A bibliometric analysis of highly cited articles in materials science. *Curr Sci* 2014;107:1565-72.
  27. Wang CC, Ho YS. Research trend of metal-organic frameworks: a bibliometric analysis. *Scientometrics* 2016;109:481-513.
  28. Ho YS, Hartley J. Classic articles in Psychology in the Science Citation Index Expanded: A bibliometric analysis. *Br J Psychol* 2016;107:768-80.
  29. Chen SR, Chiu WT, Ho YS. Asthma in children: mapping the literature by bibliometric analysis. *Rev Fr Allergol Immunol Clin* 2005;45:442-6.
  30. Lin HW, Yu TC, Ho YS. A systemic review of human papillomavirus studies: global publication comparison and research trend analyses from 1993 to 2008. *Eur J Gynaecol Oncol* 2011;32:133-40.
  31. Ho YS. The top-cited research works in the Science Citation Index Expanded. *Scientometrics* 2013;94:1297-312.
  32. Han JS, Ho YS. Global trends and performances of acupuncture research. *Neurosci Biobehav Rev* 2011;35:680-7.
  33. Halper J, Kjaer M. Basic components of connective tissues and extracellular matrix: Elastin, fibrillin, fibulins, fibrinogen, fibronectin, laminin, tenascins and thrombospondins. *Adv Exp Med Biol* 2014;802:31-47.
  34. Ho YS, Satoh H, Lin SY. Japanese lung cancer research trends and performance in Science Citation Index. *Intern Med* 2010;49:2219-28.
  35. Ho YS, Hartley J. Classic articles published by American scientists (1900-2014): a bibliometric analysis. *Curr Sci* 2016;111:1156-65.
  36. Ho YS, Kahn M. A bibliometric study of highly cited reviews in the Science Citation Index expanded. *J Assoc Inf Sci Tech* 2014;65.
  37. Riesenber D, Lundberg GD. The Order of Authorship: Who's on First? *JAMA* 1990;264:1857.
  38. Li LL, Ding GH, Feng N, et al. Global stem cell research trend: Bibliometric analysis as a tool for mapping of trends from 1991 to 2006. *Scientometrics* 2009.
  39. Brandt JS, Downing AC, Howard DL, et al. Citation classics in obstetrics and gynecology: the 100 most frequently cited journal articles in the last 50 years. *Am J Obstet Gynecol* 2010;203:355.e1-7.
  40. Kelly JC, Glynn RW, O'Briain DE, et al. The 100 classic papers of orthopaedic surgery: a bibliometric analysis. *J Bone Joint Surg Br* 2010;92:1338-43.
  41. Gaeta TJ. Authorship: "Law" and order. *Acad Emerg Med* 1999;6:297-301.
  42. Mattsson P, Sundberg CJ, Laget P. Is correspondence reflected in the author position? A bibliometric study of the relation between corresponding author and byline position. *Scientometrics* 2011;87:99-105.
  43. Burman KD. "Hanging from the Masthead": Reflections on Authorship. *Ann Intern Med* 1982;97:602-5.
  44. MacRoberts MH, MacRoberts BR. Problems of citation analysis: A critical review. *J Am Soc Inf Sci Tec* 1989;40:342-9.
  45. Ho YS. Classic papers published by Taiwanese scientists in the science citation index expanded: A bibliometric study. *COLLNET Journal of Scientometrics and Information Management* 2018;12:83-95.
  46. Gauglitz GG, Korting HC, Pavicic T, et al. Hypertrophic scarring and keloids: pathomechanisms and current and

- emerging treatment strategies. *Mol Med* 2011;17:113-25.
47. Khan MA, Ho YS. Top-cited articles in environmental sciences: Merits and demerits of citation analysis. *Sci Total Environ* 2012;431:122-7.
  48. Brown T, Gutman SA, Ho YS, et al. A bibliometric analysis of occupational therapy publications. *Scand J Occup Ther* 2018;25:1-14.
  49. Wolfram D, Tzankov A, Pulz P, et al. Hypertrophic scars and keloids--a review of their pathophysiology, risk factors, and therapeutic management. *Dermatol Surg* 2009;35:171-81.
  50. Bock O, Schmid-Ott G, Malewski P, et al. Quality of life of patients with keloid and hypertrophic scarring. *Arch Dermatol Res* 2006;297:433-8.
  51. Cheng P, Zhang J, Huang J, et al. Near-infrared fluorescence probes to detect reactive oxygen species for keloid diagnosis. *Chem Sci* 2018;9:6340-7.
  52. Miao Q, Yeo DC, Wiraja C, et al. Near-Infrared Fluorescent Molecular Probe for Sensitive Imaging of Keloid. *Angew Chem Int Ed Engl* 2018;57:1256-60.
  53. Berman B, Maderal A, Raphael B. Keloids and Hypertrophic Scars: Pathophysiology, Classification, and Treatment. *Dermatol Surg* 2017;43 Suppl 1:S3-18.
  54. Plikus MV, Guerrero-Juarez CF, Ito M, et al. Regeneration of fat cells from myofibroblasts during wound healing. *Science* 2017;355:748-52.
  55. Mao N, Wang MH, Ho YS. A bibliometric study of the trend in articles related to risk assessment published in science citation index. *Hum Ecol Risk Assess* 2010;16:801-24.
  56. Rosvall M, Bergstrom CT. Mapping Change in Large Networks. *PLoS One* 2010;5:e8694.
  57. Macarak EJ, Wermuth PJ, Rosenbloom J, et al. Keloid disorder: Fibroblast differentiation and gene expression profile in fibrotic skin diseases. *Exp Dermatol* 2021;30:132-45.
  58. Jagadeesan J, Bayat A. Transforming growth factor beta (TGFbeta) and keloid disease. *Int J Surg* 2007;5:278-85.
  59. Unahabhokha T, Sucontphunt A, Nimmannit U, et al. Molecular signalings in keloid disease and current therapeutic approaches from natural based compounds. *Pharm Biol* 2015;53:457-63.
  60. He S, Liu X, Yang Y, et al. Mechanisms of transforming growth factor beta(1)/Smad signalling mediated by mitogen-activated protein kinase pathways in keloid fibroblasts. *Br J Dermatol* 2010;162:538-46.
  61. Daian T, Ohtsuru A, Rogounovitch T, et al. Insulin-like growth factor-I enhances transforming growth factor-beta-induced extracellular matrix protein production through the P38/activating transcription factor-2 signaling pathway in keloid fibroblasts. *J Invest Dermatol* 2003;120:956-62.
  62. Liu S, Xu SW, Blumbach K, et al. Expression of integrin beta1 by fibroblasts is required for tissue repair in vivo. *J Cell Sci* 2010;123:3674-82.
  63. Al-Attar A, Mess S, Thomassen JM, et al. Keloid Pathogenesis and Treatment. *Plast Reconstr Surg* 2006;117:286-300.
  64. Behera B, Kumari R, Thappa DM, et al. Therapeutic efficacy of intralesional steroid with carbon dioxide laser versus with cryotherapy in treatment of keloids: a randomized controlled trial. *Dermatol Surg* 2016;42:1188-98.
  65. Wang J, Wu J, Xu M, et al. Combination therapy of refractory keloid with ultrapulse fractional carbon dioxide (CO<sub>2</sub>) laser and topical triamcinolone in Asians-long-term prevention of keloid recurrence. *Dermatol Ther* 2020;33:e14359.
  66. Asilian A, Darougheh A, Shariati F. New combination of triamcinolone, 5-Fluorouracil, and pulsed-dye laser for treatment of keloid and hypertrophic scars. *Dermatol Surg* 2006;32:907-15.
  67. Rossi A, Lu R, Frey MK, et al. The use of the 300 microsecond 1064 nm Nd:YAG laser in the treatment of keloids. *J Drugs Dermatol* 2013;12:1256-62.
  68. Limmer EE, Glass DA, 2nd. A Review of Current Keloid Management: Mainstay Monotherapies and Emerging Approaches. *Dermatol Ther (Heidelb)* 2020;10:931-48.
  69. Ekstein SF, Wyles SP, Moran SL, et al. Keloids: a review of therapeutic management. *Int J Dermatol* 2021;60:661-71.
  70. Sullivan T, Smith J, Kermode J, et al. Rating the burn scar. *J Burn Care Rehabil* 1990;11:256-60.
  71. Draaijers LJ, Tempelman FRH, Botman YAM, et al. The Patient and Observer Scar Assessment Scale: A Reliable and Feasible Tool for Scar Evaluation. *Plast Reconstr Surg* 2004;113:1960-5.
  72. Verhaegen PD, van der Wal MBA, Middelkoop E, et al. Objective scar assessment tools: a clinimetric appraisal. *Plast Reconstr Surg* 2011;127:1561-70.

**Cite this article as:** Chong Y, Long X, Ho YS. Scientific landscape and trend analysis of keloid research: a 30-year bibliometric review. *Ann Transl Med* 2021;9(11):945. doi: 10.21037/atm-21-508