



Diagnosis, workup, and risk stratification of upper tract urothelial carcinoma

Andrew J. Zganjar^{1^}, David D. Thiel¹, Timothy D. Lyon^{1,2}

¹Department of Urology, Mayo Clinic, Jacksonville, FL, USA; ²Robert D. and Patricia E. Kern Center for the Science of Health Care Delivery, Mayo Clinic, Rochester, MN, USA

Contributions: (I) Conception and design: All authors; (II) Administrative support: AJ Zganjar; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: AJ Zganjar, TD Lyon; (V) Data analysis and interpretation: None; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Andrew J. Zganjar, MD. Department of Urology, Mayo Clinic, 4500 San Pablo Road, Jacksonville, FL 32224, USA.

Email: zganjar.andrew@mayo.edu.

Abstract: Upper tract urothelial carcinoma (UTUC) is a relatively rare disease that presents unique challenges to urologists from both a diagnostic and management standpoint. UTUC is a clinically heterogeneous disease with a varied natural history, and given its location in the upper urinary tract, treatment has the potential to cause or worsen chronic kidney disease. Therefore, physicians caring for patients with UTUC must be facile with multiple diagnostic and therapeutic strategies in order to provide optimal patient care. We present an overview of the epidemiology, histology, risk factors, and contemporary approach to the diagnosis, laboratory evaluation, imaging, and risk stratification for patients with UTUC. Computerized tomographic urography, thorough endoscopic evaluation, and biopsy (endoscopically or percutaneously) remain the standard of care for the diagnosis and staging of patients with suspected UTUC. Preoperative nomograms are vital to select patients more optimally for preoperative systemic chemotherapy and facilitate clinical trial enrollment. A thorough understanding of the various diagnostic challenges, imaging/pathologic limitations, biomarkers, and risk stratification tools will allow us as a field to develop new modalities to improve our diagnostic capabilities and reduce the risk of under diagnosis and over treatment for our patients.

Keywords: Upper tract urothelial carcinoma (UTUC); ureteroscopy; biopsy; risk stratification

Submitted Jan 18, 2023. Accepted for publication Jul 26, 2023. Published online Aug 04, 2023.

doi: 10.21037/tau-23-45

View this article at: <https://dx.doi.org/10.21037/tau-23-45>

Introduction

Urothelial carcinoma of the ureter, renal pelvis, and intrarenal collecting system comprises about 5–10% of all urothelial carcinomas and is collectively termed upper tract urothelial carcinoma (UTUC) (1-3). UTUC is a clinically heterogeneous disease with a varied natural history, and given its location in the upper urinary tract, treatment has the potential to cause or worsen chronic kidney disease. Accordingly, physicians caring for UTUC patients must be

facile with multiple diagnostic and therapeutic strategies in order to provide optimal patient care. The objective of this review is to outline contemporary approaches to diagnosis, workup, and risk stratification for patients with UTUC.

Epidemiology and risk factors

Epidemiology

The incidence of UTUC has been estimated at 2 cases per

[^] ORCID: 0000-0002-7150-9125.

100,000 persons in the Western Hemisphere (2,4,5). The incidence is slowly rising over the past 3 decades, and there is also recent evidence of a migration towards higher stage disease at diagnosis, with the largest increase observed for *in situ* disease (2,6,7). Tumors of the renal pelvis are twice as common as ureteral tumors, and although the majority of tumors occur in a single renal unit, about 5% of UTUCs are bilateral at diagnosis (8,9). Concurrent bladder tumors are common and are found in 17% of cases (10).

The median age at diagnosis of UTUC is 73 years (4). There is a well-described sex disparity in the occurrence of UTUC, with a 2:1 male to female incidence ratio (4,11). While the majority of patients diagnosed with UTUC in the United States are Caucasian, incidence is increasing among racial minorities. In a study utilizing the Surveillance, Epidemiology, and End Results database examining trends in UTUC from 1973 to 2005, incidence was noted to increase among Black patients (3.4% to 4.3%) as well as those of 'Other' race (4% to 7.5%), including those of Asian, American Indian/Alaskan Native, or Hawaiian/Pacific Islander descent (4). Following treatment, the most frequent site of UTUC recurrence is in the bladder, which occurs in about 1/3 of patients (12,13). In those patients with low-risk disease managed endoscopically, ipsilateral recurrence rate approaches 35% (14). Recurrence in the contralateral upper tract is much less common, occurring in 5% of cases (15).

Risk factors

Tobacco exposure is the primary risk factor for developing UTUC, with an estimated 25–60% of cases attributable to smoking (16). There is also a dose-response relationship between tobacco exposure and UTUC. The relative risk for developing UTUC among smokers in the United States is around 3, but this increases to a relative risk of 7.2 among those with greater than 45 pack-years of tobacco exposure (16,17). Importantly, this risk is also modifiable, as smoking cessation is associated with a 60% to 70% reduction in risk 10 or more years after quitting (16). Urothelial carcinogenesis is thought to be potentiated by N-hydroxylamine, which is a metabolite of several aromatic amines contained in tobacco smoke. Susceptibility to the carcinogenic effects of smoking is likely influenced by genetic polymorphisms in enzymes that degrade N-hydroxylamine, which may also lead to increased susceptibility within families (18).

Several additional environmental exposures are also

implicated in risk of UTUC. Occupational exposure to aromatic amines in petroleum, industrial dyes, textiles, plastic manufacturing and coal are linked to developing UTUC (19). Exposure to aristocholic acid is similarly associated with an increasing risk of UTUC. This is thought to underlie the dramatically increased incidence of UTUC that occurred in rural areas of the Balkans (Serbia, Bulgaria, Romania, Croatia and Bosnia and Herzegovina) in the mid-20th century. Termed Balkan Endemic Nephropathy, residents here had a risk 60–100 times higher than those living in other parts of the world (19). Flour made from wheat contaminated with aristocholic acid from *Aristocholia clematitis*, a native Balkan plant, likely played a role (20). However, cases in the Balkans are decreasing over more recent years (21). Aristocholic acid is also found in some Chinese herbal products and teas, and indeed a higher incidence of UTUC is observed in Chinese mainland and Taiwan (19). Chronic exposure to arsenic in drinking water, usually due to contaminated wells, is also associated with developing UTUC (22). Use of the analgesic phenacetin has been associated with nephrotoxicity and risk of UTUC (23). Although routine use of phenacetin has been abandoned, analgesic abuse more broadly, including codeine, acetaminophen, and salicylates, has retained an association with increased risk of upper tract tumors (24). Chronic urinary tract inflammation and history of exposure to alkylating chemotherapies, such as cyclophosphamide, are further linked to risk of UTUC (19,25).

A history of urothelial carcinoma of the bladder is another well-known risk factor for UTUC, with rates of upper tract recurrence ranging from 1–6% following radical cystectomy in modern series (26,27). Although there is a theoretical risk of seeding the upper urinary tract if a ureteral stent is placed at the time of endoscopic resection of a bladder tumor, the available data are mixed regarding whether this is a clinically meaningful phenomenon. Miest and colleagues undertook a multi-institutional review of 1,049 patients who underwent radical cystectomy at one of two centers and observed no difference in the rate of subsequent upper tract recurrence between patients managed with a ureteral stent or nephrostomy tube preoperatively (28). However, Sountoulides *et al.* undertook a systematic review of the risk of metachronous UTUC recurrence after a bladder tumor, and found that among patients without preoperative hydronephrosis, subsequent UTUC was more common in those who had a ureteral stent placed after resection versus those who did not [odds ratio (OR) 3.37, 95% confidence interval (CI): 1.49–7.63] (29).

Notably, this was not found among those with preoperative hydronephrosis, as no significant difference in upper tract recurrence was seen between those managed with a ureteral stent or a nephrostomy tube in this population. Authors concluded that prophylactic ureteral stenting after ureteral orifice resection should be avoided, when possible, in patients without preoperative hydronephrosis due to the potential for upper tract seeding (29).

Lynch syndrome related cancers

A hereditary predisposition to UTUC is known to occur in the context of Lynch syndrome, also termed hereditary nonpolyposis colorectal carcinoma (HNPCC) (30). UTUC is the 3rd most common tumor in HNPCC, occurring in about 5% of patients; colon (63%) and endometrial (9%) cancers also frequently occur (30). Lynch syndrome has an autosomal dominant mode of transmission and is caused by a mutation in one of several DNA mismatch repair genes, including *MLH1*, *MSH2*, *MSH6* and *PMS2* (30). The potential for a hereditary UTUC should be suspected in patients who present at a young age (<60 years), who have a personal history of another cancer in the HNPCC spectrum (including colon, endometrial, pancreatic, ovarian and gastric cancers), or among those with two or more first-degree relatives with a Lynch spectrum cancer (30,31). Tissue-based testing for HNPCC includes polymerase chain reaction testing for high microsatellite instability within the tumor, a hallmark of Lynch syndrome, as well as with immunohistochemistry evaluating for loss of *MLH1*, *MSH2*, *MSH6* and *PMS2* (31). Suspected or identified individuals should be referred for genetic counseling. For individuals known to have Lynch syndrome, screening for UTUC with an annual urinalysis is recommended, along with consideration for periodic upper tract imaging (31).

Histology

Urothelial carcinomas are the most common tumors of the upper urinary tract, representing over 90% of cases (32). Approximately 24% of cases include histologic variants, including squamous, glandular, sarcomatoid, micropapillary or neuroendocrine differentiation (33). Primary squamous cell carcinomas or adenocarcinomas can occur in the upper urinary tract but are very rare, representing <1% of upper tract tumors (23,34).

Benign lesions of the upper tract are also known to occur, including papillomas, inverted papillomas, and von Brunn

nests. Concurrent or subsequent malignancy has been observed in 15–20% of patients with inverted papillomas of the upper tract, and therefore upper tract surveillance is recommended for these patients for at least 2 years (35–37). Other benign lesions that can occur in the upper tract include fibroepithelial polyps and neurofibromas, which are best treated with local excision (38,39).

Diagnosis and workup

Presenting symptoms

The most common symptom of UTUC is gross or microscopic hematuria, which is present in 75–80% of cases (32). Gross hematuria is predictive of higher pathologic stage at initial presentation compared to microscopic hematuria (40). Flank pain due to renal obstruction occurs in 20% of patients (32). Occasionally, UTUC is diagnosed incidentally in otherwise asymptomatic patients with evidence of a collecting system mass on cross-sectional imaging. Hydronephrosis is often present on imaging (up to 50% of cases), and multiple studies have shown that hydronephrosis is associated with higher stage at diagnosis (41,42). Constitutional symptoms, including weight loss, fever, night sweats, anorexia, or hemoptysis are associated with a worse prognosis and should prompt a complete evaluation for the presence metastatic disease (32).

Laboratory evaluation

Basic laboratory workup should include microscopic urinalysis to detect microscopic hematuria and rule out urinary tract infection, along with hemoglobin level and renal function panel. Cytology remains a reliable diagnostic method for UTUC, however voided urinary cytology has high false negative rates approaching 50–90% (43). Selective cytology of the affected upper tract has shown improved diagnostic accuracy compared to voided cytology, with meta-analysis demonstrating a sensitivity of 53.1% and specificity of 90% (43). Newer classification methods in cytologic grading (Paris System) were introduced in 2016 to improve diagnostic accuracy and focus more on high-grade diagnosis (44). Subsequent studies have demonstrated a significantly higher rate of surgical pathologic concordance when compared with traditional cytology in UTUC (45). Additionally, high rates of sensitivity (71–78.6%) and specificity (86–100%) have been found (46,47); while others report a decrease in sensitivity when compared to traditional

cytology (48). Despite these diagnostic challenges, both the European Association of Urology (EAU) and National Comprehensive Cancer Network (NCCN) recommend the addition of cytology to the initial evaluation of a patient with suspected UTUC (49,50).

Given the current limitations with non-invasive means for diagnosis of UTUC, the role of alternative urinary biomarkers has been an active area of investigation. Fluorescence in situ hybridization (FISH) testing involves the use of fluorescent probes to identify abnormalities in specific chromosomes (3, 7, 17, and 9p21) (51). While FISH is commonly used adjunctly to aid in the diagnosis of bladder cancer, efforts have been made to apply it to the diagnosis of UTUC with mixed results. When used to analyze voided urine without a concurrent bladder tumor, sensitivity and specificity have varied widely, with reported rates of 54–76.7% and 78–94.7%, respectively (52,53). The detection of DNA mutations and methylation markers in urine has gained some attention as another potential biomarker for UTUC, with preliminary studies showing promise. Monteiro-Reis *et al.* studied 22 patients with confirmed UTUC and found that by analyzing voided samples for methylation of the *GDF15*, *TMEFF2*, and *VIM* promoters they were able to detect UTUC with 91% sensitivity and 100% specificity (54). Similarly, Guo *et al.* examined the same panel with the addition of the *CDH1*, *HSPA2*, and *RASSF1A* genes, and found a lower sensitivity of 82% and specificity of 68% (55). Neither study evaluated whether methylation could distinguish between upper and lower tract lesions. Several gene mutations have been linked to urothelial carcinoma, and mutations in hotspots of the *TERT* and *FGFR3* genes have been studied as potential urinary biomarkers for UTUC. One study found that when examining *TERT* and *FGFR3* mutations in combination with cytology, a sensitivity of 78.6% and specificity of 96% was demonstrated (56). Despite sometimes promising results, urinary biomarkers for UTUC have failed to achieve mainstream support or guideline endorsement.

Imaging

Computerized tomographic urography (CTU) is the dominant imaging modality for evaluating the upper urinary tract. It has been shown to be highly accurate for diagnosing UTUC, with sensitivity of 92% and specificity of 95% according to a recent meta-analysis of 13 studies involving 1,233 patients (57). Additionally, it is widely available, able to detect lesions as small as 5 mm (58),

and can offer additional staging information—with the presence of enlarged lymph nodes being highly predictive for metastasis in UTUC (59). Current radiological methods cannot diagnose invasiveness unless the tumor is advanced. Magnetic resonance urography (MRU) may be used in patients who cannot undergo CTU, typically when iodinated contrast media is contraindicated because of severe allergy or poor renal function. However, MRU has a lower sensitivity (69–75%) for the diagnosis of UTUC compared to CTU, and its utility can be limited by motion/peristalsis artifact, increased study time, and cost (60–62). Retrograde pyelogram at the time of cystourethroscopy is also an option for patients with contraindications to CTU/MRU, however the sensitivity for UTUC is lower (25–100%) and provides no staging information (63). CT remains the preferred diagnostic modality for complete staging and detecting metastasis, however the use of ¹⁸F-fluorodeoxyglucose positron emission tomography/computed tomography (FDG-PET/CT) has shown to be effective in detecting nodal metastasis with a sensitivity and specificity of 82% and 84%, respectively (64). Currently, only the EAU have specific imaging guidelines for the diagnosis of UTUC—with CTU being the preferred modality for diagnosis and staging (49).

Endoscopic evaluation

Cystourethroscopy remains an integral initial part of UTUC evaluation to rule out concomitant bladder cancer. A retrospective review of 673 patients diagnosed and treated for UTUC found that 17% of patients presented with both UTUC and bladder cancer—with the location of the UTUC (renal pelvis *vs.* upper ureter *vs.* lower ureter) being the only predictive factor for concomitant bladder tumor on multivariate analysis (OR: 1.7; 95% CI: 1.007–2.906, *P*=0.047) (10). This can be done at the time of ureteroscopy or in the office pending the clinical scenario and imaging findings.

The use of ureteroscopy with biopsy is currently the primary method for diagnosing UTUC if the imaging and cytology are not sufficient for the diagnosis and/or risk-stratification of the tumor (49,50). Flexible ureteroscopy allows visualization of the ureter, renal pelvis and collecting system and enables selective urinary cytology as well as biopsies for suspicious lesions or positive urinary cytology with negative enhanced cystoscopy. The specific size, appearance, and multifocality of UTUC can also be fully characterized—especially with the emergence of thinner

flexible ureteroscopes with digital technology and image enhancement techniques (65,66). Talso *et al.* compared seven different flexible ureteroscope systems (Olympus V, Olympus V2, Olympus P6, Wolf Cobra vision, Storz Flex XC, Storz Flex X2, and Lithovue) and found that the newer digital scopes offered better image quality than fiber optics (65). Narrow-band imaging (NBI), which has shown higher detection of bladder tumors with greater sensitivity on meta-analysis compared to white-light cystoscopy (67), has been evaluated in UTUC in two small prospective studies with promising but preliminary results (66,68). Confocal laser endomicroscopy (CLE) enables real-time *in vivo* visualization of tissue microarchitecture and cellular morphology using a low-energy laser light source and has been shown in small case series to have high concordance with definitive histology (69,70).

While there have been improvements in the visualization of UTUC, the endoscopic yield of biopsies remains a key factor in the reliability of risk stratification for UTUC. Acquisition of adequate samples of a sufficient quality and size for accurate pathologic interpretation is a technical challenge—mainly limited by the size of current ureteroscopes while maintaining their necessary maneuverability. There are numerous ureteroscopic biopsy devices that have been evaluated for their efficacy and utility—the 3F cold-cup biopsy forceps (Piranha), 6F BIGopsy[®] backloading cold-cup biopsy forceps, and 2.2/2.4F stone wire basket are the most common (71–73). Retrospective cohort studies have found diagnostic rates of 74.9–79% for 3F cold-cup forceps, 81.9–90% for BIGopsy[®], and 87–100% for 2.2F or 2.4F wire baskets (71–73)—with small size of sampled tissue being the main driver of false negatives (74,75). Forceps (either 3F forceps or BIGopsy[®]) are generally preferable for sampling smaller, flat, or non-papillary lesions, while the basket can be used to debulk large papillary tumors by avulsing large pieces of tumor. A newer ureteroscopic biopsy technique called the “form tackle” was recently described by Klett *et al.* (76) and was compared to standard biopsy techniques in a case series of 12 patients. In each patient they performed a “standard biopsy” with a 3F cold-cup biopsy forceps and a second biopsy utilizing the same biopsy forceps and the new “form tackle” technique. This new biopsy technique involves engaging the jaws of the forceps into the base of the tumor and advancing both the ureteroscope and biopsy forceps “forward, up, and away” from the mucosal wall until the tissue is released. Using the new technique, UTUC was diagnosed in an additional 4 patients (2 low-grade, 2 high-grade) in the setting of a

benign “standard technique” biopsy.

While using larger biopsy devices, different techniques, or taking multiple biopsies can enhance the quality/size of the biopsy, it has been suggested that addition of cytology should be mandatory to improve grading accuracy (75). Several studies have demonstrated the grade concordance of ureteroscopic biopsy with radical nephroureterectomy pathologic specimens is estimated between 68–77% (72,73). This discordance is also influenced by the overall size of tumor, as multiple small biopsies may only provide pathologic data for a small region and potentially miss high-grade morphology. Breda *et al.* found that only in tumors less than 2 cm is there a high concordance between ureteroscopic biopsy grade and final pathology (72). Likewise, staging information is limited by the depth of urothelial tissue able to be sampled endoscopically (77). Vashistha *et al.* retrospectively reviewed 120 endoscopic biopsies for UTUC and found poor stage concordance (58.6%) between ureteroscopic biopsy specimen and final pathology at time of nephroureterectomy (74). Similarly, Lama and colleagues found agreement between the biopsy grade and final pathologic stage occurred in 58% of patients in their multi-institutional retrospective collaboration with 145 patients (73). A recent comprehensive systemic review and meta-analysis was performed utilizing 23 studies (almost 3,600 patients) to evaluate the accuracy of ureteroscopic biopsies in predicting the stage and grade of UTUC (78). Interestingly, they found a substantial correlation between tumor grade at ureteroscopic biopsy and the final pathology (66%, 95% CI: 55–77%) for low-grade tumors and 97% (95% CI: 94–98%) for high-grade tumors. Additionally, the stage-to-stage match between ureteroscopic biopsy/final pathology showed a positive predictive value (PPV) for cT1+/muscle-invasive disease of 94% (95% CI: 84–100%) and a NPV for cTa-Tis/non-muscle-invasive disease of 60% (95% CI: 52–68%). These data demonstrate that invasion of the subepithelial connective tissue in the ureteroscopy specimen has a strong correlation with pathologic stage, however biopsy of the subepithelium is one of the main challenges of ureteroscopic biopsy (77). Nevertheless, a certain risk of undergrading and understaging should be assumed.

Additionally, various clinical scenarios can present significant challenges for the use of diagnostic ureteroscopy (previous urinary diversion, difficult calyceal anatomy, severe hematuria, etc.) leading some authors to investigate the safety, diagnostic yield, and histologic concordance of a percutaneous biopsy approach for diagnosing UTUC.

Huang *et al.* found that 85% of patients undergoing percutaneous biopsy for various indications (nondiagnostic ureteroscopy, urinary diversion, or being poor surgical candidates) received a diagnosis of UTUC—with 86% grade concordance with final pathology (79). However, similar to ureteroscopic biopsy, percutaneous biopsy has limited ability to stage the disease. The largest series of percutaneous biopsy for UTUC was published by Joseph and colleagues in 2020 and included 42 patients from 2009–2017 undergoing radical nephroureterectomy. They demonstrated 95% histologic concordance and 90% grade concordance relative to final nephroureterectomy pathology. Additionally, at a median follow-up of 28.2 months no cases of radiographic tract seeding were identified (80). The lack of tract seeding in this series [and Huang *et al.* (79)] was felt to be due to improved biopsy technique and incorporation of coaxial needle sets—which utilize a larger outer introducing needle for the initial insertion with multiple biopsies taken through the introducer needle to minimize the risk of tract seeding.

In addition to low risk of tract seeding with percutaneous biopsy, this approach may also decrease the risk of intravesical recurrence. A recent meta-analysis by Sharma *et al.*, comparing ureteroscopy *vs.* no ureteroscopy prior to nephroureterectomy found that 8 out of 12 studies demonstrated an increased risk for intravesical recurrence if ureteroscopy was performed. On multivariate analysis, ureteroscopy with biopsy was found to increase the risk of intravesical recurrence [hazard ratio (HR) 1.40, $P=0.04$], while ureteroscopy alone and percutaneous biopsy did not (HR 1.07, $P=0.87$; HR 1.15, $P=0.54$ respectively) (81). Similarly, Douglawi and colleagues retrospectively reviewed 143 patients who underwent radical nephroureterectomy for UTUC and found higher bladder recurrence rates for patients who had ureteroscopy prior *vs.* no ureteroscopy [30.8% *vs.* 7.7%, respectively ($P=0.02$)] (82). Interestingly, a lower recurrence rate was noted in patients whom a ureteral access sheath was utilized (11.5%) *vs.* those with no access sheath (39.7%, $P=0.01$)—potentially suggesting a protective effect of the ureteral access sheath. Multivariable analysis confirmed a significant increase in bladder recurrence if ureteroscopy was performed prior to radical nephroureterectomy [HR 5.6 (1.7–18.5), $P<0.004$], however, this effect was mitigated if a ureteral access sheath was used [HR 1.3 (0.3–6.4), $P=0.76$]. These data—along with other studies demonstrating the clonal relatedness of UTUC/subsequent bladder tumors (83), a decreased risk of bladder recurrence after nephroureterectomy with a single instillation

of postoperative intravesical chemotherapy (12,13), and extrapolation from the post-TURBT setting (84)—support the rationale for use of prophylactic intravesical chemotherapy after endoscopic biopsy to prevent bladder seeding, however this approach needs to be validated with future studies.

Risk stratification

Patient and tumor-related factors

UTUC is more common in men than women, and previous research has suggested that female gender may be associated with a worse oncologic outcome. However, after taking other prognostic factors into account (such as age, pT stage, neo/adjuvant chemotherapy, etc), no significant difference in cancer-specific survival or overall survival was found between genders on systematic review and meta-analysis (85). The relationship between ethnicity and UTUC outcomes is also uncertain, with some studies finding differences in survival between ethnic groups and others finding no significant differences (86). Whether these potential differences are related to biology or access to care is unknown. Older age at the time of radical nephroureterectomy has been shown to be independently associated with decreased cancer specific survival (87). Similarly, smoking is a well-established risk factor for UTUC, with long-term smokers at higher risk of more advanced disease stage, recurrence after radical nephroureterectomy, and cancer-specific mortality (88). Various medical comorbidities (chronic kidney disease, obesity), preoperative laboratory abnormalities (hypoalbuminemia, elevated C-reactive protein, elevated fibrinogen) along with poor performance status are also associated with worse survival outcomes across disease stages (89–91). Constitutional symptoms, including weight loss, fever, night sweats, anorexia, or hemoptysis are associated with a worse prognosis and should prompt a complete evaluation for the presence metastatic disease (32).

While there are certainly some patient related factors that are prognostic, the most accurate independent factors for predicted outcome in UTUC remain tumor grade and stage (87,92–94). In a study of UTUC patients in the Netherlands, 5-year cancer-specific survival was 86% for non-muscle-invasive tumors, 70% for muscle-invasive, organ-confined tumors, and 44% for locally-advanced tumors (95). Another contemporary analysis of radical nephroureterectomies for UTUC showed that 5-year cancer-specific survival was 86% for T1N0, 77%

for T2N0, 63% for T3N0, and 39% for T4N0/T any N1-3 (96). Hydronephrosis is often present on imaging (up to 50% of cases), and multiple studies have shown that hydronephrosis is associated with higher stage at diagnosis (41,42). Additionally, after adjusting for tumor stage, patients with ureteral and/or multifocal tumors appear to have a worse prognosis than patients with renal pelvic tumors (97,98). Larger tumors are more likely to be muscle-invasive and/or non-organ-confined in both ureteral and renal pelvis UTUC—with 2 cm being the best cut-off in identifying patient's with an increased risk of pT2 UTUC or higher (99). Variant histology in UTUC has also been found to be associated with worse recurrence free survival, cancer specific survival, and overall survival on a recent systematic review and meta-analysis of 26 studies with over 1,200 patients—with micropapillary and squamous and/or glandular variant histologies being associated with the worst cancer specific survival (100). Additionally, while previous studies have found mixed results with respect to previous/concomitant bladder cancer and CSS in patients with UTUC, a recent Surveillance, Epidemiology, and End Results (SEER) database review of 8,500 patients found that UTUC patients with previous or concomitant bladder cancer had a worse prognosis and higher risk of intravesical recurrence after radical nephroureterectomy (101). Various pre-operative serum and urine-based biomarkers have been investigated to help predict the prognosis of UTUC, however none of the investigated markers have been validated yet to support their introduction into routine clinical decision-making.

Prognostic tools

Several models have been developed to predict high-grade, muscle-invasive, or non-organ confined disease in UTUC based on various prognostic factors. Brien and colleagues utilized a combination of hydronephrosis, positive urinary cytology, and high-grade disease on biopsy to predict muscle invasive carcinoma at the time of radical nephroureterectomy. All three variables were independently associated with non-organ confined disease and abnormality of all three had 89% and 73% positive predictive value for muscle invasive and nonorgan confined UTUC, respectively, but when all tests were normal, the negative predictive value was 100% (42). In the same year, Margulis *et al.* proposed a different model to predict non-organ confined disease with an accuracy of 77% based

on tumor grade, architecture, and location (102). In a subsequent follow-up multi-institutional retrospective study, a new multivariable model was developed and demonstrated that clinical stage (OR 14.0, $P < 0.01$), biopsy tumor grade (OR 3.3, $P = 0.01$), tumor architecture (OR 2.65, $P = 0.09$), and hemoglobin (OR 0.8, $P = 0.02$) level were independently associated with non-organ confined disease. A preoperative nomogram incorporating these 4 variables achieved 82% accuracy, 48% sensitivity, and 95% specificity in predicting non-organ confined UTUC (103). These preoperative nomograms can be used to select patients more optimally for preoperative systemic chemotherapy and facilitate clinical trial enrollment—although some caution should be used as none have benefited from external validation.

Because clinical stage is so difficult to determine preoperatively, the EAU guidelines for UTUC have proposed stratifying patients as either low or high risk for progression in order to identify patients who are more likely to benefit from kidney-sparing treatment and those who should be treated with radical surgery (49). Based on the previously mentioned risk data and prognostic tools, the EAU proposed a set of parameters for pretreatment risk stratification of UTUC to support clinical decision-making. A low-risk tumor is defined as unifocal disease, lesion < 2 cm, low-grade cytology/biopsy, and no invasive aspect on imaging (all must be present). High-risk tumors include those presenting with hydronephrosis or multifocal disease, any high-grade status (on cytology or biopsy), or lesions > 2 cm. A history of radical cystectomy or variant histology are also criteria for high-risk tumors. Any of these high-risk features make it a high risk UTUC (*Table 1*). The predictive ability of the EAU risk stratification system to identify \geq pT2/N+ stage patients at radical nephroureterectomy was recently evaluated and compared to a new preoperative risk model utilizing over 1,200 patients in a multi-institutional retrospective cohort. This new model utilized age, high-grade biopsy, cT1+ biopsy, preoperative hydronephrosis, tumor size, invasion on imaging, and sessile tumor architecture to achieve an accuracy of 75%—compared to 71% for the EAU risk stratification system (104). Additionally, several new risk stratification strategies have been evaluated in comparison to the EAU risk grouping—including a tumour grade-and stage-based model (105) and a three-tier risk stratification model (i.e., low-, intermediate-, and high-risk) (106). Both strategies demonstrated a potential improvement on the current EAU risk stratification system but require further external validation.

Table 1 EAU risk stratification of non-metastatic UTUC

Low-risk UTUC (all criteria must be present)
Unifocal disease
Tumor size <2 cm
Low-grade cytology
Low-grade biopsy
No invasive aspect on imaging
High-risk UTUC (any criteria may be present)
Hydronephrosis
Tumor size >2 cm
High-grade cytology
High-grade biopsy
Multifocal disease
Previous radical cystectomy for bladder cancer
Variant histology
EAU, European Association of Urology; UTUC, upper tract urothelial carcinoma.

Conclusions

UTUC is a relatively rare disease that presents unique challenges to urologists from both a diagnostic and management standpoint. It is imperative to understand the epidemiology, natural history, genetics and known risk factors for this malignancy. Additionally, a thorough understanding of the diagnostic challenges, imaging/pathologic limitations, patient comorbidities, and risk stratification tools will allow us as a field to develop new modalities to improve our diagnostic capabilities and reduce the risk of under diagnosis and over treatment for our patients.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the Guest Editors (Ram A. Pathak and Ashok K. Hemal) for the series “Upper Tract Urothelial Cancer” published in *Translational Andrology and Urology*. The article has undergone external peer review.

Peer Review File: Available at <https://tau.amegroups.com/article/view/10.21037/tau-23-45/prf>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://tau.amegroups.com/article/view/10.21037/tau-23-45/coif>). The series “Upper Tract Urothelial Cancer” was commissioned by the editorial office without any funding or sponsorship. TDL has served as a consultant for Bristol Myers Squibb and ImmunityBio. DDT has received support from the Grail Research Support for Mayo Clinic, support to present at the Southeastern Section of the American Urological Association meeting; is the President-Elect of Florida Urological Society; and owns stock in Auris Robotics. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Siegel RL, Miller KD, Fuchs HE, et al. Cancer statistics, 2022. *CA Cancer J Clin* 2022;72:7-33.
2. Lughezzani G, Jeldres C, Isbarn H, et al. Temporal stage and grade migration in surgically treated patients with upper tract urothelial carcinoma. *BJU Int* 2010;105:799-804.
3. Munoz JJ, Ellison LM. Upper tract urothelial neoplasms: incidence and survival during the last 2 decades. *J Urol* 2000;164:1523-5.
4. Raman JD, Messer J, Sielatycki JA, et al. Incidence and survival of patients with carcinoma of the ureter and renal pelvis in the USA, 1973-2005. *BJU Int* 2011;107:1059-64.
5. Rouprêt M, Zigeuner R, Palou J, et al. European guidelines for the diagnosis and management of upper

- urinary tract urothelial cell carcinomas: 2011 update. *Eur Urol* 2011;59:584-94.
6. David KA, Mallin K, Milowsky MI, et al. Surveillance of urothelial carcinoma: stage and grade migration, 1993-2005 and survival trends, 1993-2000. *Cancer* 2009;115:1435-47.
 7. Soria F, Moschini M, Haitel A, et al. HER2 overexpression is associated with worse outcomes in patients with upper tract urothelial carcinoma (UTUC). *World J Urol* 2017;35:251-9.
 8. Raman JD, Ng CK, Scherr DS, et al. Impact of tumor location on prognosis for patients with upper tract urothelial carcinoma managed by radical nephroureterectomy. *Eur Urol* 2010;57:1072-9.
 9. Holmäng S, Johansson SL. Synchronous bilateral ureteral and renal pelvic carcinomas: incidence, etiology, treatment and outcome. *Cancer* 2004;101:741-7.
 10. Cosentino M, Palou J, Gaya JM, et al. Upper urinary tract urothelial cell carcinoma: location as a predictive factor for concomitant bladder carcinoma. *World J Urol* 2013;31:141-5.
 11. Shariat SF, Favaretto RL, Gupta A, et al. Gender differences in radical nephroureterectomy for upper tract urothelial carcinoma. *World J Urol* 2011;29:481-6.
 12. Ito A, Shintaku I, Satoh M, et al. Prospective randomized phase II trial of a single early intravesical instillation of pirarubicin (THP) in the prevention of bladder recurrence after nephroureterectomy for upper urinary tract urothelial carcinoma: the THP Monotherapy Study Group Trial. *J Clin Oncol* 2013;31:1422-7.
 13. O'Brien T, Ray E, Singh R, et al. Prevention of bladder tumours after nephroureterectomy for primary upper urinary tract urothelial carcinoma: a prospective, multicentre, randomised clinical trial of a single postoperative intravesical dose of mitomycin C (the ODMIT-C Trial). *Eur Urol* 2011;60:703-10.
 14. Foerster B, D'Andrea D, Abufaraj M, et al. Endocavitary treatment for upper tract urothelial carcinoma: A meta-analysis of the current literature. *Urol Oncol* 2019;37:430-6.
 15. Li WM, Shen JT, Li CC, et al. Oncologic outcomes following three different approaches to the distal ureter and bladder cuff in nephroureterectomy for primary upper urinary tract urothelial carcinoma. *Eur Urol* 2010;57:963-9.
 16. McLaughlin JK, Silverman DT, Hsing AW, et al. Cigarette smoking and cancers of the renal pelvis and ureter. *Cancer Res* 1992;52:254-7.
 17. Crivelli JJ, Xylinas E, Kluth LA, et al. Effect of smoking on outcomes of urothelial carcinoma: a systematic review of the literature. *Eur Urol* 2014;65:742-54.
 18. Hung RJ, Boffetta P, Brennan P, et al. GST, NAT, SULT1A1, CYP1B1 genetic polymorphisms, interactions with environmental exposures and bladder cancer risk in a high-risk population. *Int J Cancer* 2004;110:598-604.
 19. Colin P, Koenig P, Ouzzane A, et al. Environmental factors involved in carcinogenesis of urothelial cell carcinomas of the upper urinary tract. *BJU Int* 2009;104:1436-40.
 20. Debelle FD, Vanherweghem JL, Nortier JL. Aristolochic acid nephropathy: a worldwide problem. *Kidney Int* 2008;74:158-69.
 21. Markovic N, Ignjatovic I, Cukuranovic R, et al. Decreasing incidence of urothelial cancer in a Balkan endemic nephropathy region in Serbia. A surgery based study from 1969 to 1998. *Pathol Biol (Paris)* 2005;53:26-9.
 22. Tan LB, Chen KT, Guo HR. Clinical and epidemiological features of patients with genitourinary tract tumour in a blackfoot disease endemic area of Taiwan. *BJU Int* 2008;102:48-54.
 23. Stewart JH, Hobbs JB, McCredie MR. Morphologic evidence that analgesic-induced kidney pathology contributes to the progression of tumors of the renal pelvis. *Cancer* 1999;86:1576-82.
 24. De Broe ME, Elseviers MM. Analgesic nephropathy. *N Engl J Med* 1998;338:446-52.
 25. Brenner DW, Schellhammer PF. Upper tract urothelial malignancy after cyclophosphamide therapy: a case report and literature review. *J Urol* 1987;137:1226-7.
 26. Miest T, Khanna A, Sharma V, et al. Development and Validation of a Risk-Adapted Scoring Model for Metachronous Upper Tract Urothelial Carcinoma following Radical Cystectomy. *J Urol* 2022;207:284-92.
 27. Naser-Tavakolian A, Ghodoussipour S, Djaladat H. Upper urinary tract recurrence following bladder cancer therapy: a review of surveillance and management. *Curr Opin Urol* 2019;29:189-97.
 28. Miest TS, Sharma V, Boeri L, et al. Does Ureteral Stent Drainage Prior to Cystectomy Increase the Risk of Subsequent Upper Tract Urothelial Carcinoma and Ureteral Complications? *Urology* 2021;153:215-20.
 29. Sountoulides P, Pyrgidis N, Brookman-May S, et al. Does Ureteral Stenting Increase the Risk of Metachronous Upper Tract Urothelial Carcinoma in Patients with Bladder Tumors? A Systematic Review and Meta-analysis. *J Urol* 2021;205:956-66.
 30. Rouprêt M, Yates DR, Comperat E, et al. Upper urinary

- tract urothelial cell carcinomas and other urological malignancies involved in the hereditary nonpolyposis colorectal cancer (lynch syndrome) tumor spectrum. *Eur Urol* 2008;54:1226-36.
31. Mork M, Hubosky SG, Rouprêt M, et al. Lynch Syndrome: A Primer for Urologists and Panel Recommendations. *J Urol* 2015;194:21-9.
 32. Rouprêt M, Babjuk M, Compérat E, et al. European Association of Urology Guidelines on Upper Urinary Tract Urothelial Carcinoma: 2017 Update. *Eur Urol* 2018;73:111-22.
 33. Rink M, Robinson BD, Green DA, et al. Impact of histological variants on clinical outcomes of patients with upper urinary tract urothelial carcinoma. *J Urol* 2012;188:398-404.
 34. Stein A, Sova Y, Lurie M, et al. Adenocarcinoma of the renal pelvis. Report of two cases, one with simultaneous transitional cell carcinoma of the bladder. *Urol Int* 1988;43:299-301.
 35. Asano K, Miki J, Maeda S, et al. Clinical studies on inverted papilloma of the urinary tract: report of 48 cases and review of the literature. *J Urol* 2003;170:1209-12.
 36. Chevillat JC, Wu K, Sebo TJ, et al. Inverted urothelial papilloma: is ploidy, MIB-1 proliferative activity, or p53 protein accumulation predictive of urothelial carcinoma? *Cancer* 2000;88:632-6.
 37. Grainger R, Gikas PW, Grossman HB. Urothelial carcinoma occurring within an inverted papilloma of the ureter. *J Urol* 1990;143:802-4.
 38. Blank C, Lissmer L, Kaneti J, et al. Fibroepithelial polyp of the renal pelvis. *J Urol* 1987;137:962-3.
 39. Varela-Duran J, Urdiales-Viedma M, Taboada-Blanco F, et al. Neurofibroma of the ureter. *J Urol* 1987;138:1425-6.
 40. Qi N, Zhang J, Chen Y, et al. Microscopic hematuria predicts lower stage in patients with upper tract urothelial carcinoma. *Cancer Manag Res* 2018;10:4929-33.
 41. Ng CK, Shariat SF, Lucas SM, et al. Does the presence of hydronephrosis on preoperative axial CT imaging predict worse outcomes for patients undergoing nephroureterectomy for upper-tract urothelial carcinoma? *Urol Oncol* 2011;29:27-32.
 42. Brien JC, Shariat SF, Herman MP, et al. Preoperative hydronephrosis, ureteroscopic biopsy grade and urinary cytology can improve prediction of advanced upper tract urothelial carcinoma. *J Urol* 2010;184:69-73.
 43. Potretzke AM, Knight BA, Vetter JM, et al. Diagnostic Utility of Selective Upper Tract Urinary Cytology: A Systematic Review and Meta-analysis of the Literature. *Urology* 2016;96:35-43.
 44. Barkan GA, Wojcik EM, Nayar R, et al. The Paris System for Reporting Urinary Cytology: The Quest to Develop a Standardized Terminology. *Acta Cytol* 2016;60:185-97.
 45. McIntire PJ, Snow JT, Robinson BD, et al. Improved correlation of urinary cytology specimens using The Paris System in biopsy-proven upper tract urothelial carcinomas. *Cancer Cytopathol* 2018;126:498-504.
 46. Xing J, Monaco SE, Pantanowitz L. Utility of The Paris System for Reporting Urinary Cytology in upper urinary tract specimens. *J Am Soc Cytopathol* 2018;7:311-7.
 47. Zheng X, Si Q, Du D, et al. The Paris System for urine cytology in upper tract urothelial specimens: A comparative analysis with biopsy and surgical resection. *Cytopathology* 2018;29:184-8.
 48. Simon CT, Skala SL, Magers MJ, et al. The utility of upper urinary tract urine cytology before and after application of the Paris system. *Diagn Cytopathol* 2019;47:421-7.
 49. EAU Guidelines on Upper Urinary Tract Urothelial Cell Carcinoma - DIAGNOSIS - Uroweb. Uroweb - European Association of Urology. [cited 2022 Dec 18]. Available online: <https://uroweb.org/guidelines/upper-urinary-tract-urothelial-cell-carcinoma/chapter/diagnosis>
 50. Guidelines Detail. NCCN. [cited 2021 Nov 16]. Available online: <https://www.nccn.org/guidelines/guidelines-detail>
 51. Sokolova IA, Halling KC, Jenkins RB, et al. The development of a multitarget, multicolor fluorescence in situ hybridization assay for the detection of urothelial carcinoma in urine. *J Mol Diagn* 2000;2:116-23.
 52. Marín-Aguilera M, Mengual L, Ribal MJ, et al. Utility of fluorescence in situ hybridization as a non-invasive technique in the diagnosis of upper urinary tract urothelial carcinoma. *Eur Urol* 2007;51:409-15; discussion 415.
 53. Johannes JR, Nelson E, Bibbo M, et al. Voided urine fluorescence in situ hybridization testing for upper tract urothelial carcinoma surveillance. *J Urol* 2010;184:879-82.
 54. Monteiro-Reis S, Leça L, Almeida M, et al. Accurate detection of upper tract urothelial carcinoma in tissue and urine by means of quantitative GDF15, TMEFF2 and VIM promoter methylation. *Eur J Cancer* 2014;50:226-33.
 55. Guo RQ, Xiong GY, Yang KW, et al. Detection of urothelial carcinoma, upper tract urothelial carcinoma, bladder carcinoma, and urothelial carcinoma with gross hematuria using selected urine-DNA methylation biomarkers: A prospective, single-center study. *Urol Oncol* 2018;36:342.e15-23.
 56. Hayashi Y, Fujita K, Matsuzaki K, et al. Diagnostic potential of TERT promoter and FGFR3 mutations in

- urinary cell-free DNA in upper tract urothelial carcinoma. *Cancer Sci* 2019;110:1771-9.
57. Janisch F, Shariat SF, Baltzer P, et al. Diagnostic performance of multidetector computed tomographic (MDCTU) in upper tract urothelial carcinoma (UTUC): a systematic review and meta-analysis. *World J Urol* 2020;38:1165-75.
 58. Caoili EM, Cohan RH, Inampudi P, et al. MDCT urography of upper tract urothelial neoplasms. *AJR Am J Roentgenol* 2005;184:1873-81.
 59. Millán-Rodríguez F, Palou J, de la Torre-Holguera P, et al. Conventional CT signs in staging transitional cell tumors of the upper urinary tract. *Eur Urol* 1999;35:318-22.
 60. Takahashi N, Glockner JF, Hartman RP, et al. Gadolinium enhanced magnetic resonance urography for upper urinary tract malignancy. *J Urol* 2010;183:1330-65.
 61. Razavi SA, Sadigh G, Kelly AM, et al. Comparative effectiveness of imaging modalities for the diagnosis of upper and lower urinary tract malignancy: a critically appraised topic. *Acad Radiol* 2012;19:1134-40.
 62. Froemming A, Potretzke T, Takahashi N, et al. Upper tract urothelial cancer. *Eur J Radiol* 2018;98:50-60.
 63. Mohapatra A, Vemana G, Bhayani S, et al. Trends in the utilization of imaging for upper tract urothelial carcinoma. *Urol Oncol* 2016;34:236.e23-8.
 64. Voskuilen CS, Schweitzer D, Jensen JB, et al. Diagnostic Value of (18)F-fluorodeoxyglucose Positron Emission Tomography with Computed Tomography for Lymph Node Staging in Patients with Upper Tract Urothelial Carcinoma. *Eur Urol Oncol* 2020;3:73-9.
 65. Talso M, Proietti S, Emiliani E, et al. Comparison of Flexible Ureterorenoscope Quality of Vision: An In Vitro Study. *J Endourol* 2018;32:523-8.
 66. Traxer O, Geavlete B, de Medina SG, et al. Narrow-band imaging digital flexible ureteroscopy in detection of upper urinary tract transitional-cell carcinoma: initial experience. *J Endourol* 2011;25:19-23.
 67. Li K, Lin T, Fan X, et al. Diagnosis of narrow-band imaging in non-muscle-invasive bladder cancer: a systematic review and meta-analysis. *Int J Urol* 2013;20:602-9.
 68. Hao YC, Xiao CL, Liu K, et al. Application of narrow-band imaging flexible ureteroscopy in the diagnosis, treatment and follow-up of upper tract urothelial carcinomas. *Zhonghua Wai Ke Za Zhi* 2018;56:222-6.
 69. Breda A, Territo A, Guttilla A, et al. Correlation Between Confocal Laser Endomicroscopy (Cellvizio®) and Histological Grading of Upper Tract Urothelial Carcinoma: A Step Forward for a Better Selection of Patients Suitable for Conservative Management. *Eur Urol Focus* 2018;4:954-9.
 70. Villa L, Cloutier J, Cotè JF, et al. Confocal Laser Endomicroscopy in the Management of Endoscopically Treated Upper Urinary Tract Transitional Cell Carcinoma: Preliminary Data. *J Endourol* 2016;30:237-42.
 71. Kleinmann N, Healy KA, Hubosky SG, et al. Ureteroscopic biopsy of upper tract urothelial carcinoma: comparison of basket and forceps. *J Endourol* 2013;27:1450-4.
 72. Breda A, Territo A, Sanguedolce F, et al. Comparison of biopsy devices in upper tract urothelial carcinoma. *World J Urol* 2019;37:1899-905.
 73. Lama DJ, Safiullah S, Patel RM, et al. Multi-institutional Evaluation of Upper Urinary Tract Biopsy Using Backloaded Cup Biopsy Forceps, a Nitinol Basket, and Standard Cup Biopsy Forceps. *Urology* 2018;117:89-94.
 74. Vashistha V, Shabsigh A, Zynger DL. Utility and diagnostic accuracy of ureteroscopic biopsy in upper tract urothelial carcinoma. *Arch Pathol Lab Med* 2013;137:400-7.
 75. Margolin EJ, Matulay JT, Li G, et al. Discordance between Ureteroscopic Biopsy and Final Pathology for Upper Tract Urothelial Carcinoma. *J Urol* 2018;199:1440-5.
 76. Klett DE, Alom M, Wymer K, et al. A novel upper tract ureteroscopic biopsy technique: the "form tackle". *Int Braz J Urol* 2022;48:367-8.
 77. Skolarikos A, Griffiths TR, Powell PH, et al. Cytologic analysis of ureteral washings is informative in patients with grade 2 upper tract TCC considering endoscopic treatment. *Urology* 2003;61:1146-50.
 78. Subiela JD, Territo A, Mercadé A, et al. Diagnostic accuracy of ureteroscopic biopsy in predicting stage and grade at final pathology in upper tract urothelial carcinoma: Systematic review and meta-analysis. *Eur J Surg Oncol* 2020;46:1989-97.
 79. Huang SY, Ahrar K, Gupta S, et al. Safety and diagnostic accuracy of percutaneous biopsy in upper tract urothelial carcinoma. *BJU Int* 2015;115:625-32.
 80. Joseph JP, Potretzke TA, Packiam V, et al. Percutaneous Image-guided Core Needle Biopsy for Upper Tract Urothelial Carcinoma. *Urology* 2020;135:95-100.
 81. Sharma V, Miest TS, Juvet TS, et al. The Impact of Upper Tract Urothelial Carcinoma Diagnostic Modality on Intravesical Recurrence after Radical Nephroureterectomy: A Single Institution Series and Updated Meta-Analysis. *J Urol* 2021;206:558-67.

82. Douglawi A, Ghoreifi A, Lee R, et al. Bladder Recurrence Following Diagnostic Ureteroscopy in Patients Undergoing Nephroureterectomy for Upper Tract Urothelial Cancer: Is Ureteral Access Sheath Protective? *Urology* 2022;160:142-6.
83. Audenet F, Isharwal S, Cha EK, et al. Clonal Relatedness and Mutational Differences between Upper Tract and Bladder Urothelial Carcinoma. *Clin Cancer Res* 2019;25:967-76.
84. Messing EM, Tangen CM, Lerner SP, et al. Effect of Intravesical Instillation of Gemcitabine vs Saline Immediately Following Resection of Suspected Low-Grade Non-Muscle-Invasive Bladder Cancer on Tumor Recurrence: SWOG S0337 Randomized Clinical Trial. *JAMA* 2018;319:1880-8.
85. Mori K, Mostafaei H, Enikeev DV, et al. Differential Effect of Sex on Outcomes after Radical Surgery for Upper Tract and Bladder Urothelial Carcinoma: A Systematic Review and Meta-Analysis. *J Urol* 2020;204:58-62.
86. Matsumoto K, Novara G, Gupta A, et al. Racial differences in the outcome of patients with urothelial carcinoma of the upper urinary tract: an international study. *BJU Int* 2011;108:E304-9.
87. Collà Ruvolo C, Nocera L, Stolzenbach LF, et al. Incidence and Survival Rates of Contemporary Patients with Invasive Upper Tract Urothelial Carcinoma. *Eur Urol Oncol* 2021;4:792-801.
88. Rink M, Xylinas E, Margulis V, et al. Impact of smoking on oncologic outcomes of upper tract urothelial carcinoma after radical nephroureterectomy. *Eur Urol* 2013;63:1082-90.
89. Aziz A, Fritsche HM, Gakis G, et al. Comparative analysis of comorbidity and performance indices for prediction of oncological outcomes in patients with upper tract urothelial carcinoma who were treated with radical nephroureterectomy. *Urol Oncol* 2014;32:1141-50.
90. Kluth LA, Xylinas E, Kent M, et al. Predictors of survival in patients with disease recurrence after radical nephroureterectomy. *BJU Int* 2014;113:911-7.
91. Mori K, Janisch F, Mostafaei H, et al. Prognostic value of preoperative blood-based biomarkers in upper tract urothelial carcinoma treated with nephroureterectomy: A systematic review and meta-analysis. *Urol Oncol* 2020;38:315-33.
92. Margulis V, Shariat SF, Matin SF, et al. Outcomes of radical nephroureterectomy: a series from the Upper Tract Urothelial Carcinoma Collaboration. *Cancer* 2009;115:1224-33.
93. Clements T, Messer JC, Terrell JD, et al. High-grade ureteroscopic biopsy is associated with advanced pathology of upper-tract urothelial carcinoma tumors at definitive surgical resection. *J Endourol* 2012;26:398-402.
94. Mbeutcha A, Rouprêt M, Kamat AM, et al. Prognostic factors and predictive tools for upper tract urothelial carcinoma: a systematic review. *World J Urol* 2017;35:337-53.
95. van Doeveren T, van der Mark M, van Leeuwen PJ, et al. Rising incidence rates and unaltered survival rates for primary upper urinary tract urothelial carcinoma: a Dutch population-based study from 1993 to 2017. *BJU Int* 2021;128:343-51.
96. Rosiello G, Palumbo C, Knipper S, et al. Contemporary conditional cancer-specific survival after radical nephroureterectomy in patients with nonmetastatic urothelial carcinoma of upper urinary tract. *J Surg Oncol* 2020;121:1154-61.
97. Kaczmarek K, Lemiński A, Gołąb A, et al. Survival differences of patients with ureteral versus pelvicalyceal tumours: a systematic review and meta-analysis. *Arch Med Sci* 2019;17:603-12.
98. Chromecki TF, Cha EK, Fajkovic H, et al. The impact of tumor multifocality on outcomes in patients treated with radical nephroureterectomy. *Eur Urol* 2012;61:245-53.
99. Foerster B, Abufaraj M, Mari A, et al. The Performance of Tumor Size as Risk Stratification Parameter in Upper Tract Urothelial Carcinoma (UTUC). *Clin Genitourin Cancer* 2021;19:272.e1-e7.
100. Mori K, Janisch F, Parizi MK, et al. Prognostic Value of Variant Histology in Upper Tract Urothelial Carcinoma Treated with Nephroureterectomy: A Systematic Review and Meta-Analysis. *J Urol* 2020;203:1075-84.
101. Zeng S, Ying Y, Yu X, et al. Impact of previous, simultaneous or intravesical recurrence bladder cancer on prognosis of upper tract urothelial carcinoma after nephroureterectomy: a large population-based study. *Transl Androl Urol* 2021;10:4365-75.
102. Margulis V, Youssef RF, Karakiewicz PI, et al. Preoperative multivariable prognostic model for prediction of nonorgan confined urothelial carcinoma of the upper urinary tract. *J Urol* 2010;184:453-8.
103. Petros FG, Qiao W, Singla N, et al. Preoperative multiplex nomogram for prediction of high-risk nonorgan-confined upper-tract urothelial carcinoma. *Urol Oncol* 2019;37:292.e1-9.
104. Foerster B, Abufaraj M, Matin SF, et al. Pretreatment Risk Stratification for Endoscopic Kidney-sparing Surgery

- in Upper Tract Urothelial Carcinoma: An International Collaborative Study. *Eur Urol* 2021;80:507-15.
105. Katayama S, Mori K, Schuettfort VM, et al. Accuracy and Clinical Utility of a Tumor Grade- and Stage-based Predictive Model in Localized Upper Tract Urothelial Carcinoma. *Eur Urol Focus* 2022;8:761-8.
106. Marcq G, Foerster B, Abufaraj M, et al. Novel Classification for Upper Tract Urothelial Carcinoma to Better Risk-stratify Patients Eligible for Kidney-sparing Strategies: An International Collaborative Study. *Eur Urol Focus* 2022;8:491-7.

Cite this article as: Zganjar AJ, Thiel DD, Lyon TD. Diagnosis, workup, and risk stratification of upper tract urothelial carcinoma. *Transl Androl Urol* 2023;12(9):1456-1468. doi: 10.21037/tau-23-45