

POSTER PRESENTATION

Open Access

NAC regulation of embryo development in conifers

Emma Larsson*, Folke Sitbon, Jens Sundström, Sara von Arnold

From IUFRO Tree Biotechnology Conference 2011: From Genomes to Integration and Delivery
Arraial d'Ajuda, Bahia, Brazil. 26 June - 2 July 2011

Background

In most gymnosperms the cotyledons develop as a crown surrounding the incipient shoot apical meristem (SAM), which maintains the radial symmetry of the plant throughout embryogenesis. This is in contrast to the *Arabidopsis* embryo in which a symmetry-breaking event (from radial to bilateral symmetry) is associated with the emergence of the SAM and the two cotyledons [1]. We have previously shown that the differentiation of cotyledons and the establishment of the SAM and the root apical meristem (RAM) in somatic embryos of Norway spruce (*Picea abies*) is dependent on polar auxin transport (PAT) [2]. In *Arabidopsis* PAT restricts the expression of several genes belonging to the large plant specific NAC family of transcription factors [3-5], which in turn induce developmental processes involved in both shoot and root formation. Members of the NAC family have a conserved DNA-binding domain in the N-terminal region known as the NAC domain, and a more diverse domain in the C-terminal region, proposed to be responsible for the trans-activation capacity of the protein [6]. Despite the diversity of the NAC gene family, and the importance of NAC proteins in fundamental processes, no NAC genes have to our knowledge been characterized in conifers.

Methods

Phylogenetic analyses of NAC domain containing genes from *Picea glauca*, *Arabidopsisthaliana*, *Medicago truncatula* and *Physcomitrella patens* were performed using PAUP* and MrBayes. Full-length cDNAs of two NAC genes were isolated from maturing somatic embryos of Norway spruce and cloned using standard protocols. The promoters were amplified using the Genome

Walker™ kit (Clontech). The motif discovery tool MEME was used to find conserved motifs in the C-terminal regions and in the promoters of the NAC sequences from different species. The expression pattern of the two isolated NAC genes was analyzed in Norway spruce somatic embryos using quantitative real-time PCR. Their expression was also analyzed in embryos that had been treated with the PAT inhibitor NPA as described earlier [2]. Promoter::GUS fusions were inserted into embryogenic cultures of Norway spruce using *Agrobacterium* mediated transformation.

Results and discussion

We identified several different NAC sequences from white spruce in GenBank and deployed them to different sub-clades within the NAC gene family. Two full-length cDNAs (*gene 1* and *gene 2*) were cloned from Norway spruce. Both genes have elements in their promoter region which are significant for specific NAC genes. Furthermore, *gene 1* harbors previously characterized motifs that have been shown to have functional importance in NAC genes in *Arabidopsis*.

The relative expression of *gene 1* increased as early embryos differentiated, and remained at a high level until separated cotyledons were clearly visible. However, the up-regulation was blocked in embryos that formed fused cotyledons and lacked distinct SAM and RAM after being treated with NPA. The relative expression of *gene 2* was high in proembryogenic masses and early differentiating embryos but decreased as the embryos developed and matured. Interestingly the down-regulation occurred at about the same time as the cotyledons were delineated. The expression of *gene 2* was not dependent on PAT. Spatial expression analysis using promoter::GUS fusions revealed that the spruce genes have specific expression patterns, distinct from the

* Correspondence: emma.larsson@vbsg.slu.se
Swedish University of Agricultural Sciences, Department of Plant Biology and Forest Genetics, PO. Box 7080, 75007 Uppsala, Sweden

expression patterns of their most closely related *Arabidopsis* counterparts.

Conclusion

Taken together, our results show that NAC genes with distinct signature motifs existed before the separation of angiosperms and gymnosperms approximately 300 million years ago. However, through evolution the expression of closely related genes has diversified leading to induction of distinct developmental programs in angiosperms and gymnosperms.

Published: 13 September 2011

References

1. Chandler J: Cotyledon organogenesis. *J Exp Bot* 2008, **59**:2917-2931.
2. Larsson E, Sitbon F, Ljung K, von Arnold S: Inhibited polar auxin transport results in aberrant embryo development in Norway spruce. *New Phytol* 2008, **177**:356-366.
3. Aida M, Vernoux X, Furutani M, Traas J, Tasaka M: Roles of PIN-FORMED1 and MONOPTEROS in pattern formation of the apical region of the *Arabidopsis* embryo. *Development* 2002, **129**:3965-3974.
4. Furutani M, Vernoux T, Traas J, Kato T, Tasaka M, Aida M: PIN-FORMED1 and PINOID regulate boundary formation and cotyledon development in *Arabidopsis* embryogenesis. *Development* 2004, **131**:5021-5030.
5. Xie Q, Frugis G, Colgan D, Chua N-H: *Arabidopsis* NAC1 transduces auxin signal downstream of TIR1 to promote lateral root development. *Genes Dev* 2000, **14**:3024-3036.
6. Taoka K, Yanagimoto Y, Daimon Y, Hibara K, Aida M, Tasaka M: The NAC domain mediates functional specificity of CUP-SHAPED COTYLEDON proteins. *Plant J* 2004, **40**:462-473.

doi:10.1186/1753-6561-5-S7-P67

Cite this article as: Larsson et al.: NAC regulation of embryo development in conifers. *BMC Proceedings* 2011 **5**(Suppl 7):P67.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

