



Original Article

Crocin Promotes Apoptosis in Human EBV-Transformed B-Lymphocyte via Intrinsic Pathway

Abdolreza Sotoodeh Jahromi¹, Mohammad Kargar^{1*}, Farshid Kafilzadeh¹, Marzieh Jamalidoust² and Maliheh Moradzadeh^{3*}.

¹ Department of Microbiology, Jahrom Branch, Islamic Azad University, Jahrom, Iran

² Department of Clinical Virology, Clinical Microbiology Research Center, Shiraz University of Medical Sciences, Namazi Hospital, Shiraz, Iran.

³ Golestan Rheumatology Research Center, Sayad Shirazi Hospital, Golestan University of Medical Sciences, Gorgan, Iran.

*Both corresponding authors contributed and supervised equally to this research work and manuscript.

Competing interests: The authors declare no conflict of Interest.

Abstract. Background: As a major carotenoid in saffron, crocin demonstrates potent anti-cancer impacts. However, its anti-lymphoma effects remain vague, especially in the human EBV-associated B-cell lymphoproliferative disorders. This study examined crocin's apoptogenic potential and its underlying mechanism in CO 88BV59-1 cell line vs. normal human peripheral blood B cells.

Methods: CO 88BV59-1 cells were treated with crocin alone or in combination with vincristine for up to 72 h. The cell viability was examined using a resazurin assay. Flow cytometry using annexin V and propidium iodide labeling was performed to detect apoptotic cells. Also, the expression levels of genes and proteins involved in apoptosis (CASP3, CASP8, CASP9, P53, Bax, and Bcl-2) were respectively determined via real-time PCR and Western blot analysis.

Results: Crocin concentration-dependently reduced cell viability in CO 88BV59-1 cells with no significant toxicity toward normal B cells. Similar to vincristine, crocin significantly increased apoptosis in these cells during 72 h of incubation. Furthermore, the combination of crocin (80 µM) and vincristine (1 µM) enhanced apoptosis in CO 88BV59-1 cells. Therefore, this synergistic effect was detected in human EBV-transformed B-lymphocyte. CASP3, CASP9, P53, and Bax/Bcl-2 ratio expressions were significantly raised in CO 88BV59-1 cells, whereas CASP8 was unaltered. It was proposed that crocin promoted apoptosis in CO 88BV59-1 cells in a time- and concentration-dependent manner via the induction of the intrinsic pathway.

Conclusion: The results suggest that crocin may serve as a good alternative/coadjuvant to vincristine in EBV-associated B-cell lymphoproliferative disorders.

Keywords: Crocin; CO 88BV59-1 cells; EBV-associated B-cell lymphoproliferative disorders; Apoptosis.

Citation: Jahromi A.S., Kargar M., Kafilzadeh F., Jamalidoust M., Moradzadeh M. Crocin promotes apoptosis in human EBV-transformed B-lymphocyte via intrinsic pathway. *Mediterr J Hematol Infect Dis* 2021, 13(1): e2021049, DOI: <http://dx.doi.org/10.4084/MJHID.2021.049>

Published: July 1, 2021

Received: February 13, 2021

Accepted: June 18, 2021

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Correspondence to: Mohammad Kargar, Department of biology, Jahrom Branch, Islamic Azad University, Jahrom, Iran Tel: 09173149203 Fax: 071 5437 2001 Email: microkargar@gmail.com

Maliheh Moradzadeh, Golestan Rheumatology Research Center, Sayad Shirazi Hospital, Gorgan, Iran Telefax: + 98 17 32239791 Email: Moradzadeh63@yahoo.com

Introduction. B-cell lymphomas associated with Epstein-Barr virus (EBV) infection are common in patients who have immunodeficiency states, e.g., organ transplantation and human immunodeficiency virus (HIV) infection. Reports show that 1.8% of global cancer deaths in 2010 are related to EBV-attributable malignancies (Approximately 142,979).¹ The three main types of B cell malignancy associated with EBV are the Burkitt, Hodgkin, and diffuse large B-cell lymphomas (respectively BL, HL, and DLBCL). These pre-neoplastic/neoplastic lesions appearing in transplant patients are collectively referred to as post-transplant lymphoproliferative disorders (PTLD). The majority of PTLD cases are B cell, but 5-10% T and NK cell lymphoma have been described.² EBV inflicts >95% of the world's population and leads to lifelong asymptomatic infection. Its ability to induce oncogenesis may occur because of the suppression of the immune system or uncontrolled proliferation. The virus increases the activity of the B cell lymphoma 2 (Bcl-2) gene, which promotes cell proliferation by curbing apoptosis. The upregulation of the Bcl-2 gene can decrease the activity of tumor suppressor genes, including P53.³

Apoptosis is an evolutionary conserved, intrinsic program of cell death that occurs in various physiological and pathological states. The underlying mechanisms for initiating an apoptosis response upon cytotoxic therapy may depend on the individual stimulus and damage to DNA; however, damage to DNA or other critical molecules is considered a common initial event propagated by the cellular stress response. Apoptosis pathways can be initiated through different entry sites, for example, at the plasma membrane by death receptor ligation (named receptor or extrinsic pathway) or at the mitochondria (mitochondrial or intrinsic pathway). Bcl-2, P53, Bax, Caspase-9 (CASP9) are involved in the intrinsic pathway, and CASP8 in the extrinsic pathway.⁴

Introducing rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone (R-CHOP) to treat B-cell lymphoma can markedly improve patients' survival rate. Nevertheless, resistance to these drugs and their toxicity are significant obstacles to the course of treatment. Therefore, there is an urgent need for the development of novel therapeutic medications.⁵

Researchers held that dietary phytochemical agents might affect chemotherapy and contribute to the treatment of cancer patients.⁶ According to some studies, phytochemicals isolated from medicinal plants inhibited cell proliferation and induced apoptosis.⁷ At present, some of these plant-derived compounds are widely applied in chemotherapy for cancer treatment. For instance, taxol analogs, vinca alkaloids (vincristine and vinblastine), and podophyllotoxin analogs have greatly contributed to cancer treatment.⁸ Plant-derived compounds such as carotenoids play a pivotal role against cancer, providing a valuable source of anti-

cancer agents. Increasing evidence suggests that crocin, a carotenoid isolated from the saffron plant (*Crocus sativus* L), has anti-cancer effects in various types of cancer.⁹ Like vincristine, crocin's anti-proliferative activity involves targeting microtubules¹⁰ or p53-dependent and -independent mechanisms in cancer cells.^{11,12} In our laboratory, we observed that crocetin, a hydrolyzed form of crocin, exerts anti-proliferative, pro-apoptotic, and pro-differentiating impacts on human leukemia cells by inhibiting protein kinase B (Akt)-mediated pro-survival cascades, raising the intracellular Bcl-2-like protein 4 (Bax)/Bcl-2 ratio, and reducing Tyrosyl-DNA phosphodiesterase 1 (TDP1) enzyme activity and the expressions of promyelocytic leukemia-retinoic acid receptor alpha (PML-RAR α), Histone deacetylase 1 (HDAC1), and Multidrug resistance (MDR)-associated proteins.^{13,14}

Our laboratory has been interested in crocin's application in lymphoma and leukemia. Still, the exact mechanism of its action against EBV-associated B-cell lymphomas remains unknown. Thus, herein, a series of experiments were designed to examine crocin's apoptogenic potential and its underlying mechanisms in CO 88BV59-1 EBV-transformed B-lymphocyte vs. normal human B cells.

Materials and Methods

Cell line and reagents. Human CO 88BV59-1 EBV-transformed B-lymphocyte ([CRL-10624TM](#)) was purchased from ATCC (USA). The high-glucose Roswell Park Memorial Institute medium (RPMI 1640), penicillin-streptomycin, and fetal bovine serum (FBS) were obtained from Gibco BRL Life Technologies (USA). Moreover, 7-hydroxy-3H-phenoxazin-3-one-10-oxide (resazurin), crocin (>95%), vincristine (>95%), Fluorescein isothiocyanate (FITC) annexin V antibody, and propidium iodide (PI) were procured from Sigma-Aldrich (USA). TRIzol was obtained from Invitrogen (USA). A real-time PCR Master Mix and a cDNA synthesis Kit were also purchased from Roche Diagnostic (Switzerland) and Fermentas (Lithuania). Moreover, an enhanced chemiluminescence (ECL) detection kit and polyvinylidene difluoride (PVDF) membranes were purchased from GE Healthcare (UK) and Millipore (USA), in that order. Primary antibodies for β -actin, Bcl-2, Bax, P53, cleaved caspases (3, 8, and 9), and secondary antibodies were obtained from Cell Signaling Technology (USA). Fluorescein isothiocyanate-conjugated antibody against CD19 was also purchased from BioLegend (USA). Finally, a human B cell isolation kit was obtained from Miltenyi Biotec (Germany).

Human normal B cell isolation and cell culture. Human peripheral blood mononuclear cells (PBMCs) were obtained from fresh blood samples taken from healthy




Figure 1. Normal B cells purification using MACS technique as evaluated by flow cytometry. **A)** Representative histogram of the fluorescence intensity of CD19-labeled normal B cells. In a series of three experiments, B cells were enriched from PBMC, to give populations of 94-98% CD19-positive B cells. **B)** IgG isotype control was used as a negative control to differentiate the background noise of the flow cytometry analysis.

volunteers by Ficoll-density (Pharmacia, Sweden) gradient centrifugations. Subsequently, B-cells were isolated from PBMC via a B-cell isolation kit. Then, the purity of the cellular preparation was tested via FITC-conjugated anti-human CD19 antibody staining in the MACS analysis, and it was found to be >97% pure (**Figure 1**).¹⁵ CO 88BV59-1 and normal B cells were also cultured in the RPMI medium containing 10% (v/v) FBS, 100 units/ml penicillin, and 100 µg/ml streptomycin maintained at 37 °C in a humidified atmosphere (90%) containing 5% CO₂. The cells were subsequently incubated with different concentrations of vincristine (0.05-50 µM) and crocin (0.2-200 µM) up to 72 h. All the treatments were conducted in triplicate.

Cell viability assay. We determined cell viability by the resazurin reagent. For this purpose, CO 88BV59-1 and normal B cells (1×10^5) were added to each well in 96-well culture plates treated with vincristine (0.05-50 µM) and crocin (0.2-200 µM) up to 72 h. Next, 20 µl of the resazurin reagent was added to each well, and the plates were incubated for 4 h. The fluorescence intensity of the product resorufin, proportional to the number of viable cells per well, was measured via a fluorescence Victor X5 2030 Multilabel Plate Reader (Perkin Elmer, Shelton, Connecticut) with excitation at 530 nm and emission at 590 nm.¹⁶

Cell apoptosis assay. Crocin's apoptosis effect on CO 88BV59-1 cells was assessed by FITC annexin V/PI staining. The cells were treated with crocin (175, 112, and 80 µM) and vincristine (35, 23, and 1 µM) according

to IC50 values for different durations (24, 48, and 72 h), respectively. Also, we assessed the combination of crocin (80 µM) and vincristine (1 µM) on these cells for 72 h. After the treatment, the cells were incubated with the FITC annexin V antibody and analyzed by a flow cytometer (BD Biosciences, USA). The FlowJo software (TreeStar Inc.) was employed for data analysis.¹⁷

Real-time PCR quantification with SYBR Green. The CO 88BV59-1 cells were treated with crocin (175, 112, and 80 µM) alone or in combination with vincristine (35, 23, and 1 µM) up to 72 h., and then RNA extraction was done using TRIzol according to the manufacturer's instruction. RNA concentration and purity were evaluated via spectrophotometry. For each sample, the complementary DNA (cDNA) was synthesized from the total RNA (100 ng) via a cDNA synthesis kit with the random hexamer primer. Primers (Bcl-2, Bax, P53, CASP3, CASP8, and CASP9 genes) were designed using the Beacon software (Applied Biosystems; **Table 1**). Gene expression changes were determined using SYBR Green-based real-time PCR technology by the Applied Biosystems Step One Plus Detection System (ABI, USA). The reaction mixture comprised 1 µl of the primers (100 pmol), 2 µl of cDNA (250-400 ng), 10 µl of 2x master mix, and dH₂O to bring the volume to 20 µl. The optimized parameters utilized for the thermocycler included a short hot-start at 95 °C for 15 min, followed by 40 cycles, each consisting of denaturing at 95 °C for 15 secs, annealing at 60 °C for 1 min, and extension at 72 °C for 20 sec. Melting curves were used from 60 to 90 °C rising by 0.3 °, as the final step of the SYBR Green

Table 1: Sequences of primer selected for real-time PCR quantification using SYBR Green.

GENE	FORWARD PRIMER	REVERSE PRIMER	PRODUCT SIZE (BP)
Bcl-2	CCAAGAAAGCAGGAAACC	GGATAGCAGCACAGGATT	170
Bax	GCCTCCTCTCCTACTTTG	CTCAGCCCATCTTCTTCC	102
P53	GGAAGTCAAGGATGCCAG	CAAGAAGTGGAGAATGTCAGTC	155
CASP3	AGAACTGGACTGTGGCATT	GCTTGTGCGCATACTGTTT	191
CASP8	TGTTGGAGGAAAGCAATCTG	CCTGGTGTCTGAAGTTCCCT	124
CASP9	CTTTGTGTCCTACTCTACTTTCC	AACAGCATTAGCGACCCTA	151
GAPDH	GAAGTCAGGTGGAGCGAGG	TGGGTGGAATCATATTGGAACAT	200

real-time PCR. Gene expressions were normalized to Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) as the housekeeping gene. The samples were run in triplicate, and the fold difference of expression in the treated and untreated samples was calculated using the $2^{-\Delta\Delta Ct}$ method.¹⁸

Western blot analysis. Following 72 h of treatment with crocin (80 μM) alone or in combination with vincristine (1 μM), the cells were lysed using the lysis buffer and centrifuged at 18000 g, at 4 °C for one h, and then the supernatant was collected. Next, the lysates were run on 10% sodium dodecyl sulfate (SDS)-polyacrylamide gel and then transferred onto nitrocellulose membranes. After blocking with 2% Bovine serum albumin (BSA), the blots were exposed to the primary antibody for one h at room temperature. In the next step, they were washed and incubated with the corresponding horseradish peroxidase-conjugated to the secondary antibody for two h. Finally, membrane visualization was performed using an ECL detection kit. The reactions were revealed and documented by Gel-Doc (Syngene, Cambridge, UK), and the images were quantified using the Image J software (version 1.46).¹⁹

Statistical analysis. The data are represented as mean \pm

SD and were analyzed using one-way analysis of variance (ANOVA) with Tukey's multiple comparisons post-hoc test in the Graph Pad PRISM software (Version 6, Graph Pad Software, CA). A p-value <0.05 was considered statistically significant.

Results.

Crocin concentration-dependently reduced the viability of CO 88BV59-1 cells. Crocin at concentrations of 100 and 200 μM significantly decreased the viability of CO 88BV59-1 cells at 48 and 72 h ($p < 0.05$; **Figure 2A**). Similarly, a significant drop in viability was observed in these cells incubated for 48 and 72 h with 25 and 50 μM of vincristine ($p < 0.05$). However, crocin did not affect the viability of normal B cells at concentrations of 0.2-200 μM (**Figure 2B**). **Table 2** presents the IC₅₀ values of crocin and vincristine in CO 88BV59-1 cells for 24, 48, and 72 h of incubation.

Table 2: IC₅₀ values of crocin and vincristine in CO 88BV59-1 cell line up to 72 h incubation.

IC ₅₀	24 h	48 h	72 h
Crocin (μM)	174.7 \pm 0.08	112.0 \pm 0.14	79.2 \pm 0.12
Vincristine (μM)	34.9 \pm 0.17	23.2 \pm 0.07	1.2 \pm 0.15




Figure 2. Effects of crocin on the viability of CO 88BV59-1 cells. **A)** The cells were treated with various concentrations of crocin (0.2-200 μM) and vincristine (0.05-50 μM) up to 72 h. **B)** Normal B cells were treated with various concentrations of crocin (0.2-200 μM) up to 24 h. Cell viability was determined using the resazurin assay. The data are expressed as the mean \pm SEM of three independent experiments performed in triplicate. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. untreated control cells (concentration of 0).




Figure 3. Effects of crocin alone or in combination with vincristine on the apoptosis of CO 88BV59-1 cells as evaluated by annexin V and propidium iodide double-staining. The representative dot blot of the fluorescence intensity of annexin V and PI double-stained cells treated with crocin (175, 112 and 80 μM) and vincristine (35, 23 and 1 μM) based on their IC₅₀ value for 24, 48, and 72 h. Also, these cells were treated with combination of crocin (80 μM) and vincristine (1 μM) for 72 h. Quantitative analysis was performed in the Flow Jo software. The data are presented as the mean \pm SEM of three independent experiments performed in triplicate. *** $p < 0.001$ vs. untreated control cells (concentration of 0)




Figure 4. Effects of crocin alone or in combination with vincristine on the expression of apoptotic and anti-apoptotic genes in CO 88BV59-1 cells. The cells were treated with vincristine (35, 23 and 1 μ M) or crocin (175, 112 and 80 μ M) based on their IC₅₀ value up to 72 h. Also, these cells were treated with combination of crocin (80 μ M) and vincristine (1 μ M) for 72 h. Next, the expression levels of apoptotic (Bax, P53, CASP3, CASP8, and CASP9) and anti-apoptotic (Bcl-2) genes were determined by real-time PCR. The data are expressed as the mean \pm SEM of three independent experiments performed in triplicate. * p < 0.05, ** p < 0.01, *** p < 0.001 vs. untreated control cells (concentration of 0).

Crocin time-dependently induced apoptosis in CO 88BV59-1 cells. **Figure 3** displays the impacts of crocin on the apoptosis of CO 88BV59-1 cells, assessed by annexin V and PI double staining. Similar to vincristine, crocin (175, 112, and 80 μ M) significantly and time-dependently spiked the apoptosis rate of these cells (p < 0.001). Furthermore, the combination of crocin (80 μ M) and vincristine (1 μ M) remarkably enhanced apoptosis by up to 68% in CO 88BV59-1 cells (p < 0.001).

Crocin modulated genes involved in survival and apoptosis in CO 88BV59-1 cells. **Figure 4** depicts the effects of crocin alone or in combination with vincristine on the expression of genes involved in survival (Bcl-2) and apoptosis (P53, Bax, CASP3, CASP8, and CASP9) in CO 88BV59-1 cells up to 72 h. The expressions of P53, CASP3, CASP8, and CASP9 were significantly raised in these cells treated with either crocin or vincristine (p <

0.001 compared with control cells). In addition, crocin alone or in combination with vincristine significantly increased the Bax/Bcl-2 ratio to 4.4 ± 0.16 and 5.7 ± 0.20 -fold in the cells during 72 h, respectively (p < 0.001 compared with control cells; **Figure 4**).

To further assess the impact of crocin on apoptotic genes, their protein levels were evaluated by western blot analysis (**Figure 5**). Treatment of cells with crocin increased the protein expressions of P53, Bax, CASP3, and CASP9 but reduced the Bcl-2 expression. On the contrary, the expression of CASP8 protein did not change in CO 88BV59-1 cells following treatment with crocin (**Figure 5**); however, crocin combined with vincristine raised the expression of CASP8 (p < 0.001). The levels of Bax/Bcl-2 protein was significantly increased in crocin (80 μ M) alone or in combination with vincristine (1 μ M)-treated cells compared to untreated cells (4.09 ± 0.18 and 5.4 ± 0.11 , respectively; p < 0.001), thereby confirming the synergistic effect of crocin on CO 88BV59-1 cells. A similar surge was also observed in the cells incubated with 1 μ M of vincristine (4.2 ± 0.11 , p < 0.001). It was demonstrated that crocin significantly increased Bax/Bcl-2 ratio, approximately equal to vincristine.

Discussion. This study was the first to examine the mechanism of apoptotic cell death induced by crocin in EBV-transformed B-lymphocyte (CO 88BV59-1 cell line), compared to the standard anti-lymphoma drug, vincristine, during short-term treatment. Crocin effectively inhibited cell proliferation and induced apoptosis at high concentrations in CO 88BV59-1 cells during 72 h of treatment. The cytotoxic effect of crocin was more noticeable against CO 88BV59-1 cells than against normal human B cells, and these effects were comparable with those of vincristine. Furthermore, crocin significantly up-regulated the expression P53, CASP3, CASP9, and Bax/Bcl-2 ratio in CO 88BV59-1 cells at the mRNA and protein level, whereas the CASP8 protein remained unchanged. The present study's





Figure 5: Effects of crocin alone or in combination with vincristine on the expression of apoptotic and anti-apoptotic proteins in CO 88BV59-1 cells. The cells were treated with crocin (80 μ M) alone or in combination with vincristine (1 μ M) for 72 h. Then, the expression levels of apoptotic (Bax, P53, CASP3, CASP8, and CASP9) and anti-apoptotic (Bcl-2) proteins were determined by the Western blot test. Quantitative analysis was performed in the Image J software, and the mean \pm SEM of the three independent experiments are given. ** p < 0.01, *** p < 0.001 vs. untreated control cells (concentration of 0).

findings revealed that crocin induces apoptosis via the intrinsic pathway in a concentration- and time-dependent manner in CO 88BV59-1 cells. Interestingly, the combination of crocin (80 μ M) and vincristine (1 μ M) induced apoptosis in these cells via both pathways (intrinsic and extrinsic). Therefore, this synergistic apoptotic effect of crocin and vincristine was detected in these cells.

Patients who have acquired or inherited immune incompetence demonstrate a high incidence of lymphoma. A common factor in these patients seems to be the impairment of immunoregulatory mechanisms involved in neoplastic and/or viral surveillance.²⁰ The incidence of EBV among B-cell lymphoma patients is <5% in the United States and European countries, but 10-15% in Latin American and Asian countries.^{21,22} However, EBV-related B-cell lymphomas in transplant recipients show some additional characteristics; most notably, a large proportion of these tumors tend to regress spontaneously upon immunosuppression withdrawal or reduction, even though they are almost universally fatal if they remain untreated.²³

Several studies reported that nontoxic natural agents could be useful either alone or in combination with conventional therapeutics to prevent tumor progression and/or treat human malignancies.²⁴

The fact that crocin is abundantly available in large quantities in food products and is reportedly nontoxic makes its anti-cancer effect even more attractive.²⁵ Furthermore, since it also possesses immunosuppressive characteristics, it can exert potent anti-inflammatory effects in autoimmune diseases via inhibiting cytokines.²⁶⁻²⁹ Consistent with our findings, a study explored the effect of crocin on the proliferation and differentiation of HL-60 cells during long-term (5 days) exposure.³⁰ In another research, crocin inhibited proliferation and induced apoptosis in leukemic cell lines (K562, HL-60, L1210, and P388):³¹ crocin also inhibited the proliferation of Jurkat and HL60 cells by reducing cell growth and induced apoptosis by raising the Bax/Bcl-2 ratio.^{32,33}

Some studies showed that the anti-proliferative activity of crocin like vincristine involves targeting microtubules¹⁰ and p53-dependent and -independent mechanisms in colon cancer cells.^{11,12}

Crocetin triggers apoptosis by increasing the Bax/Bcl-2 ratio and caspase activation in human gastric adenocarcinoma without affecting human normal fibroblast skin cells.³⁴ Also, Luo et al. reported that the combination of crocin with cisplatin exerts growth suppression and apoptosis in gastric carcinoma cells.³⁵ Crocetin (hydrolyzed crocin) induced p53-mediated cell death in functional p53-expressing cancer cells through Bax and P53-induced protein with a death domain (PIDD) caspase-2-t-BH3 interacting-domain death (BID) pathway.³⁶ Another study showed that dimethyl-

crocetin and crocin induced cytotoxicity on HL60 cells but did not affect K562 cells. They suggested that dimethyl-crocetin could disrupt DNA-protein interactions (e.g., topoisomerase II) and inhibit nucleic acid synthesis.³⁷ Controversially, the other study showed that crocetin, unlike silymarin, retinoic acid, and other drugs, was unable to prevent the neoplastic transformation of rat tracheal epithelial cells by Benzopyrene.³⁸ Another study presented that crocin suppressed multiple human myeloma growth through inhibition of STAT3-mediated gene products, including BAX, Bcl-2, vascular endothelial growth factor (VEGF), CXC Chemokine Receptor 4 (CXCR4), and cell cycle regulator (cyclin D1).³⁹ Xu et al. showed that crocin could block HL-60 cells in the G₀/G₁ phase and inhibit their proliferation. The suggested mechanism in these cells may be related to the inhibition of Bcl-2 and activation of Bax.⁴⁰

The studies investigated the effect of crocin on the proliferation and immune function of dendritic cells (DC) derived from the bone marrow of children with acute leukemia. They concluded that crocin could synergically promote the maturity of DC cooperating with recombinant human granulocyte-macrophage colony-stimulating factor (them-CSF), recombinant human IL-4 (rhIL-4), and recombinant human TNF- α (rhTNF- α). The DC induced by crocin can particularly enhance the proliferation of T cells.^{41,42}

According to Molnar et al., crocin and crocetin were ineffective in reversing multidrug resistance of lymphoma cells but inhibited the early tumor antigen expression of adenovirus-infected mouse lymphoma cells.⁴³ It is noteworthy that crocin demonstrates a significant antiviral activity against HSV-1 and also HIV-1.⁴⁴ Another study also showed that crocin concentration- and time-dependently inhibited the proliferation and prolonged the lifespan of Dalton's lymphoma-bearing animals through significant effects on hematological parameters.⁴⁵ On the other hand, Khavari et al. concluded that a combination of DNA vaccine with crocin did not potentiate protective and therapeutic effects compared to mono-therapies for controlling papillomavirus-infected tumors.⁴⁶ Based on another study, crocin exhibited low cytotoxic effects on the MOLT-4 cell line, which might be mediated through the escalation of DNA fragmentation.⁴⁷ Also, crocin significantly and concentration-dependently promoted T cell proliferation and IL-2 and IL-4 secretion. Crocin itself caused no significant damage to T cells but curbed DNA damage in T cells treated with cytarabine.⁴⁸ Clinical experiments reported that healthy volunteers treated with saffron tablets (200 mg/kg) did not demonstrate hematological or biochemical toxicity.⁴⁹ In addition, a pharmacokinetic study suggested that orally administered crocins are hydrolyzed to crocetin before or during intestinal absorption.⁵⁰ The LD50 of crocin has

been reported to be >3 g/kg.⁵¹

The present study had some limitations partially due to a reduced financial budget. This study was carried out on only one EBV-transformed B cell, and only six genes and proteins (apoptotic and anti-apoptotic) were evaluated in this research work. It should be better to utilize more than two EBV-transformed B cells and study other apoptotic and anti-apoptotic genes.

Conclusions. The results showed that crocin promotes apoptosis in CO 88BV59-1 cells in a time- and concentration-dependent manner via the induction of the P53-dependent intrinsic pathway. Furthermore, crocin

and vincristine have a synergistic effect on these cells. Thus, it is suggested from these preclinical studies to evaluate the effect of crocin alone or in combination with vincristine in EBV-associated B-cell lymphoproliferative disorders.

Ethical consideration. This study was approved by the Research Ethics Committee of Jahrom University of Medical Sciences (ethic code: IR.JUMS.REC.1399.026).

Acknowledgments. We appreciate the insightful comments of Dr. Saiedeh Erfanian, which greatly enhanced an early version of this paper.

References:

1. Pei Y, Lewis AE and Robertson ES. Current progress in EBV-associated B-cell lymphomas. In: editors. Infectious Agents Associated Cancers: Epidemiology and Molecular Biology. Springer; 2017. p. 57-74. https://doi.org/10.1007/978-981-10-5765-6_5 PMID:29052132 PMCID:PMC6053051
2. Shannon-Lowe C, Rickinson AB and Bell AI. Epstein-Barr virus-associated lymphomas. Philosophical Transactions of the Royal Society B: Biological Sciences 2017; 372: 20160271. <https://doi.org/10.1098/rstb.2016.0271> PMID:28893938 PMCID:PMC5597738
3. Fu Q, He C and Mao ZR. Epstein-Barr virus interactions with the Bcl-2 protein family and apoptosis in human tumor cells. J Zhejiang Univ Sci B 2013; 14: 8-24. <https://doi.org/10.1631/jzus.B1200189> PMID:23303627 PMCID:PMC3542954
4. Fulda S and Debatin K-M. Extrinsic versus intrinsic apoptosis pathways in anti-cancer chemotherapy. Oncogene 2006; 25: 4798-4811. <https://doi.org/10.1038/sj.onc.1209608> PMID:16892092
5. Coiffier B and Sarkozy C. Diffuse large B-cell lymphoma: R-CHOP failure-what to do? Hematology 2014, the American Society of Hematology Education Program Book 2016; 2016: 366-378. <https://doi.org/10.1182/asheducation-2016.1.366> PMID:27913503 PMCID:PMC6142522
6. Wegiera M, Smolarz HD, Jedruch M, Korczak M and Koproń K. Cytotoxic effect of some medicinal plants from Asteraceae family on J-45.01 leukemic cell line-pilot study. Acta poloniae pharmaceutica 2012; 69: 263-268.
7. Shu L, Cheung K-L, Khor TO, Chen C and Kong A-N. Phytochemicals: cancer chemoprevention and suppression of tumor onset and metastasis. Cancer and Metastasis Reviews 2010; 29: 483-502. <https://doi.org/10.1007/s10555-010-9239-y> PMID:20798979
8. Saklani A and Kutty SK. Plant-derived compounds in clinical trials. Drug discovery today 2008; 13: 161-171. <https://doi.org/10.1016/j.drudis.2007.10.010> PMID:18275914
9. Khorasanchi Z, Shafiee M, Kermanshahi F, Khazaei M, Ryzhikov M, Parizadeh MR, Kermanshahi B, Ferns GA, Avan A and Hassanian SM. Crocus sativus a natural food coloring and flavoring has potent anti-tumor properties. Phytomedicine 2018; 43: 21-27. <https://doi.org/10.1016/j.phymed.2018.03.041> PMID:29747750
10. Hire RR, Srivastava S, Davis MB, Kumar Konreddy A and Panda D. Antiproliferative Activity of Crocin Involves Targeting of Microtubules in Breast Cancer Cells. Scientific reports 2017; 7: 44984-44984. <https://doi.org/10.1038/srep44984> PMID:28337976 PMCID:PMC5364484
11. Zhong Y-j, Shi F, Zheng X-l, Wang Q, Yang L, Sun H, He F, Zhang L, Lin Y, Qin Y, Liao L-c and Wang X. Crocetin induces cytotoxicity and enhances vincristine-induced cancer cell death via p53-dependent and -independent mechanisms. Acta pharmacologica Sinica 2011; 32: 1529-1536. <https://doi.org/10.1038/aps.2011.109> PMID:21986580 PMCID:PMC4010206
12. Harchegani AB, Khor A, Niha MM, Kaboutaraki HB, Shirvani H and Shahriari A. The hepatoprotective and antioxidative effect of saffron stigma alcoholic extract against vincristine sulfate induced toxicity in rats. Interdiscip Toxicol 2019; 12: 186-191. <https://doi.org/10.2478/intox-2019-0023> PMID:32461722 PMCID:PMC7247369
13. Moradzadeh M, Ghorbani A, Erfanian S, Mohaddes S, Rahimi H, Karimiani E, Mashkani B, Chiang S, El-Khamisy S and Tabarraei A. Study of the mechanisms of crocetin-induced differentiation and apoptosis in human acute promyelocytic leukemia cells. Journal of cellular biochemistry 2018; <https://doi.org/10.1002/jcb.27489> PMID:30203596
14. Moradzadeh M, Tabarraei A, Ghorbani A, Hosseini A and Sadeghnia HR. Short - term in vitro exposure to crocetin promotes apoptosis in human leukemic HL - 60 cells via intrinsic pathway. Acta Poloniae Pharm Drug Res 2018; 75: 445-451.
15. Moore DK, Motaung B, du Plessis N, Shabangu AN, Loxton AG and Consortium S-I. Isolation of B-cells using Miltenyi MACS bead isolation kits. PloS one 2019; 14: e0213832. <https://doi.org/10.1371/journal.pone.0213832> PMID:30893384 PMCID:PMC6426237
16. Mashkani B, Tanipour MH, Saadatmandzadeh M, Ashman LK and Griffith R. FMS-like tyrosine kinase 3 (FLT3) inhibitors: Molecular docking and experimental studies. European journal of pharmacology 2016; 776: 156-166. <https://doi.org/10.1016/j.ejphar.2016.02.048> PMID:26896780
17. Rangarajan P, Dharmalingam Subramaniam SP, Kwatra D, Palaniyandi K, Islam S, Harihar S, Ramalingam S, Gutheil W, Putty S and Pradhan R. Crocetin acid inhibits hedgehog signaling to inhibit pancreatic cancer stem cells. Oncotarget 2015; 6: 27661. <https://doi.org/10.18632/oncotarget.4871> PMID:26317547 PMCID:PMC4695016
18. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic acids research 2001; 29: e45. <https://doi.org/10.1093/nar/29.9.e45> PMID:11328886 PMCID:PMC55695
19. Moradzadeh M, Tabarraei A, Sadeghnia HR, Ghorbani A, Mohamadkhani A, Erfanian S and Sahebkar A. Kaempferol increases apoptosis in human acute promyelocytic leukemia cells and inhibits multidrug resistance genes. Journal of Cellular Biochemistry 2018; 119: 2288-2297. <https://doi.org/10.1002/jcb.26391> PMID:28865123
20. Marques-Piubelli ML, Salas YI, Pachas C, Becker-Hecker R, Vega F and Miranda RN. Epstein-Barr virus-associated B-cell lymphoproliferative disorders and lymphomas: a review. Pathology 2020; 52: 40-52. <https://doi.org/10.1016/j.pathol.2019.09.006> PMID:31706670
21. Gibson SE and Hsi ED. Epstein-Barr virus-positive B-cell lymphoma of the elderly at a United States tertiary medical center: an uncommon aggressive lymphoma with a nongermline center B-cell phenotype. Human pathology 2009; 40: 653-661. <https://doi.org/10.1016/j.humpath.2008.10.007>

- PMid:19144386
22. Hoeller S, Tzankov A, Pileri SA, Went P and Dirnhofer S. Epstein-Barr virus-positive diffuse large B-cell lymphoma in elderly patients is rare in Western populations. *Human pathology* 2010; 41: 352-357. <https://doi.org/10.1016/j.humpath.2009.07.024> PMid:19913281
 23. Rossi AP and Klein CL. Posttransplant malignancy. *Surgical Clinics* 2019; 99: 49-64. <https://doi.org/10.1016/j.suc.2018.09.004> PMid:30471741
 24. Alavizadeh SH and Hosseinzadeh H. Bioactivity assessment and toxicity of crocin: a comprehensive review. *Food and Chemical Toxicology* 2014; 64: 65-80. <https://doi.org/10.1016/j.fct.2013.11.016> PMid:24275090
 25. Moradzadeh M, Kalani MR and Avan A. The antileukemic effects of saffron (*Crocus sativus* L.) and its related molecular targets: A mini review. *Journal of cellular biochemistry* 2019; 120: 4732-4738. <https://doi.org/10.1002/jcb.27525> PMid:30644127
 26. Li X, Jiang C and Zhu W. Crocin reduces the inflammation response in rheumatoid arthritis. *Bioscience, Biotechnology, and Biochemistry* 2017; 81: 891-898. <https://doi.org/10.1080/09168451.2016.1263145> PMid:28388359
 27. Nam KN, Park Y-M, Jung H-J, Lee JY, Min BD, Park S-U, Jung W-S, Cho K-H, Park J-H and Kang I. Anti-inflammatory effects of crocin and crocetin in rat brain microglial cells. *European journal of pharmacology* 2010; 648: 110-116. <https://doi.org/10.1016/j.ejphar.2010.09.003> PMid:20854811
 28. Tamaddonfard E, Farshid A-A, Eghdami K, Samadi F and Erfanparast A. Comparison of the effects of crocin, saffranal and diclofenac on local inflammation and inflammatory pain responses induced by carrageenan in rats. *Pharmacological Reports* 2013; 65: 1272-1280. [https://doi.org/10.1016/S1734-1140\(13\)71485-3](https://doi.org/10.1016/S1734-1140(13)71485-3)
 29. Li K, Li Y, Ma Z and Zhao J. Crocin exerts anti-inflammatory and anti-catabolic effects on rat intervertebral discs by suppressing the activation of JNK. *International journal of molecular medicine* 2015; 36: 1291-1299. <https://doi.org/10.3892/ijmm.2015.2359> PMid:26648423 PMCID:PMC4601741
 30. Tarantilis PA, Morjani H, POLISSIOU M and MANFAIT M. Inhibition of growth and induction of differentiation of promyelocytic leukemia (HL-60) by carotinoids from *Crocus sativus* 1994; 1913-1918.
 31. Morjani H, Tarantilis P, Polissiou M and Manfait M. Growth inhibition and induction of erythroid differentiation activity by crocin, dimethylcrocetin and β -carotene on K562 tumor cells. *Anticancer Res* 1990; 10: 1398-1406.
 32. Sun Y, Wang Z, Wang L, Wang L, Zang C and Sun L. The Effect and Mechanisms of Proliferative Inhibition of Crocin on Human Leukaemia Jurkat Cells. *The West Indian medical journal* 2015; 64: 473. <https://doi.org/10.7727/wimj.2016.053> PMid:27398676 PMCID:PMC4961334
 33. Sun Y, Xu H-J, Zhao Y-X, Wang L-Z, Sun L-R, Wang Z and Sun X-F. Crocin exhibits antitumor effects on human leukemia HL-60 cells in vitro and in vivo. *Evidence-Based Complementary and Alternative Medicine* 2013; 2013: <https://doi.org/10.1155/2013/690164> PMid:23573146 PMCID:PMC3615578
 34. Hoshyar R, Bathaie SZ and Sadeghizadeh M. Crocin triggers the apoptosis through increasing the Bax/Bcl-2 ratio and caspase activation in human gastric adenocarcinoma, AGS, cells. *DNA and cell biology* 2013; 32: 50-57. <https://doi.org/10.1089/dna.2012.1866> PMid:23347444
 35. Luo Y, Cui S, Tang F, Shen C, Qi Y, Lu D, Ma L, Yang Y, Li Y and Chen R. The combination of crocin with cisplatin suppresses growth of gastric carcinoma cell line BGC-823 and promotes cell apoptosis. *Pakistan journal of pharmaceutical sciences* 2017; 30:
 36. Ray P, Guha D, Chakraborty J, Banerjee S, Adhikary A, Chakraborty S, Das T and Sa G. Crocetin exploits p53-induced death domain (PIDD) and FAS-associated death domain (FADD) proteins to induce apoptosis in colorectal cancer. *Scientific reports* 2016; 6: 1-11. <https://doi.org/10.1038/srep32979> PMid:27622714 PMCID:PMC5020693
 37. Beljebbar A, Sockalingum G, Morjani H, Angiboust J, Polissiou M and Manfait M. Differential Interaction Modes of Dimethylcrocetin in K562 and HL60 Tumor Cells As Probed by Near Infrared FT-Raman Microspectroscopy. In: editors. *Spectroscopy of Biological Molecules*. Springer; 1995. p. 475-476. https://doi.org/10.1007/978-94-011-0371-8_217
 38. Steele VE, Kelloff GJ, Wilkinson BP and Arnold JT. Inhibition of transformation in cultured rat tracheal epithelial cells by potential chemopreventive agents. *Cancer research* 1990; 50: 2068-2074.
 39. Kim B, Lee KY and Park B. Crocin suppresses constitutively active STAT3 through induction of protein tyrosine phosphatase SHP - 1. *Journal of cellular biochemistry* 2017; 118: 3290-3298. <https://doi.org/10.1002/jcb.25980> PMid:28295507
 40. Xu H-J, Zhong R, Zhao Y-X, Li X-R, Lu Y, Song A-Q, Pang X-Y, Yao R-Y and Sun L-R. Proliferative inhibition and apoptotic induction effects of crocin on human leukemia HL-60 cells and their mechanisms. *Zhongguo shi yan xue ye xue za zhi* 2010; 18: 887-892.
 41. Xu H-J, Zhang K-P, Zhong R, Zhao Y-X, Li X-R, Lu Y, Song A-Q, Pang X-Y and Sun L-R. Influence of crocin on proliferation in vitro and function of dendritic cells derived from bone marrow of children with acute leukemia. *Zhongguo shi yan xue ye xue za zhi* 2012; 20: 57-61.
 42. Zhang K, Zhong R, Xu H, ZHAO Y-x, LI X-r, LU Y, SONG A-q and SUN L-r. Effect of crocin on culture and proliferation of dendritic cells derived from children acute leukemia blood marrow in vitro. *Prog Mod Biomed* 2011; 24: 035.
 43. Molnar J, Szabo D, Pusztai R, Mucsi I, Berek L, Ocsovszki I, Kawata E and Shoyama Y. Membrane associated antitumor effects of crocine-, ginsenoside- and cannabinoid derivatives. *Anti-cancer research* 2000; 20: 861-867.
 44. Soleymani S, Zabihollahi R, Shahbazi S, Bolhassani A. Antiviral Effects of Saffron and its Major Ingredients. *Curr Drug Deliv*. 2018;15(5):698-704. doi: 10.2174/15672018146666171129210654. <https://doi.org/10.2174/15672018146666171129210654> PMid:29189153
 45. Bakshi HA, Sam S, Feroz A, Ravesh Z, Shah GA and Sharma M. Crocin from Kashmiri saffron (*Crocus sativus*) induces in vitro and in vivo xenograft growth inhibition of Dalton's lymphoma (DLA) in mice. *Asian Pac J Cancer Prev* 2009; 10: 887-890.
 46. Khavari A, Bolhassani A, Alizadeh F, Bathaie SZ, Balaram P, Agi E and Vahabpour R. Chemo-immunotherapy using saffron and its ingredients followed by E7-NT (gp96) DNA vaccine generates different anti-tumor effects against tumors expressing the E7 protein of human papillomavirus. *Archives of virology* 2015; 160: 499-508. <https://doi.org/10.1007/s00705-014-2250-9> PMid:25395243
 47. Rezaee R, Mahmoudi M, Abnous K, Zamani Taghizadeh Rabe S, Tabasi N, Hashemzaei M and Karimi G. Cytotoxic effects of crocin on MOLT-4 human leukemia cells. *Journal of Complementary and Integrative Medicine* 2013; 10: 105-112. <https://doi.org/10.1515/jcim-2013-0011> PMid:23934514
 48. Zhang K, Wang L, Si S, Sun Y, Pei W, Ming Y and Sun L. Crocin improves the proliferation and cytotoxic function of T cells in children with acute lymphoblastic leukemia. *Biomedicine & Pharmacotherapy* 2018; 99: 96-100. <https://doi.org/10.1016/j.biopha.2018.01.042> PMid:29329036
 49. Modagheh M-H, Shahabian M, Esmaceli H-A, Rajbai O and Hosseinzadeh H. Safety evaluation of saffron (*Crocus sativus*) tablets in healthy volunteers. *Phytomedicine* 2008; 15: 1032-1037. <https://doi.org/10.1016/j.phymed.2008.06.003> PMid:18693099
 50. Xi L, Qian Z, Du P and Fu J. Pharmacokinetic properties of crocin (crocetin digentiobiose ester) following oral administration in rats. *Phytomedicine* 2007; 14: 633-636. <https://doi.org/10.1016/j.phymed.2006.11.028> PMid:17215113
 51. Bostan HB, Mehri S and Hosseinzadeh H. Toxicology effects of saffron and its constituents: a review. *Iranian journal of basic medical sciences* 2017; 20: 110.