




# Characteristics of Induced-Sputum Inflammatory Phenotypes in Adults with Asthma: Predictors of Bronchial Eosinophilia

Astrid Crespo-Lessmann <sup>1-3,\*</sup>, Elena Curto <sup>1-3,\*</sup>, Eder Freddy Mateus Medina <sup>1-3</sup>, Esther Palones <sup>1-3</sup>, Alicia Belda Soler <sup>1-3</sup>, Soraya Sánchez Maza <sup>1-3</sup>, Lorena Soto-Retes <sup>1-3</sup>, Vicente Plaza <sup>1-3</sup>

<sup>1</sup>Servicio de Neumología y Alergia, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain; <sup>2</sup>Institut d'Investigació Biomèdica Sant Pau, Barcelona, Spain; <sup>3</sup>Departament de Medicina, Universitat Autònoma de Barcelona, Barcelona, Spain

\*These authors contributed equally to this work

Correspondence: Astrid Crespo-Lessmann, Servicio de Neumología y Alergia, Hospital de la Santa Creu i Sant Pau, Carrer Mas Casanovas 90, Barcelona, 08041, Spain, Tel +34-935565972, Fax +34 935565601, Email [acrespo@santpau.cat](mailto:acrespo@santpau.cat)

**Purpose:** The objectives of this study were, for patients attending a specialist asthma clinic at a tertiary care hospital, to determine, from sputum induction (SI), proportions of bronchial inflammatory phenotypes, demographic, clinical and functional characteristics of each phenotype, and the most accessible non-invasive inflammatory marker that best discriminates between phenotypes.

**Patients and Methods:** Included were 96 patients with asthma, attending a specialist asthma clinic at a tertiary care hospital, who underwent testing as follows: SI, spirometry, fractional exhaled nitric oxide (FeNO), blood eosinophilia, total immunoglobulin E (IgE), and a skin prick test.

**Results:** SI phenotypes were 46.9% eosinophilic, 33.3% paucigranulocytic, 15.6% neutrophilic, and 4.2% mixed. No significantly different clinical or functional characteristics were observed between the phenotypes. A positive correlation was observed between SI eosinophilia and both emergency visits in the last 12 months ( $p = 0.041$ ;  $r = 0.214$ ) and FeNO values ( $p = 0.000$ ;  $r = 0.368$ ). Blood eosinophilia correlated with SI eosinophilia ( $p = 0.001$ ;  $r = 0.362$ ) and was the best predictor of bronchial eosinophilia, followed by FeNO, and total blood IgE (area under the receiver operating characteristic curve (AUC-ROC) 72%, 65%, and 53%, respectively), although precision was only fair.

**Conclusion:** In consultations for severe asthma, the most frequent phenotype was eosinophilic. Peripheral blood eosinophilia is a reliable marker for discriminating between different bronchial inflammatory phenotypes, is useful in enabling doctors to select a suitable biologic treatment and so prevent asthma exacerbation, and is a better predictor of bronchial eosinophilia than FeNO and IgE values.

**Keywords:** asthma, sputum induction, phenotype, eosinophilia

## Introduction

Sputum induction (SI), the gold standard for evaluating bronchial inflammation in patients with asthma, is a non-invasive, standardized, and validated test<sup>1,2</sup> that distinguishes between 4 bronchial inflammatory phenotypes: eosinophilic, paucigranulocytic, neutrophilic, and mixed.<sup>3</sup> This technique, however, is not available in all hospitals as it requires trained personnel and a suitable infrastructure; therefore, other more accessible markers are used in current clinical practice, such as eosinophil count in peripheral blood and fractional exhaled nitric oxide (FeNO) measurement.

Especially important for severe asthma is phenotype identification, as it enables an individualized approach to treatment.<sup>4,5</sup> Several studies have confirmed that eosinophilic airway inflammation predicts response to anti-inflammatory treatment with both inhaled corticosteroids<sup>6,7</sup> and biologics.<sup>8-13</sup> Indeed, the main clinical guidelines for asthma management propose using SI to evaluate severe asthma<sup>14,15</sup> and to manage severe uncontrolled asthma treatment

if the patient is clinically followed up in suitably equipped centers.<sup>14</sup> However, the fact that SI is laborious and requires experienced personnel explains current attempts to identify non-invasive markers that discriminate between different bronchial inflammatory phenotypes in a simple and cost-effective way. While peripheral blood eosinophilia is a marker that predicts airway eosinophilia,<sup>16</sup> a common cut-off point has not been established, and correlations between blood and SI eosinophilia vary widely.<sup>16–18</sup> Studies have also reported a relationship between blood eosinophilia and the risk of severe exacerbations, decreased lung function, responsiveness to corticosteroid treatment, and predicted efficacy of some biologic treatments.<sup>10–13,19–21</sup> While FeNO is a non-invasive marker that reflects eosinophilic inflammation,<sup>22,23</sup> certain variables may modify its value,<sup>24,25</sup> and its correlation with bronchial and peripheral blood eosinophilia is also variable.<sup>26–29</sup>

Our study objectives, for a population of patients with asthma attending a specialist asthma clinic attached to a tertiary care hospital, were as follows: (a) to determine, from an SI test, the different proportions of bronchial inflammatory phenotypes and their demographic, clinical, and functional characteristics and (b) to identify an accessible non-invasive inflammatory marker used in routine clinical practice that discriminates between the different bronchial inflammatory phenotypes.

## Materials and Methods

Our cross-sectional descriptive study included 96 patients, aged 18–80 years old. Patients were consecutively enrolled from our tertiary care university hospital's severe asthma outpatient unit (located in Spain) for evaluation in 2018 and 2019. All the patients complied with asthma diagnostic criteria according to Global Initiative for Asthma (GINA) guidelines.<sup>15</sup> Excluded were smokers and patients who had experienced respiratory infections or required oral corticosteroids in the previous month.

Demographic, clinical, and functional data were collected for the 96 patients, and on the same day, the following procedures were carried out: skin prick test, total blood immunoglobulin E (IgE), absolute eosinophil count, SI inflammatory cell count, and forced expiratory volume in 1 second (FEV1), forced vital capacity (FVC), and FeNO measurements. Patients were also asked to complete the Asthma Control Test (ACT)<sup>30</sup> and Asthma Quality of Life Questionnaire (AQLQ).<sup>31</sup> The skin prick test, performed for common local aeroallergens according to the standard procedure,<sup>32</sup> was considered positive when papule diameter was >3 mm. Total blood IgE was determined using the enzyme-linked immunosorbent assay (ELISA) method (UNICAP, Pharmacia, Uppsala, Sweden) and was considered increased for values >160 IU/mL. Used to assess asthma control was the ACT, a self-assessment questionnaire validated in Spanish,<sup>30</sup> with good control considered to be >20 points. Quality of life (QoL) was assessed using the self-administered AQLQ, likewise validated in Spanish.<sup>32</sup> Spirometry measurements were made with a Datospir-600 device (Sibelmed SA, Barcelona, Spain) by an experienced technician and following the 2005 recommendations of the American Thoracic Society/European Respiratory Society (ATS/ERS);<sup>33</sup> FEV1 >80% was considered to be in the reference range of the theoretical value.<sup>34</sup> FeNO, following ATS/ERS 2005 recommendations<sup>35</sup> and using an electrochemical analyzer (NO Vario Analyzer, Filt Lungen- und Thoraxdiagnostik GmbH, Berlin, Germany), was measured at a flow of 50 mL/s and was considered to be significantly increased when values were >50 ppb.<sup>36</sup> Sputum samples were obtained and processed according to the method described by Djukanović et al,<sup>1</sup> and patients were classified according to cell counts as follows: eosinophilic if eosinophils  $\geq 3\%$ , neutrophilic if neutrophils  $\geq 61\%$ , paucigranulocytic if neutrophils <61% and eosinophils <3%, and mixed if neutrophils  $\geq 61\%$  and eosinophils  $\geq 3\%$ .<sup>6</sup> An absolute eosinophil count of  $\geq 300$  cells/ $\mu\text{L}$  was taken to define blood eosinophilia.<sup>14</sup>

## Ethical and Legal Aspects

The study complied with the principles of the Declaration of Helsinki (18th World Medical Assembly, 1964) and was approved by the Clinical Research Ethics Committee of Hospital Santa Creu i Sant Pau in Barcelona. The patients provided their written consent prior to participation in the study and all study data were anonymized.

## Statistical Analysis

Categorical variables were expressed as absolute and relative frequencies and quantitative variables as mean and standard deviation (SD) values. Between-group comparisons were analyzed using analysis of variance (ANOVA) for quantitative variables, and the chi-square or McNemar test for categorical variables, as appropriate. The area under the receiver operating characteristic curve (AUC-ROC) of the biomarkers used to detect the eosinophilic inflammatory phenotype was calculated using a combined impact model (general linear model; GLM), and Pearson's test was used for correlation analyses of the studied population.

The results were considered significant for  $p < 0.05$ . The analysis was performed with SPSS version 26.0 (SPSS Inc., Chicago, IL, USA).

## Results

### SI Inflammatory Phenotype Proportions and Characteristics

Of the 96 patients who underwent the SI test, almost half were eosinophilic ( $n = 45$ ; 46.9%), around a third were paucigranulocytic ( $n = 32$ ; 33.3%), and the remainder were neutrophilic ( $n = 15$ ; 15.6%) or mixed ( $n = 4$ ; 4.2%).

Demographic, clinical, and functional characteristics are summarized in Table 1. Overall mean age was 50 years. No significant differences were observed regarding sex, body mass index (BMI), asthma severity, disease control, emergency

**Table 1** Demographic, Clinical, and Functional Characteristics for 4 Inflammatory Phenotypes Identified by SI ( $n = 96$ )

	Eosinophilic (n=45)	Paucigranulocytic (n=32)	Neutrophilic (n=15)	Mixed (n=4)	p
Age, mean (SD) years	52.1 (14.7)	51.2 (14)	53.2 (17.9)	57 (26.2)	0.900
Women, %	71.1%	62.5%	46.7%	25%	0.143
Childhood asthma diagnosis, %	24.4%	21.9%	20%	25%	0.984
Severe persistent asthma, %	53.3%	34.4%	33.3%	50%	0.674
GINA 2021 asthma treatment steps, %	STEP 1–2: 6.7%	STEP 1–2: 12.9%	STEP 1–2: 15.9%	STEP 1–2: 50%	0.671
	STEP 3: 28.9%	STEP 3: 22.5%	STEP 3: 21%	STEP 3: 0%	
	STEP 4: 35.5%	STEP 4: 38.8%	STEP 4: 21%	STEP 4: 0%	
	STEP 5: 28.8%	STEP 5: 25.8%	STEP 5: 21%	STEP 5: 50%	
Poor asthma control (ACT <20), %	15.6%	12.5%	6.7%	50%	0.371
ED visits in previous 12 months, mean (SD)	1.4 (2.2)	1.1 (1.7)	1.3 (2.6)	3.5 (5.7)	0.291
AQLQ, mean (SD)	3.3 (2.3)	3.1 (3.2)	1.8 (2.1)	8.1	0.162
BMI, mean (SD) kg/m <sup>2</sup>	27.1 (4.2)	27.6 (5.6)	24.8 (3.6)	24.8 (5.7)	0.201
Nasal polyposis, %	24.4%	9.4%	13.3%	0%	0.250
Rhinitis, %	68.9%	68.8%	53.3%	25%	0.243
FEV1, mean (SD) %	83.7 (21)	103.9 (98.9)	80.9 (17.5)	77.5 (16)	0.433
BDT, %	26.7%	15.6%	33%	50%	0.327
Prick test +, %	68.9%	71.9%	60%	100%	0.476
FeNO, mean (SD) ppb	46.1 (37.2)	33.4 (26.5)	29.8 (27.4)	30.2 (12.9)	0.207

(Continued)

Table 1 (Continued).

	Eosinophilic (n=45)	Paucigranulocytic (n=32)	Neutrophilic (n=15)	Mixed (n=4)	p
Total IgE, mean (SD) IU/mL	460.4 (770.8)	276 (301.4)	241.9 (503)	239.2 (349.1)	0.454
Blood eosinophils, mean (SD) cells/ $\mu$ L	360 (300)	230 (100)	230 (100)	320 (800)	0.057 <sup>a</sup> 0.027 <sup>b</sup>
Eosinophils, mean (SD) %	5.4 (4)	3.5 (2.1)	3.5 (1.9)	3.7 (1)	0.041 <sup>a</sup> 0.021 <sup>b</sup>
Dose of inhaled corticosteroids, %	Medium 37.8% and high 28.9%	Medium 37.5% and high 28.1%	Medium 20% and high 40%	High 75%	0.587
Eosinophils, mean (SD) %	12.4 (12.3)	0.79 (0.7)	1.17 (0.9)	9.2 (5.8)	0.000
Neutrophils, mean (SD) %	38 (17.7)	36.6 (18)	76.2 (7.6)	72 (3.4)	0.000
Macrophages, mean (SD) %	48.35 (18.7)	59.8 (17.8)	19.8 (8.7)	17 (8.5)	0.000
Lymphocytes, mean (SD) %	0.89 (0.6)	0.93 (0.6)	0.91 (0.4)	1.58 (1.4)	0.207

Notes: <sup>a</sup>Significance comparing the 4 groups. <sup>b</sup>Significance comparing eosinophilic, paucigranulocytic, and neutrophilic phenotype groups.

Abbreviations: ACT, asthma control test; AQLQ, Asthma Quality of Life Questionnaire; BDT, bronchodilator test; BMI, body mass index; ED, emergency department; FeNO, fractional exhaled nitric oxide; FEV1, forced expiratory volume in 1 second; GINA, Global Initiative for Asthma; IgE, immunoglobulin E; SD, standard deviation; SI, sputum induction.

department (ED) visits in the past 12 months, QoL as measured by AQLQ, bronchial obstruction (FEV1), associated rhinitis or nasal polyposis, total blood IgE, FeNO, or inhaled corticosteroid dose. In contrast, significant differences were observed between the different phenotypes in peripheral blood eosinophil percentages and counts, which were significantly higher in the eosinophilic group.

## Asthma Severity

In the analysis by asthma severity, the predominant inflammatory phenotypes were as follows: paucigranulocytic with intermittent asthma, 46.6%; eosinophilic with mild persistent asthma, 45%; eosinophilic with moderate persistent asthma, 36%; and eosinophilic with severe persistent asthma, 57.14% ( $p = 0.674$ ).

## Variable Correlations for the Studied Population


For the 96 patients, positive correlations were observed between SI eosinophilia and ED visits in the previous 12 months ( $p = 0.041$ ;  $r = 0.214$ ), between SI eosinophilia and FeNO values ( $p = 0.000$ ;  $r = 0.368$ ), and between SI eosinophilia and peripheral blood eosinophilia ( $p = 0.001$ ;  $r = 0.362$ ). We interpreted discriminatory capacity as follows (see Figure 1):  $r = 0.5$ , equivalent to a coin toss;  $r = 0.5-0.6$ , poor;  $r = 0.6-0.75$ , fair;  $r = 0.75-0.9$ , good; and  $r = 0.9-0.97$ , very good.

## SI Eosinophilia $\geq 3\%$ Detection in the Combined Model

The AUC-ROC values that detected SI eosinophilia  $\geq 3\%$  were as follows: absolute blood eosinophils (EOS), 72% ( $p = 0.000$ ); FeNO, 65% ( $p = 0.014$ ); and total blood IgE, 53% ( $p = 0.590$ ).

## Discussion

In patients attending our specialist asthma clinic at a tertiary care hospital, the eosinophilic bronchial inflammatory phenotype was predominant, and there were no significant differences in clinical and functional characteristics for the various bronchial inflammatory phenotypes. Furthermore, our study supports the following findings: (a) peripheral blood eosinophilia is a marker that enables the eosinophilic inflammatory phenotype to be differentiated from other bronchial inflammatory phenotypes, although note that precision is only fair according to the AUC-ROC; and (b) while positive



**Figure 1** AUC-ROC plot from EOS, FeNO, and total IgE values in a combined model (n=96).

**Abbreviations:** AUC-ROC, area under the receiver operating characteristic curve; EOS, absolute eosinophils; FeNO, fractional exhaled nitric oxide; IgE, immunoglobulin E.

correlations exist between SI eosinophilia and both FeNO and blood eosinophil values, they were not strong, suggesting a possible activation of various inflammatory pathways in patients with asthma. This would point to the need for a more comprehensive approach to asthma management that goes beyond mere biomarker threshold positivity.

Our AUC-ROC value for detecting SI eosinophilia  $\geq 3\%$  was slightly higher than documented in the literature. Hastie et al<sup>39</sup> reported a value of 69% for detection of SI eosinophilia  $\geq 2\%$  and concluded that while there was an association between blood eosinophilia and SI eosinophilia, precision in terms of correct classification of patients with and without an eosinophilic phenotype was poor, and, furthermore, that this poor precision persisted when biomarkers such as FeNO and IgE were also considered.

The usefulness of blood eosinophilia is supported by several studies. Wagener et al<sup>16</sup> showed that blood eosinophilia was an accurate biomarker of eosinophilic airway inflammation in 2 independent cohorts of patients with asthma (AUC 89%;  $p < 0.001$ ; sensitivity 78% and specificity 91%), while Schleich et al<sup>3</sup> reported that blood eosinophilia  $\geq 220/\text{mm}^3$  enabled SI eosinophilia  $\geq 3\%$  to be detected with 77% sensitivity and 70% specificity (AUC 0.79,  $p < 0.0001$ ).

There is no consensus regarding the blood eosinophilia cutoff value to define the eosinophilic phenotype, and, although current evidence points to 150–300 cells/ $\mu\text{L}$ , that range of values is still a matter of debate.<sup>16,18</sup> Note that, in patients with severe asthma, part of the variability reported in different studies is explained by possible variations in blood and SI eosinophilia depending on doses of inhaled or systemic corticosteroids.<sup>4,37</sup> Variability may also result from factors such as smoking (OR = 6.44;  $p = 0.013$ ) and having had a recent asthma exacerbation (OR = 5.84;  $p = 0.022$ ).<sup>38</sup>

While we found positive correlations between SI eosinophilia and both ED visits in the previous 12 months ( $p = 0.041$ ;  $r = 0.214$ ) and FeNO values ( $p = 0.000$ ;  $r = 0.368$ ) and between SI eosinophilia and peripheral blood eosinophilia ( $p = 0.001$ ;  $r = 0.362$ ), none of those correlations were sizeable, indicating no linear relationship between the variables. Another study found better correlation for patients with asthma when the comparison was based on  $\geq 300$  cells/ $\mu\text{L}$  in peripheral blood and SI eosinophilia  $\geq 2\%$  ( $p = 0.0002$ ;  $r = 0.5235$ ).<sup>40</sup>

The growing importance attached to blood eosinophilia is because it is the most relevant marker for both the choice of, and response to, biologic treatments for severe uncontrolled eosinophilic asthma. The fact that the vast majority of studies use peripheral blood eosinophilia and not SI eosinophilia as the biomarker of choice for a biologic is because not all hospitals have the facilities necessary for SI cell counting.<sup>9,41</sup>

Peripheral blood eosinophilia has been demonstrated to be a marker of a better response to biologics. For instance, it is the key biomarker for measuring response to mepolizumab, a monoclonal antibody against interleukin-5 (IL-5), with exacerbations greatly reduced (73%) in patients with blood eosinophils  $\geq 500/\mu\text{L}$ .<sup>9,41</sup> Efficacy of another intravenously administered antibody against IL-5, reslizumab, has also been demonstrated for patients with blood eosinophilia  $\geq 400/$

$\mu\text{L}$ .<sup>42</sup> Finally, benralizumab, an antibody that acts against the IL-5 receptor through apoptosis of eosinophils and basophils, has been shown to reduce exacerbation rates in patients with blood eosinophilia  $\geq 300/\mu\text{L}$ .<sup>9,20</sup>

Blood eosinophilia can also predict response to treatment with both inhaled and systemic corticosteroids<sup>14</sup> and can help adjust oral corticosteroid dosage for patients with severe asthma, as demonstrated by Wark et al,<sup>43</sup> who reported that blood eosinophilia maintained at  $< 200$  cells/ $\mu\text{L}$  prevented exacerbations, improved asthma control, and enabled lower oral corticosteroid doses.

In our series, the FeNO value was less correlated with SI eosinophilia (AUC-ROC 65%) than in the review by Korevaar et al,<sup>44</sup> which included 12 studies with a combined sensitivity of 66% and specificity of 76% in detecting SI eosinophilia  $\geq 3\%$  (AUC 0.74; 95% confidence interval (CI): 0.70–0.78). This difference is possibly explained by the multiple factors that modify FeNO values, such as allergic rhinitis, upper and lower respiratory tract viral infections, age, tobacco use, and atopy;<sup>25–28</sup> it may also be due to the higher sensitivity but lower specificity of FeNO, resulting in high negative predictive values but low positive predictive values for eosinophilic inflammation.<sup>45</sup>

However, it is important to understand the complexity of inflammatory mechanisms in type 2 inflammation in different patients, so it is recommended to simultaneously measure several biomarkers (EOS, IgE, FeNO) to identify potential targets for treatment with biologics. A post-hoc analysis of the QUEST study<sup>46</sup> that assessed dupilumab efficacy by biomarker subgroups, as defined by GINA,<sup>15</sup> found that reference blood eosinophilia and FeNO levels clearly pointed to similar disease severity at the outset in all subpopulations (EOS  $\geq 150$  cells/ $\mu\text{L}$ , FeNO  $\geq 20$  ppb, and both EOS  $\geq 150$  cells/ $\mu\text{L}$  and FeNO  $\geq 20$  ppb).<sup>47</sup> In a study of 110 patients published by our working group, we reported a dissociation between increased FeNO ( $\geq 50$  ppb) and SI eosinophilia in 42% of patients; that study identified 2 groups with discordant values: a younger group mainly associated with a paucigranulocytic phenotype and atopy, with high FeNO and no SI eosinophilia, and with better FEV1, and an older group mainly associated with a non-allergic eosinophilic phenotype, with low FeNO and high SI eosinophilia, and accounting for more ED visits in the previous 12 months.<sup>48</sup> Those data support the existence of different activation patterns in underlying inflammatory pathways in patients with asthma, suggesting the need for a more comprehensive and more personalized approach to management that goes beyond mere biomarker threshold positivity.

Use of IgE as a biomarker of eosinophilia is poorly supported. In our study, total IgE was the weakest predictor of SI eosinophilia, corroborating other studies<sup>45–47</sup> reporting low sensitivity, specificity, and AUC values for IgE. Demarche et al,<sup>49</sup> in particular, indicated that IgE alone does not adequately predict SI eosinophilic status. Westerhof et al<sup>17</sup> proposed joint use of blood eosinophilia and FeNO to improve airway predictions of eosinophilia (AUC 0.87;  $p = 0.027$ ), while Demarche et al<sup>49</sup> proposed the joint use of blood eosinophilia, FeNO, and IgE, as an approach that, in their study, identified 58% of patients with a high or low probability of having SI eosinophilia  $\geq 3\%$ , and that correctly classified 87% of those patients.

Regarding the distribution of inflammatory phenotypes in our study, the eosinophilic phenotype predominated (almost half), followed by the paucigranulocytic phenotype (around a third); this finding corroborates another large series,<sup>3</sup> but contradicts other studies that reported predominance of the paucigranulocytic phenotype.<sup>50</sup> The difference is possibly explained by the fact that the studies in which the paucigranulocytic phenotype predominated were of patients whose asthma was less severe than that of patients recruited in specialist asthma clinics.

We found no significant differences in clinical and functional characteristics between the bronchial inflammatory phenotypes. This finding differs from that of Schleich et al,<sup>3</sup> who reported that the eosinophilic phenotype was associated with atopy, bronchial hyperresponsiveness, poorer control, a reduced FEV1/FVC ratio, increased FeNO and IgE values, and nasal polyposis.<sup>51</sup> The difference may reflect sample size: 96 in our series compared to 508 in the study by Schleich et al.<sup>3</sup> Note that there is probably a significant overlap in biomarker positivity in patients with asthma<sup>52</sup> that may suggest no differentiating characteristics according to inflammatory phenotype. This issue needs to be addressed through more studies, as relevant pathogenic knowledge is required for an era of biologic monoclonals and more personalized medicine.

Another result to highlight from our study was the positive correlation between SI eosinophilia and ED visits in the previous 12 months, possibly comparable to the poorer asthma control of the eosinophilic phenotype reported by



Schleich et al,<sup>3</sup> and in line with the established fact that eosinophilia is a predictor of exacerbations in patients with asthma.<sup>52</sup>

As limitations, our cross-sectional descriptive study may incur possible selection bias; all our patients were required to have undergone specific testing for inclusion, and were patients with predominantly moderate-severe persistent asthma attending a specialist clinic in a tertiary care hospital. Another limitation is the small sample size compared to other studies (for instance, those by Schleich et al<sup>3</sup> and Abdo et al<sup>53</sup>), and, within our sample, the fact that the patients with a mixed inflammatory phenotype were so few that we were unable to characterize them; note, however, that the mixed phenotype prevalence rate in our study reflects that of other studies.<sup>3</sup> A strength of our study is that inflammatory phenotypes were identified on the basis of SI, and comparisons were possible with biomarkers used in typical asthma consultations, such as peripheral eosinophilia and FeNO.

The main conclusions of our study of 96 patients with asthma attending our specialist asthma clinic at a tertiary care hospital are as follows: (a) the predominant bronchial inflammatory phenotype was eosinophilic, and there were no significant differences in clinical and functional characteristics between the 4 different bronchial inflammatory phenotypes; and (b) peripheral eosinophilia detected SI eosinophilia  $\geq 3\%$  with greater diagnostic accuracy than markers such as FeNO and total IgE and was also the only marker that distinguished the eosinophilic phenotype from the other inflammatory phenotypes.

While the SI cell count is the gold standard for non-invasive evaluation of bronchial inflammation in patients with severe asthma and a useful test to guide the choice of biologic treatment, our study would suggest peripheral blood eosinophilia as an alternative when this test is not available, given that, as a good marker for detection of the eosinophilic inflammatory phenotype, it can be potentially useful for doctors to select a suitable biologic treatment and prevent asthma exacerbations.

## Abbreviations

ACT, Asthma Control Test; ANOVA, analysis of variance; ATS, American Thoracic Society; AQLQ, Asthma Quality of Life Questionnaire; AUC, area under the curve; BMI, body mass index; CI, confidence interval; ED, emergency department; ELISA, enzyme-linked immunosorbent assay; EOS, absolute eosinophils; ERS, European Respiratory Society; FeNO, fractional exhaled nitric oxide; FEV1, forced expiratory volume in 1 second; FVC, forced vital capacity; GEMA, Spanish Asthma Management Guidelines; GINA, Global Initiative for Asthma; IgE, immunoglobulin E; IL-5, interleukin-5; QoL, quality of life; ROC, receiver operating characteristic; SD, standard deviation; SI, sputum induction.

## Acknowledgments

The authors would like to thank Ailish M J Maher for translating and reviewing the article.

## Disclosure

AC-L has received fees in the last 3 years for talks at meetings sponsored by AstraZeneca, Bial, Boehringer Ingelheim, Chiesi, Ferrer, GlaxoSmithKline, MSD, Novartis, Orion Pharma, and Sanofi, has received travel and attendance expenses for conferences from Bial, Gebro, GlaxoSmithKline, Novartis, and TEVA, and has received funds/grants for research projects from several state agencies, non-profit foundations, and AstraZeneca and GlaxoSmithKline. EC reports non-financial support from ALK, AstraZeneca, Novartis, and Menarini, personal fees from Boehringer-Ingelheim and TEVA, and personal fees/non-financial support from Chiesi outside the submitted work. This paper is part of the doctoral thesis of EC. EP has received travel and attendance expenses for conferences from Gebro Pharma, Chiesi, FAES Farma, Rovi, GlaxoSmithKline, and Sanofi, and has received funds/grants for research projects from state agencies, non-profit foundations, and Alpha Bioresearch. LS-R has received fees in the last 3 years for talks at meetings sponsored by AstraZeneca, Diater, Chiesi, and GlaxoSmithKline, has received travel and attendance expenses for conferences from Sanofi, Allergy-Therapeutics, Hal Allergy, and FAES Farma, has acted as a consultant for Sanofi, Stallergenes-Greer, GlaxoSmithKline, and AstraZeneca, and has received funds/grants for research projects from Spanish Allergy and Clinical Immunology Society (SEAIC), a non-profit foundation. VP has received fees in the last 3 years for talks at meetings sponsored by AstraZeneca, Boehringer-Ingelheim, Merck Sharp & Dohme, and Chiesi, has received travel and

attendance expenses for conferences from AstraZeneca, Chiesi, and Novartis, has acted as a consultant for ALK, AstraZeneca, Boehringer, Merck Sharp & Dohme, MundiPharma, and Sanofi, and has received funds/grants for research projects from several state agencies, non-profit foundations, and AstraZeneca, Chiesi and Menarini. EFMM and ABS and SSM declare no conflicts of interest in this work.

## References

- Djukanovic R, Sterk PJ, Fahy JV, Hargreave FE. Standardized methodology of sputum induction and processing. *Eur Respir J*. 2002;20 (Supplement 37):1–52, 4. doi:10.1183/09031936.02.00000102
- Spanevello A, Confalonieri M, Sulotto F, et al. Induced sputum cellularity: reference values and distribution in normal volunteers. *Am J Respir Crit Care Med*. 2000;162(3):1172–1174. doi:10.1164/ajrccm.162.3.9908057
- Schleich FN, Manise M, Sele J, Henket M, Seidel L, Louis R. Distribution of sputum cellular phenotype in a large asthma cohort: predicting factors for eosinophilic vs neutrophilic inflammation. *BMC Pulm Med*. 2013;13:11. doi:10.1186/1471-2466-13-11
- Robinson D, Humbert M, Buhl R, et al. Revisiting type 2-high and type 2-low airway inflammation in asthma: current knowledge and therapeutic implications. *Clin Exp Allergy*. 2017;47:1365–2222. doi:10.1111/cea.12880
- Aziz-Ur-Rehman A, Dasgupta A, Kjarsgaard M, Hargreave FE, Nair P. Sputum cell counts to manage prednisone-dependent asthma: effects on FEV1 and eosinophilic exacerbations. *Allergy Asthma Clin Immunol*. 2017;13(1):17. doi:10.1186/s13223-017-0190-0
- Green RH, Brightling CE, Woltmann G, Parker D, Wardlaw AJ, Pavord ID. Analysis of induced sputum in adults with asthma: identification of subgroup with isolated sputum neutrophilia and poor response to inhaled corticosteroids. *Thorax*. 2002;57(10):875–879.
- Park SY, Kang SY, Song WJ, Kim JH. Evolving concept of severe asthma: transition from diagnosis to treatable traits. *Allergy Asthma Immunol Res*. 2022;14(5):447–464. doi:10.4168/aaair.2022.14.5.447
- Haldar P, Brightling CE, Hargadon B, et al. Mepolizumab and exacerbations of refractory eosinophilic asthma. *N Engl J Med*. 2009;360 (10):973–984. doi:10.1056/NEJMoa0808991
- Holguin F, Cardet JC, Chung KF, et al. Management of severe asthma: a European respiratory society/American thoracic society guideline. *Eur Respir J*. 2020;55(1):1900588. doi:10.1183/13993003.00588-2019
- Bleecker ER, FitzGerald JM, Chanez P, et al. Efficacy and safety of benralizumab for patients with severe asthma uncontrolled with high-dosage inhaled corticosteroids and long-acting  $\beta_2$ -agonists (SIROCCO): a randomised, multicentre, placebo-controlled Phase 3 trial. *Lancet*. 2016;388 (10056):2115–2127. doi:10.1016/S0140-6736(16)31324-1
- FitzGerald JM, Bleecker ER, Nair P, et al. Benralizumab, an anti-interleukin-5 receptor  $\alpha$  monoclonal antibody, as add-on treatment for patients with severe, uncontrolled, eosinophilic asthma (CALIMA): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet*. 2016;388 (10056):2128–2141. doi:10.1016/S0140-6736(16)31322-8
- Domingo RC, Carrillo DT, Blanco AM, Martínez ME, Banas CD, Sánchez HMG; REDES Study Group. Correction to: rEal world effectiveness and safety of mepolizumab in a multicentric Spanish cohort of asthma patients stratified by Eosinophils: the REDES study. *Drugs*. 2021;81 (16):1949–1951. doi:10.1007/s40265-021-01622-x
- Ortega HG, Yancey SW, Mayer B, et al. Severe eosinophilic asthma treated with mepolizumab stratified by baseline eosinophil thresholds: a secondary analysis of the DREAM and MENSA studies. *Lancet Respir Med*. 2016;4(7):549–556. doi:10.1016/S2213-2600(16)30031-5
- GEMA. Guía Española para el Manejo del Asma (GEMA 5.2). Available from: <http://gemasma.com>. Accessed January 9, 2023.
- Global Initiative for Asthma (GINA). Global strategy for asthma management and prevention; 2021. Available from: <https://ginasthma.org/>. Accessed January 9, 2023.
- Wagener AH, de Nijs SB, Lutter R, et al. External validation of blood eosinophils, FeNO and serum periostin as surrogates for sputum eosinophils in asthma. *Thorax*. 2015;70:115–120. doi:10.1136/thoraxjnl-2014-205634
- Westerhof GA, Korevaar DA, Amelink M, et al. Biomarkers to identify sputum eosinophilia in different adult asthma phenotypes. *Eur Respir J*. 2015;46(3):688–696. doi:10.1183/09031936.00012415
- Heffler E, Terranova G, Chessari C, et al. Point-of-care blood eosinophil count in a severe asthma clinic setting. *Ann Allergy Asthma Immunol*. 2017;119(1):16–20. doi:10.1016/j.anaai.2017.05.016
- Sinz H, Renz H, Skevaki C. Cellular and noncellular bloodborne biomarkers in asthma. *Ann Allergy Asthma Immunol*. 2017;118(6):672–679. doi:10.1016/j.anaai.2017.04.016
- Menzella F, Biava M, Bagnasco D, et al. Efficacy and steroid-sparing effect of benralizumab: has it an advantage over its competitors? *Drugs in Context*. 2019;8:212580. doi:10.7573/dic.212580
- Tiotiu A. Biomarkers in asthma: state of the art. *Asthma Res Pract*. 2018;4(1):10. doi:10.1186/s40733-018-0047-4
- Lex C, Ferreira F, Zacharasiewicz A, et al. Airway eosinophilia in children with severe asthma: predictive values of non-invasive tests. *Am J Respir Crit Care Med*. 2006;174(12):1286–1291. doi:10.1164/rccm.200603-352OC
- Lim S, Jatakanon A, Meah S, Oates T, Chung KF, Barnes PJ. Relationship between exhaled nitric oxide and mucosal eosinophilic inflammation in mild to moderately severe asthma. *Thorax*. 2000;55(3):184–188. doi:10.1136/thorax.55.3.184
- Kharitonov SA, Robbins RA, Yates D, Keatings V, Barnes PJ. Acute and chronic effects of cigarette smoking on exhaled nitric oxide. *Am J Respir Crit Care Med*. 1995;152(2):609–612. doi:10.1164/ajrccm.152.2.7543345
- Olin AC, Rosengren A, Thelle DS, Lissner L, Bake B, Torén K. Height, age, and atopy are associated with fraction of exhaled nitric oxide in a large adult general population sample. *Chest*. 2006;130(5):1319–1325. doi:10.1378/chest.130.5.1319
- Strunk RC, Szeffler SJ, Phillips BR, et al. Relationship of exhaled nitric oxide to clinical and inflammatory markers of persistent asthma in children. *J Allergy Clin Immunol*. 2003;112(5):883–892. doi:10.1016/j.jaci.2003.08.014
- Jatakanon A, Lim S, Kharitonov SA, Chung KF, Barnes PJ. Correlation between exhaled nitric oxide, sputum eosinophils, and methacholine responsiveness in patients with mild asthma. *Thorax*. 1998;53(2):91–95. doi:10.1136/thx.53.2.91
- Jones SL, Kittelson J, Cowan JO, et al. The predictive value of exhaled nitric oxide measurements in assessing changes in asthma control. *Am J Respir Crit Care Med*. 2001;164(5):738–743. doi:10.1164/ajrccm.164.5.2012125



29. Berry MA, Shaw DE, Green RH, et al. The use of exhaled nitric oxide concentration to identify eosinophilic airway inflammation: an observational study in adults with asthma. *Clin Exp Allergy*. 2005;35(9):1175–1179. doi:10.1111/j.1365-2222.2005.02314.x
30. Vega JM, Badia X, Badiola C, et al. Validation of the Spanish version of the asthma control test (ACT). *J Asthma*. 2007;44(10):867–872. doi:10.1080/02770900701752615
31. Perpiñá M, Belloch A, Pascual LM, de Diego A, Compte L. The quality of life in asthma: an evaluation of the AQLQ questionnaire for its use on a Spanish population. Asthma quality of life questionnaire. *Arch Bronconeumol*. 1995;31(5):211–218. doi:10.1016/S0300-2896(15)30926-1
32. Bernstein IL, Li JT, Bernstein DI, et al. Allergy diagnostic testing: an updated practice parameter. *Ann Allergy Asthma Immunol*. 2008;100(3 Suppl 3):S1–148. doi:10.1016/s1081-1206(10)60305-5
33. Miller MR, Crapo R, Hankinson J, et al; ATS/ERS Task Force. General considerations for lung function testing. *Eur Respir J*. 2005;1:153–161. doi:10.1183/09031936.05.00034505
34. Roca J, Sanchis J, Agusti-Vidal A, et al. Spirometric reference values from a Mediterranean population. *Bull Eur Physiopathol Respir*. 1986;22(3):217–224.
35. American Thoracic Society/European Respiratory Society (ATS/ERS). ATS/ERS recommendations for standardized procedures for the online and offline measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide, 2005. *Am J Respir Crit Care Med*. 2005;171(8):912–930. doi:10.1164/rccm.200406-710ST
36. Dweik RA, Boggs PB, Erzurum SC, et al. American thoracic society committee on interpretation of exhaled nitric oxide levels (FeNO) for clinical applications. An official ATS clinical practice guideline: interpretation of exhaled nitric oxide levels (FeNO) for clinical applications. *Am J Respir Crit Care Med*. 2011;5:602–615. doi:10.1164/rccm.9120-11ST
37. Yancey SW, Keene ON, Albers FC, et al. Biomarkers for severe eosinophilic asthma. *J Allergy Clin Immunol*. 2017;140(6):1509–1518. doi:10.1016/j.jaci.2017.10.005
38. Suárez-Cuartín G, Crespo A, Mateus E, et al. Variability in asthma inflammatory phenotypes in induced sputum. Frequency and causes. *Arch Bronconeumol*. 2016;52(2):76–81. doi:10.1016/j.arbres.2015.03.007
39. Hastie AT, Moore WC, Li H, et al; National Heart, Lung, and Blood Institute's Severe Asthma Research Program. Biomarker surrogates do not accurately predict sputum eosinophil and neutrophil percentages in asthmatic subjects. *J Allergy Clin Immunol*. 2013;132(1):72–80. doi:10.1016/j.jaci.2013.03.044
40. Rial MJ, Álvarez-puebla MJ, Arismendi E, et al. Clinical and inflammatory characteristics of patients with asthma in the Spanish MEGA project cohort. *Clin Transl Allergy*. 2021;11:e12001. doi:10.1002/ct2.12001
41. Pavord ID, Korn S, Howarth P, et al. Mepolizumab for severe eosinophilic asthma (DREAM): a multicentre, double-blind, placebo-controlled trial. *Lancet*. 2012;380(9842):651–659. doi:10.1016/S0140-6736(12)60988-X
42. Brussino L, Heffler E, Bucca C, Nicola S, Rolla G. Eosinophils target therapy for severe asthma: critical points. *Biomed Res Int*. 2018;2018:7582057. doi:10.1155/2018/7582057
43. Wark PA, McDonald VM, Gibson PG. Adjusting prednisone using blood eosinophils reduces exacerbations and improves asthma control in difficult patients with asthma. *Respirology*. 2015;20(8):1282–1284. doi:10.1111/resp.12602
44. Korevaar DA, Westerhof GA, Wang J, et al. Diagnostic accuracy of minimally invasive markers for detection of airway eosinophilia in asthma: a systematic review and meta-analysis. *Lancet Respir Med*. 2015;3(4):290–300. doi:10.1016/S2213-2600(15)00050-8
45. Taylor DR. Advances in the clinical applications of exhaled nitric oxide measurements. *J Breath Res*. 2012;6(4):047102. doi:10.1088/1752-7155/6/4/047102
46. Pelaia C, Vatrella A, Gallelli L, et al. Dupilumab for the treatment of asthma. *Expert Opin Biol Ther*. 2017;17(12):1565–1572. doi:10.1080/14712598.2017.1387245
47. Busse WW, Maspero JF, Rabe KF, et al. Liberty asthma QUEST: phase 3 randomized, double-blind, placebo-controlled, parallel-group study to evaluate dupilumab efficacy/safety in patients with uncontrolled, moderate-to-severe asthma. *Adv Ther*. 2018;35(5):737–748. doi:10.1007/s12325-018-0702-4
48. Crespo A, Giner J, Torrejón M, et al. Clinical and inflammatory features of asthma with dissociation between fractional exhaled nitric oxide and eosinophils in induced sputum. *J Asthma*. 2016;53(5):459–464. doi:10.3109/02770903.2015.1116086
49. Demarche SF, Schleich FN, Paulus VA, Henket MA, Van Hees TJ, Louis RE. Is it possible to claim or refute sputum eosinophils  $\geq 3\%$  in asthmatics with sufficient accuracy using biomarkers? *Respir Res*. 2017;18(1):133. doi:10.1186/s12931-017-0615-9
50. Ntontsi P, Loukides S, Bakakos P, et al. Clinical, functional and inflammatory characteristics in patients with paucigranulocytic stable asthma: comparison with different sputum phenotypes. *Allergy*. 2017;72(11):1761–1767.
51. Laidlaw T, Mullol J, Woessner K, Amin N, Mannent L. Chronic rhinosinusitis with nasal polyps and asthma. *J Allergy Clin Immunol Pract*. 2021;9(3):1133–1141. doi:10.1016/j.jaip.2020.09.063
52. Denton E, Price D, Tran T, et al. Cluster analysis of inflammatory biomarker expression in the international severe asthma registry. *J Allergy Clin Immunol Pract*. 2021;9(7):2680–2688. doi:10.1016/j.jaip.2021.02.059
53. Abdo M, Pedersen F, Kirsten A-M, et al. Longitudinal impact of sputum inflammatory phenotypes on small airway dysfunction and disease outcomes in asthma. *J Allergy Clin Immunol Pract*. 2022;10(6):1545–1553.