

Complete Genome Sequence of the *Pseudomonas fluorescens* Bacteriophage UFV-P2

Monique R. Eller,^a Rafael L. Salgado,^b Pedro M. P. Vidigal,^c Maura P. Alves,^d Roberto S. Dias,^a Leandro L. de Oliveira,^a Cynthia C. da Silva,^a Antônio F. de Carvalho,^d Sérgio O. De Paula^a

Laboratory of Molecular Immunovirology, Department of General Biology,^a Department of Biochemistry and Molecular Biology,^b Bioinformatics Laboratory, Institute of Applied Biotechnology to Agriculture (BIOAGRO),^c and Department of Food Technology,^d Federal University of Viçosa, Minas Gerais, Brazil

Milk proteolysis caused by *Pseudomonas fluorescens* is a serious problem in the dairy industries as a result of its ability to grow under refrigeration. The use of phages to control contaminants in food has been considered an alternative to traditional methods; therefore, a thorough understanding of such organisms is vital for their use. In this study, we show the complete genome sequence and analysis of a *P. fluorescens* phage isolated from wastewater of a dairy industry in Brazil.

Received 2 October 2012 Accepted 18 October 2012 Published 15 January 2013

Citation Eller MR, Salgado RL, Vidigal PMP, Alves MP, Dias RS, de Oliveira LL, da Silva CC, de Carvalho AF, De Paula SO. 2013. Complete genome sequence of the *Pseudomonas fluorescens* bacteriophage UFV-P2. *Genome Announc.* 1(1):e00006-12. doi:10.1128/genomeA.00006-12.

Copyright © 2013 Eller et al. This is an open-access article distributed under the terms of the [Attribution 3.0 Unported Creative Commons License](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Sérgio O. De Paula, depaula@ufv.br.

Pseudomonas fluorescens is the major microorganism associated with milk deterioration by proteolysis (1–3). This bacterium produces heat-stable proteases and lipases responsible for the gelation of ultra-high-temperature (UHT) milk and destabilization of casein micelles and is associated with off-flavors and yield loss of dairy products (2–5). Because the use of antibiotics and other broad-range antimicrobial agents can be discouraged as a result of the need to maintain milk indigenous microbiota, the control of milk proteolysis using specific bacteriophages has been suggested as a strategic alternative (6, 7). However, its use in food would be considered only after a thorough examination to ensure its safety and effectiveness. Therefore, it was essential to determine the complete genome sequence of phage UFV-P2, a *P. fluorescens* phage with a high ability to reduce casein proteolysis in milk.

Phage UFV-P2 was isolated and purified from the wastewater of a dairy industry in Brazil, and its genomic DNA was extracted and sequenced using an Illumina Genome Analyzer II by CD Genomics (New York, NY). The viral genome was assembled and analyzed using CLC Genomics Workbench version 5.1 (CLC bio). The reads were assembled in contigs that considered more stringent parameters, in which 90% of each read had to cover the other read with 90% identity. This assembly produced the UFV-P2 genome sequence with coverage of 30,655-fold. Approximately 92 open reading frames (ORFs) were predicted using the Bacterial Genetic Code (NCBI translation Table 11) and alternative start codons (AUG, CUG, and UUG). All predicted ORFs were functionally annotated using BlastX searches against GenBank (<http://www.ncbi.nlm.nih.gov/genbank>) and UniProt (<http://www.uniprot.org>) databases. Only 41 ORFs (44.57%) presented significant similarities to known proteins and were considered in genome annotation. Additionally, the presence of tRNA genes was predicted using tRNAscan-SE version 1.21 (8).

The phage UFV-P2 has a linear 45,517-bp DNA genome with no tRNA genes, a GC content of 51.5%, and 41 ORFs (19 positive- and 22 negative-stranded), representing a gene density of 0.9008/

kb. The ORFs analyzed were annotated to five different protein groups, one of them containing 14 hypothetical proteins with unknown function (34.1%). The remaining groups consisted of one chaperone and four constitutive and seven structural protein genes, including a major head, a portal, and a hypothetical tail collar domain. Finally, there were 15 ORF hits (36.6%) with genes encoding enzymes, including one lysozyme, the terminase small and large subunits, an exonuclease, an endonuclease, a primase/helicase and two parts of the DNA polymerase. The bioinformatics analyses showed 53.61% identity to the genome of the temperate *Pseudomonas* phage PaP3 (9). Knowledge of this group of phages is still limited, and further analyses are needed to confirm the safety of UFV-P2 and its potential as an agent for biocontrol of milk contaminants.

Nucleotide sequence accession number. The complete genome sequence of *P. fluorescens* phage UFV-P2 has been deposited in GenBank under accession number [JX863101](https://www.ncbi.nlm.nih.gov/nuccore/JX863101).

ACKNOWLEDGMENTS

We thank Cristina Dantas Vanetti of the Federal University of Viçosa for providing materials and technical support.

This work was supported by grants from the Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). The funders had no role in the study design, data collection, analysis, decision to publish, or preparation of the manuscript.

REFERENCES

1. Arcuri EF, Silva PDL, Brito MAVP, Brito JRF, Lange CC, Magalhães MMA. 2008. Counts, isolation and characterization of psychrotrophic bacteria from refrigerated raw milk. *Cienc. Rural* 38:2250–2255.
2. Mu Z, Du M, Bai Y. 2009. Purification and properties of a heat-stable enzyme of *Pseudomonas fluorescens* Rm12 from raw milk. *Eur. Food Res. Technol.* 228:725–734.

3. Nörnberg MFBL, Friedrich RSC, Weiss RDN, Tondo EC, Brandelli A. 2010. Proteolytic activity among psychrotrophic bacteria isolated from refrigerated raw milk. *Int. J. Dairy Technol.* **63**:41–46.
4. Dufour D, Nicodème M, Perrin C, Driou A, Brusseau E, Humbert G, Gaillard JL, Dary A. 2008. Molecular typing of industrial strains of *Pseudomonas* spp. isolated from milk and genetical and biochemical characterization of an extracellular protease produced by one of them. *Int. J. Food Microbiol.* **125**:188–196.
5. Kives J, Guadarrama D, Orgaz B, Rivera-Sen A, Vazquez J, SanJose C. 2005. Interactions in biofilms of *Lactococcus lactis* ssp. *cremoris* and *Pseudomonas fluorescens* cultured in cold UHT milk. *J. Dairy Sci.* **88**:4165–4171.
6. Thiel K. 2004. Old dogma, new tricks—21st century phage therapy. *Nat. Biotechnol.* **22**:31–36.
7. Dixon B. 2004. New dawn for phage therapy. *Lancet Infect. Dis.* **4**:186.
8. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* **25**:955–964.
9. Tan Y, Zhang K, Rao X, Jin X, Huang J, Zhu J, Chen Z, Hu X, Shen X, Wang L, Hu F. 2007. Whole genome sequencing of a novel temperate bacteriophage of *P. aeruginosa*: evidence of tRNA gene mediating integration of the phage genome into the host bacterial chromosome. *Cell. Microbiol.* **9**:479–491.