



Molecular mechanisms of electron transfer employed by native proteins and biological-inorganic hybrid systems



Michael Lienemann

VTT Technical Research Centre of Finland Ltd., 02150 Espoo, Finland

ARTICLE INFO

Article history:

Received 15 September 2020

Received in revised form 3 December 2020

Accepted 5 December 2020

Available online 14 December 2020

Keywords:

Electron transfer

Electron bifurcation

Redox potential

Redox protein

Enzymatic electrosynthesis

Protein engineering

ABSTRACT

Recent advances in enzymatic electrosynthesis of desired chemicals in biological-inorganic hybrid systems has generated interest because it can use renewable energy inputs and employs highly specific catalysts that are active at ambient conditions. However, the development of such innovative processes is currently limited by a deficient understanding of the molecular mechanisms involved in electrode-based electron transfer and biocatalysis. Mechanistic studies of non-electrosynthetic electron transferring proteins have provided a fundamental understanding of the processes that take place during enzymatic electrosynthesis. Thus, they may help explain how redox proteins stringently control the reduction potential of the transferred electron and efficiently transfer it to a specific electron acceptor. The redox sites at which electron donor oxidation and electron acceptor reduction take place are typically located in distant regions of the redox protein complex and are electrically connected by an array of closely spaced cofactors. These groups function as electron relay centers and are shielded from the surrounding environment by the electrically insulating apoprotein. In this matrix, electrons travel via electron tunneling, i.e. hopping between neighboring cofactors, over impressive distances of up to several nanometers and, as in the case of the *Shewanella oneidensis* Mtr electron conduit, traverse the bacterial cell wall to extracellular electron acceptors such as solid ferrihydrite. Here, the biochemical strategies of protein-based electron transfer are presented in order to provide a basis for future studies on the basis of which a more comprehensive understanding of the structural biology of enzymatic electrosynthesis may be attained.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of Research Network of Computational and Structural Biotechnology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1. Introduction	206
2. Microbial electron transfer systems	208
2.1. Nitrogen reduction to ammonia by the <i>Azotobacter vinelandii</i> Mo-nitrogenase	208
2.2. Electron transfer by multiheme cytochromes <i>c</i> .	208
2.3. Flavin-based electron bifurcation in the NADH-dependent ferredoxin:NADP reductase Nfn.	210
3. Concluding remarks	211
Declaration of Competing Interest	211
Acknowledgements	211
References	211

1. Introduction

Survival of each biological species is highly dependent on its ability to harness the energy contained in the chemical bonds in

its environment. These reactions are catalyzed by enzymes and power other reactions that ensure the survival and propagation of the cell. Instrumental in this process are electron transfers between intracellular and extracellular compounds, which has

<https://doi.org/10.1016/j.csbj.2020.12.004>

2001-0370/© 2020 The Author(s). Published by Elsevier B.V. on behalf of Research Network of Computational and Structural Biotechnology.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).




Fig. 1. Electron transfer in the apoproteins presented in this review is mediated by various cofactors including the covalently bound iron-sulfur clusters [2Fe-2S] (A) and [4Fe-4S] (B), the P-cluster (C), FeMo cofactor (FeMo-co; D), heme *c* (E) as well as the non-covalently bound flavin adenine dinucleotide (FAD; F). Amino acids contributing to the covalent cofactor attachment to the apoprotein are shown in grey. Catalytically active iron atoms are colored blue and FAD nitrogen atoms transferring electrons during electron bifurcations are displayed in red. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

been fittingly captured by Nobel Prize-winning physiologist Albert Szent-Györgyi in his famous quote “Life is nothing but an electron looking for a place to rest”. As with most biochemical processes, electron transfer inside proteins is an electron tunneling reaction during which the electron hops from one electron relaying redox center to another, in a stepping stone fashion, until it is transferred to an external electron acceptor (cofactors discussed in this review as redox centers are presented in Fig. 1). A theoretical framework for electron transfer is provided by the Marcus theory with which tunneling rates can be directly predicted for a wide range of reactions [1,2]. According to this theory, the rate of electron transfer depends on five parameters, i.e., the free reaction energy, ΔG^0 , the free activation energy, ΔG^\ddagger , the reorganization energy, λ , the electronic coupling of the electron donor–acceptor pair, H_{DA} , and the temperature. H_{DA} is a measure of the probability at which electron tunneling occurs between degenerate states of the donor and acceptor. A large H_{DA} is characteristic for an adiabatic electron transfer with a low energy barrier ΔG^\ddagger when compared to non-adiabatic (low H_{DA}) electron transfers and strong coupling between the nuclei and electron motion. ΔG^0 is directly determined by the redox potentials of the electron donor–acceptor couple, while λ is the energy required to reorganize the nuclei of the surrounding physical medium and increases with donor–acceptor separation. In agreement with the Marcus theory, it was demonstrated by M. Kuss-Petermann et al. that electron transfer reactions are fastest when ΔG^0 and λ are of equal magnitude and that reactions are slowed down by an excessively large ΔG^0 [3]. The latter case is referred to as the “inverted regime” and has been subsequently experimentally confirmed using synthetic molecules that perform intramolecular electron transfer [4]. A theory of long-range electron transfer by tunneling in biological systems was developed already in 1974 by J. J. Hopfield [5] and, eight years later, validated

experimentally by J. R. Winkler and H. B. Gray. They designed a photochemical system consisting of a photoactive Ru complex as redox center that is covalently bound to the surface of a cytochrome or a blue copper protein [6,7]. By systematic variation of the position of Ru complex attachment, they were able to estimate H_{DA} and λ , and determine the pathways along which electrons travel between the redox centers [8,9]. In the case of electron transfer within proteins, edge-to-edge distances between two individual redox centers do not typically exceed 14 Å thereby ensuring that electron tunneling rates are faster than the common millisecond bond-breaking at the active sites of enzymes [10].

According to a recent survey, iron sulfur clusters and hemes are the most common electron-transferring cofactors in proteins [10], within which these are arranged as precisely spaced arrays that are shielded from the surrounding aqueous environment by an insulating protein matrix [11]. Electron transfer by proteins was initially described for the four membrane complexes of the mitochondrial respiratory chain that transfers two electrons from the reduction equivalents NADH and succinate to a terminal electron acceptor, such as oxygen, using a set of different redox cofactors including flavins, iron sulfur clusters and hemes [12]. The latter two are employed by Complex III and will be discussed in this minireview to illustrate the principles of electron bifurcation.

The possibility of driving enzyme reactions by an electric current at coulombic efficiencies of up to 99% has already been demonstrated for several oxidases and reductases [13] and a vast range of substrates [14]. Here, the catalytic center of the enzyme exchanges electrons with the electrode either directly or indirectly, utilizing stably protein-bound redox cofactors or soluble redox shuttles, respectively. From the perspective of practical application, the direct process appears more appealing since no redox mediator needs to be replenished after removal of the reaction




Fig. 2. Schematic representation of the *Azotobacter vinelandii* Mo-nitrogenase, displaying the locations of redox cofactors involved electron transfer from soluble electron donors to N_2 .

product. However, the stability and efficiency of most tested enzyme systems have been found to be suboptimal for practical applications [13] and would require further development of this technology, possibly with improvement by rational design. This approach requires a comprehensive understanding of the structure–function relationships of electron transferring proteins to which published research has already made many important contributions. This review discusses the concepts and current knowledge of the molecular mechanisms relevant to electron transfer in proteins and highlights some examples of previously described enzyme–electrode hybrid systems.

2. Microbial electron transfer systems

2.1. Nitrogen reduction to ammonia by the *Azotobacter vinelandii* Mo-nitrogenase

The reduction of nitrogen is a biologically important process that converts the inert gas into an activated form that is accessible to cell as N-source for the biosynthesis of a wide range of biomolecules such as proteins, DNA and carbohydrates. However, breaking of the triple bond in the nitrogen molecule requires a high amount of chemical energy. In order to perform the highly endergonic nitrogen reduction, nitrogen fixing microorganisms employ the nitrogenase enzyme complex, which consists of a reductase (Fe protein) and an N_2 -reducing subunit (MoFe protein) (Fig. 2). The Fe protein harbors a catalytic cofactor containing a bound iron atom and an additional eponymous metal atom, either Mo, V or Fe. The catalytic mechanism of enzymatic N_2 reduction has been studied in detail using the Mo-nitrogenase of the model nitrogen-fixing microbe *A. vinelandii* but is still a matter of debate. During the reduction of one molecule of N_2 , at least one equivalent of H_2 is produced. In the case of productive H_2 evolution, this reaction supports N_2 reduction by promoting the binding of the N_2 molecule and its partial reduction [15]. In addition to this so-called productive H_2 production, H_2 can be produced unproductively without concomitant electron transfer to N_2 , as discussed in detail elsewhere [16]. Thereby, the reduction of one N_2 molecule requires the passage of a minimum of eight electrons through the nitrogenase complex, during which the sequence of events needs to take place eight times (once per electron). The currently debated reaction mechanism is called “Fe protein cycle” and can be described as a ping-pong mechanism. Here, the Fe protein interacts with the β -unit of the MoFe protein through electrostatic interactions and then separates from the MoFe protein to associate with a soluble electron donor (Fid/Flx) using the same surface area. This

sequence of events is preceded by a step-wise modulation of the $[4Fe-4S]^{2+/1+}$ cofactor reduction pair potential through ATP binding ($-0.30 \rightarrow -0.43$ V vs. SHE) and association of the Fe protein with the MoFe protein ($-0.43 \rightarrow -0.62$ V vs. SHE), thus providing the reduction power required to eventually transfer an electron to the catalytic MoCo cofactor [17,18]. Then, a single electron is transferred to the MoCo cofactor by an initial abstraction from the P-cluster followed by an electron transfer from the $[4Fe-4S]^{1+}$, hydrolysis of 2 ATP, the release of 2 P_i , and a subsequent electron transfer from ferredoxin or flavodoxin to the Fe-protein cofactor $[4Fe-4S]^{2+}$ [19,20]. Owing to the soluble electron donor being oxidized as a terminal step, this process is called the “deficit-spend ing-mechanism” and has inspired engineering efforts, aiming at reducing the P-cluster of the catalytically active protein (i.e., MoFe, VFe or FeFe proteins) by either addition of soluble mediators, covalent attachment of photoelectrocatalytic materials or direct electron transfer from an electrode [21]. These strategies are attractive from an application point of view because they simplify and accelerate the electron transfer reaction by rendering it independent from the Fe-protein-catalyzed ATP hydrolysis and P_i release as the rate-limiting step in N_2 reduction [22].

The role of amino acids in the proximity of the P-cluster and the FeMo cofactor of the MoFe protein of the *A. vinelandii* Mo-nitrogenase has been investigated using mutational analysis [23]. The electroenzymatic reduction of hydrazine (N_2H_4) to NH_3 was quantified using a polyaminocarboxylate-ligated Eu(II) as an electron transfer mediator. The authors found that independent Tyr/His and Phe/His substitutions endowed the enzyme with the ability to electrocatalytically reduce hydrazine while N_2 reduction was not observed. In 2018, D. P. Hickey *et al.* reported the MoFe-based electroreduction of N_2 by non-mediated electron transfer to an electrode-bound MoFe protein with concomitant H_2 reduction [24]. This reaction produced 1.1 $\mu\text{mol } NH_3$ per mg of MoFe protein, which highlights the potential of direct electron transfer for the development of enzymatic electrosynthesis applications.

2.2. Electron transfer by multiheme cytochromes c

Among the known redox proteins, c-type multiheme cytochromes (cyts c) stand out because, unlike iron-sulfur clusters, they are insensitive to oxygen and are the only known cofactor allowing electron transfer proteins to directly reduce extracellular solid-phase electron acceptors such as Fe(III) and Mn(IV) minerals [25,26]. The redox cofactor used by multiheme cytochromes is heme c, in which a redox-active iron atom is coordinated by a porphyrin molecule in a tetradentate complex (Fig. 1). In cytochromes this iron is present either in the ferrous (Fe^{II}) or ferric state (Fe^{III}), which have characteristic UV–vis absorption maxima that can be monitored spectroscopically in order to follow cytochrome-mediated electron transfer [27,28]. The porphyrin ring of heme c is covalently bound to the apoprotein through two thioether bonds formed with cysteine residues of the characteristic CXXCH heme binding motif of cyts c [29].

Numerous studies have investigated the extracellular electron transfer of the Gram-negative bacterium *Shewanella oneidensis*, establishing it as a model species and providing detailed insights into the molecular mechanisms of this process [26,30]. According to current understanding, the transmembrane electron transfer proceeds by a multistep incoherent hopping mechanism between redox centers that is initiated at the cytoplasmic face of the inner membrane by extraction of a single electron from NAD(P)H by NAD dehydrogenase (NAD DH). Then, the electron is transferred to a membrane-soluble menaquinone, which in turn reduces the membrane-bound menaquinol dehydrogenase CymA (Fig. 3). CymA passes the electron on either to the periplasmic fumarate reductase (FccA) or to the periplasmic small tetraheme cytochrome




Fig. 3. Electron conduit in the cell envelope of *Shewanella oneidensis* connecting the cytosolic face of the inner membrane (IM) and the extracellular face of the outer membrane (OM). Electrons are exchanged between redox cofactors (standard reduction potentials indicated) that are either functional as single-molecule units (menaquinone/menaquinol; MQ/MQH₂) or bound in soluble cytochromes CymA, FccA and STC (yellow) or membrane-bound electron transfer proteins (grey). Figure modified from [31]. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

(STC), which are soluble proteins and are thereby able to interact with the inner and outer membrane. Upon contact with the inner face of the outer membrane, the reduced periplasmic cytochrome transfers an electron to the transmembrane complex MtrABC. In addition to cytochromes MtrA and MtrC, this complex contains the funnel-shaped porin MtrB which does not actively participate in electron transfer but orients the 185 Å-long heme array of the complex perpendicular to the outer membrane and serves as an insulating sheet to the membrane-spanning MtrA [31]. Both MtrB functions are important for the function of the Mtr complex by, respectively, facilitating optimal electron transfer away from the cell, and shielding the MtrA hemes from membrane-soluble molecules such as oxygen, the reduction of which may give rise to reactive oxygen species causing oxidative damage to the cell. The hemes in the Mtr complex function as stepping stones for the transferred electron, with reduction potentials ranging from 0 to –400 mV. This heterogeneity of the electron transferring cofactors does not, however, restrict the electron transfer in a single direction as it is dependent on the relative redox potentials at the faces of the outer membrane. This was exemplified with purified Mtr complexes in liposome membranes, in which an inward electron transfer occurs when sodium dithionite ($E^{0'} = -660$ mV) is present in the exterior and methyl viologen ($E^{0'} = -446$ mV) in the interior space [32]. MtrC is bound to the extracellular face of MtrA from which electrons are channeled into the decaheme MtrC by a single heme and transferred to three surface-accessible hemes. This transfer is facilitated by 10 bis-His coordinated hemes that are arranged as a “staggered cross”. The electron transfer to the solid-phase electron acceptor is believed to occur at heme C10 since it is located furthest away from the hydrophobic bilayer core of the OM (~90 Å) and in the vicinity of a putative binding site with a PTPD amino acid sequence motif [33,28][33,28]. The remaining surface-accessible hemes C2 and C7 have higher redox potentials than the remaining MtrC hemes and are thus suggested to function as junctions through which electrons are passed on to neighboring cytochromes and facilitate long-range electron transport parallel to the cell membrane [34].

As is valid for all metalloproteins, cytochromes have evolved such that their redox potential corresponds to that of their respective redox partners and can be modulated through alterations of its primary structure. Various factors contribute to the cytochrome redox potential including (in order of decreasing magnitude) the

protein matrix, 1st coordination sphere ligands, type of heme deformation and attachment sequence, heme type, heme accessibility, 2nd coordination sphere ligands and surface charges [35]. The magnitude with which these factors modulate the cytochrome redox potential is, however, strongly dependent on the experimental conditions, and is a factor that complicates the estimation of combined effects. Among the mentioned factors, their modulation by stabilization of the heme cofactor in the hydrophobic heme pocket has been investigated using cyts *c*. Here, the low dielectric constant of the protein matrix relative to aqueous solution destabilizes the charged ferric state of the heme iron (Fe^{III}) over the neutral ferrous state (Fe^{II}) [36]. Cyts *c* span a broad redox potential spectrum that extends to positive redox potentials, which are not reached by cyts with non-covalently bound hemes, such as heme *b*. This is believed to be due to the 10⁶ times weaker binding of the highly charged oxidized heme to the apoprotein when compared to its reduced state, which results in solvation and probably loss of the cofactor [37]. Such dislocation of heme *c* is prevented in cyts *c* by two thioether linkages originating from two Cys residues of the aforementioned CXXCH motif that also directly interacts with the heme Fe^{II/III} atom through the His as proximal ligand. The residues of the highly variable “XX” peptide sequence participate in H-bonding at the axial His, where loss of interactions reduces the histidinate character and increases the redox potential [38]. A counteracting effect of the H-bond network, in terms of the redox potential, is its stabilization of a distorted heme conformation, with respect to a perfectly planar tetrapyrrole ring, that has been demonstrated in case of saddled porphyrins to decrease the reduction potential of the Fe(II)/Fe(III) couple [39,40]. Hydrophobic and H-bonding interactions involving cyt *c* residues of the 2nd coordination sphere are considered to be crucial determinants of the redox potential and ET reorganization energy, as outlined for the CXXCH motif. Notably, the most prominent structural change observed for cyts *c* upon transition from the ferric to the ferrous state is the reorganization to the H-bond network that extends to the axial Met ligand, as demonstrated for the Tyr67 of yeast *iso-1*-cytochrome *c* [41–43]. The 1st coordination sphere of the heme iron of natural cyts *c* contains Met at the axial distal ligand position, which influences the redox potential through a combination of endergonic and entropic effects [36]. When considering cytochromes in general, this position can either remain vacant or be occupied by endogenous ligands (e.g., Met or His) or exogenous

ones (e.g., H_2O or OH^-). Most cyts contain endogenous axial ligands, among which a 100–150 mV higher redox potential is found for Met ligation when compared to bis-His complexes [44–46]. The thioether bond of the axial Met ligand has a π -electron-acceptor character with which it stabilizes the ferrous form of the heme iron and is mainly responsible for the high redox potentials of class I cyts *c* [36,47]. A comparison of microperoxidase-8 and cyt *c* with an unoccupied distal position and a distal Met ligand, respectively, showed that the cyt *c* protein fold efficiently excludes the solvent and thereby renders entropic effects negligible [47].

Multiheme cytochromes are not catalytically active and therefore have not yet been applied for electrosynthetic applications. It is, however, noteworthy that attachment and electrocatalysis has been demonstrated at cathodes and anodes using different heme-containing dehydrogenase and peroxidase enzymes, respectively [14]. Structural studies of cellobiose dehydrogenase revealed that its heme transfers electrons between an external electron donor and the catalytic FAD cofactor and thereby may function as an in-built redox mediator [48,49]. A comprehensive survey of cytochrome applications involving direct electron transfer is provided in [50].

2.3. Flavin-based electron bifurcation in the NADH-dependent ferredoxin:NADP reductase *Nfn*

Electron bifurcation is a unique case of electron transfer, as it splits the electron influx derived from substrate oxidation to yield one product with a lower and another with a higher reduction potential than the substrate. This is achieved by two half-reactions, which are energetically coupled to drive the endergonic one at the expense of a preceding exergonic transfer [51]. The concept of electron bifurcation was first presented in 1975 to rationalize the electron transfer performed by the quinone-based respiratory chain component complex III (cytochrome *bc*₁) in the cristall membrane of mitochondria [52]. Essential to understanding the mechanisms of electron bifurcation are the electron transfer reactions yielding and oxidizing the highly energized semiquinone (SQ). Initially, two electrons are transferred from a membrane-soluble hydroquinone (QH_2) to a complex-III-bound quinone (Q) to yield a bound QH_2 ($E^0 = +90$ mV). The exergonic 1-electron transfer from the bound QH_2 to soluble cytochrome *c* ($E^0 = +250$ mV) via a bound iron sulfur cluster is followed by a movement of the iron sulfur cluster away from the bound SQ that increases the distance between both redox centers to > 20 Å. This structural change favors the second 1-electron transfer “uphill” along the endergonic path from SQ to the low-potential acceptor cytochrome *b*_L ($E^0 = -60$ mV) [53]. The second electron is then transferred and eventually recycled by transfer to a membrane-soluble Q or QH to

yield QH_2 . The spatial separation of SQ and the iron sulfur cluster is a common mechanism among bifurcating flavoproteins by which the high fidelity of electron bifurcation is ensured [54–56]. In addition, all bifurcating enzymes contain a bifurcating cofactor exhibiting “crossed-over potentials” by which bifurcated electrons are directed into the exergonic and endergonic path at a 1:1 distribution [51]. As opposed to the “normal mode”, where both electrons are transferred to high-potential acceptors, during electron bifurcation, the first transferred electron has a higher potential than the second transferred electron. Here, the first electron transfer produces a highly reactive low-potential SQ radical as a reaction intermediate that is able to reduce a nearby low-potential acceptor.

Until the year 2008, quinones were the only known bifurcating cofactors. This was changed by the discovery of flavin-based electron bifurcation [57,58]. The bacterial ferredoxin:NADP reductase (*NfnAB*) is the best-characterized electron bifurcating enzyme complex (Fig. 4) and differs from complex III in that no major conformational changes occur during electron transfer [59,60]. According to the current understanding of the *NfnAB*-catalyzed process, electron bifurcation is initiated in the large subunit by a hydride transfer from soluble NADPH to b-FAD. The fully reduced bifurcating cofactor b-FADH is oxidized in a thermodynamically uphill 1-electron transfer to b-FAD^{•−} ($E_{m,\text{SQ/HQ}} = -359$ mV) by the high-potential $[2\text{Fe-2S}]^{2+}$ cluster ($E_m = +80$ mV) [51]. This is followed by an immediate transfer of a second electron from the highly reactive semiquinone FAD^{•−} ($E_{m,\text{SQ/Q}} = -911$ mV) to the proximal $[4\text{Fe-4S}]^{2+}$ cluster ($E_m = +80$ mV) from which the electron is passed on to the distal $[4\text{Fe-4S}]^{2+}$ cluster ($E_m = -513$ mV) and transferred further to ferredoxin ($E_m = -500$ to -400 mV) as the terminal electron acceptor of the endergonic path. The final steps of the electron bifurcation process occur in the small subunit NfnB, where the $[2\text{Fe-2S}]^{1+}$ cluster is oxidized by a-FAD ($E_m = -276$ mV) in an endergonic reaction yielding a-FAD^{•−} after an additional completion of the aforementioned sequence of electron transfers following the NADPH oxidation. Ultimately, two electrons are transferred from a-FAD^{•−} to the soluble electron acceptor NAD^+ yielding NADH. Thereby, the first electron removal from the reduced flavin b-FADH generates the thermodynamic driving force for the transfer of the second electron into the endergonic path of the *NfnAB* complex. Furthermore, the five bound redox centers of *Nfn* illustrate the “14 Å rule” according to which neighboring reduction centers need to be ≤ 14 Å apart to allow for effective electron transfer to the catalytic sites and not limit substrate turnover, which typically takes place in redox enzymes in about one millisecond [10].

The recently published structure of the heterodisulfide reductase/hydrogenase MvhADG – HdrABC from the methanogenic archaeon *Methanothermococcus thermolithotrophicus* provided




Fig. 4. Electron transfer in NADH-dependent ferredoxin:NADP reductase (*NfnAB*). Edge-to-edge distances between cofactors and mid-point potentials of the enzyme-bound redox cofactors and soluble electron donor and acceptor molecules participating in the electron transfer are indicated as reported in [51].




Fig. 5. Proposed pathways along which electrons are transferred in *Methanothermococcus thermolithotrophicus* heterodisulfide reductase/hydrogenase MvhADG – HdrABC during oxidation of H₂ and bifurcated to reduce the CoM–CoB heterodisulfide and ferredoxin. Reproduced from [61] with permission from the American Association for the Advancement of Science.

detailed information on the mechanism of electron bifurcation during methanogenesis utilizing an extended array of iron sulfur clusters [61], (Fig. 5). This enzyme oxidizes H₂ and transfers the extracted high-potential electron to coenzymes M and B and the low-potential electron to ferredoxin using a bifurcating flavin and an array of 14 iron sulfur clusters. According to the available structural data, the two isoalloxazine rings of the flavin are separated > 30 Å from the closest cofactor of the electron transfer route originating at the site of H₂ oxidation ([2Fe–2S] cluster MD). The apparent MD–FAD distancing violates the 14 Å rule since it exceeds the permissible length for electron tunneling and, based on high B factors, conformational rearrangements have been proposed to occur during electron transfer that reduce the MD–FAD spacing. Another noteworthy heterodisulfide reductase-containing complex is the *Methanococcus maripaludis* heterodisulfide reductase supercomplex, which contains two electron transfer routes spanning the heterodisulfide reductase HdrABC, with the bifurcating FAD bound to HdrA, and an attached hydrogenase domain VhuD as a constant component. The electron transfer routes are extended by either a formate dehydrogenase FdhAB or an auxiliary hydrogenase VhuAG that are incorporated during growth under H₂ limitation or H₂ excess, respectively [62,63]. Enzymatic electrosynthesis of formate at 90% efficiency was demonstrated using the (Fdh)₂ homocomplex, but the redox cofactors and molecular mechanisms involved in this reaction require further investigation [64].

3. Concluding remarks

The discussed electron-transferring proteins allow their host cells to perform vital anabolic and catabolic reactions by accessing distant electron acceptors, conservation of chemical energy or accessing essential nutrients that require chemical activation. As apparent from the electron bifurcating flavoproteins and the wire-like multiheme cytochromes, dynamic changes of the protein structure can support the catalysis of redox reactions but were not essential for electron transfer over long distances. In addition, the transient association of the nitrogenase protein complex components effectively facilitated the gradual supply of chemical energy required for the highly endergonic electron transfer from flavodoxin/ferredoxin to the inert gaseous substrate. Furthermore, the structure-aided engineering of this complex facilitating electrosyn-

thesis of H₂ and NH₃ serves as a proof-of-principle for the possibility to employ engineering approaches for the advancement of the enzymatic electrosynthesis technology. The knowledge gained until now on the functionality of the protein structure for electron transfer on a molecular level motivates further studies employing experimental and computational methods for the design of novel biological-inorganic hybrid systems.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was supported by an Academy Research Fellowship grant awarded by the Academy of Finland (Decision No. 321723). Peter Richard is thanked for providing helpful advice.

References

- [1] Marcus RA. On the theory of oxidation–reduction reactions involving electron transfer. I. *J Chem Phys* 1956;24(5):966–78. <https://doi.org/10.1063/1.1742723>.
- [2] Marcus RA, Sutin N. Electron transfers in chemistry and biology. *BBA Rev Bioenerg* 1985;811:265–322.
- [3] Kuss-Petermann M, Wenger OS. Electron transfer rate maxima at large donor–acceptor distances. *J Am Chem Soc* 2016;138(4):1349–58. <https://doi.org/10.1021/jacs.5b11953.s001>.
- [4] Miller JR, Calcaterra LT, Closs GL. Intramolecular long-distance electron transfer in radical anions. The effects of free energy and solvent on the reaction rates. *J Am Chem Soc* 1984;106(10):3047–9. <https://doi.org/10.1021/ja00322a058>.
- [5] Hopfield JJ. Electron transfer between biological molecules by thermally activated tunneling. *Proc Natl Acad Sci* 1974;71(9):3640–4. <https://doi.org/10.1073/pnas.71.9.3640>.
- [6] Gray HB, Winkler JR. Electron flow through metalloproteins. *Biochim Biophys Acta (BBA) – Bioenerg* 2010;1797(9):1563–72. <https://doi.org/10.1016/j.bbabi.2010.05.001>.
- [7] Winkler JR, Gray HB. Electron flow through metalloproteins. *Chem Rev* 2014;114(7):3369–80. <https://doi.org/10.1021/cr4004715>.
- [8] Winkler JR, Gray HB. Electron transfer in ruthenium-modified proteins. *Chem Rev* 1992;92(3):369–79. <https://doi.org/10.1021/cr00011a001>.
- [9] Wuttke DS, Bjerrum MJ, Winkler JR, Gray HB. Electron-tunneling pathways in cytochrome c. *Science* 1992;256(5059):1007–9. <https://doi.org/10.1126/science.256.5059.1007>.

- [10] Page C, Moser CC, Dutton PL. Mechanism for electron transfer within and between proteins. *Curr Opin Chem Biol* 2003;7:551–6.
- [11] H. Beinert Iron-Sulfur Clusters: Nature's Modular, Multipurpose Structures 277 5326 653 659 10.1126/science.277.5326.653.
- [12] Berg JM, Tymoczko JL, Stryer L. *Biochemistry*. 5th ed. New York: W H Freeman; 2002.
- [13] Wu R, Ma C, Zhu Z. Enzymatic electrosynthesis as an emerging electrochemical synthesis platform. *Curr Opin Electrochem* 2020;19:1–7. <https://doi.org/10.1016/j.coelec.2019.08.004>.
- [14] Dominguez-Benetton X, Srikanth S, Satyawali Y, Vanbroekhoven K, Pant D. Enzymatic electrosynthesis: an overview on the progress in enzyme-electrodes for the production of electricity, fuels and chemicals. *J Microb Biochem Technol* 2013.
- [15] Lukoyanov D, Barney BM, Dean DR, Seefeldt LC, Hoffman BM. Connecting nitrogenase intermediates with the kinetic scheme for N₂ reduction by a relaxation protocol and identification of the N₂ binding state. *Proc Natl Acad Sci* 2007;104(5):1451–5. <https://doi.org/10.1073/pnas.0610975104>.
- [16] Hoffman BM, Dean DR, Seefeldt LC. Climbing nitrogenase: toward a mechanism of enzymatic nitrogen fixation. *Acc Chem Res* 2009;42(5):609–19. <https://doi.org/10.1021/ar8002128>.
- [17] Lanzilotta WN, Seefeldt LC. Changes in the Midpoint Potentials of the Nitrogenase Metal Centers as a Result of Iron Protein–Molybdenum–Iron Protein Complex Formation †. *Biochemistry* 1997;36(42):12976–83. <https://doi.org/10.1021/bi9715371>.
- [18] Lough S, Burns A, Watt GD. Redox reactions of the nitrogenase complex from *Azotobacter vinelandii*. *Biochemistry* 1983;22(17):4062–6. <https://doi.org/10.1021/bi00286a011>.
- [19] Danyal K, Dean DR, Hoffman BM, Seefeldt LC. Electron Transfer within Nitrogenase: Evidence for a Deficit-Spending Mechanism. *Biochemistry* 2011;50(43):9255–63. <https://doi.org/10.1021/bi201003a>.
- [20] Duval S, Danyal K, Shaw S, Lytle AK, Dean DR, Hoffman BM, Antony E, Seefeldt LC. Electron transfer precedes ATP hydrolysis during nitrogenase catalysis. *Proc Natl Acad Sci* 2013;110(41):16414–9. <https://doi.org/10.1073/pnas.1311218110>.
- [21] Gu W, Milton RD. Natural and engineered electron transfer of nitrogenase. *Chemistry (Easton)* 2020;2:322–46.
- [22] Yang Z-Y, Ledbetter R, Shaw S, Pence N, Tokmina-Lukaszewska M, Eilers B, Guo Q, Pokhrel N, Cash VL, Dean DR, Antony E, Bothner B, Peters JW, Seefeldt LC. Evidence that the P i release event is the rate-limiting step in the nitrogenase catalytic cycle. *Biochemistry* 2016;55(26):3625–35. <https://doi.org/10.1021/acs.biochem.6b00421.s001>.
- [23] Danyal K, Rasmussen AJ, Keable SM, Inglet BS, Shaw S, Zadornyy OA, Duval S, Dean DR, Raugei S, Peters JW, Seefeldt LC. Fe protein-independent substrate reduction by nitrogenase MoFe protein variants. *Biochemistry* 2015;54(15):2456–62. <https://doi.org/10.1021/acs.biochem.5b00140>.
- [24] Hickey DP, Lim K, Cai R, Patterson AR, Yuan M, Sahin S, Abdellaoui S, Minter SD. Pyrene hydrogel for promoting direct bioelectrochemistry: ATP-independent electroenzymatic reduction of N₂. *Chem Sci* 2018;9(23):5172–7. <https://doi.org/10.1039/C8SC01638K>.
- [25] Edwards MJ, Richardson DJ, Paquette CM, Clarke TA. Role of multiheme cytochromes involved in extracellular anaerobic respiration in bacteria. *Protein Sci* 2020;29(4):830–42. <https://doi.org/10.1002/pro.3787>.
- [26] Beblawy S, Bursac T, Paquette C, Louro R, Clarke TA, Gescher J. Extracellular reduction of solid electron acceptors by *Shewanella oneidensis*. *Mol Microbiol* 2018;109(5):571–83. <https://doi.org/10.1111/mmi.14067>.
- [27] Yamanaka T. *The biochemistry of bacterial cytochromes*. New York: Springer-Verlag; 1992.
- [28] Rosso KM, Blumberger J, Butt JN. Multi-haem cytochromes in *Shewanella oneidensis* MR-1: structures, functions and opportunities. *J R Soc Interface* 2015;12:20141117. <https://doi.org/10.1098/rsif.2014.1117>.
- [29] Liu J, Chakraborty S, Hosseinzadeh P, Yu Y, Tian S, Petrik I, Bhagi A, Lu Yi. Metalloproteins containing cytochrome, iron-sulfur, or copper redox centers. *Chem Rev* 2014;114(8):4366–469. <https://doi.org/10.1021/cr400479b>.
- [30] Pirbadian S, El-Naggar MY. Multistep hopping and extracellular charge transfer in microbial redox chains. *Phys Chem Chem Phys* 2012;14(40):13802. <https://doi.org/10.1039/c2cp41185g>.
- [31] Edwards MJ, White GF, Butt JN, Richardson DJ, Clarke TA. The crystal structure of a biological insulated transmembrane molecular wire. *Cell* 2020;181(3):665–673.e10. <https://doi.org/10.1016/j.cell.2020.03.032>.
- [32] White GF, Shi Z, Shi L, Wang Z, Dohnalkova AC, Marshall MJ, Fredrickson JK, Zachara JM, Butt JN, Richardson DJ, Clarke TA. Rapid electron exchange between surface-exposed bacterial cytochromes and Fe(III) minerals. *Proc Natl Acad Sci* 2013;110(16):6346–51. <https://doi.org/10.1073/pnas.1220074110>.
- [33] Lower BH, Lins RD, Oestreicher Z, Straatsma TP, Hochella Jr MF, Shi L, Lower SK. In vitro evolution of a peptide with a hematite binding motif that may constitute a natural metal-oxide binding archetype. *Environ Sci Technol* 2008;42(10):3821–7. <https://doi.org/10.1021/es702688c>.
- [34] Jiang X, Burger B, Gajdos F, Bortolotti C, Futera Z, Breuer M, Blumberger J. Kinetics of trifurcated electron flow in the decaheme bacterial proteins MtrC and MtrF. *Proc Natl Acad Sci USA* 2019;116(9):3425–30. <https://doi.org/10.1073/pnas.1818003116>.
- [35] Alvarez-Paggi D, Hannibal L, Castro MA, Oviedo-Rouco S, Demicheli V, Tórtora V, Tomasina F, Radi R, Murgida DH. Multifunctional cytochrome c : learning new tricks from an old dog. *Chem Rev* 2017;117(21):13382–460. <https://doi.org/10.1021/acs.chemrev.7b00257>.
- [36] Tezcan FA, Winkler JR, Gray HB. Effects of ligation and folding on reduction potentials of heme proteins. *J Am Chem Soc* 1998;120(51):13383–8. <https://doi.org/10.1021/ja982536e>.
- [37] Zheng Z, Gunner MR. Analysis of the electrochemistry of hemes with E_m s spanning 800 mV. *Proteins* 2009;75(3):719–34. <https://doi.org/10.1002/prot.22282>.
- [38] Can M, Zoppellaro G, Andersson KK, Bren KL. Modulation of ligand-field parameters by heme ruffling in cytochromes c revealed by EPR spectroscopy. *Inorg Chem* 2011;50(23):12018–24. <https://doi.org/10.1021/ic201479g>.
- [39] Olea Jr C, Kuriyan J, Marletta MA. Modulating heme redox potential through protein-induced porphyrin distortion. *J Am Chem Soc* 2010;132(37):12794–5. <https://doi.org/10.1021/ja106252b>.
- [40] Grinstaff MW, Hill MG, Birnbaum ER, Schaefer WP, Labinger JA, Gray HB. Structures, electronic properties, and oxidation-reduction reactivity of halogenated iron porphyrins. *Inorg Chem* 1995;34(19):4896–902. <https://doi.org/10.1021/ic00123a026>.
- [41] Berghuis AM, Guillemette JG, Smith M, Brayer GD. Mutation of tyrosine-67 to phenylalanine in cytochrome c significantly alters the local heme environment. *J Mol Biol* 1994;235(4):1326–41. <https://doi.org/10.1006/jmbi.1994.1086>.
- [42] Berghuis AM, Guillemette JG, McLendon G, Sherman F, Smith M, Brayer GD. The role of a conserved internal water molecule and its associated hydrogen bond network in cytochrome c. *J Mol Biol* 1994;236(3):786–99. <https://doi.org/10.1006/jmbi.1994.1189>.
- [43] Lett CM, Berghuis AM, Frey HE, Lepock JR, Guillemette JG. The role of a conserved water molecule in the redox-dependent thermal stability of iso-1-cytochrome c. *J Biol Chem* 1996;271(46):29088–93. <https://doi.org/10.1074/jbc.271.46.29088>.
- [44] Miller GT, Zhang B, Hardman JK, Timkovich R. Converting a c-type to a b-type cytochrome: Met61 to His61 mutant of *Pseudomonas* cytochrome c-551. *Biochemistry* 2000;39:9010–7.
- [45] Hay S, Wydrzynski T. Conversion of the *Escherichia coli* cytochrome b₅₆₂ to an archetype cytochrome b: a mutant with bis-histidine ligation of heme iron. *Biochemistry* 2005;44:431–9.
- [46] Raphael AL, Gray HB. Axial ligand replacement in horse heart cytochrome c by semisynthesis. *Proteins* 1989;6(3):338–40. <https://doi.org/10.1002/prot.340060316>.
- [47] Battistuzzi G, Borsari M, Cowan JA, Ranieri A, Sola M. Control of cytochrome c redox potential: axial ligation and protein environment effects. *J Am Chem Soc* 2002;124:5315–24.
- [48] Tan T-C, Kracher D, Gandini R, Sygmond C, Kittl R, Haltrich D, Hällberg BM, Ludwig R, Divne C. Structural basis for cellobiose dehydrogenase action during oxidative cellulose degradation. *Nat Commun* 2015;6(1). <https://doi.org/10.1038/ncomms8542>.
- [49] Scheiblbrandner S, Ludwig R. Cellobiose dehydrogenase: bioelectrochemical insights and applications. *Bioelectrochemistry* 2020;131:107345. <https://doi.org/10.1016/j.bioelechem.2019.107345>.
- [50] Ma Su, Ludwig R. Direct electron transfer of enzymes facilitated by cytochromes. *ChemElectroChem* 2019;6(4):958–75. <https://doi.org/10.1002/celec.201801256>.
- [51] Buckel W, Thauer RK. Flavin-based electron bifurcation, a new mechanism of biological energy coupling. *Chem Rev* 2018;118(7):3862–86. <https://doi.org/10.1021/acs.chemrev.7b00707>.
- [52] Mitchell P. The protonmotive Q cycle: a general formulation. *FEBS Lett* 1975;59:137–9.
- [53] Zhu J, Egawa T, Yeh S-R, Yu L, Yu C-A. Simultaneous reduction of iron-sulfur protein and cytochrome b_L during ubiquinol oxidation in cytochrome bc₁ complex. *Proc Natl Acad Sci* 2007;104(12):4864–9. <https://doi.org/10.1073/pnas.0607812104>.
- [54] Garcia Costas AM, Poudel S, Miller A-F, Schut GJ, Ledbetter RN, Fixen KR, Seefeldt LC, Adams MWW, Harwood CS, Boyd ES, Peters JW, Becker A. Defining electron bifurcation in the electron-transferring flavoprotein family. *J Bacteriol* 2017;199(21). <https://doi.org/10.1128/JB.00440-17>.
- [55] Demmer JK, Bertsch J, Öppinger C, Wohlers H, Kayastha K, Demmer U, Ermiler U, Müller V. Molecular basis of the flavin-based electron-bifurcating caffeoyl-CoA reductase reaction. *FEBS Lett* 2018;592(3):332–42. <https://doi.org/10.1002/1873-3468.12971>.
- [56] Demmer JK, Pal Chowdhury N, Selmer T, Ermiler U, Buckel W. The semiquinone swing in the bifurcating electron transferring flavoprotein/butyryl-CoA dehydrogenase complex from *Clostridium difficile*. *Nat Commun* 2017;8(1). <https://doi.org/10.1038/s41467-017-01746-3>.
- [57] Herrmann G, Jayamani E, Mai G, Buckel W. Energy conservation via electron-transferring flavoprotein in anaerobic bacteria. *JB* 2008;190(3):784–91. <https://doi.org/10.1128/JB.01422-07>.
- [58] Li F, Hinderberger J, Seedorf H, Zhang J, Buckel W, Thauer RK. Coupled ferredoxin and crotonyl coenzyme A (CoA) reduction with NADH catalyzed by the butyryl-CoA dehydrogenase/Etf Complex from *Clostridium kluyveri*. *J Bacteriol* 2008;190(3):843–50. <https://doi.org/10.1128/JB.01417-07>.
- [59] Lubner CE, Jennings DP, Mulder DW, Schut GJ, Zadornyy OA, Hoben JP, Tokmina-Lukaszewska M, Berry L, Nguyen DM, Lipscomb GL, Bothner B, Jones AK, Miller A-F, King PW, Adams MWW, Peters JW. Mechanistic insights into energy conservation by flavin-based electron bifurcation. *Nat Chem Biol* 2017;13(6):655–9. <https://doi.org/10.1038/nchembio.2348>.
- [60] Berry L, Poudel S, Tokmina-Lukaszewska M, Colman DR, Nguyen DMN, Schut GJ, Adams MWW, Peters JW, Boyd ES, Bothner B. H/D exchange mass

- spectrometry and statistical coupling analysis reveal a role for allostery in a ferredoxin-dependent bifurcating transhydrogenase catalytic cycle. *Biochimica et Biophysica Acta (BBA) - General Subjects* 2018;1862(1):9–17. <https://doi.org/10.1016/j.bbagen.2017.10.002>.
- [61] Wagner T, Koch J, Ermler U, Shima S. Methanogenic heterodisulfide reductase (HdrABC-MvhAGD) uses two noncubane [4Fe-4S] clusters for reduction. *Science* 2017;357(6352):699–703. <https://doi.org/10.1126/science.aan0425>.
- [62] Milton RD, Ruth JC, Deutzmann JS, Spormann AM. *Methanococcus maripaludis* employs three functional heterodisulfide reductase complexes for flavin-based electron bifurcation using hydrogen and formate. *Biochemistry* 2018;57(32):4848–57. <https://doi.org/10.1021/acs.biochem.8b00662.s001>.
- [63] Costa KC, Wong PM, Wang T, Lie TJ, Dodsworth JA, Swanson I, et al. Protein complexing in a methanogen suggests electron bifurcation and electron delivery from formate to heterodisulfide reductase. *Proc Natl Acad Sci* 2010;107:11050–5. <https://doi.org/10.1073/pnas.1003653107>.
- [64] Lienemann M, Deutzmann JS, Milton RD, Sahin M, Spormann AM. Mediator-free enzymatic electrosynthesis of formate by the *Methanococcus maripaludis* heterodisulfide reductase supercomplex. *Bioresour Technol* 2018;254:278–83. <https://doi.org/10.1016/j.biortech.2018.01.036>.