

SCIENTIFIC REPORTS



OPEN

Measuring the Energy of Ventilation and Circulation during Human Walking using Induced Hypoxia

Masahiro Horiuchi¹, Yoshiyuki Fukuoka², Yoko Handa¹, Daijiro Abe³ & Herman Pontzer^{4,5}

Energy expenditure (EE) during walking includes energy costs to move and support the body and for respiration and circulation. We measured EE during walking under three different oxygen concentrations. Eleven healthy, young, male lowlanders walked on a treadmill at seven gait speeds ($0.67\text{--}1.83\text{ m s}^{-1}$) on a level gradient under normobaric normoxia (room air, 21% O_2), moderate hypoxia (15% O_2), and severe hypoxia (11% O_2). By comparing the hypoxia-induced elevation in heart rate (HR [bpm]), ventilation (V_E [L min^{-1}]) with the change in energy expenditure (EE [W]) at each speed, we were able to determine circulatory and respiratory costs. In a multivariate model combining HR and V_E , respiratory costs were $0.44 \pm 0.15\text{ W}$ per each L min^{-1} increase in V_E , and circulatory costs were $0.24 \pm 0.05\text{ W}$ per each bpm increase in HR (model adjusted $r^2 = 0.97$, $p < 0.001$). These V_E costs were substantially lower than previous studies that ignored the contribution of HR to cardiopulmonary work. Estimated HR costs were consistent with, although somewhat higher than, measures derived from catheterization studies. Cardiopulmonary costs accounted for 23% of resting EE, but less than 5% of net walking costs (i.e., with resting EE subtracted).

Humans' bipedal walking gait is a defining feature of our lineage, and the metabolic costs of walking have figured prominently in discussions of human evolution and ecology^{1–3}. The energy cost of walking must include energy expenditure (EE) to move and support the body and to fuel physiological changes, such as increased ventilation and circulation, needed to meet increased metabolic demand. To date, studies of walking cost have largely ignored the costs of ventilation and circulation, focusing instead on the factors known to affect the costs of movement such as walking speed, internal and external work^{4,5}, gradients^{6,7}, body mass^{8–11}, and load carriage^{12–15}. Locomotor cost is of central importance in human ecology^{16,17}, behavior^{18,19}, evolution^{1–3}, and biomechanics research^{4,5,7,12}, and a full accounting of human walking cost and its determinants requires an understanding of the contribution of cardiopulmonary work to total metabolic cost.

Respiratory costs have been measured in humans and other vertebrates during rest and exercise^{20–30}. Early studies of respiratory cost at rest, measured by having the subject breathe through a long tube to increase the dead space and thus the work needed to breathe, found that ventilation costs in humans at rest were quite low ($<1\text{ mlO}_2$ per L V_E)^{21,27}. Subsequent studies reported substantially greater respiratory costs during exercise, though still less than 10% of total exercise energy expenditure^{20,22,23,25,29,30}. However, these studies have generally used an experimental design in which respiratory costs are not measured during exercise, but rather during voluntary hyperpnea in which subjects match their respiratory frequency and tidal volume to levels observed during exercise²⁰. The contribution of increased heart rate (HR) is generally ignored, although HR can increase substantially during hyperpnea³¹ and could contribute substantially to the measured increase in energy expenditure. Further, these studies have typically focused on running and other moderate- or high-intensity exercise. The contribution of cardiopulmonary work to walking cost may be higher, as the total cost of walking is considerably lower³².

Measuring walking costs under hypoxic conditions provides a potential experimental design for measuring cardiopulmonary costs during exercise. At high altitude, and under normobaric hypoxia, a reduction of alveolar

¹Division of Human Environmental Science, Mt. Fuji Research Institute, Kami-yoshida 5597-1, Fuji-yoshida-city, Yamanashi, 4030005, Japan. ²Faculty of Health and Sports Science, Doshisha University, Tatara 1-3, Kyotanabe, Kyoto, 6100394, Japan. ³Center for Health and Sports Science, Kyushu Sangyo University, Matsukadai 2-3-1, Higashi-ku, Fukuoka-city, Fukuoka, 8130003, Japan. ⁴Department of Anthropology, Hunter College, The City University of New York, New York, NY, 10065, USA. ⁵New York Consortium for Evolutionary Primatology, New York, NY, 10065, USA. Correspondence and requests for materials should be addressed to M.H. (email: mhoriuchi@mfri.pref.yamanashi.jp)

	Normoxia	Moderate	Severe	
EE, Watts			Δ EE	Δ EE
Rest	103 ± 18	100 ± 9	-3	106 ± 12
0.67 ms ⁻¹	211 ± 23	205 ± 23	-6	218 ± 20
0.86 ms ⁻¹	230 ± 28	224 ± 23	-6	236 ± 23
1.06 ms ⁻¹	252 ± 34	253 ± 28	1	261 ± 33
1.25 ms ⁻¹	281 ± 34	285 ± 31	4	288 ± 39
1.44 ms ⁻¹	328 ± 50	332 ± 36	4	338 ± 49
1.64 ms ⁻¹	412 ± 64	415 ± 59	3	424 ± 59
1.83 ms ⁻¹	495 ± 83	499 ± 79	4	513 ± 78
V_E, L min⁻¹			Δ V _E	Δ V _E
Rest	10.9 ± 0.8	11.2 ± 1.4	0.3	12.6 ± 1.4
0.67 ms ⁻¹	18.5 ± 2.5	18.1 ± 2.3	-0.4	21.4 ± 3.2
0.86 ms ⁻¹	20.2 ± 2.8	19.7 ± 2.7	-0.5	23.9 ± 4.0
1.06 ms ⁻¹	21.8 ± 3.4	22.0 ± 3.4	0.2	26.7 ± 4.9
1.25 ms ⁻¹	24.9 ± 3.1	25.5 ± 4.3	0.6	30.9 ± 6.8
1.44 ms ⁻¹	28.0 ± 3.2	28.9 ± 3.9	0.9	36.7 ± 7.0
1.64 ms ⁻¹	33.3 ± 4.3	34.9 ± 5.9	1.6	45.6 ± 9.4
1.83 ms ⁻¹	40.5 ± 5.3	42.2 ± 8.5	1.7	58.8 ± 15.7
HR, bpm			Δ HR	Δ HR
Rest	80 ± 11	83 ± 10	3	93 ± 11
0.67 ms ⁻¹	78 ± 10	83 ± 8	5	98 ± 11
0.86 ms ⁻¹	82 ± 10	85 ± 9	3	103 ± 11
1.06 ms ⁻¹	84 ± 11	89 ± 9	5	107 ± 11
1.25 ms ⁻¹	89 ± 11	94 ± 9	5	114 ± 10
1.44 ms ⁻¹	94 ± 10	101 ± 9	7 [‡]	123 ± 12
1.64 ms ⁻¹	101 ± 11	109 ± 8	8 [‡]	136 ± 12
1.83 ms ⁻¹	109 ± 12	119 ± 9	10 [‡]	146 ± 14

Table 1. Energy expenditure (EE), pulmonary ventilation (V_E), and heart rate (HR) at rest and each gait speed under three oxygen concentrations. Changes in these variables (Δ) under moderate and severe hypoxia versus normoxia are shown. Values are mean ± standard deviation. bpm; beats per minute. * $P < 0.05$ between normoxia and severe hypoxia, † $P < 0.05$ between moderate and severe hypoxia, ‡ $P < 0.05$ between normoxia and moderate hypoxia.

PO₂ limits pulmonary O₂ diffusion capability, which induces a decrease in arterial O₂ saturation (SpO₂)^{33, 34}. Under these conditions, pulmonary ventilation (V_E) increases within minutes³⁵, followed by increases in HR³⁶. These responses are generally considered part of the compensatory mechanisms against hypoxia with reduced oxygen delivery to the peripheral tissues. These results suggest that if hypoxic-induced increases in energy cost of ventilation and/or circulation are observed even at resting conditions, differential energy costs during walking under hypoxia would also be observed.

We developed and tested an experimental model to evaluate the energy cost of ventilation and circulation during walking using acute hypoxic exposure. We used pulmonary ventilation (V_E) and heart rate (HR) as indices of respiratory and circulatory work, respectively. Healthy male adults, not acclimated to hypoxic or high-altitude conditions, walked over a range of speeds under three different oxygen concentrations (normoxia, moderate, and severe hypoxia) at level gradient. This design enabled us to vary the cardiopulmonary work done while keeping the energy costs to support and move the body (i.e., the mechanical cost of walking) constant across conditions. Our objective was to determine the costs of ventilation and circulation during walking, and to compare values measured during walking to those measured during standing at rest.

Results

Mean (±SD) values for EE, V_E, and HR at rest and each walking speed, and changes in these variables with hypoxia are given in Table 1. All three change variables, Δ EE, Δ V_E, and Δ HR, increased with speed and degree of hypoxia, with severe hypoxia having a much stronger effect than moderate hypoxia (Fig. 1). Both walking speed and the degree of hypoxia were significant predictors for Δ EE (model adjusted $r^2 = 0.82$, $p < 0.001$), Δ V_E (adj. $r^2 = 0.66$, $p < 0.001$), and Δ HR (adj. $r^2 = 0.92$, $p < 0.001$) in multiple regression.

Bivariate analyses of Δ EE revealed significant correlations with both Δ V_E and Δ HR (Fig. 2). With the intercept set at 0, EE increased with ventilatory effort as Δ EE = $1.10 \pm 0.12 \Delta$ V_E ($r^2 = 0.84$, $p < 0.001$), or roughly 1 Watt for every 1 L min⁻¹ increase in V_E. This cost is equivalent to 3.3 mL O₂ per L V_E, assuming 20.1 J per mL O₂. Energy expenditure increased with circulation as Δ EE = $0.36 \pm 0.04 \Delta$ HR ($r^2 = 0.81$, $p < 0.001$), or roughly 1 Watt for every 3 beats per minute increase in HR. Slopes changed less than 3% when the three hypoxia-speed conditions with negative Δ EE values were excluded from analysis (Fig. 2).




Figure 1. Mean values of differences in energy expenditure (EE: top panel), pulmonary ventilation (V_E : middle panel), and heart rate (HR: bottom panel) between normoxic trial and moderate hypoxia (open circles), and between normoxic trial and severe hypoxia (filled circles) at rest and each walking speed. The number of 0.0 on the X axis indicate at rest.

ΔV_E and ΔHR were strongly correlated ($r^2 = 0.96$, second order polynomial regression, Fig. 2), making it difficult to determine their independent effects on EE. When ΔV_E and ΔHR were entered as independent variables in a multiple regression, ΔV_E achieved significance ($t(14) = 2.33$, $p = 0.036$) but ΔHR did not ($t(14) = 1.42$, $p = 0.178$). However, when the three hypoxia-speed conditions with negative ΔEE values were excluded from the analysis, both ΔV_E ($t(11) = 2.87$, $p = 0.015$) and ΔHR ($t(11) = 4.65$, $p < 0.001$) were significant, and the fit of the model was very strong ($\Delta EE = 0.44 \pm 0.15 \Delta V_E + 0.24 \pm 0.05 \Delta HR$, adj. $r^2 = 0.97$, $p < 0.001$). Notably, this multivariate model reduces the estimated individual costs of ventilation and circulation compared to bivariate analyses. Ventilatory costs are approximately 1 Watt for every 2.3 L min^{-1} increase in V_E ; circulatory costs are approximately 1 Watt for every 4 beats per minute increase in HR. The fit of this predictive model for all data is shown in Fig. 3.




Figure 2. Changes in ΔEE versus in ΔV_E (top panel), and changes in ΔEE versus in ΔHR (middle panel). Ordinary least squares linear regression equations, with intercept = 0, are shown. Changes in ΔV_E versus in ΔHR (bottom panel). Second order polynomial regression equation, with intercept = 0, are shown.

We used the combined model with ΔV_E and ΔHR (i.e., $EE = 0.44V_E + 0.24HR$) along with V_E and HR values at each condition to estimate the energy cost of cardiopulmonary work at rest and during walking under normoxia, moderate hypoxia, and severe hypoxia. At rest, the estimated energy cost of breathing and circulation accounted for approximately one-fourth of total EE across conditions, rising slightly under severe hypoxia compared to normoxia and moderate hypoxia. As walking speed and EE increased, the percentage of EE attributed to cardiopulmonary work decreased under all conditions. This change was similar in moderate hypoxia and severe hypoxia (Table 2).

Next, we subtracted EE and estimated cardiopulmonary expenditure during resting from values measured during walking in order to calculate net walking EE and net walking cardiopulmonary work. Net cardiopulmonary energy expenditure accounted for 3% of net walking EE at 0.67 m s^{-1} , rising to 5% at 1.83 m s^{-1} under both




Figure 3. Observed ΔEE plotted against that predicted ΔEE from the least squares regression with V_E and HR: $\Delta EE = 0.44\Delta V_E + 0.24\Delta HR$; see text. Line indicates $y = x$. This predictive fit model for data was calculated using the hypoxia-speed conditions with only positive ΔEE values (filled circles), but excluding the hypoxia-speed conditions with negative ΔEE values (open circles).

	Normoxia	Moderate	Severe
Watts, W			
0.67 ms ⁻¹	26.5	27.5	32.5
0.86 ms ⁻¹	28.2	28.7	34.8
1.06 ms ⁻¹	29.4	30.6	36.9
1.25 ms ⁻¹	31.9	33.3	40.4
1.44 ms ⁻¹	34.4	36.5	45.1
1.64 ms ⁻¹	38.4	41.0	52.0
1.83 ms ⁻¹	43.4	46.5	60.2
%EE, %			
0.67 ms ⁻¹	12.6	13.4	14.9
0.86 ms ⁻¹	12.3	12.8	14.7
1.06 ms ⁻¹	11.7	12.1	14.2
1.25 ms ⁻¹	11.4	11.7	14.0
1.44 ms ⁻¹	10.5	11.0	13.3
1.64 ms ⁻¹	9.3	9.9	12.3
1.83 ms ⁻¹	8.8	9.3	11.7
% walk net, %			
0.67 ms ⁻¹	2.6	2.9	4.5
0.86 ms ⁻¹	3.6	3.4	5.6
1.06 ms ⁻¹	3.8	4.0	6.1
1.25 ms ⁻¹	4.6	4.8	7.1
1.44 ms ⁻¹	4.8	5.2	7.6
1.64 ms ⁻¹	4.8	5.2	7.7
1.83 ms ⁻¹	5.0	5.5	8.0

Table 2. Mean values of EE, the percentage EE to total cardiopulmonary work, and the percentage of net cardiopulmonary EE at rest and each walking speed under all conditions.




Figure 4. Contribution of V_E (gray segments) and HR (white segments) to cardiopulmonary energy expenditure at rest and at each walking speed under normoxia (N), moderate hypoxia (M), and severe hypoxia (S) conditions.

normoxia and moderate hypoxia. Under severe hypoxia, these values were slightly higher compared to normoxia and moderate hypoxia. (Table 2).

Finally, we applied the multivariate model with ΔV_E and ΔHR to the V_E and HR values at each condition to calculate the relative contributions of ventilatory and circulatory work to cardiopulmonary energy expenditure. Across normoxic and both hypoxic conditions, the estimated energy cost of circulation exceeded that of ventilation. Heart beat accounted for 80% of cardiopulmonary work at rest in all three conditions. The proportion of cardiopulmonary work attributable to HR decreased regularly with walking speed under all conditions (Fig. 4).

Discussion

Both breathing and circulation have measurable effects on EE during both resting and walking. While our findings are consistent with studies using hyperpnea at rest to mimic the costs of ventilation during exercise^{20, 22, 23, 25, 29, 30}, results here also indicate that previous work has greatly underestimated the contribution of the heart to cardiopulmonary costs. Our bivariate analysis of V_E and EE (Fig. 2) gives a ventilation cost of 3.3 mlO₂ per L V_E (assuming 20.1 J per mlO₂), consistent with hyperpnea studies reporting 2–4 mlO₂ per L V_E during exercise^{20, 26, 30}. However, HR and V_E increase together during exercise (Fig. 2) and during enforced hyperventilation at rest³¹. Consequently, analyses that ignore HR and attribute cardiopulmonary costs entirely to breathing will artificially inflate the estimated cost of ventilation. This inflation may be substantial. Ventilation costs decreased 60%, to 1.3 mlO₂ per L V_E , and the heart accounted for more than half (60–80%) of cardiopulmonary energy cost, when ventilatory and circulatory costs were considered together in our multivariate model. Future studies using forced hyperpnea to estimate ventilation costs should include simultaneous measures of HR in order to assess the potential contribution of circulatory costs.

Measuring cardiopulmonary costs by induced hypoxia, as done here, enabled us to measure the energy expenditure of the heart non-invasively. To our knowledge, the only previous *in vivo* measures of the heart's energy expenditure were conducted by monitoring oxygen concentrations in arterial blood supply and venous return of a catheterized heart³⁷. Catheterization methods produce estimates of 7 W for heart energy expenditure in adult males lying supine, at rest³⁷. Assuming a resting HR of ~60 bpm, this estimate translates to ~7 J per beat. HR cost in our multivariate model was 0.24 ± 0.05 W per bpm, which translates to 14 ± 3 J per beat, twice that of catheterization methods. Given the very different experimental designs, these estimates are reasonably similar. One reason that our approach may produce higher estimates is that our measurements were taken primarily during exercise, which increases blood pressure and stroke volume compared to resting, and presumably increases the energy cost per heartbeat. More work is needed to refine non-invasive estimates of HR cost and to reconcile them with catheterization studies.

The present study focused on walking, rather than the running or cycling tests typical of previous work on cardiopulmonary cost, e.g., refs 20, 22–24, 26, 30. As a result, we examined a relatively low range of V_E . Maximum V_E in the present study was 59 L min⁻¹, which falls at the very low end of the range examined in most previous works. The lower range of V_E in this study may explain why we did not detect the exponential increase in V_E cost at higher ventilation rates reported by those studies^{26, 30}.

Cardiopulmonary effort accounts for a substantial portion of resting EE, approximately 23% in normoxic conditions. In contrast, cardiopulmonary energy expenditure accounts for only 3 to 5% of net walking EE (i.e., with resting EE subtracted). The small contribution of cardiopulmonary effort to net walking cost is in line with previous work in humans and a range of other vertebrates²⁵. These results lend support to analyses of locomotor

cost that focus on musculoskeletal energy expenditure, ignoring cardiopulmonary work, e.g., refs 3, 6, 7, 38–40. Even under severe hypoxia, cardiopulmonary energy cost never exceeded 8% of net walking EE, suggesting that the primary cost of walking at altitude remains the musculoskeletal effort needed to support and propel the body.

Of note, we found small (~3%) reductions in EE with moderate hypoxia compared to normoxia at rest and during walking at slow speeds (0.67 and 0.83 m s⁻¹). In our original study⁴¹, we found that VO₂ and VCO₂ in these conditions trended lower (without achieving statistical significance; see Table 1 and Fig. 1 in Horiuchi *et al.*⁴¹). Similarly, the small reductions reported here for EE, which is calculated from VO₂ and VCO₂, do not reach statistical significance. These results are nonetheless unexpected and we can only speculate regarding the cause of this phenomenon, but there are three general possibilities. One is that these small differences are spurious, reflecting measurement error and the limits of precision in our experimental set up. A second possibility is that the body fueled some small percentage of physiological activity using anaerobic metabolism in these conditions. In this case, the aerobic measures of EE employed here would fail to capture some small portion of total energy expenditure in these (and possibly other) conditions, leading to an underestimation of EE and ΔEE. Third, the body may be adjusting to hypoxic conditions by reducing EE to some unspecified set of physiological activities. Given the small size of the effect (~3% or less of EE), we do not expect that this phenomenon greatly affects the results of the currently analyses. Nonetheless, additional research into this phenomenon is warranted.

Another analytical concern here is the effect of CO₂ stores at the onset of exercise on our measures of VCO₂, and hence EE. As the generation of CO₂ increases at the beginning of exercise, initial production will be absorbed in the bloodstream and accumulate there, prior to being exchanged in the lungs and expired. Consequently, measures of expired air, such as those employed here, may underestimate VCO₂ at the beginning of an exercise bout. However, previous work has demonstrated that the CO₂ stores were in equilibrium by ~2 min of exercise⁴². As we discarded the first 3 minutes of each trial and used only the last 1 min data of VO₂ and VCO₂ at each gait speed in our analyses, the effect of CO₂ accumulation in the blood should be negligible for this study.

Limitations of this study include the absence of women in our sample and the relatively small range of V_E, HR, and EE examined. Our multivariate analyses also underscore the inherent difficulties and limitations in parsing the effects of HR and V_E on EE, given the physiological linkage between HR and V_E. Additional work, examining a broader range of subjects and activities, is needed to improve our estimates of HR and V_E cost and to determine whether these costs vary with body size, sex, age, or other factors.

Imposing greater cardiopulmonary effort through induced hypoxia provides a non-invasive approach for measuring cardiac and ventilatory costs during locomotion. The advantage of this approach is the ability to measure V_E and HR during a given activity, non-invasively, rather than mimicking these costs separately during forced hyperpnea. Results here suggest that previous studies may have overestimated the cost of respiration during locomotion by ignoring the cost of increased HR. Induced hypoxia experiments may prove useful in determining respiratory and circulatory costs across a broader range of activities and conditions.

Methods

Participants. The present report represents additional data from investigations that were published previously⁴¹, examining the interaction of speed and hypoxia on walking cost. Of the 12 subjects who completed the entire protocol in the previous study, 11 completed the experiment in this study. We used the data from these 11 subjects for this study. Subjects (n = 11) were male athletes (sprinters, middle-distance runners, and soccer and baseball players) who engaged in strenuous daily training (2 hours per day, 5–6 days per week). Their mean age, height, and body weight were 24 ± 8 years, 174 ± 6 cm, and 70 ± 9 kg, respectively (values are mean ± standard deviation [SD]). Before this study, we explained all procedures, possible risks, and benefits of participation to the subjects, and all subjects signed an informed consent form. They were asked to refrain from intense physical activity 2 days before testing and to refrain from drinking any alcohol or caffeinated beverages the day before testing. This study was approved by an ethical committee at the Mount Fuji Research Institute according to the Declaration of Helsinki (No: ECMFRI-03-2014).

Exercise Protocols. All experiments were carried out on a motor-driven treadmill, 2.21 m long and 0.88 m wide (T7000, Johnson Health Tech. Co., Ltd., Taichung Hsein, Taiwan). During an experiment, the subject was allowed to choose his stride and step frequency freely during walking. Subjects wore underwear, shirts, socks, shorts, and lightweight training shoes⁴³. They practiced at least three times at several gait speeds to familiarize themselves with treadmill walking while wearing a gas-collection mask. Inspired oxygen concentrations were set at normobaric normoxia (21% room air), moderate hypoxia (FiO₂; 15%), and severe hypoxia (FiO₂; 11%). Each trial was performed on a different day in random order with a single-blind study design. First, each subject sat on a chair for 10 minutes, then stood for 5 minutes on the treadmill for baseline measurement. Thereafter, he began to walk on the treadmill. As described in our previous study, seven speeds were set incrementally at 0.67, 0.86, 1.06, 1.25, 1.44, 1.64, and 1.83 m s⁻¹, and each speed lasted for 4 minutes^{41, 44}.

Measurements. Pulmonary ventilation (V_E) and gas-exchange variables were measured with a computerized breath-by-breath system (AE-310S, Minato Ltd., Osaka, Japan). The standard, known gases (O₂ 15.23%, CO₂ 4.999%, and N₂ balance) and room air were used to calibrate the gas analyzer. Inspired and expired gas volumes were measured with a hot wire respiratory flow system. Flow signals were electrically integrated for the duration of each breath to calculate minute ventilation. The expired fractions of O₂ and CO₂ were analyzed with a zirconium solid electrolyte oxygen analyzer and an infrared carbon dioxide analyzer, respectively. Each oxygen concentration gas was supplied via a 200-L Douglas bag with a hypoxic gas generator system (Everest summit II, Will Co. Ltd., Tokyo, Japan). HR was measured throughout the study with a commercial HR monitor (POLAR RC800X, POLAR Electro, Tokyo, Japan).

SpO₂ was monitored with a pulse oximeter on the left middle finger every 1 minute throughout the study (TM-2564G, A&D, Tokyo, Japan).

Data Analysis. To calculate energy expenditure, EE, at rest and at each gait speed, VO₂ and VCO₂ were measured with the following equation.

$$\text{Energy Expenditure (Watts)} = [(3.869 \times \text{VO}_2/1000) + (1.195 \times \text{VCO}_2/1000)] \times 4.186/60 \times 1000^{45,46}.$$

At baseline, all physiological values (i.e., gas exchange variables, HR, and SpO₂) were averaged for the last 2 minutes of standing prior to the start of walking. During walking, a single sample with an average final 1 minute of physiological data was also obtained.

Calculating Ventilatory and Heart Rate Costs. To evaluate respiratory and circulatory cost, total walking energy expenditure at each speed (including rest) under normoxia (EE_{norm}) was subtracted from values under moderate or severe hypoxia (EE_{hyp}) to calculate a ΔEE value. We then compared ΔEE to the change in ventilation rate, $\Delta V_E = (V_{E\text{hyp}} - V_{E\text{norm}})$ or change in heart rate, $\Delta\text{HR} = (\text{HR}_{\text{hyp}} - \text{HR}_{\text{norm}})$ between conditions. First, we examined the change in EE with respect to V_E and HR separately, in a bivariate regression model (Ventilation Cost: $\Delta\text{EE} \sim \Delta V_E$. Circulation Cost: $\Delta\text{EE} \sim \Delta\text{HR}$). Next we used multiple regression to test for the independent, additive effects of V_E and HR on EE (Cardiopulmonary Cost: $\Delta\text{EE} \sim \Delta V_E + \Delta\text{HR}$). These regressions were evaluated with the intercept set at 0. This approach assumes that any change in EE between conditions at a given speed is attributable to the change in cardiopulmonary work. This design yielded $n = 16$ comparisons (1 rest + 7 walking speeds for 2 hypoxic conditions).

Statistics. All data are presented as mean \pm SD. A two-way repeated ANOVA (oxygen level \times walking speed) was conducted for comparison in EE, V_E , and HR. As stated above, a bivariate regression model was used for ventilation or circulation cost. Further, multiple regression analysis was also used to test for the independent, additive effects of V_E and HR on EE. All statistical analysis were performed using R ver. 2.13.1 (R Core Team, 2014). A P value less than 0.05 was considered statistically significant.

References

- Pontzer, H., Raichlen, D. A. & Rodman, P. S. Bipedal and quadrupedal locomotion in chimpanzees. *J Hum Evol* **66**, 64–82 (2014).
- Rodman, P. S. & McHenry, H. M. Bioenergetics and the origin of hominid bipedalism. *Am J Phys Anthropol* **52**, 103–106 (1980).
- Sockol, M. D., Raichlen, D. A. & Pontzer, H. Chimpanzee locomotor energetics and the origin of human bipedalism. *Proc Natl Acad Sci USA* **104**, 12265–12269 (2007).
- Cavagna, G. A. & Kaneko, M. Mechanical work and efficiency in level walking and running. *J Physiol* **268**, 467–481 (1977).
- Willems, P. A., Cavagna, G. A. & Heglund, N. C. External, internal and total work in human locomotion. *J Exp Biol* **198**, 379–393 (1995).
- Abe, D., Muraki, S. & Yasukouchi, A. Ergonomic effects of load carriage on energy cost of gradient walking. *Appl Ergon* **39**, 144–149 (2008).
- Horiuchi, M., Endo, J., Horiuchi, Y. & Abe, D. Comparisons of energy cost and economical walking speed at various gradients in healthy active younger and older adults. *J Exerc Sci Fit* **13**, 79–85 (2015).
- Browning, R. C., Baker, E. A., Herron, J. A. & Kram, R. Effects of obesity and sex on the energetic cost and preferred speed of walking. *J Appl Physiol* **100**, 390–398 (2006).
- Browning, R. C. & Kram, R. Energetic cost and preferred speed of walking in obese vs. normal weight women. *Obes Res* **13**, 891–899 (2005).
- Peyrot, N. *et al.* Why does walking economy improve after weight loss in obese adolescents? *Med Sci Sports Exerc* **44**, 659–665 (2012).
- Wall-Scheffler, C. M. & Myers, M. J. Reproductive costs for everyone: how female loads impact human mobility strategies. *J Hum Evol* **64**, 448–456 (2013).
- Abe, D., Muraki, S. & Yasukouchi, A. Ergonomic effects of load carriage on the upper and lower back on metabolic energy cost of walking. *Appl Ergon* **39**, 392–398 (2008).
- Bastien, G. J., Schepens, B., Willems, P. A. & Heglund, N. C. Energetics of load carrying in Nepalese porters. *Science* **308**, 1755 (2005).
- Bastien, G. J., Willems, P. A., Schepens, B. & Heglund, N. C. Effect of load and speed on the energetic cost of human walking. *Eur J Appl Physiol* **94**, 76–83 (2005).
- Maloij, G. M., Heglund, N. C., Prager, L. M., Cavagna, G. A. & Taylor, C. R. Energetic cost of carrying loads: have African women discovered an economic way? *Nature* **319**, 668–669 (1986).
- Bornstein, M. H. & Bornstein, H. G. The pace of life. *Nature* **259**, 557–559 (1976).
- Raichlen, D. A. *et al.* Evidence of Levy walk foraging patterns in human hunter-gatherers. *Proc Natl Acad Sci USA* **111**, 728–733 (2014).
- Shadmehr, R., Huang, H. J. & Ahmed, A. A. A Representation of Effort in Decision-Making and Motor Control. *Curr Biol: CB* **26**, 1929–1934 (2016).
- Selinger, J. C., O'Connor, S. M., Wong, J. D. & Donelan, J. M. Humans Can Continuously Optimize Energetic Cost during Walking. *Curr Biol: CB* **25**, 2452–2456 (2015).
- Aaron, E. A., Johnson, B. D., Seow, C. K. & Dempsey, J. A. Oxygen cost of exercise hyperpnea: measurement. *J Appl Physiol* **72**, 1810–1817 (1992).
- Campbell, E. J., Westlake, E. K. & Cherniack, R. M. Simple methods of estimating oxygen consumption and efficiency of the muscles of breathing. *J Appl Physiol* **11**, 303–308 (1957).
- Coast, J. R. *et al.* Ventilatory work and oxygen consumption during exercise and hyperventilation. *J Appl Physiol* **74**, 793–798 (1993).
- Levison, H. & Cherniack, R. M. Ventilatory cost of exercise in chronic obstructive pulmonary disease. *J Appl Physiol* **25**, 21–27 (1968).
- Margarita, R., Milic-Emili, G., Petit, J. M. & Cavagna, G. Mechanical work of breathing during muscular exercise. *J Appl Physiol* **15**, 354–358 (1960).
- Markley, J. S. & Carrier, D. R. The cost of ventilation in birds measured via unidirectional artificial ventilation. *Comp Biochem Physiol A, Mol Integr Physiol* **155**, 146–153 (2010).
- Marks, D. *et al.* Oxygen cost of ventilation and its effect of the VO₂ plateau. *J. Exerc. Physiol. Online* **8**, 1–12 (2005).
- Milic-Emili, G. & Petit, J. M. Mechanical efficiency of breathing. *J Appl Physiol* **15**, 359–362 (1960).
- Murray, J. F. Oxygen cost of voluntary hyperventilation. *J Appl Physiol* **14**, 187–190 (1959).

29. Scano, G., Grazzini, M., Stendardi, L. & Gigliotti, F. Respiratory muscle energetics during exercise in healthy subjects and patients with COPD. *Respir Med* **100**, 1896–1906 (2006).
30. Vella, C. A., Marks, D. & Robergs, R. A. Oxygen cost of ventilation during incremental exercise to VO_2 max. *Respirology* **11**, 175–181 (2006).
31. Alexopoulos, D. *et al.* Hemodynamic response to hyperventilation test in healthy volunteers. *Clin Cardiol* **18**, 636–641 (1995).
32. Rubenson, J. *et al.* Reappraisal of the comparative cost of human locomotion using gait-specific allometric analyses. *J Exp Biol* **210**, 3513–3524 (2007).
33. Calbet, J. A. & Lundby, C. Air to muscle O_2 delivery during exercise at altitude. *High Alt Med Biol* **10**, 123–134 (2009).
34. Schoene, R. B. Limits of human lung function at high altitude. *J Exp Biol* **204**, 3121–3127 (2001).
35. Dempsey, J. A. & Forster, H. V. Mediation of Ventilatory Adaptations. *Physiol Rev* **62**, 262–346 (1982).
36. Naeije, R. Physiological adaptation of the cardiovascular system to high altitude. *Prog Cardiovasc Dis* **52**, 456–466 (2010).
37. Elia, M. Organ and tissue contribution to metabolic rate. In *Energy Metabolism: Tissue Determinants and Cellular Corollaries*. J. M. Kinney and H. N. Tucker eds pp. 19–60, Raven Press, New York, NY, USA, (1992).
38. Hubel, T. Y. & Usherwood, J. R. Children and adults minimise activated muscle volume by selecting gait parameters that balance gross mechanical power and work demands. *J Exp Biol* **218**, 2830–2839 (2015).
39. Pontzer, H. A unified theory for the energy cost of legged locomotion. *Biol Lett* **12**, 20150935 (2016).
40. Usherwood, J. R. Constraints on muscle performance provide a novel explanation for the scaling of posture in terrestrial animals. *Biol Lett* **9**, 20130414 (2013).
41. Horiuchi, M., Handa, Y., Abe, D. & Fukuoka, Y. Walking economy at simulated high altitude in human healthy young male lowlanders. *Biol Open* **5**, 1408–1414 (2016).
42. Chuang, M. L. *et al.* Aerobically generated CO_2 stored during early exercise. *J Appl Physiol* **87**, 1048–1058 (1999).
43. Abe, D., Yanagawa, K. & Niihata, S. Effects of load carriage, load position, and walking speed on energy cost of walking. *Appl Ergon* **35**, 329–335 (2004).
44. Abe, D., Fukuoka, Y. & Horiuchi, M. Economical Speed and Energetically Optimal Transition Speed Evaluated by Gross and Net Oxygen Cost of Transport at Different Gradients. *PLoS One* **10**, e0138154 (2015).
45. Brouwer, E. On simple formulae for calculating the heat expenditure and the quantities of carbohydrate and fat oxidized in metabolism of men and animals, from gaseous exchange (Oxygen intake and carbonic acid output) and urine-N. *Acta Physiol Pharmacol Neerl* **6**, 795–802 (1957).
46. Masschelein, E. *et al.* Dietary nitrate improves muscle but not cerebral oxygenation status during exercise in hypoxia. *J Appl Physiol* **113**, 736–745 (2012).

Author Contributions

M.H. and H.P. conceived and designed the research. M.H. and Y.H. performed the experiments. M.H., Y.H., D.A., Y.F. and H.P. analyzed and interpreted data. Y.H. prepared tables and figures. M.H. and H.P. wrote the first manuscript. All authors edited, revised and approved the final version of this manuscript.

Additional Information

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017