



Relative substituent orientation in the structure of *cis*-3-chloro-1,3-dimethyl-*N*-(4-nitrophenyl)-2-oxocyclopentane-1-carboxamideMatthias Zeller,<sup>a</sup> Jonas Warneke<sup>b</sup> and Vladimir Azov<sup>b\*</sup>Received 31 July 2014  
Accepted 1 August 2014<sup>a</sup>Department of Chemistry, Youngstown State University, 1 University Plaza, Youngstown, Ohio 44555, USA, and  
<sup>b</sup>University of Bremen, Department of Chemistry, Leobener Str. NW 2C, D-28359 Bremen, Germany. \*Correspondence  
e-mail: vazov@uni-bremen.deEdited by P. C. Healy, Griffith University,  
Australia**Keywords:** crystal structure; hydrogen bonds;  $\pi$ - $\pi$  stacking; methacryloyl chloride dimer; Diels–Alder reaction**CCDC reference:** 1017486**Supporting information:** this article has supporting information at journals.iucr.org/e


The structure of the title compound,  $C_{14}H_{15}ClN_2O_4$ , prepared by reaction of a methacryloyl dimer with nitroaniline, was determined to establish the relative substituent orientation on the cyclopentanone ring. In agreement with an earlier proposed reaction mechanism, the amide group and the methyl group adjacent to the chloro substituent adopt equatorial positions and relative *cis* orientation, whereas the Cl substituent itself and the methyl group adjacent to the amide have axial orientations relative to the mean plane of the five-membered ring. The conformation of the molecule is stabilized by one classical N–H $\cdots$ O (2.18 Å) and one non-classical C–H $\cdots$ O (2.23 Å) hydrogen bond, each possessing an *S*(6) graph-set motif. The crystal packing is defined by several non-classical intramolecular hydrogen bonds, as well as by partial stacking of the aromatic rings.

## 1. Chemical context

The title compound, *cis*-3-chloro-1,3-dimethyl-*N*-(4-nitrophenyl)-2-oxocyclopentane-1-carboxamide, (1), was prepared in the course of study of the formation and reactivity of methacryloyl chloride dimers (2), (3) and (4) (Warneke *et al.*, 2014). The scheme below shows the reactivity of methacryloyl dimers and the synthesis of the title compound (1) (LA = Lewis acid).



Dimer (2) forms in the oxa-Diels–Alder reaction of two methacryloyl chloride molecules and, in the presence of a Lewis acid (LA, such as  $AlCl_3$  or  $TiCl_4$ ), rearranges to cyclopentanone derivatives (3) (kinetic product) and (4) (thermodynamic product). Compounds (3) and (4) show similar  $^1H$  and  $^{13}C$  NMR spectra, making the direct assignment of the relative orientation of the cyclopentanone substituents almost impossible. The crystal structure of (1), as well as the crystal structure of another aromatic amide, *cis*-3-chloro-*N*-(3,5-dichlorophenyl)-1,3-dimethyl-2-oxocyclopentanecarboxamide,




**Figure 1**  
Plot of the title molecule, (1), with the atom-numbering scheme. Displacement ellipsoids are represented at 50% probability levels.

solved and reported earlier (Warneke *et al.*, 2014), were crucial for the determination of the substituent orientation of the cyclopentanone ring after the isolation and derivatization of (4). For the X-ray structures of related *trans*-3-chloro-*N*-(3,5-dichlorophenyl)-1,3-dimethyl-2-oxocyclopentanecarboxamide with *cis* orientation of two methyl groups, see Fischer *et al.* (1985).

## 2. Structural commentary

The molecular structure of the title compound with atom numbering is shown in Fig. 1. All bond lengths and angles may be considered normal. The crystal structure shows the *cis* disposition of the two methyl substituents of the cyclopentanone ring. The C1 and C7 substituents adopt equatorial, whereas the C8 and C11 substituents have axial orientations relative to the mean plane of the five-membered ring. The 4-nitroanilide group is essentially planar, with a maximum deviation of fitted atoms from the least-square plane, which is defined by atoms C9–C14, N1, N2, O1 and O2, of 0.0139 (9) Å for N1. The conformation of the amide is stabilized by one classical N1–H1···O1 (2.18 Å) and one non-classical C10–H10···O2 (2.23 Å) hydrogen bonds (Fig. 2), both with an *S*(6) graph-set motif (Bernstein *et al.*, 1995).



**Figure 2**  
Plot of compound (1) depicting one classical N1–H1···O1 and one non-classical C10–H10···O2 intramolecular hydrogen bond (blue), as well as intermolecular interactions with distances shorter than van der Waals contacts (red).


**Table 1**  
Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N1–H1···O1	0.88	2.18	2.8536 (12)	134
C10–H10···O2	0.95	2.23	2.8467 (14)	122
C7–H7A···O1 <sup>i</sup>	0.98	2.53	3.3577 (13)	142
C7–H7C···O4 <sup>ii</sup>	0.98	2.54	3.4898 (15)	165
C10–H10···C11 <sup>iii</sup>	0.95	2.86	3.5362 (10)	129
C14–H14···C1 <sup>iv</sup>	0.95	2.96	3.9034 (10)	171
C6–H6B···O3 <sup>v</sup>	0.99	2.68	3.1440 (14)	109

Symmetry codes: (i)  $x, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (ii)  $x + 1, y, z + 1$ ; (iii)  $-x + 1, -y + 1, -z + 1$ ; (iv)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (v)  $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$ .

## 3. Supramolecular features

The crystal packing is governed by several short contacts, which may be classified as non-classical hydrogen bonds (for reviews on weak non-classical hydrogen bonding, see Desiraju & Steiner, 1999; Steiner, 2002; Desiraju, 2005), and by partial stacking of the aromatic rings. Molecules of the title compound form columns with alternating enantiomeric molecules along the *c* axis. Although no tight stacking of the aromatic rings can be established [distance between the ring centroids of 4.3719 (6) Å], the aromatic rings of neighboring molecules show partial stacking with several short contacts centered near their nitro-substituent: C14···C13<sup>i</sup> [3.3843 (15) Å; symmetry code: (i)  $x, -y + \frac{1}{2}, z + \frac{1}{2}$ ], C14···C12<sup>i</sup> [3.2483 (15) Å], and C13···N2<sup>i</sup> [3.1860 (14) Å]. The C7–H7A···O1<sup>i</sup> hydrogen bond (2.53 Å) provides additional cohesion between neighboring enantiomeric molecules in the columns (Table 1; Fig. 3). Along the *b* axis, parallel columns are interconnected by C10–H10···C11<sup>iii</sup> [2.86 Å; symmetry code: (iii)  $-x + 1, -y + 1, -z + 1$ ], and along the *a* axis by C7–H7C···O4<sup>ii</sup> [2.54 Å; symmetry code: (ii)  $x + 1, y, z + 1$ ] non-classical hydrogen bonds (Fig. 4). Although the C6–H6B···O3<sup>v</sup> [2.68 Å; symmetry code: (v)  $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$ ] contact also lies below the sum of van der Waals radii, its classification as a hydrogen bond is disputable due to an unfavorable angle of 108°.



**Figure 3**  
Plot of the pair of enantiomeric molecules of (1), showing short contacts between two aromatic rings and the C7–H7A···O1 hydrogen bond.

**Table 2**  
Experimental details.

Crystal data	
Chemical formula	C <sub>14</sub> H <sub>15</sub> ClN <sub>2</sub> O <sub>4</sub>
<i>M<sub>r</sub></i>	310.73
Crystal system, space group	Monoclinic, <i>P</i> 2 <sub>1</sub> / <i>c</i>
Temperature (K)	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	11.4117 (4), 16.1679 (7), 7.8201 (3)
$\beta$ (°)	103.382 (2)
<i>V</i> (Å <sup>3</sup> )	1403.66 (10)
<i>Z</i>	4
Radiation type	
$\mu$ (mm <sup>-1</sup> )	Mo <i>K</i> $\alpha$
Crystal size (mm)	0.28 × 0.18 × 0.16
Data collection	
Diffractometer	Bruker D8 Quest CMOS
Absorption correction	Multi-scan ( <i>SADABS</i> ; Bruker, 2014)
<i>T<sub>min</sub></i> , <i>T<sub>max</sub></i>	0.681, 0.747
No. of measured, independent and observed [ <i>I</i> > 2 $\sigma$ ( <i>I</i> )] reflections	15441, 6627, 5116
<i>R<sub>int</sub></i>	0.028
( <i>sin</i> $\theta$ / $\lambda$ ) <sub>max</sub> (Å <sup>-1</sup> )	0.862
Refinement	
<i>R</i> [ <i>F</i> <sup>2</sup> > 2 $\sigma$ ( <i>F</i> <sup>2</sup> )], <i>wR</i> ( <i>F</i> <sup>2</sup> ), <i>S</i>	0.046, 0.114, 1.06
No. of reflections	6627
No. of parameters	192
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\max}$ , $\Delta\rho_{\min}$ (e Å <sup>-3</sup> )	0.58, -0.36


Computer programs: *APEX2* and *SAINT* (Bruker, 2014), *SHELXS97* and *SHELXL2013* (Sheldrick, 2008), *SHELXL* (Hübschle *et al.*, 2011), *ORTEP-3 for Windows* (Farrugia, 2012), *Mercury* (Macrae *et al.*, 2006), *pubCIF* (Westrip, 2010) and *enCIFer* (Allen *et al.*, 2004).

#### 4. Synthesis and crystallization

The title compound was prepared as described by Warneke *et al.* (2014) by reaction of 4-nitroaniline and *cis*-3-chloro-1,3-dimethyl-2-oxocyclopentanecarbonyl chloride in the presence of Et<sub>3</sub>N in THF. The product was purified by column chromatography on SiO<sub>2</sub> (CHCl<sub>3</sub>) and readily afforded large transparent X-ray quality crystals upon slow evaporation of CHCl<sub>3</sub>/heptane solution (m.p. 402–403 K). <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>):  $\delta$  8.89 (*bs*, 1H), 8.26–8.16 (*m*, 2H), 7.78–7.70 (*m*, 2H), 2.91–2.78 (*m*, 1H), 2.49–2.40 (*m*, 1H), 2.12–2.05 (*m*, 1H), 2.05–1.98 (*m*, 1H), 1.75 (*s*, 3H), 1.51 (*s*, 3H). <sup>13</sup>C NMR (90 MHz, CDCl<sub>3</sub>):  $\delta$  212.4, 168.9, 143.7, 143.3, 125.0, 119.3, 69.7, 55.0, 35.6, 29.4, 25.0, 24.1. MS (EI): *m/z* (%) 310 (85) [*M*]<sup>+</sup>, 173 (85) [*M*-NHAr]<sup>+</sup>. HRMS (EI): *m/z* [*M*]<sup>+</sup> calculated for C<sub>14</sub>H<sub>15</sub>ClN<sub>2</sub>O<sub>4</sub> 310.07203, found 310.07170.

#### 5. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. H atoms were included at calculated positions using a riding model, with aromatic, methyl and amide C–H bond lengths of 0.99, 0.98 and 0.95 Å, respectively, and amide N–H bond lengths of 0.88 Å. The *U*<sub>iso</sub>(H) values



**Figure 4**  
Crystal packing of (1), viewed along the *c* axis. C10–H10...Cl1 contacts are shown as blue dashed lines and C7–H7C...O4 contacts as green dashed lines.

were fixed at 1.5*U*<sub>eq</sub>(C) for methyl H atoms, and 1.2*U*<sub>eq</sub>(C,N) for all other carrier atoms.

#### Acknowledgements

MZ acknowledges the US National Science Foundation for grant 1337296 for the purchase of the X-ray diffractometer.

#### References

- Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed.* **34**, 1555–1573.
- Bruker (2014). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Desiraju, G. R. (2005). *Chem. Commun.* pp. 2995–3001.
- Desiraju, G. R. & Steiner, T. (1999). *The Weak Hydrogen Bond in Structural Chemistry and Biology*. New York: Oxford University Press Inc.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Fischer, W., Belluš, D., Adler, A., Francotte, E. & Roloff, A. (1985). *Chimia*, **39**, 19–20.
- Hübschle, C. B., Sheldrick, G. M. & Dittrich, B. (2011). *J. Appl. Cryst.* **44**, 1281–1284.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Steiner, T. (2002). *Angew. Chem. Int. Ed.* **41**, 48–76.
- Warneke, J., Wang, Z., Zeller, M., Leibfritz, D., Plaumann, M. & Azov, V. A. (2014). *Tetrahedron* **70**, 6515–6521.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2014). E70, 121-123 [doi:10.1107/S1600536814017711]

## Relative substituent orientation in the structure of *cis*-3-chloro-1,3-dimethyl-*N*-(4-nitrophenyl)-2-oxocyclopentane-1-carboxamide

Matthias Zeller, Jonas Warneke and Vladimir Azov

### Computing details

Data collection: *APEX2* (Bruker, 2014); cell refinement: *SAINTE* (Bruker, 2014); data reduction: *SAINTE* (Bruker, 2014); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008) and *SHELXL* (Hübschle *et al.*, 2011); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010) and *enCIFer* (Allen *et al.*, 2004).

### *cis*-3-Chloro-1,3-dimethyl-*N*-(4-nitrophenyl)-2-oxocyclopentane-1-carboxamide

#### Crystal data

C<sub>14</sub>H<sub>15</sub>ClN<sub>2</sub>O<sub>4</sub>

*M<sub>r</sub>* = 310.73

Monoclinic, *P*2<sub>1</sub>/*c*

*a* = 11.4117 (4) Å

*b* = 16.1679 (7) Å

*c* = 7.8201 (3) Å

β = 103.382 (2)°

*V* = 1403.66 (10) Å<sup>3</sup>

*Z* = 4

*F*(000) = 648

*D<sub>x</sub>* = 1.470 Mg m<sup>-3</sup>

Melting point: 402 K

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 8547 reflections

θ = 2.5–37.8°

μ = 0.29 mm<sup>-1</sup>

*T* = 100 K

Block, colourless

0.28 × 0.18 × 0.16 mm

#### Data collection

Bruker D8 Quest CMOS

diffractometer

Radiation source: I-μ-S microsource X-ray

tube

'laterally graded multilayer (Goebel) mirror'

monochromator

ω and phi scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2014)

*T<sub>min</sub>* = 0.681, *T<sub>max</sub>* = 0.747

15441 measured reflections

6627 independent reflections

5116 reflections with *I* > 2σ(*I*)

*R<sub>int</sub>* = 0.028

θ<sub>max</sub> = 37.8°, θ<sub>min</sub> = 2.2°

*h* = -19→17

*k* = -21→27

*l* = -12→9

#### Refinement

Refinement on *F*<sup>2</sup>

Least-squares matrix: full

*R*[*F*<sup>2</sup> > 2σ(*F*<sup>2</sup>)] = 0.046

*wR*(*F*<sup>2</sup>) = 0.114

*S* = 1.06

6627 reflections

192 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0482P)^2 + 0.5524P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.58 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.64522 (9)	0.42645 (6)	0.37961 (14)	0.01331 (17)
C2	0.76549 (9)	0.42549 (6)	0.51672 (14)	0.01232 (17)
C3	0.77769 (9)	0.35857 (6)	0.65852 (13)	0.01139 (16)
C4	0.83111 (9)	0.39694 (6)	0.83990 (13)	0.01175 (16)
C5	0.87160 (10)	0.48290 (6)	0.79734 (15)	0.01608 (19)
H5A	0.9561	0.4818	0.7857	0.019*
H5B	0.8649	0.5228	0.8906	0.019*
C6	0.78602 (10)	0.50660 (6)	0.62191 (14)	0.01600 (19)
H6A	0.7091	0.5285	0.6412	0.019*
H6B	0.8231	0.5489	0.5596	0.019*
C7	0.92411 (9)	0.34331 (6)	0.96019 (14)	0.01468 (18)
H7A	0.8894	0.2888	0.9723	0.022*
H7B	0.9483	0.3695	1.0760	0.022*
H7C	0.9947	0.3370	0.9100	0.022*
C8	0.86666 (10)	0.40959 (7)	0.41801 (16)	0.0181 (2)
H8A	0.8532	0.3561	0.3578	0.027*
H8B	0.9449	0.4091	0.5024	0.027*
H8C	0.8658	0.4536	0.3315	0.027*
C9	0.50005 (9)	0.33261 (6)	0.18533 (13)	0.01140 (16)
C10	0.43082 (10)	0.39401 (6)	0.08226 (14)	0.01465 (18)
H10	0.4525	0.4506	0.1006	0.018*
C11	0.33034 (9)	0.37168 (6)	-0.04677 (14)	0.01470 (18)
H11	0.2830	0.4129	-0.1173	0.018*
C12	0.29953 (9)	0.28902 (6)	-0.07204 (14)	0.01272 (17)
C13	0.36802 (9)	0.22669 (6)	0.02663 (14)	0.01327 (17)
H13	0.3463	0.1702	0.0063	0.016*
C14	0.46849 (9)	0.24892 (6)	0.15510 (14)	0.01253 (17)
H14	0.5165	0.2073	0.2233	0.015*
N1	0.60174 (8)	0.35022 (5)	0.31966 (12)	0.01282 (15)
H1	0.6424	0.3073	0.3715	0.015*
N2	0.19374 (8)	0.26654 (6)	-0.20886 (13)	0.01633 (17)
O1	0.75274 (7)	0.28588 (5)	0.63432 (10)	0.01533 (15)
O2	0.59793 (8)	0.49203 (5)	0.32621 (13)	0.02455 (19)
O3	0.16813 (9)	0.19301 (6)	-0.23155 (13)	0.0270 (2)
O4	0.13478 (8)	0.32247 (6)	-0.29704 (12)	0.02274 (18)
Cl1	0.69827 (2)	0.40657 (2)	0.93120 (4)	0.01568 (6)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0149 (4)	0.0129 (4)	0.0118 (4)	-0.0017 (3)	0.0024 (4)	0.0009 (3)
C2	0.0143 (4)	0.0112 (4)	0.0107 (4)	-0.0023 (3)	0.0015 (3)	0.0011 (3)
C3	0.0117 (4)	0.0114 (4)	0.0113 (4)	0.0001 (3)	0.0031 (3)	0.0001 (3)
C4	0.0125 (4)	0.0121 (4)	0.0106 (4)	-0.0012 (3)	0.0026 (3)	-0.0003 (3)
C5	0.0191 (5)	0.0129 (4)	0.0147 (5)	-0.0057 (3)	0.0008 (4)	-0.0001 (3)
C6	0.0217 (5)	0.0106 (4)	0.0140 (5)	-0.0040 (3)	0.0007 (4)	0.0010 (3)
C7	0.0123 (4)	0.0166 (4)	0.0143 (5)	0.0007 (3)	0.0013 (4)	0.0018 (3)
C8	0.0159 (5)	0.0241 (5)	0.0151 (5)	-0.0027 (4)	0.0049 (4)	0.0018 (4)
C9	0.0112 (4)	0.0124 (4)	0.0106 (4)	0.0001 (3)	0.0025 (3)	0.0002 (3)
C10	0.0147 (4)	0.0129 (4)	0.0150 (5)	0.0009 (3)	0.0007 (4)	0.0015 (3)
C11	0.0133 (4)	0.0154 (4)	0.0145 (5)	0.0019 (3)	0.0014 (4)	0.0019 (3)
C12	0.0102 (4)	0.0174 (4)	0.0108 (4)	0.0005 (3)	0.0028 (3)	-0.0004 (3)
C13	0.0131 (4)	0.0137 (4)	0.0132 (4)	-0.0002 (3)	0.0034 (4)	-0.0004 (3)
C14	0.0126 (4)	0.0124 (4)	0.0124 (4)	0.0009 (3)	0.0025 (3)	0.0004 (3)
N1	0.0128 (4)	0.0109 (3)	0.0131 (4)	0.0001 (3)	-0.0003 (3)	0.0006 (3)
N2	0.0128 (4)	0.0224 (4)	0.0131 (4)	-0.0007 (3)	0.0016 (3)	-0.0007 (3)
O1	0.0203 (4)	0.0104 (3)	0.0146 (4)	-0.0009 (2)	0.0026 (3)	-0.0009 (2)
O2	0.0275 (4)	0.0131 (3)	0.0261 (5)	0.0012 (3)	-0.0081 (4)	0.0022 (3)
O3	0.0256 (4)	0.0228 (4)	0.0265 (5)	-0.0073 (3)	-0.0064 (4)	-0.0012 (3)
O4	0.0163 (4)	0.0281 (4)	0.0204 (4)	0.0043 (3)	-0.0025 (3)	0.0038 (3)
C11	0.01539 (11)	0.01668 (11)	0.01600 (12)	0.00173 (8)	0.00574 (9)	-0.00161 (8)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

C1—O2	1.2189 (13)	C8—H8A	0.9800
C1—N1	1.3701 (13)	C8—H8B	0.9800
C1—C2	1.5338 (15)	C8—H8C	0.9800
C2—C3	1.5324 (14)	C9—C10	1.4017 (14)
C2—C6	1.5369 (14)	C9—N1	1.4025 (13)
C2—C8	1.5515 (15)	C9—C14	1.4061 (13)
C3—O1	1.2136 (12)	C10—C11	1.3884 (15)
C3—C4	1.5390 (14)	C10—H10	0.9500
C4—C7	1.5169 (14)	C11—C12	1.3844 (15)
C4—C5	1.5253 (14)	C11—H11	0.9500
C4—C11	1.8257 (10)	C12—C13	1.3938 (14)
C5—C6	1.5371 (15)	C12—N2	1.4616 (14)
C5—H5A	0.9900	C13—C14	1.3856 (14)
C5—H5B	0.9900	C13—H13	0.9500
C6—H6A	0.9900	C14—H14	0.9500
C6—H6B	0.9900	N1—H1	0.8800
C7—H7A	0.9800	N2—O3	1.2273 (13)
C7—H7B	0.9800	N2—O4	1.2373 (13)
C7—H7C	0.9800		
O2—C1—N1	124.66 (10)	C4—C7—H7C	109.5

O2—C1—C2	120.13 (9)	H7A—C7—H7C	109.5
N1—C1—C2	115.13 (8)	H7B—C7—H7C	109.5
C3—C2—C1	115.45 (8)	C2—C8—H8A	109.5
C3—C2—C6	103.77 (8)	C2—C8—H8B	109.5
C1—C2—C6	111.49 (8)	H8A—C8—H8B	109.5
C3—C2—C8	106.84 (8)	C2—C8—H8C	109.5
C1—C2—C8	107.60 (9)	H8A—C8—H8C	109.5
C6—C2—C8	111.64 (9)	H8B—C8—H8C	109.5
O1—C3—C2	126.31 (9)	C10—C9—N1	123.06 (9)
O1—C3—C4	124.24 (9)	C10—C9—C14	119.77 (9)
C2—C3—C4	109.42 (8)	N1—C9—C14	117.17 (9)
C7—C4—C5	116.90 (9)	C11—C10—C9	119.64 (9)
C7—C4—C3	114.34 (8)	C11—C10—H10	120.2
C5—C4—C3	103.98 (8)	C9—C10—H10	120.2
C7—C4—C11	109.35 (7)	C12—C11—C10	119.64 (9)
C5—C4—C11	109.18 (7)	C12—C11—H11	120.2
C3—C4—C11	101.95 (7)	C10—C11—H11	120.2
C4—C5—C6	105.06 (8)	C11—C12—C13	121.84 (10)
C4—C5—H5A	110.7	C11—C12—N2	118.95 (9)
C6—C5—H5A	110.7	C13—C12—N2	119.20 (9)
C4—C5—H5B	110.7	C14—C13—C12	118.54 (9)
C6—C5—H5B	110.7	C14—C13—H13	120.7
H5A—C5—H5B	108.8	C12—C13—H13	120.7
C2—C6—C5	104.58 (8)	C13—C14—C9	120.55 (9)
C2—C6—H6A	110.8	C13—C14—H14	119.7
C5—C6—H6A	110.8	C9—C14—H14	119.7
C2—C6—H6B	110.8	C1—N1—C9	127.61 (9)
C5—C6—H6B	110.8	C1—N1—H1	116.2
H6A—C6—H6B	108.9	C9—N1—H1	116.2
C4—C7—H7A	109.5	O3—N2—O4	123.15 (10)
C4—C7—H7B	109.5	O3—N2—C12	118.38 (9)
H7A—C7—H7B	109.5	O4—N2—C12	118.47 (9)
O2—C1—C2—C3	140.81 (11)	C1—C2—C6—C5	154.86 (9)
N1—C1—C2—C3	-42.27 (13)	C8—C2—C6—C5	-84.75 (10)
O2—C1—C2—C6	22.73 (14)	C4—C5—C6—C2	-37.41 (11)
N1—C1—C2—C6	-160.35 (9)	N1—C9—C10—C11	-179.12 (10)
O2—C1—C2—C8	-100.01 (12)	C14—C9—C10—C11	1.15 (16)
N1—C1—C2—C8	76.91 (11)	C9—C10—C11—C12	0.10 (16)
C1—C2—C3—O1	47.32 (14)	C10—C11—C12—C13	-1.21 (16)
C6—C2—C3—O1	169.62 (10)	C10—C11—C12—N2	-179.78 (10)
C8—C2—C3—O1	-72.28 (13)	C11—C12—C13—C14	1.02 (16)
C1—C2—C3—C4	-134.50 (9)	N2—C12—C13—C14	179.59 (9)
C6—C2—C3—C4	-12.19 (11)	C12—C13—C14—C9	0.26 (15)
C8—C2—C3—C4	105.90 (9)	C10—C9—C14—C13	-1.34 (15)
O1—C3—C4—C7	39.21 (14)	N1—C9—C14—C13	178.91 (9)
C2—C3—C4—C7	-139.02 (9)	O2—C1—N1—C9	2.35 (18)
O1—C3—C4—C5	167.84 (10)	C2—C1—N1—C9	-174.41 (10)

C2—C3—C4—C5	-10.39 (11)	C10—C9—N1—C1	4.49 (17)
O1—C3—C4—C11	-78.65 (11)	C14—C9—N1—C1	-175.77 (10)
C2—C3—C4—C11	103.11 (8)	C11—C12—N2—O3	179.12 (11)
C7—C4—C5—C6	156.09 (9)	C13—C12—N2—O3	0.51 (15)
C3—C4—C5—C6	29.04 (11)	C11—C12—N2—O4	-0.39 (15)
C11—C4—C5—C6	-79.18 (9)	C13—C12—N2—O4	-179.00 (10)
C3—C2—C6—C5	29.97 (11)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1	0.88	2.18	2.8536 (12)	134
C10—H10...O2	0.95	2.23	2.8467 (14)	122
C7—H7A...O1 <sup>i</sup>	0.98	2.53	3.3577 (13)	142
C7—H7C...O4 <sup>ii</sup>	0.98	2.54	3.4898 (15)	165
C10—H10...C11 <sup>iii</sup>	0.95	2.86	3.5362 (10)	129
C14—H14...C11 <sup>iv</sup>	0.95	2.96	3.9034 (10)	171
C6—H6B...O3 <sup>v</sup>	0.99	2.68	3.1440 (14)	109

Symmetry codes: (i)  $x, -y+1/2, z+1/2$ ; (ii)  $x+1, y, z+1$ ; (iii)  $-x+1, -y+1, -z+1$ ; (iv)  $x, -y+1/2, z-1/2$ ; (v)  $-x+1, y+1/2, -z+1/2$ .